

Pacific Journal of Mathematics

NONCOINCIDENCE INDEX OF MANIFOLDS

MIKE HOFFMAN

NONCOINCIDENCE INDEX OF MANIFOLDS

MICHAEL HOFFMAN

For a connected topological manifold M we define the noncoincidence index of M , a topological invariant reflecting the abundance of fixed-point-free self-maps of M . We give some theorems on noncoincidence index and compute the noncoincidence index of the homogeneous manifold $U(n)/H$, where H is conjugate to $U(1)^k \times U(n-k)$.

1. Introduction. Let M be a manifold (connected locally Euclidean Hausdorff space). We define the *noncoincidence index* of M , $\#M$, as follows. If M admits k fixed-point-free self-maps, no pair of which has a coincidence, set $\#M \geq k + 1$. If $\#M \geq i$ for all i , put $\#M = \infty$; otherwise, $\#M$ is the greatest number i with $\#M \geq i$. (This definition is inspired by [5].)

Evidently a manifold has noncoincidence index 1 if and only if it has the fixed-point property. On the other hand, if a group G acts freely on M , then $\#M \geq \text{card } G$ if G is finite and $\#M = \infty$ if G is infinite. In particular, any connected nontrivial Lie group has noncoincidence index ∞ .

As we see in §2, many manifolds besides Lie groups have noncoincidence index ∞ . In §3 we show how the Lefschetz coincidence theorem can be used to put a finite upper bound on $\#M$ for certain compact oriented manifolds M . These results are used in §4 to compute the noncoincidence index of the homogeneous space $U(n)/H$ for H conjugate to $U(1)^k \times U(n-k)$. Section 5 is devoted to proving a classification theorem for endomorphisms of $H^*(U(n)/H; \mathbf{Q})$ which is needed in §4.

I thank my colleague W. Homer for greatly improving Lemma 5.3, and I thank A. Dold for some helpful observations.

2. Sufficient conditions for $\#M = \infty$. In this section we give some sufficient conditions for a manifold M to have $\#M = \infty$. The following result gives some easily checked homological conditions.

THEOREM 2.1. *Let M be a compact manifold. Then $\#M = \infty$ if either of the following is true:*

1. M has nonzero first Betti number, or
2. $\chi(M) = 0$.

Proof. For (1), see Corollary 5.1 of [5]. Now suppose $\chi(M) = 0$. By [4], there is a map $s: [0, 1] \times M \rightarrow M$ with $s(0, \cdot) = \text{id}_M$ and $s(t, \cdot): M \rightarrow M$ fixed-point-free for $t > 0$. Let d be a metric for M , and set

$$N(t) = \inf_{x \in M} d(s(t, x), x), \quad F(t) = \sup_{x \in M} d(s(t, x), x).$$

Then $F(t) \geq N(t) > 0$ for $t > 0$, and $F(t), N(t) \rightarrow 0$ as $t \rightarrow 0$. Choose $0 < t_k < t_{k-1} < \cdots < t_1 \leq 1$ so that $F(t_i) < N(t_{i-1})$: then

$$x \rightarrow s(t_i, x), \quad 1 \leq i \leq k,$$

is a set of k fixed-point-free, noncoincident maps. Since we can do this for any k , $\#M = \infty$.

From the preceding result, we see that any odd-dimensional compact manifold has noncoincidence index ∞ . It also follows that $\#M = \infty$ for any compact surface M , except $M = S^2$ and $M = \mathbf{R}P^2$ (of course $\#\mathbf{R}P^2 = 1$, and we see in the next section that $\#S^2 = 2$).

The next result gives another useful sufficient condition for $\#M = \infty$.

THEOREM 2.2. *Let M be a compact manifold which admits a fixed-point-free nonsurjective self-map. Then $\#M = \infty$.*

Proof. Let $f: M \rightarrow M$ be fixed-point-free and nonsurjective. By Theorem 1.11 of [3], there is a path field nonsingular on the image of f , i.e. a map $s: [0, 1] \times M \rightarrow M$ such that $s(0, \cdot) = \text{id}_M$ and $s(t, \cdot)$ fixes no point of $f(M)$ for $t > 0$. Let d be a metric for M and take $\varepsilon > 0$ so that $d(f(x), x) \geq \varepsilon$ for $x \in M$. Then there is some $t_0 > 0$ so that

$$\sup_{x \in M} d(s(t, x), x) < \varepsilon$$

for $t < t_0$. Now proceed as in the proof of 2.1; set

$$N(t) = \inf_{x \in f(M)} d(s(t, x), x), \quad F(t) = \sup_{x \in f(M)} d(s(t, x), x)$$

(note $f(M)$ is compact) and choose $0 < t_k < t_{k-1} < \cdots < t_1 < t_0$ such that $F(t_i) < N(t_{i-1})$. Then there are k fixed-point-free noncoincident self-maps of M given by

$$x \rightarrow s(t_i, f(x)), \quad 1 \leq i \leq k.$$

Since k is arbitrary, $\#M = \infty$.

3. The Lefschetz coincidence theorem. In this section we show how the Lefschetz coincidence theorem can be used to put a finite upper bound on the noncoincidence index in some cases. As we see in the next

section, such an upper bound combined with constructions of fixed-point-free maps often gives the noncoincidence index exactly.

Throughout this section, M will be a compact oriented n -manifold. We shall use the following version of the Lefschetz coincidence theorem: for a more general statement, see [9].

THEOREM 3.1. *For maps $f, g: M \rightarrow M$, set*

$$L(f, g) = \sum_{i=0}^n (-1)^i \operatorname{Tr}(\Phi_i^{-1} g_* \Phi_i f^*),$$

where $\Phi_i: H^i(M; \mathbf{Q}) \rightarrow H_{n-i}(M; \mathbf{Q})$ is the Poincaré duality isomorphism. If $L(f, g) \neq 0$, then f and g have a coincidence.

REMARKS. 1. It is immediate that $L(f, \operatorname{id}) = L(f)$, the ordinary Lefschetz number of f , so this result implies the Lefschetz fixed-point theorem for M .

2. It follows from properties of trace that $L(f, g) = (-1)^n L(g, f)$.

Let g be a self-map of M . We define the degree of g by $g_*[M] = (\deg g)[M]$, where $[M] \in H_n(M; \mathbf{Q})$ is the fundamental class of M . The following result is useful in computing the Lefschetz coincidence number.

PROPOSITION 3.2. *If g is a self-map of M with $\deg g \neq 0$, then $g^*: H^*(M; \mathbf{Q}) \rightarrow H^*(M; \mathbf{Q})$ has an inverse \bar{g}^* and*

$$L(f, g) = (\deg g) L(\bar{g}^* f^*)$$

for any other self-map f of M .

Proof. If $\deg g \neq 0$, it follows from consideration of Poincaré duality that g^* is injective. Then g^* is an automorphism, since each vector space $H^i(M; \mathbf{Q})$ is finite-dimensional. For $u \in H^i(M; \mathbf{Q})$,

$$\begin{aligned} \Phi_1^{-1} g_* \Phi_i f^*(u) &= \Phi_i^{-1} g_*(g^* \bar{g}^* f^*(u) \cap [M]) \\ &= \Phi_i^{-1}(\bar{g}^* f^*(u) \cap g_*[M]) = (\deg g) \bar{g}^* f^*(u), \end{aligned}$$

and the conclusion follows from the definition of $L(f, g)$.

By Theorem 3.1, any fixed-point-free self-map f of M must have $L(f) = 0$, and any pair f, g of self-maps without a coincidence must have $L(f, g) = 0$. We put

$$LZ(M) = \{f^* | f: M \rightarrow M \text{ and } L(f) = 0\}$$

and say $f^*, g^* \in LZ(M)$ are *compatible* if $L(f, g) = 0$. If $\chi(M) \neq 0$ and $LZ(M)$ consists of automorphisms of $H^*(M; \mathbf{Q})$, we call M *L-rigid*. We then have the following result.

PROPOSITION 3.3. *Suppose M is L-rigid. If $\#M \geq k + 1$, then $LZ(M)$ contains a subset of k pairwise compatible elements.*

Proof. By the hypothesis, there is a set S of k pairwise noncoincident fixed-point-free self-maps of M . Let $f, g \in S$. Then f^* and g^* are compatible elements of $LZ(M)$. We have $f^* \neq g^*$, since otherwise

$$L(f, g) = L(f, f) = (\deg f)L(\text{id}) = (\deg f)\chi(M) \neq 0.$$

Thus, $\{f^* | f \in S\}$ is a set of k pairwise compatible elements of $LZ(M)$.

REMARK. Note that if M is *L-rigid*, then any pair $f^*, g^* \in LZ(M)$ is compatible if and only if $L(\bar{g}^*f^*) = 0$.

It follows immediately from 3.3 that

$$(1) \quad \#M \leq \text{card } LZ(M) + 1$$

when M is *L-rigid* and $LZ(M)$ is finite. Thus we have, e.g., $\#S^{2n} \leq 2$ for any even sphere S^{2n} (and in fact $\#S^{2n} = 2$, since the antipodal map is fixed-point-free). As we see in the next section, however, 3.3 sometimes gives a sharper upper bound than (1).

4. Noncoincidence index of some flag manifolds. Let $F(1^k, n)$ denote the homogeneous space $U(n+k)/H$, where H is conjugate to $U(1)^k \times U(n)$. (We can assume $n = 0$ or $n \geq 2$: in the former case we write $F(1^k)$ instead of $F(1^k, 0)$.) It is proved in [7] that $\#F(1^k) = k!$. In this section we compute $\#F(1^k, n)$ for all k and n .

The manifold $F(1^k, n)$ can be thought of as in the space of k -tuples of orthogonal lines in \mathbf{C}^{n+k} . Thus, there are maps

$$\pi_i: F(1^k, n) \rightarrow \mathbf{C}P^{n+k-1}, \quad 1 \leq i \leq k,$$

given by picking out the i th line. If we let $t \in H^2(\mathbf{C}P^{n+k-1}; \mathbf{Q})$ be the first Chern class of the canonical line bundle over $\mathbf{C}P^{n+k-1}$ and put $t_i = \pi_i^*(t)$, we have the following description of $H^*(F(1^k, n); \mathbf{Q})$ [1].

$$H^*(F(1^k, n); \mathbf{Q}) = \mathbf{Q}[t_1, t_2, \dots, t_k] / \{h_i | n + 1 \leq i \leq n + k\},$$

where h_i is the i th complete symmetric function in t_1, t_2, \dots, t_k , i.e.

$$h_i = \sum_{p_1 + \dots + p_k = i} t_1^{p_1} t_2^{p_2} \dots t_k^{p_k}.$$

There is a free action of the symmetric group Σ_k on $F(1^k, n)$ by permutation of lines, and this action evidently permutes the t_i in cohomology.

For any $m \in \mathbf{Q}$ and $\sigma \in \Sigma_k$, let h_m^σ denote the endomorphism of $H^*(F(1^k, n); \mathbf{Q})$ given by

$$h_m^\sigma(t_i) = mt_{\sigma(i)}.$$

The following classification theorem for endomorphisms of $H^*(F(1^k, n); \mathbf{Q})$ is proved in §5.

THEOREM 4.1. *Unless $k = 2$ and n is a positive even number, all endomorphisms of $H^*(F(1^k, n); \mathbf{Q})$ are of the form h_m^σ for some $m \in \mathbf{Q}$ and $\sigma \in \Sigma_k$. If $k = 2$ and $n \geq 2$ is even, the only additional endomorphisms are*

$$t_i \rightarrow (-1)^i mt_q, \quad i = 1, 2,$$

for $q \in \{1, 2\}$ and $m \in \mathbf{Q}$.

The next result gives a formula for $L(h_m^\sigma)$.

THEOREM 4.2. *Let $\lambda_1 \geq \lambda_2 \geq \dots$ be the cycle-type of $\sigma \in \Sigma_k$ (so λ_1 is the length of the longest cycle in σ , λ_2 is the length of the next longest cycle, etc.). Then $h_m^\sigma: H^*(F(1^k, n); \mathbf{Q}) \rightarrow H^*(F(1^k, n); \mathbf{Q})$ has Lefschetz number*

$$(1) \quad L(h_m^\sigma) = \frac{(1 - m^{n+1})(1 - m^{n+2}) \dots (1 - m^{n+k})}{(1 - m^{\lambda_1})(1 - m^{\lambda_2}) \dots}.$$

Proof. For $h_m^\sigma: H^*(F(1^k, n); \mathbf{Q}) \rightarrow H^*(F(1^k, n); \mathbf{Q})$, let $P_{k,n}(\sigma, m)$ denote $L(h_m^\sigma)$. From [7] we have the formula

$$(2) \quad P_{k,0}(\sigma, m) = \frac{(1 - m)(1 - m^2) \dots (1 - m^k)}{(1 - m^{\lambda_1})(1 - m^{\lambda_2}) \dots}.$$

Now the spectral sequence of the fibration

$$F(1^k) \rightarrow F(1^k, n) \rightarrow G_k(\mathbf{C}^{n+k}),$$

where $G_k(\mathbf{C}^{n+k})$ is the Grassmannian of k -planes in \mathbf{C}^{n+k} , collapses for degree reasons. Thus

$$H^*(F(1^k, n); \mathbf{Q}) \cong H^*(F(1^k); \mathbf{Q}) \otimes H^*(G_k(\mathbf{C}^{n+k}); \mathbf{Q})$$

additively. Now $H^*(G_k(\mathbf{C}^{n+k}); \mathbf{Q})$ can be regarded as the invariant subring of $H^*(F(1^k, n); \mathbf{Q})$ under the Σ_k -action, and $H^*(F(1^k); \mathbf{Q})$ is a quotient of $H^*(F(1^k, n); \mathbf{Q})$ (the projection is the obvious map sending $t_i \in H^2(F(1^k, n); \mathbf{Q})$ to $t_i \in H^2(F(1^k); \mathbf{Q})$). Any endomorphism h_m^σ of

$H^*(F(1^k, n); \mathbf{Q})$ restricts to the endomorphism of $H^*(G_k(\mathbf{C}^{n+k}); \mathbf{Q})$ which multiplies dimension $2i$ by m^i , and gives rise to the corresponding h_m^σ on $H^*(F(1^k); \mathbf{Q})$. Since trace is multiplicative on tensor products,

$$(3) \quad P_{k,n}(\sigma, m) = P_{k,0}(\sigma, m) \sum_{i \geq 0} m^i \dim H^{2i}(G_k(\mathbf{C}^{n+k}); \mathbf{Q}).$$

It is well known (see e.g. [1]) that

$$\sum_{i \geq 0} m^i \dim H^{2i}(G_k(\mathbf{C}^{n+k}); \mathbf{Q}) = \frac{(1 - m^{n+1})(1 - m^{n+2}) \cdots (1 - m^{n+k})}{(1 - m)(1 - m^2) \cdots (1 - m^k)},$$

and this together with (2) and (3) implies the conclusion.

We can use Theorems 4.1 and 4.2 to show that many of the manifolds $F(1^k, n)$ are L -rigid.

PROPOSITION 4.3. *Suppose $k \neq 2$, n is odd, or $n = 0$. Then $F(1^k, n)$ is L -rigid. Further, if Γ is the set of products of $\lfloor k/2 \rfloor$ disjoint transpositions in Σ_k , then*

$$LZ(F(1^k, n)) = \begin{cases} \{h_1^\sigma | \sigma \neq \text{id}\} \cup \{h_{-1}^\sigma | \sigma \in \Sigma_k\}, & kn \text{ odd,} \\ \{h_1^\sigma | \sigma \neq \text{id}\} \cup \{h_{-1}^\sigma | \sigma \notin \Gamma\}, & kn \text{ even.} \end{cases}$$

Proof. By 4.1, an element of $LZ(F(1^k, n))$ must be an endomorphism h_m^σ of $H^*(F(1^k, n); \mathbf{Q})$ with $L(h_m^\sigma) = 0$. The above list follows from consideration of (1) (it is easy to see that any h_m^σ , $m = \pm 1$, is induced by a self-map of $F(1^k, n)$). Now

$$\chi(F(1^k, n)) = k! \binom{n+k}{k}$$

and clearly h_m^σ is an automorphism for $m \neq 0$ (in fact $\deg h_m^\sigma = m^d \text{sgn } \sigma$, where $d = \dim_{\mathbf{C}} F(1^k, n)$), so $F(1^k, n)$ is L -rigid.

Now we can give an upper bound for $\#F(1^k, n)$ when n and k satisfy the hypothesis of the preceding result.

PROPOSITION 4.4. *Suppose $k \neq 2$, n is odd, or $n = 0$. Then $\#F(1^k, n) \leq k!$ if kn is even, and $\#F(1^k, n) \leq 2k!$ if kn is odd.*

Proof. The statement about $\#F(1^k, n)$ for kn odd follows immediately from 3.3, since $LZ(F(1^k, n))$ has $2k! - 1$ elements by 4.3. Again by 3.3, to prove the statement about $\#F(1^k, n)$ for kn even it suffices to show that any set of pairwise compatible elements of $LZ(F(1^k, n))$ has at most $k! - 1$ elements in this case. Suppose kn even and let $S \subset LZ(F(1^k, n))$ be a set of compatible elements. Let

$$H = \{ \sigma \in \Sigma_k | h_1^\sigma \in S \} \cup \{ \text{id} \},$$

$$K = \{ \sigma \in \Sigma_k | h_{-1}^\sigma \in S \}.$$

Then $\text{card } S = \text{card } H + \text{card } K - 1$. Since

$$L(h_{-1}^\tau, h_1^\sigma) = (\text{deg } h_1^\sigma)L(h_1^{\sigma^{-1}}h_{-1}^\tau) = \pm L(h_{-1}^{\sigma^{-1}\tau}),$$

we must have $\sigma^{-1}\tau \notin \Gamma$ for $\tau \in K$ and $\sigma \in H$ (here Γ is as in 4.3). But then, if we take $\rho \in \Gamma$, we have $\sigma\rho \notin K$ for every $\sigma \in H$: hence $\text{card } H + \text{card } K \leq k!$, and the conclusion follows.

Next we show that the inequalities of 4.4 are equalities. To do this, we construct fixed-point-free, noncoincident maps. For any even number $2r$, define $J: \mathbf{C}^{2r} \rightarrow \mathbf{C}^{2r}$ by

$$J(z_1, z_2, \dots, z_{2r-1}, z_{2r}) = (-\bar{z}_2, \bar{z}_1, \dots, -\bar{z}_{2r}, \bar{z}_{2r-1}).$$

Then J is a conjugate-linear map of \mathbf{C}^{2r} with $J^2 = -\text{id}$. Under the evident identification $\mathbf{C}^{2r} \cong \mathbf{H}^r$, we can regard J as multiplication by the quaternion j . Any subspace of \mathbf{C}^{2r} invariant under J can be given the structure of a quaternionic vector space, and thus must be even-dimensional (cf. the proof of Theorem 1 of [6]). Further, if $\langle \ , \ \rangle$ denotes inner product,

$$(4) \quad \langle Jv, Jw \rangle = \langle w, v \rangle \quad \text{for } v, w \in \mathbf{C}^{2r}.$$

Thus J preserves orthogonality.

THEOREM 4.5. *Unless $k = 2$ and n is a positive even number,*

$$\#F(1^k, n) = \begin{cases} 2k!, & kn \text{ odd,} \\ k!, & kn \text{ even.} \end{cases}$$

Proof. Since there is a free Σ_k -action on $F(1^k, n)$ (i.e., permutation of lines), we have $\#F(1^k, n) \geq k!$; together with 4.4, this disposes of the case kn even. Now suppose kn is odd. Then $n + k$ is even, and we have the map $J: \mathbf{C}^{n+k} \rightarrow \mathbf{C}^{n+k}$ defined above. Consider the $2k! - 1$ self-maps of $F(1^k, n)$ defined by

$$(5) \quad (l_1, l_2, \dots, l_k) \rightarrow (l_{\pi(1)}, l_{\pi(2)}, \dots, l_{\pi(k)}), \quad \pi \in \Sigma_k - \{\text{id}\}$$

and

$$(6) \quad (l_1, l_2, \dots, l_k) \rightarrow (Jl_{\pi(1)}, Jl_{\pi(2)}, \dots, Jl_{\pi(k)}), \quad \pi \in \Sigma_k.$$

We claim these maps are fixed-point-free and pairwise noncoincident. Clearly the maps in (5) are fixed-point-free and pairwise noncoincident, and the maps in (6) are pairwise noncoincident. Suppose now we have a fixed point of a map in (6) or a coincidence between a map in (5) and one in (6), i.e., an element (l_1, l_2, \dots, l_k) of $F(1^k, n)$ with

$$l_{\pi(i)} = Jl_{\sigma(i)}, \quad 1 \leq i \leq k,$$

for some $\pi, \sigma \in \Sigma_k$. Then J fixes $l_1 \oplus l_2 \oplus \dots \oplus l_k$. But this is impossible, since J cannot fix an odd-dimensional subspace of \mathbf{C}^{n+k} .

Finally, we dispose of the case $k = 2$ and $n \geq 2$ even.

THEOREM 4.6. *If $n \geq 2$ is even, then $\#F(1^2, n) = \infty$.*

Proof. Let $J: \mathbf{C}^{n+2} \rightarrow \mathbf{C}^{n+2}$ be as defined above. Note that for $v \in \mathbf{C}^{n+2}$,

$$\langle Jv, v \rangle = \langle Jv, J^2v \rangle = -\langle Jv, v \rangle$$

by (4) above; thus $\langle Jv, v \rangle = 0$, and Jl is orthogonal to l for any line l . Define $\psi: F(1^2, n) \rightarrow F(1^2, n)$ by $\psi(l_1, l_2) = (Jl_1, l_1)$: then ψ is fixed-point-free and nonsurjective, and the conclusion follows by 2.2.

5. Proof of the endomorphism theorem. This section is devoted to a proof of Theorem 4.1. We use the notation of the previous section.

Since $t_i \in H^2(F(1^k, n); \mathbf{Q})$ is pulled back from $H^2(\mathbf{C}P^{n+k-1}; \mathbf{Q})$, we have $t_i^{n+k} = 0$. The next result gives a converse: it is proved in [8] for $k \leq n$, and in [2] without restriction.

THEOREM 5.1. *If $u \in H^2(F(1^k, n); \mathbf{Q})$ and $u^{n+k} = 0$, then u is of the form at_i for some $a \in \mathbf{Q}$ and $1 \leq i \leq k$.*

Now suppose f is an endomorphism of $H^*(F(1^k, n); \mathbf{Q})$. Then $f(t_i)^{n+k} = f(t_i^{n+k}) = 0$ for $1 \leq i \leq k$, and it follows from 5.1 that

$$f(t_i) = m_i t_{p(i)}, \quad 1 \leq i \leq k,$$

for some function $p: \{1, 2, \dots, k\} \rightarrow \{1, 2, \dots, k\}$ and rational numbers m_i . We shall prove that p is a permutation and all the m_i are equal unless $k = 2$ and n is even.

First we prove a technical lemma. The expression $h_i(x_1, x_2, \dots, x_r)$ denotes the i th complete symmetric function in x_1, x_2, \dots, x_r .

LEMMA 5.2. *Let $n \geq 0$ be an integer, a_1, a_2, \dots, a_r real numbers, and suppose*

$$h_{n+i}(a_1, a_2, \dots, a_r) = 0, \quad 1 \leq i \leq r - 1.$$

Then unless $r = 2$ and n is even, $a_1 = a_2 = \dots = a_r = 0$. If $r = 2$ and n is even, $a_2 = -a_1$.

Proof. Suppose first that $r = 2$. Then we have

$$(1) \quad h_{n+1}(a_1, a_2) = a_1^{n+1} + a_1^n a_2 + \dots + a_1 a_2^n + a_2^{n+1} = 0.$$

If $a_1 = a_2$, this evidently implies $a_1 = a_2 = 0$. If $a_1 \neq a_2$, then (1) is

$$\frac{a_1^{n+2} - a_2^{n+2}}{a_1 - a_2} = 0,$$

from which it follows that $a_2 = -a_1$ and n is even.

Now suppose $r = 3$. We have

$$h_{n+1}(a_1, a_2, a_3) = h_{n+2}(a_1, a_2, a_3) = 0.$$

Then from the relations

$$h_{n+2}(a_1, a_2, a_3) = a_1^{n+2} + a_1^{n+1}h_1(a_2, a_3) + \cdots + a_1h_{n+1}(a_2, a_3) + h_{n+2}(a_2, a_3)$$

and

$$a_1h_{n+1}(a_1, a_2, a_3) = a_1^{n+2} + a_1^{n+1}h_1(a_2, a_3) + \cdots + a_1h_{n+1}(a_2, a_3)$$

we get

$$(2) \quad h_{n+2}(a_2, a_3) = 0.$$

Similarly,

$$(3) \quad h_{n+2}(a_1, a_3) = 0$$

and

$$(4) \quad h_{n+2}(a_1, a_2) = 0.$$

By the argument of the preceding paragraph, these equations imply $a_1 = a_2 = a_3 = 0$ unless n is odd. In this case, (2) gives $a_3 = -a_2$, (4) gives $a_2 = -a_1$, and (3) gives $a_1 = -a_3$; but then $a_1 = a_2 = a_3 = 0$. It is now clear how to prove the result by induction for any $r > 3$.

As noted in the previous section, $H^*(F(1^k, n); \mathbf{Q})$ is the quotient of $\mathbf{Q}[t_1, \dots, t_k]$ by the ideal generated by R_1, R_2, \dots, R_k , where

$$R_i = h_{n+i}(t_1, t_2, \dots, t_k), \quad 1 \leq i \leq k.$$

For f to be a well-defined endomorphism of $H^*(F(1^k, n); \mathbf{Q})$ there must be relations

$$(5) \quad f(R_i) = N_i R_i + \sum_{1 \leq |\alpha| < i} N_i^{\alpha} t^{\alpha} R_{i-|\alpha|}, \quad 1 \leq i \leq k,$$

in $\mathbf{Q}[t_1, \dots, t_k]$, where the sum is over multi-indices $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_k)$ with $|\alpha| = \alpha_1 + \alpha_2 + \cdots + \alpha_k$ and

$$t^{\alpha} = t_1^{\alpha_1} t_2^{\alpha_2} \cdots t_k^{\alpha_k}.$$

Then we have the following result.

LEMMA 5.3. *If r of the elements t_1, t_2, \dots, t_k are missing from the image of f , then $f(R_i) = 0$ in $\mathbf{Q}[t_1, \dots, t_k]$ for $1 \leq i \leq r$.*

Proof. Permuting the t_i if necessary, we can assume that t_1, t_2, \dots, t_r are missing from the image of f . Define $\pi: F(1^k, n) \rightarrow F(1^{k-r}, n+r)$ by

$$\pi(l_1, l_2, \dots, l_k) = (l_{r+1}, \dots, l_k).$$

Then π^* sends $t_i \in H^2(F(1^{k-r}, n+r); \mathbf{Q})$ to $t_{r+i} \in H^2(F(1^k, n); \mathbf{Q})$ for $1 \leq i \leq k-r$, and is injective since the spectral sequence of the fibration

$$F(1^r, n) \rightarrow F(1^k, n) \xrightarrow{\pi} F(1^{k-r}, n+r)$$

collapses for degree reasons. Now t_1, \dots, t_r are missing from $\text{im } f$, so $\text{im } f \subset \text{im } \pi^*$ in $H^*(F(1^k, n); \mathbf{Q})$. Hence $f = \pi^*g$ for

$$g = (\pi^*)^{-1}f: H^*(F(1^k, n); \mathbf{Q}) \rightarrow H^*(F(1^{k-r}, n+r); \mathbf{Q}).$$

But the first nontrivial relation in $H^*(F(1^{k-r}, n+r); \mathbf{Q})$ is in dimension $2(n+r+1)$, so $g(R_i) = 0$ in $\mathbf{Q}[t_1, \dots, t_{k-r}]$ for $1 \leq i \leq r$ and the conclusion follows.

Suppose $n \geq 2$. For each i from 1 to k , we define the *weight* of i to be the cardinality of $\{r | p(r) = i \text{ and } m_r \neq 0\}$, i.e., the number of t_r that f maps to t_i with nonzero coefficient. The following result is the key to the proof of Theorem 4.1.

PROPOSITION 5.4. *Let $n \geq 2$. Then i has weight at most 1 for $1 \leq i \leq k$ unless $k = 2$ and n is even. If $k = 2$, n is even, and $q \in \{1, 2\}$ has weight 2, then f has the form*

$$f(t_i) = (-1)^i m t_q, \quad i = 1, 2.$$

Proof. Suppose q has weight $w > 1$. Then at least $w-1$ of the t_i are missing from the image of f , and by 5.3

$$(6) \quad f(R_i) = 0, \quad 1 \leq i \leq w-1,$$

in $\mathbf{Q}[t_1, \dots, t_k]$. We can assume t_1, t_2, \dots, t_w map to t_q with nonzero coefficient. Examine the coefficient of t_q^{n+i} in (6) to get

$$h_i(m_1, m_2, \dots, m_w) = 0, \quad 1 \leq i \leq w-1.$$

Then unless $w = 2$ and n is even, $m_1 = m_2 = \dots = m_w = 0$ by 5.2, a contradiction. If $w = 2$ and n is even, 5.2 gives $m_2 = -m_1 \neq 0$. In this case, $f(R_1) = 0$ and $f(R_2) \neq 0$ in $\mathbf{Q}[t_1, \dots, t_k]$; but then no i can have weight 1 and no more than one t_i can be missing from the image of f , from which follows $k = 2$.

REMARK. The case $n = 0$ is disposed of in [7], where it is proved that all endomorphisms of $H^*(F(1^k); \mathbf{Q})$ have the form h_m^σ .

By the preceding result, the function p of $\{1, 2, \dots, k\}$ can be assumed a permutation if $k \neq 2$ or n is odd. To finish the proof of 4.1, we need only show all the m_i are equal in this case. Now in $f(R_1)$ the coefficient of t_r^{n+1} is m_s^{n+1} , where $p(s) = r$. The coefficient of t_r^{n+1} on the right-hand side of (5) (with $i = 1$) is N_1 . Thus $m_s^{n+1} = N_1$ for $1 \leq s \leq k$. For n even, this shows all the m_s are equal. For n odd, it is also necessary to inspect the coefficients of terms $t_r^n t_s$, $r \neq s$, in equation (5) with $i = 1$.

REFERENCES

- [1] A. Borel, *Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts*, Ann. of Math., **57** (1953), 115–207.
- [2] S. A. Broughton, M. Hoffman and W. Homer, *The height of two-dimensional cohomology classes of complex flag manifolds*, Canad. Math. Bull., **26** (1983), 498–502.
- [3] R. F. Brown, *Path fields on manifolds*, Trans. Amer. Math. Soc., **118** (1965), 180–191.
- [4] R. F. Brown and E. Fadell, *Nonsingular path fields on compact topological manifolds*, Proc. Amer. Math. Soc., **16** (1965), 1342–1349.
- [5] E. Fadell and L. Neuwirth, *Configuration spaces*, Math. Scand., **10** (1962), 111–118.
- [6] H. Glover and W. Homer, *Fixed points on flag manifolds*, Pacific J. Math., **101** (1982), 303–306.
- [7] M. Hoffman, *On fixed point free maps of the complex flag manifold*, Indiana U. Math. J., **33** (1984), 249–255.
- [8] A. Liulevicius, *Homotopy rigidity of linear actions: characters tell all*, Bull. Amer. Math. Soc., **84** (1978), 213–221.
- [9] J. Vick, *Homology Theory*, Academic Press, New York, 1973.

Received March 4, 1983 and in revised form April 29, 1983.

OHIO STATE UNIVERSITY
COLUMBUS, OH 43210

Current address: U. S. Naval Academy
Annapolis, MD 21402

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)
University of California
Los Angeles, CA 90024

J. DUGUNDJI
University of Southern California
Los Angeles, CA 90089-1113

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

C. C. MOORE
University of California
Berkeley, CA 94720

ARTHUR OGUS
University of California
Berkeley, CA 94720

HUGO ROSSI
University of Utah
Salt Lake City, UT 84112

H. SAMELSON
Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$66.00 a year to individual members of supporting institutions. Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1984 by Pacific Journal of Mathematics

Ersan Akyildiz, Gysin homomorphism and Schubert calculus	257
Marilyn Breen, Clear visibility and unions of two starshaped sets in the plane	267
Robert F. Brown, Retraction methods in Nielsen fixed point theory	277
Herbert Busemann and Bhalchandra B. Phadke, A general version of Beltrami's theorem in the large	299
Gerald Arthur Edgar and Robert Francis Wheeler, Topological properties of Banach spaces	317
Yaakov Friedman and Bernard Russo, Conditional expectation without order	351
Robert Allen Goggins, Cobordism of manifolds with strong almost tangent structures	361
Mike Hoffman, Noncoincidence index of manifolds	373
William H. Julian, ε -continuity and monotone operations	385
Gerasimos E. Ladas, Y. G. Sficas and I. P. Stavroulakis, Nonoscillatory functional-differential equations	391
Arnold William Miller and Karel Libor Prikry, When the continuum has cofinality ω_1	399
Jean-Leah Mohrherr, Density of a final segment of the truth-table degrees	409
Carl Norman Mutchler, The flat Cauchy problem for radially hyperbolic operators from a characteristic manifold of high codimension	421
Kenji Nakagawa, On the orders of automorphisms of a closed Riemann surface	435
W. Ricker, Representation of vector-valued functions by Laplace transforms	445
Jorge Donato Samur, On semigroups of convolution operators in Hilbert space	463
Joseph Gail Stampfli, One-dimensional perturbations of operators	481
Andrew George Earnest and John Sollion Hsia, Correction to: "Spinor norms of local integral rotations. II"	493