

Pacific Journal of Mathematics

ε -CONTINUITY AND MONOTONE OPERATIONS

WILLIAM H. JULIAN

ε -CONTINUITY AND MONOTONE OPERATIONS

WILLIAM JULIAN

We prove constructively in the sense of Bishop that a monotone, ε -continuous operation from $[0, 1]$ into a metric space is 2ε -uniformly continuous. We derive a suitable version of Brouwer's fan theorem.

1. Introduction. Zaslavskii [ABR, Theorem 7.3] gives an example of a real valued function on $[0, 1]$ which is continuous at each computable point but which fails to be uniformly continuous. Zaslavskii [ABR, Theorem 7.14] and Mandelkern [MND1, MND2] show constructively in the Russian and Bishop sense, respectively, that a *monotone*, continuous, real valued function on $[0, 1]$ is uniformly continuous. In this paper, we weaken the hypothesis of continuity to ε -continuity, generalize the definition of monotone so that the map can be into any metric space, and consider (non-extensional) operations instead of functions. We prove constructively [BSH] that a monotone, ε -continuous operation from $[0, 1]$ into a metric space is 2ε -uniformly continuous. Delimiting examples show that the 2ε in the conclusion is best possible.

2. Valuated fans. The *binary fan* F consists of all finite or empty strings from $\{0, 1\}$. Denote a string $a \in F$ by $a_1 a_2 \cdots a_n$ where $a_i \in \{0, 1\}$, and the empty string by \emptyset . The *length* $|a|$ of a is the cardinality n of the string a . The *descendants* of string a are strings containing a as an initial segment. The *immediate descendants* of a are $a0 = a_1 a_2 \cdots a_n 0$ and $a1 = a_1 a_2 \cdots a_n 1$. Note $\emptyset 0 = 0$ and $\emptyset 1 = 1$. A *branch* B is the set of initial segments of a countable string $B_1 B_2 \cdots$ from $\{0, 1\}$. We shall write $B \sim B_1 B_2 \cdots$.

A *valuated fan* F is the binary fan together with a function V mapping F into the set N of non-negative integers. A valuation is *sub-additive* if $V(a) \geq V(a0) + V(a1)$, for all $a \in F$.

A valuation is *branch bounded* if for any branch B of F there is an integer n so that if $a \in B$ and $|a| \geq n$, then $V(a) = 0$. A valuation is *bounded* if there is an integer m so that if $a \in F$ and $|a| \geq m$, then $V(a) = 0$.

The *valuated fan generated by* $a \in F$ consists all descendents of a but with their initial segments a deleted; the valuation is the induced valuation.

We now arrive at a theorem implied by Brouwer's fan theorem [HTG] but that is valid in the sense of Bishop [BSH].

PROPOSITION 1. *Every branch bounded, sub-additive valuation on the binary fan is bounded.*

Proof. We induct on the value $V(\emptyset)$. If $V(\emptyset) = 0$ we are done, so let $V(\emptyset) > 0$. Construct a branch B starting at \emptyset by induction. If $a \in B$ and $V(a1) = V(\emptyset)$, then append $a1$ to B . Otherwise append $a0$. Since F is sub-additive, if $a \in F$ and $V(a) = V(\emptyset)$ then $a \in B$. Since F is branch bounded, there is an integer n so that if $|a| \geq n$ and $a \in B$, then $V(\emptyset) = 0$. Hence if $|a| \geq n$ then $V(a) < V(\emptyset)$. Construct the 2^n fans F_i generated by those $a \in F$ with $|a| = n$. In each, the induced value $V_i(\emptyset)$ is strictly less than $V(\emptyset)$ in F . By induction, each is bounded: There are integers m_i such that if $|b| \geq m_i$ and $b \in F_i$, then the induced value $V_i(b) = 0$. Hence if $a \in F$ and $|a| \geq n + \max\{m_i\}$, then $V(a) = 0$. \square

3. Assigning valuations. In this section we show how an operation from $[0, 1]$ induces a valuation on the binary fan. To each $a = a_1 a_2 \cdots a_n \in F$ assign the diadic rationals

$$.a = \sum a_k 2^{-k} = .a_1 a_2 \cdots a_n \quad (\text{binary}),$$

$$.a^+ = .a + 2^{-|a|},$$

and the interval $I(a) = [.a, .a^+]$. To each branch $B \sim B_1 B_2 \cdots$ assign the real number

$$.B = \sum B_k 2^{-k} = .B_1 B_2 \cdots \quad (\text{binary}).$$

Note that if $a \in B$, then $.B \in I(a)$.

DEFINITION. We denote two subsets of $[0, 1]$ by

$$B[0, 1] = \{x \in [0, 1] \mid x \text{ has an explicit binary representation}\},$$

and

$$D[0, 1] = \{x \in [0, 1] \mid x \text{ has a terminating binary representation}\}.$$

Note that $x \in B[0, 1]$ iff $x \geq d$ or $x \leq d$ for every diadic rational $d \in D[0, 1]$. \square

DEFINITION. Let f be an operation on $B[0, 1]$ into a metric space M , d and $\varepsilon > 0$. Fix one value of $f(x)$ for each $x \in D[0, 1]$. A *valuation* on the

binary fan *induced* by f , ε is assigned so that

$$V(a) = P \text{ implies } P - 2^{-2|a|} < \rho(a) < P + 1 - 2^{-2|a|^{-1}},$$

where $\rho(a) = \varepsilon^{-1}d(f(.a^+), f(.a))$. □

Note that if $\rho(a) > P - 2^{-2|a|^{-1}}$, then $V(a) \geq P$, and if $\rho(a) < 1 - 2^{-2|a|}$, then $V(a) = 0$.

4. Monotone operations. In this section we consider what valuation is induced on the binary fan by a monotone operation into a metric space. The notion of “between” replaces “order” in the definition of monotone.

DEFINITION. Let M, d be a metric space. A point $x \in M$ is *between* a and $b \in M$ if

$$d(a, x) + d(x, b) = d(a, b).$$

In addition if x is distinct from a and b , then x is *strictly between* a and b . □

The notion of “between” has been discussed by Blumenthal [BLM]; his use of “between” corresponds to our usage of “strictly between”. We distinguish the present notions of “between” and “strictly between” in the next definition:

DEFINITION. An operation f from a metric space M_1 to a metric space M_2 is *monotone* if whenever x is strictly between a and $b \in M_1$, then $f(x)$ is between $f(a)$ and $f(b)$. □

LEMMA 1. *If x and y are between a and b then $d(x, y) \leq d(a, b)$.*

Proof. Let x and y be between a and b . Thus, adding

$$d(a, z) + d(z, b) = d(a, b)$$

for z equal to x and z equal to y , we obtain

$$2d(x, y) \leq d(a, x) + d(a, y) + d(b, x) + d(b, y) = 2d(a, b). \quad \square$$

The next lemmas and a counterexample stated without proof show how *order* and *between* are related on the real line.

LEMMA. *If x is between distinct points a and b and not strictly between them, then $x = a$ or $x = b$.* □

LEMMA. *A real number x is (strictly) between a and $b \in R$ if $(a < x < b)$ $a \leq x \leq b$, or if $(a > x > b)$ $a \geq x \geq b$.* \square

LEMMA. *If x is strictly between a and $b \in R$, then $a < x < b$ or $b < x < a$.* \square

COUNTEREXAMPLE. If x between a and $b \in R$ implies $a \leq x \leq b$ or $a \geq x \geq b$, then for all $a \in R$, either $a \geq 0$ or $a \leq 0$. \square

A real valued function which is monotone in the present sense need not be increasing or decreasing.

LEMMA. *If f is a monotone operation on $S \subset R$ to a metric space M , d and $a \leq x < b$ are in S with $d(f(a), f(x)) + d(f(x), f(b)) > d(f(a), f(b))$, then $x = a$.* \square

Next we show that a monotone operation on $[0, 1]$ induces a sub-additive valuation on the binary fan.

PROPOSITION 2. *If f is a monotone operation from $B[0, 1]$ to a metric space M , d and $\varepsilon > 0$, then the valuation induced by f , ε is sub-additive.*

Proof. Now $|a0| = |a1| = |a| + 1$, so

$$V(a0) - 2^{-2|a|-2} < \rho(a0) \quad \text{and} \quad V(a1) - 2^{-2|a|-2} < \rho(a1).$$

Noting that monotonicity of f implies that $\rho(a) = \rho(a0) + \rho(a1)$, we find

$$V(a0) + V(a1) - 2^{-2|a|-1} < \rho(a).$$

Hence $V(a) \geq V(a0) + V(a1)$ and the valuation is sub-additive. \square

5. ε -continuous operations. In this section we turn our attention to what valuation on the binary fan is induced by an ε -continuous operation.

DEFINITION. An operation f from a metric space M_1 , d_1 to a metric space M_2 , d_2 is ε -continuous if for some $\varepsilon' < \varepsilon$ then for every $x \in M_1$ there is a $\delta > 0$ such that whenever $y \in M_1$ and $d_1(x, y) < \delta$, then $d_2(f(x), f(y)) < \varepsilon'$. \square

Note that if $\varepsilon' < \varepsilon'' < \varepsilon$ then f is also ε'' -continuous. Furthermore if $x = y$ then $d_2(f(x), f(y)) < \varepsilon'$.

PROPOSITION 3. *If f is an $\epsilon/2$ -continuous operation from $B[0, 1]$ to a metric space M , d then the valuation induced by f , ϵ is branch bounded.*

Proof. Let B be a branch of the binary fan F . Choose $\epsilon' < \epsilon/2$ and $\delta > 0$ such that if $y \in B[0, 1]$ and $|y - .B| < \delta$, then $d(f(y), f(.B)) < \epsilon'$. Pick n so that $2^{-n} < \delta$ and $2\epsilon' < \epsilon(1 - 2^{-2n})$, and let $a \in B$ with $|a| \geq n$. Now $.B \in I(a)$ so $|a - .B| < \delta$ and $|.a^+ - .B| < \delta$. Hence

$$\begin{aligned} d(f(.a^+), f(.a)) &\leq d(f(.a^+), f(.B)) + d(f(.B), f(.a)) \\ &< 2\epsilon' < \epsilon(1 - 2^{-2|a|}), \end{aligned}$$

and thus $V(a) = 0$. □

6. ϵ -uniformly continuous operations. In this section we prove that a monotone, ϵ -continuous operation on $[0, 1]$ is 2ϵ -uniformly continuous.

DEFINITION. An operation f is ϵ -uniformly continuous from a metric space M_1, d_1 to a metric space M_2, d_2 if there is $\epsilon' < \epsilon$ and $\delta > 0$ such that whenever $x, y \in M_1$ and $d_1(x, y) < \delta$, then $d_2(f(x), f(y)) < \epsilon'$.

THEOREM. *If f is a monotone operation from $[0, 1]$ into a metric space M , d and $\epsilon/2$ -continuous on $B[0, 1]$ then f is ϵ -uniformly continuous on $[0, 1]$.*

Proof. Choose $\epsilon' < \epsilon$ so that f is also $\epsilon'/2$ -continuous on $B[0, 1]$. Let the binary fan F have the valuation induced by f , ϵ' . By Proposition 3 the valuation is branch bounded, and by Propositions 1 and 2, the valuation is bounded. Hence, there is an m so that if $a \in F$ and $|a| \geq m$, then $V(a) = 0$.

Consider the finite set $S = \{1\} \cup \{.a \mid a \in F \text{ and } |a| = m\}$. By $\epsilon'/2$ -continuity, choose δ in $(0, 2^{-m})$ such that if $x \in [0, 1]$, $z \in S$, and $|x - z| < \delta$, then $d(f(x), f(z)) < \epsilon'/2$. Suppose that $x, y \in [0, 1]$ and $|x - y| < \delta/3$. Either $|x - z| < \delta/2$ for some $z \in S$, or $|x - z| > \delta/3$ for each $z \in S$. In the former case $|y - z| < 5\delta/6$, and

$$d(f(x), f(y)) \leq d(f(x), f(z)) + d(f(z), f(y)) < \epsilon' < \epsilon.$$

In the latter case, pick $a \in F$ with $|a| = m$, such that x and y are strictly between $.a$ and $.a^+$; then by Lemma 1 and $V(a) = 0$:

$$d(f(x), f(y)) \leq d(f(.a), f(.a^+)) < \epsilon' < \epsilon. \quad \square$$

7. Delimiting examples. The theorem is valid with $[0, 1]$ replaced by $B[0, 1]$. The first example shows that the result stated in the theorem is

sharp with regard to ε . There is no obvious constructive example, so we give a classical one.

EXAMPLE 1 (Classical). The classical function

$$f(x) = \begin{cases} \varepsilon'/2, & \text{for } x > 1/2, \\ 0, & \text{for } x = 1/2, \\ -\varepsilon'/2, & \text{for } x < 1/2 \end{cases}$$

is monotone and $\varepsilon/2$ -continuous for all $\varepsilon > \varepsilon' > 0$, but is not ε' -uniformly continuous. \square

The next example shows that there are constructive ε -continuous operations which are neither continuous nor functions.

EXAMPLE 2 (Constructive). Let $x = .x_1x_2x_3 \cdots \in B[0, 1]$. The operation

$$g(x) = \varepsilon'(x_1 - 1/2)$$

is ε -continuous and ε -uniformly continuous on $B[0, 1]$ for any $\varepsilon > \varepsilon' > 0$.

REFERENCES

- [ABR] O. Aberth, *Computable Analysis*, McGraw-Hill, NY, 1980.
- [BSH] E. A. Bishop, *Foundations of Constructive Analysis*, McGraw-Hill, NY 1967.
- [BLM] L. Blumenthal, *Theory and Applications of Distance Geometry*, Chelsea Publishing Company, Bronx, NY 1970.
- [HTG] A. Heyting, *Intuitionism, an Introduction*, 3rd ed., North-Holland Publishing Co., Amsterdam 1966.
- [MND1] M. Mandelkern, *Continuity of monotone functions*, Pacific J. Math., **99** (1982), 413–418.
- [MND2] ———, *Constructive continuity*, Mem. Amer. Math. Soc., **42** (1983), 277.

Received April 28, 1983.

NEW MEXICO STATE UNIVERSITY
LAS CRUCES, NM 88003

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DONALD BABBITT (Managing Editor)
University of California
Los Angeles, CA 90024

J. DUGUNDJI
University of Southern California
Los Angeles, CA 90089-1113

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

C. C. MOORE
University of California
Berkeley, CA 94720

ARTHUR OGUS
University of California
Berkeley, CA 94720

HUGO ROSSI
University of Utah
Salt Lake City, UT 84112

H. SAMELSON
Stanford University
Stanford, CA 94305

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$66.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1984 by Pacific Journal of Mathematics

Ersan Akyildiz, Gysin homomorphism and Schubert calculus	257
Marilyn Breen, Clear visibility and unions of two starshaped sets in the plane	267
Robert F. Brown, Retraction methods in Nielsen fixed point theory	277
Herbert Busemann and Bhalchandra B. Phadke, A general version of Beltrami's theorem in the large	299
Gerald Arthur Edgar and Robert Francis Wheeler, Topological properties of Banach spaces	317
Yaakov Friedman and Bernard Russo, Conditional expectation without order	351
Robert Allen Goggins, Cobordism of manifolds with strong almost tangent structures	361
Mike Hoffman, Noncoincidence index of manifolds	373
William H. Julian, ε -continuity and monotone operations	385
Gerasimos E. Ladas, Y. G. Sficas and I. P. Stavroulakis, Nonoscillatory functional-differential equations	391
Arnold William Miller and Karel Libor Prikry, When the continuum has cofinality ω_1	399
Jean-Leah Mohrherr, Density of a final segment of the truth-table degrees	409
Carl Norman Mutchler, The flat Cauchy problem for radially hyperbolic operators from a characteristic manifold of high codimension	421
Kenji Nakagawa, On the orders of automorphisms of a closed Riemann surface	435
W. Ricker, Representation of vector-valued functions by Laplace transforms	445
Jorge Donato Samur, On semigroups of convolution operators in Hilbert space	463
Joseph Gail Stampfli, One-dimensional perturbations of operators	481
Andrew George Earnest and John Sollion Hsia, Correction to: "Spinor norms of local integral rotations. II"	493