

Pacific Journal of Mathematics

**TRIDIAGONAL MATRIX REPRESENTATIONS OF CYCLIC
SELF-ADJOINT OPERATORS. II**

JOANNE MARIE DOMBROWSKI

TRIDIAGONAL MATRIX REPRESENTATIONS OF CYCLIC SELF-ADJOINT OPERATORS. II

J. DOMBROWSKI

A bounded cyclic self-adjoint operator C defined on a separable Hilbert space H can be represented as a tridiagonal matrix with respect to the basis generated by the cyclic vector. An operator J can then be defined so that $CJ - JC = -2iK$ where K also has tridiagonal form. If the subdiagonal elements of C converge to a non-zero limit and if K is of trace class then C must have an absolutely continuous part.

1. Introduction. A bounded cyclic self-adjoint operator C defined on a separable Hilbert space H has a tridiagonal matrix representation with respect to the basis generated by the cyclic vector. The spectral properties of C are studied in [1] under the assumption that the main diagonal elements in the tridiagonal representation are zeros. In this case C is the real part of a weighted shift operator, and if J is the corresponding imaginary part then $CJ - JC = 2iK$ where K is diagonal. It is shown in [1] that C has an absolutely continuous part if K is of trace class.

The purpose of this paper is to extend the above result to tridiagonal matrices with non-zero diagonal. This extension is significant for the study of systems of orthogonal polynomials which satisfy a three term recursion formula. The coefficients in the recursion formula correspond to a unique tridiagonal matrix whose spectral measure is the measure of orthogonality for the system of polynomials. If the measure is symmetric about the origin (as in the case of the normalized Legendre polynomials) then the diagonal entries of the corresponding tridiagonal matrix will be zero. See [1] for further background.

Suppose now that C is a bounded cyclic self-adjoint operator with the following tridiagonal matrix representation with respect to the basis $\{\phi_n\}$:

$$(1.1) \quad C = \begin{bmatrix} b_1 & a_1 & 0 & 0 & \cdots \\ a_1 & b_2 & a_2 & 0 & \cdots \\ 0 & a_2 & b_3 & a_3 & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix}, \quad a_n > 0.$$

Then $\phi_n = P_n(C)\phi_1$ with $P_1(\lambda) = 1$, $P_2(\lambda) = (\lambda - b_1)/a_1$ and for $n > 2$,

$$P_n(\lambda) = \frac{(\lambda - b_{n-1})P_{n-1}(\lambda) - a_{n-2}P_{n-2}(\lambda)}{a_{n-1}}.$$

It follows that C is the real part of the operator T defined by $T\phi_n = b_n\phi_n + 2a_n\phi_{n+1}$. If $J = (1/2i)(T + T^*)$ then $CJ - JC = -2iK$ where $K = [k_{ij}]$ has tridiagonal form with

$$(1.2) \quad \begin{aligned} k_{ii} &= -a_{i-1}^2 + a_i^2 \quad (a_0 = 0), \\ k_{i,i+1} &= k_{i+1,i} = \frac{1}{2}a_i(b_i - b_{i+1}). \end{aligned}$$

The aim now is to analyze the spectrum of C if $\sum |k_{ii}| < \infty$ and $\sum |k_{i,i+1}| < \infty$. Note that in this case $\lim a_n$ and $\lim b_n$ exist. If $\lim a_n = \alpha$ ($\alpha \neq 0$) and $\lim b_n = 0$ then the spectrum of C at least contains the interval $(-2\alpha, 2\alpha)$. Note also that the diagonals of K are absolutely summable whenever $K \geq 0$ so that T is hyponormal. For if $K = [k_{ij}]$ is positive then $\{a_n^2\}$ is an increasing sequence and $|k_{ij}|^2 \leq |k_{ii}||k_{jj}|$. It follows that

$$\frac{1}{2} \sum |a_i| |b_{i+1} - b_i| \leq \left(\sum |-a_{i-1}^2 + a_i^2| \right)^{1/2} \left(\sum |-a_i^2 + a_{i+1}^2| \right)^{1/2}.$$

Furthermore, the following is true:

THEOREM. *The operator K in (1.2) is of trace class if and only if $\sum |k_{ii}| < \infty$ and $\sum |k_{i,i+1}| < \infty$.*

Proof. Assume that the diagonals of K are absolutely summable. It follows from Jensen's Inequality that

$$\sum \langle (K^*K)^{1/2} \phi_n, \phi_n \rangle \leq \sum \langle K^*K \phi_n, \phi_n \rangle^{1/2}$$

and a direct computation shows that if $K^*K = [p_{ij}]$ then

$$\begin{aligned} p_{11} &= a_1^4 + \frac{1}{4}(-a_1b_1 + a_1b_2)^2, \\ p_{ii} &= \frac{1}{4}(-a_{i-1}b_{i-1} + a_{i-1}b_i)^2 + (-a_{i-1}^2 + a_i^2)^2 \\ &\quad + \frac{1}{4}(-a_ib_i + a_ib_{i+1})^2 \quad \text{for } i > 1. \end{aligned}$$

Since

$$\begin{aligned} \sum p_{ii}^{1/2} &\leq \sqrt{2} \left(a_1^2 + \frac{1}{2} |-a_1b_1 + a_1b_2| \right) \\ &\quad + \sqrt{3} \sum \left(\frac{1}{2} |-a_{i-1}b_{i-1} + a_{i-1}b_i| + |-a_{i-1}^2 + a_i^2| \right. \\ &\quad \left. + \frac{1}{2} |-a_ib_i + a_ib_{i+1}| \right), \end{aligned}$$

K is of trace class.

Suppose K is of trace class. Write $K = D + W$ where D is diagonal and W is the real part of a weighted shift operator. Since $\sum |\langle K\phi_n, \phi_n \rangle| < \infty$, D (and hence W) must be of trace class. Furthermore, W with weight

sequence $\{w_i\}$ is unitarily equivalent to W' with weight sequence $\{|w_i|\}$. If V denotes the unilateral shift operator then $W'(V + V^*)$ with diagonal elements $|w_1|$, $|w_1| + |w_2|$, $|w_2| + |w_3|, \dots$, is of trace class. Hence the diagonals of K are absolutely summable.

2. The absolutely continuous part of C . If $C = \int \lambda dE_\lambda$ let $H_a(C)$ denote the set of x in H for which $\|E_\lambda x\|^2$ is an absolutely continuous function of λ . The subspace $H_a(C)$ reduces C and the restriction of C to $H_a(C)$ is called the absolutely continuous part of C . If C is represented as a multiplication operator on $L^2(\mu)$ these ideas are related to the decomposition of the spectral measure μ .

The following lemma is needed for the proof of the main result.

LEMMA. *If*

$$S_{N+1}(\lambda) = a_1^2 P_1^2(\lambda) + \sum_{n=2}^{N+1} [(-a_{n-1}^2 + a_n^2) P_n^2(\lambda) + a_{n-1}(-b_{n-1} + b_n) P_{n-1}(\lambda) P_n(\lambda)],$$

then

$$S_{N+1}(\lambda) = a_N^2 P_N^2(\lambda) - a_N(\lambda - b_{N+1}) P_N(\lambda) P_{N+1}(\lambda) + a_{N+1}^2 P_{N+1}^2(\lambda).$$

Proof. Note that

$$\begin{aligned} S_2(\lambda) &= a_1^2 P_1^2(\lambda) + a_1(-b_1 + b_2) P_1(\lambda) P_2(\lambda) \\ &\quad - a_1(\lambda - b_1) P_2(\lambda) P_1(\lambda) + a_2^2 P_2^2(\lambda) \\ &= a_1^2 P_1^2(\lambda) + a_1(b_2 - \lambda) P_1(\lambda) P_2(\lambda) + a_2^2 P_2^2(\lambda). \end{aligned}$$

Assume that the result has been verified for $N = 1, \dots, k$ and recall that $a_k P_{k+1}(\lambda) = (\lambda - b_k) P_k(\lambda) - a_{k-1} P_{k-1}(\lambda)$. Then

$$\begin{aligned} S_{k+1}(\lambda) &= a_{k-1}^2 P_{k-1}^2(\lambda) - a_{k-1}(\lambda - b_k) P_{k-1}(\lambda) P_k(\lambda) \\ &\quad + a_k^2 P_k^2(\lambda) - [(\lambda - b_k) P_k(\lambda) - a_{k-1} P_{k-1}(\lambda)]^2 \\ &\quad + a_{k+1}^2 P_{k+1}^2(\lambda) + (-a_k b_k + a_k b_{k+1}) P_k(\lambda) P_{k+1}(\lambda) \\ &= a_k^2 P_k^2(\lambda) - (\lambda - b_k) P_k(\lambda) [(\lambda - b_k) P_k(\lambda) - a_{k-1} P_{k-1}(\lambda)] \\ &\quad + a_{k+1}^2 P_{k+1}^2(\lambda) + (-a_k b_k + a_k b_{k+1}) P_k(\lambda) P_{k+1}(\lambda) \\ &= a_k^2 P_k^2(\lambda) - a_k(\lambda - b_{k+1}) P_k(\lambda) P_{k+1}(\lambda) + a_{k+1}^2 P_{k+1}^2(\lambda). \end{aligned}$$

THEOREM. *Let $\lim a_n = 1/2$ and $\lim b_n = 0$. If $\sum | -a_{n-1}^2 + a_n^2 | < \infty$ and $\sum |a_{n-1}(-b_{n-1} + b_n)| < \infty$ then the spectral measure of C is absolutely continuous on $(-1, 1)$.*

Proof. The plan is to first consider an operator obtained from C by replacing a finite number of constants in such a way as to simplify the structure of K and make use of the absolute summability of the diagonals. Given $k > 1$ choose $R(k)$ such that

$$\sum_{R+1}^{\infty} |-a_{n-1}^2 + a_n^2| \leq \frac{1}{32k}, \quad \sum_{R+1}^{\infty} |a_{n-1}(-b_{n-1} + b_n)| < \frac{1}{32k}$$

and such that if $n > R$ then $|4a_n^2 - 1| < 1/16k$, $|b_n| < 1/16k$. Obtain C_k from C by replacing the constants b_1, \dots, b_R by 0 and the constants a_1, \dots, a_R by $1/2$. It will now be shown that the spectral measure of C_k is absolutely continuous on $(-1 + 1/k, 1 - 1/k)$. Since C is a trace class perturbation of C_k the same result will then hold for C .

Let Δ be a subinterval of $(-1 + 1/k, 1 - 1/k)$ and for $C_k = \int \lambda dE_\lambda$ let $\mu(\Delta) = \|E(\Delta)\phi_1\|^2$. With J and K defined as before,

$$\begin{aligned} (2.1) \quad & \langle KE(\Delta)\phi_1, E(\Delta)\phi_1 \rangle \\ &= a_1^2 |\langle E(\Delta)\phi_1, \phi_1 \rangle|^2 \\ &+ \sum_{n=2}^{\infty} \left[a_{n-1}(-b_{n-1} + b_n) \langle E(\Delta)\phi_1, \phi_{n-1} \rangle \langle E(\Delta)\phi_1, \phi_n \rangle \right. \\ &\quad \left. + (-a_{n-1}^2 + a_n^2) |\langle E(\Delta)\phi_1, \phi_n \rangle|^2 \right] \\ &= a_1^2 \left| \int_{\Delta} P_1 d\mu \right|^2 + \sum_{n=2}^{\infty} \left[a_{n-1}(-b_{n-1} + b_n) \int_{\Delta} P_{n-1} d\mu \int_{\Delta} P_n d\mu \right. \\ &\quad \left. + (a_{n-1}^2 + a_n^2) \left| \int_{\Delta} P_n d\mu \right|^2 \right]. \end{aligned}$$

Several cases need to be considered.

Case I. Suppose $\int_{\Delta} P_n^2 d\mu \leq \mu(\Delta) = \int_{\Delta} P_1^2 d\mu$ for each n . Then

$$\begin{aligned} & \langle KE(\Delta)\phi_1, E(\Delta)\phi_1 \rangle \\ & \geq a_1^2 |\mu(\Delta)|^2 - \mu(\Delta) \sum_{n=M+1}^{\infty} \left[|a_{n-1}(-b_{n-1} + b_n)| \left(\int_{\Delta} P_{n-1}^2 d\mu \right)^{1/2} \right. \\ & \quad \left. \times \left(\int_{\Delta} P_n^2 d\mu \right)^{1/2} + |-a_{n-1}^2 + a_n^2| \int_{\Delta} P_n^2 d\mu \right] \\ & \geq |\mu(\Delta)|^2 \left(\frac{1}{4} - \frac{1}{16k} \right). \end{aligned}$$

Case II. Suppose $\int_{\Delta} P_n^2 d\mu \leq \mu(\Delta) = \int_{\Delta} P_1^2 d\mu$ except for a finite number of n . Then there exists N such that $\int_{\Delta} P_N^2 d\mu = \max \int_{\Delta} P_n^2 d\mu$. Adding and subtracting

$$\mu(\Delta) \sum_{n=2}^N \int [a_{n-1}(-b_{n-1} + b_n)P_{n-1}P_n + (-a_{n-1}^2 + a_n^2)P_n^2] d\mu$$

to the expansion of $\langle KE(\Delta)\phi_1, E(\Delta)\phi_1 \rangle$ in (2.1) and using the expression for S_N given in the lemma it follows that

$$\begin{aligned} & \langle KE(\Delta)\phi_1, E(\Delta)\phi_1 \rangle \\ &= \mu(\Delta) \left\{ \int \left[a_{N-1}P_{N-1}(\lambda) - \frac{1}{2}(\lambda - b_N)P_N(\lambda) \right]^2 d\mu \right. \\ & \qquad \qquad \qquad \left. + \int \left[a_N^2 - \frac{1}{4}(\lambda - b_N)^2 P_N^2(\lambda) \right] d\mu \right\} \\ & - \mu(\Delta) \sum_{n=2}^N \int [a_{n-1}(-b_{n-1} + b_n)P_{n-1}P_n + (-a_{n-1}^2 + a_n^2)P_n^2] d\mu \\ & + \sum_{n=2}^{\infty} \left[a_{n-1}(-b_{n-1} + b_n) \int_{\Delta} P_{n-1} d\mu \int_{\Delta} P_n d\mu \right. \\ & \qquad \qquad \qquad \left. + (-a_{n-1}^2 + a_n^2) \left| \int_{\Delta} P_n d\mu \right|^2 \right] \\ & \geq \frac{1}{4} \mu(\Delta) \int [4a_N^2 - (\lambda - b_N)^2] P_N^2(\lambda) d\mu - \frac{1}{8k} \mu(\Delta) \int_{\Delta} P_N^2 d\mu \\ & \geq |\mu(\Delta)|^2 \frac{3}{32k}. \end{aligned}$$

Case III. Suppose there exists a subsequence $\{P_{n_j}\}$ such that $\int_{\Delta} P_{n_j}^2 d\mu \geq \mu(\Delta)$. Choose L sufficiently large such that

$$\begin{aligned} & \sum_{L+1}^{\infty} \left[a_{n-1}(-b_{n-1} + b_n) \int_{\Delta} P_{n-1} d\mu \int_{\Delta} P_n d\mu + (-a_{n-1}^2 + a_n^2) \left| \int_{\Delta} P_n d\mu \right|^2 \right] \\ & \leq \mu(\Delta) \sum_{L+1}^{\infty} [|a_{n-1}(-b_{n-1} + b_n)| + |-a_{n-1}^2 + a_n^2|] \leq \frac{1}{32k} |\mu(\Delta)|^2. \end{aligned}$$

Choose $M > L$ such that $\int_{\Delta} P_M^2 d\mu \geq \mu(\Delta)$. Then there exists N such that $\int_{\Delta} P_N^2 d\mu = \max_{1 \leq n \leq M} \int_{\Delta} P_n^2 d\mu$. Using the same approach as in Case II it

can be shown that

$$\begin{aligned}
& \langle KE(\Delta)\phi_1, E(\Delta)\phi_1 \rangle \\
& \geq \mu(\Delta) \int_{\Delta} \left[a_N^2 - \frac{1}{4}(\lambda - b_N)^2 \right] P_N^2(\lambda) d\mu \\
& \quad - \mu(\Delta) \sum_{n=2}^N \int [a_{n-1}(-b_{n-1} + b_n) P_{n-1} P_n d\mu + (-a_{n-1}^2 + a_n^2) P_n^2] d\mu \\
& \quad + \sum_{n=2}^{\infty} \left[a_{n-1}(-b_{n-1} + b_n) \int_{\Delta} P_{n-1} d\mu \int_{\Delta} P_n d\mu \right. \\
& \qquad \qquad \qquad \left. + (-a_{n-1}^2 + a_n^2) \left| \int_{\Delta} P_n d\mu \right|^2 \right] \\
& \geq \mu(\Delta) \int_{\Delta} \left[a_N^2 - \frac{1}{4}(\lambda - b_N)^2 \right] P_N^2(\lambda) d\mu \\
& \quad - \mu(\Delta) \int_{\Delta} P_N^2 d\mu \sum_2^N [|a_{n-1}(-b_{n-1} + b_n)| + |-a_{n-1}^2 + a_n^2|] \\
& \quad - \mu(\Delta) \int_{\Delta} P_N^2 d\mu \sum_{n=2}^M [|a_{n-1}(-b_{n-1} + b_n)| + |-a_{n-1}^2 + a_n^2|] \\
& \quad - \left| \sum_{M+1}^{\infty} [a_{n-1}(-b_{n-1} + b_n)] \int_{\Delta} P_{n-1} d\mu \int_{\Delta} P_n d\mu \right. \\
& \qquad \qquad \qquad \left. + (-a_{n-1}^2 + a_n^2) \left(\int_{\Delta} P_n d\mu \right)^2 \right| \\
& \geq \mu(\Delta) \frac{1}{32k} \int_{\Delta} P_N^2 d\mu - \frac{1}{32k} |\mu(\Delta)|^2 \geq \frac{1}{16k} |\mu(\Delta)|^2.
\end{aligned}$$

In each case, therefore, $\langle KE(\Delta)\phi_1, E(\Delta)\phi_1 \rangle \geq (1/16k)|\mu(\Delta)|^2$. It can also be shown (see the proof of Theorem 2.2.4 in [4]) that

$$|\langle KE(\Delta)\phi_1, E(\Delta)\phi_1 \rangle| \leq \frac{1}{2} \|J\| |\Delta| \|E(\Delta)\phi_1\|^2$$

where $|\Delta|$ denotes the length of Δ . It therefore follows that $|\mu(\Delta)| \leq 8k \|J\| |\Delta|$. Let β be a Borel subset of $(-1 + 1/k, 1 - 1/k)$ of Lebesgue measure zero. Then for any $\varepsilon > 0$ there exists a pairwise disjoint sequence of intervals $\{\Delta_j\}$ such that $\Delta_j \subset (-1 + 1/k, 1 - 1/k)$, $\beta \subset \bigcup \Delta_j$ and $\sum |\Delta_j| < \varepsilon$. Since $\mu(\beta) \leq \sum \mu(\Delta_j) \leq 8k \|J\| \sum |\Delta_j|$ it follows that $\mu(\beta) = 0$.

Hence the spectral measure of C_k is absolutely continuous on $(-1 + 1/k, 1 - 1/k)$. By the Kato-Rosenblum Theorem the same result holds for the spectral measure of C .

COROLLARY. *Let $\lim a_n = \alpha$ ($\alpha \neq 0$) and $\lim b_n = b$. If*

$$\sum | -a_{n-1}^2 + a_n^2 | < \infty \quad \text{and} \quad \sum | a_{n-1}(-b_{n-1} + b_n) | < \infty$$

then the spectral measure of C is absolutely continuous on $(-2\alpha + b, 2\alpha + b)$.

Finally it should be observed that the above theorem becomes false under the weaker condition that K is compact. To see this assume that the main diagonal elements of C are zero and define $a_{2n-1}^2 = 1$, $a_{2n}^2 = (n+1)^2/n^2$ for $n = 1, 2, \dots$. Since $P_n^2(0) = (a_{n-2}^2/a_{n-1}^2)P_{n-2}^2(0)$ it follows that $P_{2n}^2(0) = 0$ and $P_{2n+1}^2(0) = 1/(n+1)^2$. Therefore $\lambda = 0$ is an eigenvalue. (See Stone [5].)

Note. A similar result has recently been attained by Mate and Nevai in [2] with very different techniques.

REFERENCES

- [1] J. Dombrowski, *Tridiagonal matrix representations of cyclic self-adjoint operators*, to appear in *Pacific Math*.
- [2] A. Mate and P. Nevai, *Orthogonal polynomials and absolutely continuous measures*, preprint.
- [3] P. Nevai, *Orthogonal polynomials*, *Mem. Amer. Math. Soc.*, **18** (1979).
- [4] C. R. Putnam, *Commutation Properties of Hilbert space Operators and Related Topics*, *Ergebnisse der Math.*, **36**, Springer, 1967.
- [5] M. H. Stone, *Linear transformations in Hilbert space*, *Amer. Math. Soc.*, New York, 1932.

Received January 20, 1984.

WRIGHT UNIVERSITY
DAYTON, OH 45435

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN (Managing Editor)
University of California
Los Angeles, CA 90024

CHARLES R. DEPRIMA
California Institute of Technology
Pasadena, CA 91125

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

H. SAMELSON
Stanford University
Stanford, CA 94305

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Ulrich F. Albrecht , A note on locally A -projective groups	1
Marilyn Breen , A Krasnosel'skiĭ-type theorem for unions of two starshaped sets in the plane	19
Anthony Carbery, Sun-Yung Alice Chang and John Brady Garnett , Weights and $L \log L$	33
Joanne Marie Dombrowski , Tridiagonal matrix representations of cyclic self-adjoint operators. II	47
Heinz W. Engl and Werner Römisch , Approximate solutions of nonlinear random operator equations: convergence in distribution	55
P. Ghez, R. Lima and J. E. Roberts , W^* -categories	79
Barry E. Johnson , Continuity of homomorphisms of Banach G -modules ...	111
Elyahu Katz and Sidney Allen Morris , Free products of topological groups with amalgamation. II	123
Neal I. Koblitz , p -adic integral transforms on compact subgroups of C_p	131
Albert Edward Livingston , A coefficient inequality for functions of positive real part with an application to multivalent functions	139
Scott Carroll Metcalf , Finding a boundary for a Hilbert cube manifold bundle	153
Jack Ray Porter and R. Grant Woods , When all semiregular H -closed extensions are compact	179
Francisco José Ruiz and José Luis Torrea , A unified approach to Carleson measures and A_p weights. II	189
Timothy DuWayne Sauer , The number of equations defining points in general position	199
John Brendan Sullivan , Universal observability and codimension one subgroups of Borel subgroups	215
Akihito Uchiyama , Extension of the Hardy-Littlewood-Fefferman-Stein inequality	229