

Pacific Journal of Mathematics

**VECTOR BUNDLES OVER $(8k + 3)$ -DIMENSIONAL
MANIFOLDS**

TZE-BENG NG

VECTOR BUNDLES OVER $(8k + 3)$ -DIMENSIONAL MANIFOLDS

TZE BENG NG

Let M be a closed, connected, smooth and 3-connected mod 2 (i.e. $H_i(M; \mathbf{Z}_2) = 0, 0 < i \leq 3$) manifold of dimension $3 + 8k$ with $k > 1$. We obtain some necessary and sufficient condition for the span of a $(3 + 8k)$ -plane bundle η over M to be greater than or equal to 7 or 8. We obtain, for M 4-connected mod 2 and satisfying $\text{Sq}^2 \text{Sq}^1 H^{n-8}(M) = \text{Sq}^2 H^{n-7}(M)$, where $n = \dim M \equiv 11 \pmod{16}$ with $n > 11$, that $\text{span } M \geq 8$ if and only if $\chi_2(M) = 0$. Some applications to product manifolds and immersion are given.

1. Introduction. Let M be a closed, connected and smooth manifold whose dimension n is congruent to 3 mod 8 with $n \geq 11$. Let η be an n -plane bundle over M . Recall $\text{span}(\eta)$ is defined to be the maximal number of linearly independent cross sections of η . When η is the tangent bundle of M we simply write $\text{span}(M)$ for $\text{span}(\eta)$. Recall that the Kervaire mod 2 semi-characteristic of M $\chi_2(M)$, is defined by

$$\chi_2(M) = \sum_{2i < n} \dim_{\mathbf{Z}_2} H^i(M; \mathbf{Z}_2) \pmod{2}.$$

Suppose M is a 1-connected mod 2 spin manifold. Then according to Thomas [14] and Randall [11], $\text{span}(M) \geq 4$ if and only if $\chi_2(M) = 0$ and $w_{n-3}(M) = 0$, where $w_j(M)$ is the j th-mod 2 Stiefel-Whitney class of the tangent bundle of M . In this paper we shall derive some necessary and sufficient condition for $\text{span}(M) \geq 7$ or 8 when M is 3-connected mod 2.

For the rest of the paper we shall assume that M is 3-connected mod 2. Then from [14] and the methods of Mahowald [4] we have:

THEOREM 1.1 (Thomas-Mahowald). *Span* $(M) \geq 5$ if and only if $\delta w_{n-5}(M) = 0$ and $\chi_2(M) = 0$. Here δ is the Bockstein coboundary homomorphism associated with the exact sequence $0 \rightarrow \mathbf{Z} \rightarrow \mathbf{Z} \rightarrow \mathbf{Z}_2 \rightarrow 0$.

We shall consider the modified Postnikov tower for the fibration $B\text{spin}_{n-k} \rightarrow B\text{spin}_n$ for $n \geq 19$ and $k = 7$ or 8 where $B\text{spin}_j$ is the classifying space of spin j -plane bundles. Then for an n -plane bundle η over M , η is classified by a map g from M into $B\text{spin}_n$. Then η admits k -linearly independent cross-sections if and only if g lifts to $B\text{spin}_{n-k}$.

For the remainder of the paper we shall assume that $n \geq 19$. All cohomology will be ordinary mod 2 cohomology unless otherwise specified. Let \mathfrak{A} denote the mod 2 Steenrod algebra.

Let ϕ_3 be the stable secondary cohomology operation associated with the relation in \mathfrak{A} also denoted by the same symbol:

$$\phi_3: \text{Sq}^2\text{Sq}^2 + \text{Sq}^1(\text{Sq}^2\text{Sq}^1) = 0.$$

Then ϕ_3 is spin-trivial in the sense of Thomas [13].

Then η admits 6 linearly independent cross sections if and only if $w_{n-5}(\eta) = 0$, $0 \in k_1^2(\eta)$ and $0 \in k_4^2(\eta)$ where k_i^2 , $i = 1, 4$ are the k -invariants for the n -modified Postnikov tower for $\pi: B\text{spin}_{n-6} \rightarrow B\text{spin}_n$ defined by the following relations

$$\begin{aligned} k_1^2: \text{Sq}^2 w_{n-5} &= 0; \\ k_4^2: (\text{Sq}^4 + w_4)w_{n-3} &= 0. \end{aligned}$$

Let $B\text{spin}_{n-6} \xrightarrow{q} E_1 \xrightarrow{p} B\text{spin}_n$ be the first stage n -MPT for the fibration π . Let $j: B\text{spin}_{n-7} \rightarrow B\text{spin}_{n-6}$ be the obvious inclusion. Then since ϕ_3 is spin-trivial $0 \in \phi_3(w_{n-7}) \subset H^{n-4}(B\text{spin}_{n-7})$. Since j^* is an epimorphism in dimensions $\leq n$ and a monomorphism in dimension $n - 4$, we see that $0 \in \phi_3(w_{n-7}) \subset H^{n-4}(B\text{spin}_{n-6})$. Since $\text{Sq}^2 w_{n-7} = w_{n-5}$ and $\text{Sq}^2\text{Sq}^1 w_{n-7} = 0$ we conclude that the class w_{n-7} in $H^{n-7}(B\text{spin}_n)$ is a generating class (see [12]) for k_1^2 in $H^{n-4}(E_1)$ relative to the operation ϕ_3 . Suppose now $\text{Sq}^1 H^{n-5}(M) \subset \text{Sq}^2 H^{n-6}(M)$. We have by the generating class theorem that $0 \in k_1^2(\eta)$ if and only if $0 \in \phi_3(w_{n-7}(\eta))$.

Let Γ be the unstable cohomology operation of Hughes-Thomas type associated with the following relation in \mathfrak{A} on cohomology classes of dimension $\leq n$

$$\Gamma: \text{Sq}^4\text{Sq}^{n-3} + \text{Sq}^2(\text{Sq}^{n-3}\text{Sq}^2) + \text{Sq}^1(\text{Sq}^{n-3}\text{Sq}^3 + \text{Sq}^{n-1}\text{Sq}^1) = 0.$$

Let U be the Thom class of the universal n -plane bundle γ over $B\text{spin}_n$. Let U' be the Thom class of the bundle over $B\text{spin}_{n-6}$ induced by π from γ . Let $T\pi$ be the map between the Thom spaces. Then Γ is defined on $U' = (T\pi)^*U$ and is trivial on U' . Then $w_{n-3} \in H^{n-3}(B\text{spin}_n)$ is an admissible class (see Ng [8]) for k_4^2 with respect to the operation Γ . Then by the admissible class theorem since $(T\pi)^*$ is an epimorphism in dimension $2n$, $0 \in k_4^2(\eta)$ if and only if $0 \in \Gamma(U(\eta))$, where $U(\eta)$ is the Thom class of the bundle η . Hence we have proved

THEOREM 1.2. *Suppose $\text{Sq}^1 H^{n-5}(M) \subset \text{Sq}^2 H^{n-6}(M)$.*

(a) *If $w_4(\eta) \neq w_4(M)$ then $\text{span}(\eta) \geq 6$ if and only if $w_{n-5}(\eta) = 0$ and $0 \in \phi_3(w_{n-7}(\eta))$.*

(b) If $w_4(\eta) = w_4(M)$ then $\text{span}(\eta) \geq 6$ if and only if $w_{n-5}(\eta) = 0$, $0 \in \phi_3(w_{n-7}(\eta))$ and $\Gamma(U(\eta)) = 0$ modulo zero indeterminacy.

If η is the tangent bundle of M and $w_4(M) = 0$ then it can be easily deduced that for $n \equiv 11 \pmod{16}$, $w_{n-5}(M) = w_{n-7}(M) = 0$. According to [14], $\Gamma(U(\eta)) = \chi_2(M) \cdot U(\eta) \cdot \mu$ where $\mu \in H^n(M)$ is a generator. Hence we have

THEOREM 1.3. *Suppose $\text{Sq}^1 H^{n-5}(M) \subset \text{Sq}^2 H^{n-6}(M)$.*

(a) *If $n \equiv 3 \pmod{16} > 3$, then $\text{span}(M) \geq 6$ if and only if $w_{n-5}(M) = 0$, $\chi_2(M) = 0$ and $0 \in \phi_3(w_{n-7}(M))$.*

(b) *If $n \equiv 11 \pmod{16} > 11$ and $w_4(M) = 0$, then $\text{span}(M) \geq 6$ if and only $\chi_2(M) = 0$.*

2. Statement of results for the case $k = 7$. The n -MPT for $B\text{spin}_{n-7} \rightarrow B\text{spin}_n$ is given in Ng [8]. We list the result (in the relevant dimensions) as follows:

TABLE 1. The n -MPT for $B\text{spin}_{n-7} \rightarrow B\text{spin}_n$.

k -invariant	Dimension	Defining Relation
k_1^1	$n - 6$	$k_1^1 = \delta w_{n-7}$
k_2^1	$n - 5$	$k_2^1 = w_{n-5}$
k_3^1	$n - 3$	$k_3^1 = w_{n-3}$
k_1^2	$n - 5$	$\text{Sq}^2 k_1^1 = 0$
k_2^2	$n - 4$	$\text{Sq}^2 k_2^1 = 0$
k_6^2	n	$(\text{Sq}^4 + w_4)k_3^1 = 0$
k_1^3	$n - 4$	$\text{Sq}^2 k_1^2 = 0$.

Consider the following stable secondary cohomology operations associated with the following relations in \mathfrak{A} also denoted by the same symbols

$$\phi_4: \text{Sq}^2(\text{Sq}^2 \text{Sq}^1) = 0$$

$$\phi_5: (\text{Sq}^2 \text{Sq}^1)(\text{Sq}^2 \text{Sq}^1) + \text{Sq}^1(\text{Sq}^2 \text{Sq}^3) = 0$$

such that $\text{Sq}^2 \phi_4 + \text{Sq}^1 \phi_5 = 0$. Let ψ_5 be a stable tertiary cohomology operation associated with the above relation. We assume that (ϕ_4, ϕ_5) and ψ_5 are chosen to be spin trivial in the sense of Theorem 3.7 of Thomas [13].

Let $\phi_{0,0}$ and ϕ_3 be the Adams basic stable secondary cohomology operations associated with the relations:

$$\begin{aligned} \phi_{0,0}: \text{Sq}^1 \text{Sq}^1 &= 0 \quad \text{and} \\ \phi_3: \text{Sq}^2 \text{Sq}^2 + \text{Sq}^1(\text{Sq}^2 \text{Sq}^1) &= 0 \end{aligned}$$

respectively.

We shall prove the following theorem.

THEOREM 2.1. *Suppose*

$$\text{Sq}^1 H^{n-5}(M) \subset \text{Sq}^2 H^{n-6}(M), \quad \text{Sq}^2 H^{n-7}(M; \mathbf{Z}) = \text{Sq}^2 H^{n-7}(M)$$

and $\text{Indet}^{n-4}(\psi_5, M) = \text{Indet}^{n-4}(k_1^3, M)$. Then $\text{span}(M) \geq 7$ if and only if $\delta w_{n-7}(M) = 0$, $w_{n-5}(M) = 0$, $0 \in \phi_4(w_{n-9}(M))$, $0 \in \phi_3(w_{n-7}(M))$, $\chi_2(M) = 0$ and $0 \in \psi_5(w_{n-9}(M))$.

Suppose $\text{Sq}^1 H^4(M) = 0$, $\phi_{0,0} H^4(M) = 0$ and $H_6(M; \mathbf{Z})$ has no 2-torsion. Then it is easily deduced that $\text{Indet}^{n-4}(\psi_5, M) = \text{Indet}^{n-4}(k_1^3, M)$ and $\text{Sq}^3 H^{n-7}(M) = 0$. It follows from Theorem 2.1 the following theorem:

THEOREM 2.2. *Suppose $\text{Sq}^1 H^4(M) = 0$, $\phi_{0,0} H^4(M) = 0$ and $H_6(M; \mathbf{Z})$ has no 2-torsion. Then $\text{span}(M) \geq 7$ if and only if $\delta w_{n-7}(M) = 0$, $w_{n-5}(M) = 0$, $0 \in \phi_4(w_{n-9}(M))$, $0 \in \phi_3(w_{n-7}(M))$, $\chi_2(M) = 0$ and $0 \in \psi_5(w_{n-9}(M))$.*

If $n = 11 + 16k$ with $k \geq 1$, then according to Wu and 6.6 of [8], if $w_4(M) = 0$, $w_{n-9}(M) = w_{n-7}(M) = w_{n-5}(M) = 0$. Hence we have

COROLLARY 2.3. *Suppose either M satisfies the hypothesis of 2.2 or M is 4-connected mod 2 with $\text{Sq}^2 H^{n-7}(M; \mathbf{Z}) = \text{Sq}^2 H^{n-7}(M)$. If $n \equiv 11 \pmod{16} > 11$ and $w_4(M) = 0$, then $\text{span}(M) \geq 7$ if and only if $\chi_2(M) = 0$.*

3. Statement of results for the case $k = 8$. Let $B\hat{S}O_j\langle 8 \rangle$ be the classifying space for orientable j -plane bundles ξ satisfying $w_2(\xi) = w_4(\xi) = 0$. We shall consider the modified Postnikov tower for the fibration $B\hat{S}O_{n-8}\langle 8 \rangle \rightarrow B\hat{S}O_n\langle 8 \rangle$ through dimension n . The computation is done in [8]. We list here the k -invariants for the relevant dimensions only for reference. In particular the k -invariants k_1^2, k_5^2 and k_1^3 for the n -MPT for $B\text{spin}_{n-8} \rightarrow B\text{spin}_n$ are the images of k_1^2, k_6^2 and k_1^3 , respectively, of the n -MPT for $B\text{spin}_{n-7} \rightarrow B\text{spin}_n$ under the map between the towers.

TABLE 2. The n -MPT for $B\text{spin}_{n-8} \rightarrow B\text{spin}_n$.

k -invariant	Dimension	Defining relation
k_1^1	$n - 7$	$k_1^1 = w_{n-7}$
k_2^1	$n - 3$	$k_2^1 = w_{n-3}$
k_1^2	$n - 5$	$\text{Sq}^2 \text{Sq}^1 k_1^1 = 0$
k_2^2	$n - 4$	$(\chi \text{Sq}^4 + w_4)k_1^1 = 0$
k_5^2	n	$(\text{Sq}^4 + w_4)k_1^1 = 0$
k_1^3	$n - 4$	$\text{Sq}^2 k_1^2 = 0$

Consider the following stable secondary cohomology operations ϕ_6, ϕ_7 and ζ_7 associated with the following relations in \mathfrak{A} also denoted by the same symbols as for the operations:

$$\begin{aligned} \phi_6: (\text{Sq}^2 \text{Sq}^1) \chi \text{Sq}^4 + \text{Sq}^2(\text{Sq}^4 \text{Sq}^1) &= 0 \\ \phi_7: \chi \text{Sq}^4 \chi \text{Sq}^4 + \text{Sq}^2(\text{Sq}^4 \text{Sq}^2) + \text{Sq}^1(\text{Sq}^4 \text{Sq}^2 \text{Sq}^1) &= 0 \\ \zeta_7: (\text{Sq}^2 \text{Sq}^1)(\text{Sq}^4 \text{Sq}^1) &= 0. \end{aligned}$$

These operations satisfy the relation

$$(3.1) \quad \text{Sq}^2 \phi_6 + \text{Sq}^1 \zeta_7 = 0$$

Let ψ_7 be a stable tertiary cohomology operation associated with relation (3.1). Note that the operations ϕ_6, ϕ_7 and ψ_7 are chosen such that $\phi_4 \circ \text{Sq}^2 \subset \phi_6, \phi_5 \circ \text{Sq}^2 \subset \zeta_7$ and $\psi_5 \circ \text{Sq}^2 \subset \psi_7$.

Let U_j be the Thom class of the universal j -plane bundle over $B\hat{\text{S}}O_j\langle 8 \rangle$ for $j \geq 4$. Then it is easily seen that

$$U_j(\text{Sq}^2 \nu_4, \text{Sq}^3 \nu_4, 0) \in (\phi_6, \phi_7, \zeta_7)(U_j)$$

where $\nu_4 \in H^4(B\hat{\text{S}}O_j\langle 8 \rangle) \approx \mathbf{Z}_2$ is a generator. Hence

$$(3.2) \quad (0, 0, 0) \in (\phi_6, \phi_7, \zeta_7)(U_j).$$

Since $H^7(B\hat{\text{S}}O_j\langle 8 \rangle) \approx \mathbf{Z}_2$ is generated by $\text{Sq}^3 \nu_4$, trivially

$$(3.3) \quad 0 \in \psi_7(U_j).$$

Let K_4 be the Eilenberg-MacLane space of type $(\mathbf{Z}_2, 4)$. Consider the Massey-Peterson algebra $\mathfrak{A}(K_4)$. Let ι_4 be the fundamental class of K_4 . Let $\gamma = (1 \otimes \chi \text{Sq}^4 + \iota_4 \otimes 1), \alpha = (\text{Sq}^2 \text{Sq}^1 \iota_4 \otimes 1 + \iota_4 \otimes \text{Sq}^2 \text{Sq}^1 + \text{Sq}^1 \iota_4 \otimes \text{Sq}^2 + 1 \otimes \text{Sq}^4 \text{Sq}^2 \text{Sq}^1), \theta = (1 \otimes \text{Sq}^4 + \iota_4 \otimes 1)$. Then we have the following relation in $\mathfrak{A}(K_4)$:

$$\tilde{\phi}_7: \gamma \cdot \gamma + \text{Sq}^2(\theta \cdot \text{Sq}^2) + \text{Sq}^1 \alpha = 0.$$

Let $\tilde{\phi}_7$ be the twisted cohomology operation associated with this relation also denoted by the same symbols as the operation. For $j \geq 4$ let U_j be the Thom class of the universal j -plane bundle over $B\text{spin}_j$. Then $\tilde{\phi}_7$ is defined on (U_j, w_4) . Since $H^7(B\text{spin}_j)$ is generated by $w_7 = \text{Sq}^1 w_6$, trivially

$$(3.4) \quad 0 \in \tilde{\phi}_7(U_j, w_4).$$

We shall prove the following theorems.

THEOREM 3.1. *Suppose either $\text{Sq}^2 \text{Sq}^1 H^{n-8}(M) = \text{Sq}^2 H^{n-7}(M)$ and $\text{Sq}^4 H^{n-8}(M) = \text{Sq}^2 H^4(M) = \text{Sq}^1 H^4(M) = 0$ or $\text{Sq}^2 H^{n-7}(M) = 0$ and $\text{Sq}^1 H^{n-5}(M) + \text{Sq}^2 H^{n-6}(M) \subset \chi \text{Sq}^4 H^{n-8}(M)$. If $w_4(M) = 0$ and $\text{Indet}^{n-4}(\psi_7, M) = \text{Indet}^{n-4}(k_1^3, M)$, then $\text{span}(M) \geq 8$ if and only if $w_{n-7}(M) = 0$, $0 \in \phi_6(w_{n-11}(M))$, $0 \in \phi_7(w_{n-11}(M))$, $\chi_2(M) = 0$ and $0 \in \psi_7(w_{n-11}(M))$.*

THEOREM 3.2. *Suppose either $\text{Sq}^2 \text{Sq}^1 H^{n-8}(M) = \text{Sq}^2 H^{n-7}(M)$, $(\text{Sq}^4 + w_4(M) \cdot) H^{n-8}(M) = 0$ and $\text{Sq}^2 H^4(M) = \text{Sq}^1 H^4(M) = 0$ or $\text{Sq}^2 H^{n-7}(M) = 0$ and*

$$\text{Sq}^2 H^{n-6}(M) + \text{Sq}^1 H^{n-5}(M) \subset (\chi \text{Sq}^4 + w_4(M) \cdot) H^{n-8}(M).$$

If $\text{Indet}^{n-4}(\psi_5, M) = \text{Indet}^{n-4}(k_1^3, M)$ then $\text{span}(M) \geq 8$ if and only if $w_{n-7}(M) = 0$, $0 \in \phi_4(w_{n-9}(M))$, $0 \in \tilde{\phi}_7(w_{n-11}(M), w_4(M))$, $\chi_2(M) = 0$ and $0 \in \psi_5(w_{n-9}(M))$.

By the choice of operations (ϕ_6, ζ_7) and ψ_7 we can deduce that the indeterminacy $\text{Indet}^{n-4}(\psi_7, M)$ is given by the image of the stable operation ϕ_3 on classes $\text{Sq}^1 x + y$, where $x \in H^{n-8}(M)$ and $y \in H^{n-7}(M)$ satisfy $\text{Sq}^2 \text{Sq}^1 x + \text{Sq}^2 y = 0$ and $\text{Sq}^2 \text{Sq}^1 y = 0$. Note that the map from the tower of operations for $(\psi_7, (\phi_6, \zeta_7))$ on N -dimensional classes to the tower of operations for $(\psi_5, (\phi_4, \phi_5))$ on $N + 2$ dimensional classes is induced by the commutative square:

$$\begin{array}{ccc} K_N & \xrightarrow{\text{Sq}^2} & K_{N+2} \\ (\chi \text{Sq}^4, \text{Sq}^4 \text{Sq}^1) \downarrow & & \downarrow (\text{Sq}^2 \text{Sq}^1, \text{Sq}^2 \text{Sq}^3) \\ K_{N+4} \times K_{N+5} & \xrightarrow{(\text{Sq}^1 \iota_{N+4} + \iota_{N+5}, 0)} & K_{N+5} \times K_{N+7} \end{array}$$

where K_j is the Eilenberg-Maclane space of type (\mathbf{Z}_2, j) and ι_j is its fundamental class.

Now the map from the n -MPT for $B\text{spin}_{n-8} \rightarrow B\text{spin}_n$ to the n -MPT for $B\text{spin}_{n-7} \rightarrow B\text{spin}_n$ is induced by the commutative square:

$$\begin{array}{ccc}
 B\text{spin}_n & \xlongequal{\quad} & B\text{spin}_n \\
 (\omega_{n-7}, \omega_{n-3}) \downarrow & & \downarrow (\omega_{n-7}, \omega_{n-5}, \omega_{n-3}) \\
 K_{n-7} \times K_{n-3} & \xrightarrow{(\delta\iota_{n-7}, \text{Sq}^2 \iota_{n-7}, \iota_{n-3})} & K_{n-6}^* \times K_{n-5} \times K_{n-3}
 \end{array}$$

where K_{n-6}^* is the Eilenberg-MacLane space of type $(\mathbf{Z}, n - 6)$. In view of the fact that (k_1^2, k_2^2) is independent of w_{n-3} (see [15, §4]) and k_1^3 is independent of k_2^2 we can deduce that $\text{Indet}^{n-4}(k_1^3, M)$ is the image of the stable operations ϕ_3^* on classes δx , where $x \in H^{n-8}(M)$ satisfies $\text{Sq}^2 \text{Sq}^1 x = 0$ and $(\chi \text{Sq}^4 + w_4(M) \cdot)x = 0$ and ϕ_3^* is a stable cohomology operation associated with the relation $\text{Sq}^2 \text{Sq}^2 = 0$ on integral classes.

Suppose M satisfies the following conditions:

- (A) $H_6(M; \mathbf{Z})$ has no 2-torsion and
- (B) $H_7(M; \mathbf{Z})$ has no free parts and its 2-torsion elements are all of order 2.

Then $\text{Sq}^1 H^{n-8}(M) = H^{n-7}(M)$. If further

$$\text{Sq}^1 H^{n-5}(M) \subset \text{Sq}^2 H^{n-6}(M) \text{ and } (\chi \text{Sq}^4 + w_4(M) \cdot) H^{n-8}(M) = 0,$$

then $\text{Indet}^{n-4}(\psi_7, M) = \text{Indet}^{n-4}(k_1^3, M)$. Hence we have by Theorem 3.1 the following immediate corollary:

COROLLARY 3.3. *Suppose M satisfies both condition (A) and (B). If $w_4(M) = 0$, $\text{Sq}^1 H^4(M) \simeq \text{Sq}^2 H^4(M) \simeq 0$ and $\chi \text{Sq}^4 H^{n-8}(M) = 0$ then*

- (i) $\phi_4(\omega_{n-9}(M)) = \phi_6(\omega_{n-11}(M))$;
- (ii) if $\phi_{0,0} H^4(M) = 0$ then $\psi_7(\omega_{n-11}(M)) = \psi_5(\omega_{n-9}(M))$;
- (iii) $\text{span}(M) \geq 8$ if and only if $w_{n-7}(M) = 0$, $0 \in \phi_6(\omega_{n-11}(M))$, $0 \in \phi_7(\omega_{n-11}(M))$, $\chi_2(M) = 0$ and $0 \in \psi_7(\omega_{n-11}(M))$.

Similarly from Theorem 3.2 we have

COROLLARY 3.4. *Suppose M satisfies conditions A and B. Assume $\text{Sq}^1 H^4(M) = 0$, $\phi_{0,0} H^4(M) = 0$, $\text{Sq}^2 H^4(M) = 0$ and*

$$(\chi \text{Sq}^4 + w_4(M) \cdot) H^{n-8}(M) = 0.$$

Then $\text{span}(M) \geq 8$ if and only if $w_{n-7}(M) = 0$, $0 \in \phi_4(\omega_{n-9}(M))$, $0 \in \tilde{\phi}_7(\omega_{n-11}(M), w_4(M))$, $\chi_2(M) = 0$ and $0 \in \psi_5(\omega_{n-9}(M))$.

Suppose $n = 11 + 16k$ with $k \geq 1$, $w_4(M) = w_8(M) = 0$. Then by Wu $w_{n-5}(M) = w_{n-7}(M) = w_{n-9}(M) = 0$ and $w_{n-11}(M) = v_{8k}^2$, where $v_j \in H^j(M)$ is the j th Wu-class of M . Since $w_8(M) = 0$, utilizing the Adém relations for $Sq^8 Sq^{8k-4}$, $Sq^4 Sq^{8k-2}$ and $Sq^2 Sq^{8k-1}$ we can show that $Sq^4 v_{8k} = 0$. Therefore ϕ_7 is defined on v_{8k} . By a Cartan formula for ϕ_7 , (see [7])

$$\begin{aligned} \phi_7(w_{n-11}(M)) &= \phi_7(v_{8k}^2) = \phi_7(v_{8k}) \cdot v_{8k} + v_{8k} \cdot \phi_7(v_{8k}) \\ &= \{0\} \quad \text{modulo indeterminacy of } \phi_7. \end{aligned}$$

If $n = 19 + 32k$ with $k \geq 0$ and $w_4(M) = w_8(M) = 0$, then $w_{n-5}(M) = w_{n-7}(M) = w_{n-9}(M) = w_{n-11}(M) = 0$. Thus we have from Corollary 3.3 the following:

THEOREM 3.5. *Suppose $n \equiv 11 \pmod{16} > 11$ or $n \equiv 19 \pmod{32}$ and $w_4(M) = w_8(M) = 0$. Then*

- (a) *Suppose $Sq^1 H^4(M) \simeq Sq^2 H^4(M) \simeq Sq^4 H^{n-8}(M) \simeq 0$. Then if M satisfies (A) and (B), $\text{span}(M) \geq 8$ if and only if $\chi_2(M) = 0$.*
- (b) *Suppose M is 4-connected mod 2 and $Sq^2 Sq^1 H^{n-8}(M) = Sq^2 H^{n-7}(M)$. Then $\text{span}(M) \geq 8$ if and only if $\chi_2(M) = 0$.*

4. Proof of Theorem 2.1. Let w_{n-9} be the $(n - 9)$ th mod 2 universal Stiefel-Whitney class considered as in $H^{n-9}(B\text{spin}_{n-7})$. Then according to [13] we have

PROPOSITION 4.1 (E. Thomas).

- (a) $(0, 0) \in (\phi_4, \phi_5)(w_{n-9}) \subset H^{n-5}(B\text{spin}_{n-7}) \oplus H^{n-4}(B\text{spin}_{n-7})$
- (b) $0 \in \psi_5(w_{n-9}) \subset H^{n-4}(B\text{spin}_{n-7})$.

Proof. Part (a) is proved in [13, Proposition 4.2] and we shall not present it here. Let $j: B\text{spin}_{n-9} \rightarrow B\text{spin}_{n-7}$ be the inclusion map. Since ψ_5 is spin trivial $0 \in \psi_5(w_{n-9}) \subset H^{n-4}(B\text{spin}_{n-9})$. Now $\text{Indet}^{n-4}(\psi_5, B\text{spin}_{n-9})$ is determined by a stable cohomology operation Θ defined on cohomology vectors

$$(x, y) \in H^{n-7}(B\text{spin}_{n-9}) \times H^{n-5}(B\text{spin}_{n-9})$$

satisfying $Sq^2 x = 0$ and $Sq^2 Sq^1 x + Sq^1 y = 0$. Since j^* is an epimorphism in dimensions $\leq n - 4$, there are classes $x' \in H^{n-7}(B\text{spin}_{n-7})$ and $y' \in H^{n-5}(B\text{spin}_{n-7})$ such that $j^*x' = x$ and $j^*y' = y$. Since j^* is an isomorphism in dimension $n - 5$, $Sq^2 x = 0$ implies that $Sq^2 x' = 0$.

Since $\text{Ker } j^*$ in dimension $n - 4$ is generated by $w_{n-8} \cdot w_4$, $\text{Sq}^2 \text{Sq}^1 x' + \text{Sq}^1 y' = \alpha w_{n-8} \cdot w_4$ for some $\alpha \in \mathbf{Z}_2$. Hence

$$\text{Sq}^2 \text{Sq}^1 x' + \text{Sq}^1(y' + \alpha w_{n-9} \cdot w_4) = 0.$$

Thus Θ is defined on $(x', y' + \alpha w_{n-9} \cdot w_4)$. Now Θ is defined on $(0, w_{n-9} \cdot w_4) \in H^{n-7}(B\text{spin}_{n-9}) \times H^{n-5}(B\text{spin}_{n-9})$. $\Theta(0, w_{n-9} \cdot w_4) = \phi_{0,0}(w_{n-9} \cdot w_4) \subset H^{n-4}(B\text{spin}_{n-9})$ modulo indeterminacy of Θ . But

$$\phi_{0,0}(w_{n-9} \cdot w_4) = \phi_{0,0}(w_{n-9}) \cdot w_4 + w_{n-9} \cdot \phi_{0,0}(w_4) = 0,$$

since $\phi_{0,0}$ is spin-trivial and $H^5(B\text{spin}_{n-9}) \approx 0$. Thus $\Theta(0, w_{n-9} \cdot w_4) = \{0\} \subset H^{n-4}(B\text{spin}_{n-9})$. Thus

$$\begin{aligned} \Theta(x, y) &= \Theta(x, y) + \Theta(0, \alpha w_{n-9} \cdot w_4) = \Theta(x, y + \alpha w_{n-9} \cdot w_4) \\ &= \Theta(j^*x', j^*(y' + \alpha w_{n-9} \cdot w_4)) = j^*\Theta(x', y' + \alpha w_{n-9} \cdot w_4) \end{aligned}$$

since j^* is an epimorphism in dimensions $\leq n - 4$. Thus

$$\text{Indet}^{n-4}(\psi_5, B\text{spin}_{n-9}) = j^*\text{Indet}^{n-4}(\psi_5, B\text{spin}_{n-7}).$$

This clarifies an argument in [13, Proposition 4.2]. Since $w_{n-8} \cdot w_4 = \text{Sq}^1(w_{n-9} \cdot w_4)$ we conclude that $0 \in \psi_5(w_{n-9}) \subset H^{n-4}(B\text{spin}_{n-7})$.

Similarly, since ϕ_3 is spin-trivial, we have

PROPOSITION 4.2. $0 \in \phi_3(w_{n-7}) \subset H^{n-4}(B\text{spin}_{n-7})$.

Let the n -MPT for $\pi: B\text{spin}_{n-k} \rightarrow B\text{spin}_n$ for $k = 7$ or 8 be indicated by the following diagram:

$$\begin{array}{ccccc} & & & & B\text{spin}_{n-k} \\ & & & & \downarrow \pi \\ & & q_2 \swarrow & \swarrow q_1 & \\ E_2 & \xrightarrow{p_2} & E_1 & \xrightarrow{p_1} & B\text{spin}_n \end{array}$$

when $k = 8$, it is understood this tower is induced over $B\hat{\text{S}}\text{O}_n\langle 8 \rangle$. We shall still use the same symbols for the second and third stage of the tower over $B\hat{\text{S}}\text{O}_n\langle 8 \rangle$ when no confusion need arise.

Recall the definition of a generating class [12]. Then we have

PROPOSITION 4.3 (*E. Thomas*).

(a) The class w_{n-9} in $H^{n-9}(B\text{spin}_n)$ is a generating class for the pair $(k_1^2, 0)$ in $H^{n-5}(E_1) \oplus H^{n-4}(E_1)$ relative to the pair (ϕ_4, ϕ_5) .

(b) The class $p_1^*(w_{n-9})$ is a generating class for k_1^3 , relative to the operation ψ_5 .

(c) The class w_{n-7} in $H^{n-7}(B\text{spin}_n)$ is a generating class for k_2^2 in $H^{n-4}(E_1)$ relative to the operation ϕ_3 .

Now by inspection of the k -invariants for the n -MPT for π and the connectivity condition on M , together with Proposition 4.1, 4.2, 4.3, the generating class theorem of [12] and the admissible class theorem [8] we have:

THEOREM 4.4. *Let η be an n -plane bundle over M . Suppose*

$$\text{Sq}^2 H^{n-7}(M; \mathbf{Z}) = \text{Sq}^2 H^{n-7}(M), \quad \text{Sq}^1 H^{n-5}(M) \subset \text{Sq}^2 H^{n-6}(M)$$

and $\text{Indet}^{n-4}(\psi_5, M) = \text{Indet}^{n-4}(k_1^3, M)$.

(a) *Suppose $w_4(\eta) \neq w_4(M)$. Then η admits 7-linearly independent cross sections if and only if $\delta w_{n-7}(\eta) = 0$, $w_{n-5}(\eta) = 0$, $0 \in \phi_4(w_{n-9}(\eta))$, $0 \in \phi_3(w_{n-7}(\eta))$ and $0 \in \psi_5(w_{n-9}(\eta))$.*

(b) *Suppose $w_4(\eta) = w_4(M)$. Then η admits 7-linearly independent cross sections if and only if $\delta w_{n-7}(\eta) = 0$, $w_{n-5}(\eta) = 0$, $0 \in \phi_4(w_{n-9}(\eta))$, $0 \in \phi_3(w_{n-7}(\eta))$, $0 \in \psi_5(w_{n-9}(\eta))$ and $\Gamma(U(\eta)) = 0$.*

We now specialise to the case when η is the tangent bundle of M . Let $g: M \times M \rightarrow T(\tau)$ be the map that collapses the complement of a tubular neighbourhood of the diagonal in $M \times M$ to a point. Let $U = g^*(U(\tau))$. Then we have the following decomposition of Milnor and Wu:

$$U \text{ mod } 2 = \sum_{2i < n} \sum_k \alpha_i^k \otimes \beta_{n-i}^k + \sum_{2i < n} \sum_k \beta_{n-i}^k \otimes \alpha_i^k$$

where $\alpha_i^k \in H^i(M)$, $\beta_{n-i}^k \in H^{n-i}(M)$ and $\alpha_i^k \cup \beta_{n-i}^j = \delta_{kj} \mu$ where $\mu \in H^n(M)$ is a generator and δ_{kj} is the Kronecker function.

Then we have

LEMMA 4.5. *Let $A = \sum_{2i < n} \sum_k \alpha_i^k \otimes \beta_{n-i}^k$.*

(i) $\underline{U} \text{ mod } 2 = A + t^*A$ where $t^*: H^*(M \times M) \rightarrow H^*(M \times M)$ is the homomorphism induced by the map that interchanges the factors.

(ii) $A \cup t^*A = \chi_2(M)\mu \otimes \mu$.

(iii) $\text{Sq}^{n-3} A = \text{Sq}^{n-3} \text{Sq}^2 A = (\text{Sq}^{n-3} \text{Sq}^3 + \text{Sq}^{n-1} \text{Sq}^1) A = 0$.

Proof. (i) and (ii) are due to Thomas [15] (iii) is a consequence of the connectivity condition on M .

Then according to Mahowald [4] we have

THEOREM 4.6 (Mahowald-Thomas). Γ is defined on A and so on t^*A . In particular modulo zero indeterminacy,

$$\Gamma(U(\tau)) = \chi_2(M)U(\tau)\mu.$$

4.7. *Proof of Theorem 2.1.* This now follows immediately from Theorem 4.4 (b) and Theorem 4.6.

5. Proof of Theorem 3.1 and Theorem 3.2. Consider the fibration $\pi: B\hat{S}O_{n-8}\langle 8 \rangle \rightarrow B\hat{S}O_n\langle 8 \rangle$. Let U_j be the Thom class of the universal j -plane bundle over $B\hat{S}O_j\langle 8 \rangle$ for $j \geq 4$. Then from §3 we know that $(0, 0, 0) \in (\phi_6, \phi_7, \zeta_7)U_j$. Let $l: B\hat{S}O_{n-11}\langle 8 \rangle \rightarrow B\hat{S}O_{n-8}\langle 8 \rangle$ be the obvious inclusion. Then w_{n-11} in $H^*(B\hat{S}O_{n-11}\langle 8 \rangle)$ is the reduction mod 2 of the Euler class of the universal $(n - 11)$ -plane bundle over $B\hat{S}O_{n-11}\langle 8 \rangle$. Hence

$$(5.1) \quad (0, 0, 0) \in (\phi_6, \phi_7, \zeta_7)(w_{n-11}) \subset H^{n-5}(B\hat{S}O_{n-11}\langle 8 \rangle) \\ \oplus H^{n-4}(B\hat{S}O_{n-11}\langle 8 \rangle) \oplus H^{n-4}(B\hat{S}O_{n-11}\langle 8 \rangle).$$

Now it can be easily checked that $l^*: H^*(B\hat{S}O_{n-8}\langle 8 \rangle) \rightarrow H^*(B\hat{S}O_{n-11}\langle 8 \rangle)$ is an epimorphism in dimension $\leq n - 4$ for $n > 43$, while for $n = 19$ and 27 l^* is monomorphic in dimension $\leq n - 4$. This is readily derived by considering the Leray-Serre spectral sequence for $B\hat{S}O_j\langle 8 \rangle$. Also for $n > 27$, $\ker l^*$ in dimension $n - 4$ is generated by $\{w_{n-10}\text{Sq}^2\nu_4, w_{n-8} \cdot \nu_4\}$, where $\nu_4 \in H^4(B\hat{S}O_{n-8}\langle 8 \rangle) \approx \mathbf{Z}_2$ is a generator, while in dimension $n - 5$ $\ker l^*$ is generated by $\{w_{n-9} \cdot \nu_4\}$.

Now for $n \leq 43$, $\text{Indet}^{n-5, n-4, n-4}((\phi_6, \phi_7, \zeta_7), B\hat{S}O_{n-11}\langle 8 \rangle)$ on $\text{Cok } l^*$ is contained in $\text{Coker } l^*$. Therefore in view of (5.1), $(0, 0, 0) \in l^*(\phi_6, \phi_7, \zeta_7)(w_{n-11})$ modulo $l^* \text{Indet}^{n-5, n-4, n-4}((\phi_6, \phi_7, \zeta_7), B\hat{S}O_{n-8}\langle 8 \rangle)$ for $n \geq 19$. Thus by naturality there exist classes $U_1 \in H^{n-5}(B\hat{S}O_{n-8}\langle 8 \rangle)$, $U_2, U_3 \in H^{n-4}(B\hat{S}O_{n-8}\langle 8 \rangle)$ such that $(U_1, U_2, U_3) \in (\phi_6, \phi_7, \zeta_7)w_{n-11}$ and $l^*U_1 = 0, l^*U_2 = 0$ and $l^*U_3 = 0$.

Since l^* is injective for $n = 19, 27$, in dimension $n - 4, n - 5$, $(0, 0, 0) \in (\phi_6, \phi_7, \zeta_7)(w_{n-11}) \subset H^{n-5}(B\hat{S}O_{n-8}\langle 8 \rangle) \oplus H^{n-4}(B\hat{S}O_{n-8}\langle 8 \rangle) \oplus H^{n-4}(B\hat{S}O_{n-8}\langle 8 \rangle)$ for $n = 19$ and 27 . For $n > 27$, $\text{Sq}^3(w_{n-9}\nu_4) \neq 0$ for $w_{n-9}\nu_4 \in H^{n-5}(B\hat{S}O_{n-8}\langle 8 \rangle)$. But by (3.1), $\text{Sq}^3 U_1 = 0$, since $\text{Sq}^3 \phi_6 = 0$. Thus $U_1 = 0$. Since $\text{Sq}^2(w_{n-10}\nu_4) = w_{n-10}\text{Sq}^2\nu_4$ and $\text{Sq}^1(w_{n-9}\nu_4) = w_{n-8}\nu_4$, we may assume that $U_2 = 0$. Now $U_3 \neq w_{n-10}\text{Sq}^2\nu_4$ since by (3.1) $\text{Sq}^1 U_3 = \text{Sq}^2 U_1 = 0$ but $\text{Sq}^1(w_{n-10}\text{Sq}^2\nu_4) = w_{n-10}\text{Sq}^3\nu_4 \neq 0$. Now $U_3 \neq w_{n-8}\nu_4$ for $\text{Sq}^2 U_3 = \text{Sq}^2 \zeta_7(w_{n-11}) = \alpha \text{Sq}^8 \text{Sq}^1 w_{n-11}$ for some $\alpha \in \mathbf{Z}_2$ and $\text{Sq}^2(w_{n-8}\nu_4) = w_{n-8}\text{Sq}^2\nu_4 \neq w_8 w_{n-10}$. Thus $U_3 = 0$.

We have thus proved.

PROPOSITION 5.2. *Let the operation $(\phi_6, \phi_7, \zeta_7)$ be given as in §3. Then*

$$(0, 0, 0) \in (\phi_6, \phi_7, \zeta_7)(w_{n-11}) \subset H^{n-5}(B\hat{S}O_{n-8}\langle 8 \rangle) \\ \oplus H^{n-4}(B\hat{S}O_{n-8}\langle 8 \rangle) \oplus H^{n-4}(B\hat{S}O_{n-8}\langle 8 \rangle).$$

$\text{Ker } l^*$ is generated by $\{w_{n-10}, w_{n-9}\}$ as a $\mathfrak{A}(H^*(B\hat{S}O_{n-8}\langle 8 \rangle))$ -module in dimensions $\leq n + 1$. Now $\text{Indet}^{n-4}(\psi_7, B\hat{S}O_{n-11}\langle 8 \rangle)$ is determined by a cohomology operation Ω on cohomology vectors $(x, y) \in H^{n-8}(B\hat{S}O_{n-11}\langle 8 \rangle) \times H^{n-7}(B\hat{S}O_{n-11}\langle 8 \rangle)$ satisfying $\text{Sq}^2 \text{Sq}^1 x + \text{Sq}^2 y = 0$ and $\text{Sq}^2 \text{Sq}^1 y = 0$. Since l^* is onto in dimensions $\leq n - 8$ and onto in dimension $n - 7$ if $n > 43$ and since for $n \leq 43$, $\text{Sq}^2(\text{Cok } l^*)_{n-7} \subseteq (\text{Cok } l^*)_{n-5}$ and is non-zero in $(\text{Cok } l^*)_{n-5}$, there exist classes $x' \in H^{n-8}(B\hat{S}O_{n-8}\langle 8 \rangle)$ and $y' \in H^{n-7}(B\hat{S}O_{n-8}\langle 8 \rangle)$ such that $l^*x' = x$ and $l^*y' = y$. Therefore $l^*(\text{Sq}^2 \text{Sq}^1 x' + \text{Sq}^2 y') = 0$ and $l^*(\text{Sq}^2 \text{Sq}^1 y') = 0$. For $n = 19$ and 27 , l^* is a monomorphism in dimension $n - 4$ and $n - 5$ and so $\text{Sq}^2 \text{Sq}^1 x' + \text{Sq}^2 y' = 0$ and $\text{Sq}^2 \text{Sq}^1 y' = 0$.

Now consider the case $n \geq 35$. $\text{Sq}^2 \text{Sq}^1 x' + \text{Sq}^2 y' = \alpha w_{n-9} \cdot \nu_4$ for some $\alpha \in \mathbf{Z}_2$. Let $K: B\hat{S}O_j\langle 8 \rangle \rightarrow B\text{spin}_j$ be the obvious map. Then by considering for $p \leq n - 4$,

$$\begin{aligned} H^p(B\hat{S}O_{n-8}\langle 8 \rangle) &= K^*H^p(B\text{spin}_{n-8}) + \nu_4 \cdot K^*H^{p-4}(B\text{spin}_{n-8}) \\ &\quad + \text{Sq}^2 \nu_4 \cdot K^*H^{p-6}(B\text{spin}_{n-8}) \\ &\quad + \text{Sq}^3 \nu_4 \cdot K^*H^{p-7}(B\text{spin}_{n-8}) \end{aligned}$$

modulo $\{(\text{Sq}^I \nu_4)^j K^*H^*(B\text{spin}_{n-8}); I \text{ is an admissible sequence of excess } < 4, \text{ length } l(I) \geq 2 \text{ and } j \geq 1 \text{ or } l(I) \leq 1 \text{ and } j \geq 2\}$ and the fact that l^* is an isomorphism in dimensions $\leq n - 11$ we can show that

$$w_{n-9} \cdot \nu_4 \notin (\text{Im } \text{Sq}^2 \text{Sq}^1 + \text{Im } \text{Sq}^2) \cap \text{Ker } l^* \cap H^{n-5}(B\hat{S}O_{n-8}\langle 8 \rangle).$$

Thus $\alpha = 0$ and $\text{Sq}^2 \text{Sq}^1 x' + \text{Sq}^2 y' = 0$. Similarly it can be shown that $\text{Sq}^2 \text{Sq}^1 x' + \text{Sq}^2 y' = 0$ and $l^*(\text{Sq}^2 \text{Sq}^1 y') = 0$ implies that $\text{Sq}^2 \text{Sq}^1 y' = 0$. Hence Ω is defined on (x', y') . Thus $\Omega(x, y) = \Omega(l^*x', l^*y') = l^*\Omega(x', y')$ modulo $\text{Indet}^{n-4}(\Omega, B\hat{S}O_{n-11}\langle 8 \rangle)$. In view of this and the fact that ψ_7 is trivial on the Thom class of the universal $(n - 11)$ -plane bundle over $B\hat{S}O_{n-11}\langle 8 \rangle$,

$$l^*\psi_7(w_{n-11}) \subseteq \text{Sq}^2 H^{n-6}(B\hat{S}O_{n-11}\langle 8 \rangle) + \text{Sq}^1 H^{n-7}(B\hat{S}O_{n-11}\langle 8 \rangle)$$

modulo $l^* \text{Indet}^{n-4}(\psi_7, B\hat{S}O_{n-8}\langle 8 \rangle)$. Since for $n \leq 43$ $\text{Sq}^2(\text{Cok } l^*)_{n-6} + \text{Sq}^1(\text{Cok } l^*)_{n-5}$ is non-trivial in $(\text{Cok } l^*)_{n-4}$ and since l^* is an epimorphism in dimensions $\leq n - 4$ for $n > 43$, we can conclude that $l^*\psi_7(w_{n-11}) \subset l^* \text{Indet}^{n-4}(\psi_7, B\hat{S}O_{n-8}\langle 8 \rangle)$. Since $\text{Ker } l^*$ is generated by

$$\begin{aligned} \{w_{n-10} \text{Sq}^2 \nu_4, w_{n-8} \nu_4\} &\subset \text{Sq}^2 H^{n-6}(B\hat{S}O_{n-8}\langle 8 \rangle) \\ &\quad + \text{Sq}^1 H^{n-5}(B\hat{S}O_{n-8}\langle 8 \rangle), \end{aligned}$$

we deduce that $0 \in \psi_7(w_{n-11}) \subset H^{n-4}(B\hat{S}O_{n-8}\langle 8 \rangle)$.

Hence

PROPOSITION 5.3. *Let ψ_7 be the stable tertiary cohomology operation given as in §3. Then $0 \in \psi_7(w_{n-11}) \subset H^{n-4}(B\hat{S}O_{n-8}\langle 8 \rangle)$.*

We can now use the generating class theorem to realize the k -invariants for liftings. Now in $H^*(B\hat{S}O_n\langle 8 \rangle)$, $\chi Sq^4 w_{n-11} = w_{n-7}$, $Sq^4 Sq^1 w_{n-11} = 0$, $Sq^4 Sq^2 w_{n-11} = 0$ and $Sq^4 Sq^2 Sq^1 w_{n-11} = 0$. $\Pi^*: H^*(B\hat{S}O_n\langle 8 \rangle) \rightarrow H^*(B\hat{S}O_{n-8}\langle 8 \rangle)$ is an epimorphism in dimension $\leq n - 4$ for $n \geq 35$. Now for $n = 27$ or 19 , Π^* is an epimorphism in dimensions $n - 8$, $n - 6$, $n - 5$ and $n - 4$ and q_1^* is an epimorphism in dimensions $\leq n$. A computational check shows that

$$\begin{aligned} & \{ (Sq^2 x, Sq^2 Sq^1 x) : x \in H^{n-7}(B\hat{S}O_{n-8}\langle 8 \rangle) \} \\ & \subset \Pi^* \{ (Sq^2 x, Sq^2 Sq^1 x) : x \in H^{n-7}(B\hat{S}O_n\langle 8 \rangle) \}. \end{aligned}$$

Thus by the generating class theorem, Proposition 5.2 and Proposition 5.3 we have

PROPOSITION 5.4.

(a) *The class $w_{n-11} \in H^{n-11}(B\hat{S}O_n\langle 8 \rangle)$ is a generating class for $(k_1^2, k_2^2, 0)$ in $H^{n-5}(E_1) \oplus H^{n-4}(E_1) \oplus H^{n-4}(E_1)$ relative to the operation $(\phi_6, \phi_7, \zeta_7)$.*

(b) *The class $p_1^*(w_{n-11})$ in $H^{n-11}(E_1)$ is a generating class for $k_1^3 \in H^{n-4}(E_2)$ relative to the operation ψ_7 .*

Consider now $\pi: Bspin_{n-8} \rightarrow Bspin_n$ for $n \geq 19$. Then π^* is an epimorphism in dimension $\leq n$. Then by the admissible class theorem we have

$$U(E_1) \cdot \{k_6^2\} \in \Gamma(U(E_1))$$

where E_1 is the first stage of the modified Postnikov tower for π and $U(E_1)$ is the Thom class of the n -plane bundle over E_1 induced from the universal n -plane bundle over $Bspin_n$ by p_1 . Therefore by naturality we have

PROPOSITION 5.5. *$U(E_1) \cdot \{k_6^2\} \in \Gamma(U(E_1)) \subset H^{2n}(T(E_1))$ where $T(E_1)$ is the Thom space of the n -plane bundle over E_1 induced from the universal n -plane bundle over $B\hat{S}O_n\langle 8 \rangle$ by p_1 and $U(E_1)$ the corresponding Thom class.*

We also have the following proposition since $\tilde{\phi}_7$ is in the terminology of [12], of bundle type 2.

PROPOSITION 5.6.

(a) $0 \in \tilde{\phi}_7(w_{n-11}, w_4) \subset H^{n-4}(B\text{spin}_{n-8})$

(b) *The class $w_{n-11} \in H^{n-11}(B\text{spin}_n)$ is a generating class for k_2^2 in $H^{n-4}(E_1)$ relative to the operation $\tilde{\phi}_7$ with twisting given by w_4 .*

Proof. (a) is a consequence of (3.4). Since $(\chi \text{Sq}^4 + w_4)w_{n-11} = w_{n-7} \in H^{n-7}(B\text{spin}_n)$, $(\theta \cdot \text{Sq}^2)(w_{n-11}) = 0$, $\alpha(w_{n-11}) = 0$ and $\pi^*: H^*(B\text{spin}_n) \rightarrow H^*(B\text{spin}_{n-8})$ is an epimorphism in dimension $\leq n$. (b) follows.

Suppose either $\text{Sq}^2 \text{Sq}^1 H^{n-8}(M) = \text{Sq}^2 H^{n-7}(M)$ and $\text{Sq}^4 H^{n-8}(M) = \text{Sq}^2 H^4(M) = \text{Sq}^1 H^4(M) = 0$ or $\text{Sq}^2 H^{n-7}(M) = 0$ and $\text{Sq}^1 H^{n-5}(M) + \text{Sq}^2 H^{n-6}(M) \subset \chi \text{Sq}^4 H^{n-8}(M)$.

Suppose $\text{Indet}^{n-4}(\psi_7, M) = \text{Indet}^{n-4}(k_1^3, M)$. Then we have with the above assumption

THEOREM 5.7. *Let η be an n -plane bundle over M such that $w_4(\eta) = 0$.*

(a) *Suppose $w_4(M) \neq 0$. Then $\text{span}(\eta) \geq 8$ if and only if $w_{n-7}(\eta) = 0$, $0 \in \phi_6(w_{n-11}(\eta))$, $0 \in \phi_7(w_{n-11}(\eta))$ and $0 \in \psi_7(w_{n-11}(\eta))$.*

(b) *Suppose $w_4(M) = 0$. Then $\text{span}(\eta) \geq 8$ if and only if $w_{n-7}(\eta) = 0$, $0 \in \phi_6(w_{n-11}(\eta))$, $0 \in \phi_7(w_{n-11}(\eta))$, $\Gamma(U(\eta)) = 0$ and $0 \in \psi_7(w_{n-11}(\eta))$.*

Proof. Since $w_4(M) \neq 0$. $\text{Sq}^4 H^{n-4}(M) = H^n(M)$. Trivially $0 \in k_6^2(\eta)$. Part (a) then follows from Propositions 5.2, 5.3, 5.4 and the generating class theorem.

Part (b) is similar and it follows from Propositions 5.2, 5.3, 5.4, 5.5 the generating class theorem and the admissible class theorem. We leave the details to the reader.

Similarly we have a variation of Theorem 5.7.

THEOREM 5.8. *Let η be an n -plane bundle over M . Suppose either*

$$\text{Sq}^2 \text{Sq}^1 H^{n-8}(M) = \text{Sq}^2 H^{n-7}(M), \quad (\text{Sq}^4 + w_4(\eta))H^{n-8}(M) = 0$$

and

$$\text{Sq}^2 H^4(M) = \text{Sq}^1 H^4(M) = 0 \quad \text{or} \quad \text{Sq}^2 H^{n-7}(M) = 0$$

and

$$\text{Sq}^2 H^{n-6}(M) + \text{Sq}^1 H^{n-5}(M) \subset (\chi \text{Sq}^4 + w_4(\eta))H^{n-8}.$$

Assume $\text{Indet}^{n-4}(\psi_5, M) = \text{Indet}^{n-4}(k_1^3, M)$. Then

(a) If $w_4(\eta) \neq w_4(M)$, then $\text{span}(\eta) \geq 8$ if and only if $w_{n-7}(\eta) = 0$, $0 \in \phi_4(w_{n-9}(\eta))$, $0 \in \tilde{\phi}_7(w_{n-11}(\eta), w_4(\eta))$ and $0 \in \psi_5(w_{n-9}(\eta))$.

(b) If $w_4(\eta) = w_4(M)$, then $\text{span}(\eta) \geq 8$ if and only if $w_{n-7}(\eta) = 0$, $0 \in \phi_4(w_{n-9}(\eta))$, $0 \in \tilde{\phi}_7(w_{n-11}(\eta), w_4(\eta))$, $\Gamma(U(\eta)) = 0$ and $0 \in \psi_5(w_{n-9}(\eta))$.

The proof is similar to that of 5.7 using Proposition 5.6. We leave the details to the reader.

5.9. Proof of Theorem 3.1 and Theorem 3.2. Theorem 3.1 is now a consequence of Theorem 5.7(b) by taking η to be the tangent bundle of M and the fact that $\Gamma(U(\tau)) = \chi_2(M)U(\tau)\mu$ (cf. Theorem 4.6). Similarly Theorem 3.2 follows from Theorem 5.8(b) and Theorem 4.6.

6. Applications.

6.1. Let $M = S^{7+8k} \times QP^{2l+1}$ for $k \geq 0, l \geq 0$. Then we have

THEOREM. $\text{Span}(M) \geq 8$.

Proof. Plainly $H^{n-5}(M) \simeq H^{n-6}(M) \simeq 0$, $w_4(M) = 0$ and $\chi_2(M) = 0$. For $k \geq 1$, $H^{n-7}(M) \simeq 0$ and so $\text{Indet}^{n-4}(\psi_7, M) = \text{Indet}^{n-4}(k_1^3, M) \simeq 0$. For $k = 0$, $\delta H^{n-8}(M) \simeq 0$ and so $\text{Indet}^{n-4}(k_1^3, M) \simeq 0$. For $k = 0$, $H^{n-7}(M) \simeq H^{n-7}(QP^{2l+1})$ and so by naturality $\text{Indet}^{n-4}(\psi_7, M) \simeq 0$. Now $w_{n-11}(M) = 0$ if $k \geq 1$ and $w_{n-11}(M) = 1 \times U^{2l}$ if $k = 0$ where $U \in H^4(QP^{2l+1}) \simeq \mathbf{Z}_2$ is a generator. Trivially $w_{n-7}(M) = 0$. Thus if $k \geq 1$ $\text{span}(M) \geq 8$. Now for $k = 0$ $w_{n-11}(M) = j^*(U^{2l})$, where $j: M \rightarrow QP^{2l+1}$ is the projection. For dimensional reasons $(\phi_6, \phi_7, \zeta_7)(U^{2l}) = (0, 0, 0)$ and $\psi_7(U^{2l}) = 0$. Therefore by naturality $0 \in \psi_7(w_{n-11}(M))$. It follows from Theorem 3.1 that $\text{span}(M) \geq 8$. This completes the proof.

Similarly we have

THEOREM 6.2. $\text{Span}(S^{3+8k} \times QP^{2l+1} \times QP^{2j+1}) \geq 8$ for $k \geq 1, l, j \geq 0$.

We now give an application to immersion of M into euclidean spaces. Suppose $w_4(M) = 0$ and $\dim M = n \equiv 11 \pmod{16} > 11$. Then following Massey one readily deduces that $\bar{w}_{n-j}(M) = 0$ for $j = 0, 1, 2, \dots, 10, 11$.

Let ν be the stable normal bundle of M . For stable bundle we can ignore the secondary obstruction in the top dimension given by Table 1 and Table 2. Thus by Theorem 4.4 we have

THEOREM 6.3. *Suppose $\text{Sq}^1 H^4(M) = 0$, $\phi_{0,0} H^4(M) = 0$ and $H_6(M; \mathbf{Z})$ has no 2-torsion. Then if $w_4(M) = 0$ and $n \equiv 11 \pmod{16} > 11$, M immerses in \mathbf{R}^{2n-7} .*

Consequently we have by Theorem 5.8:

THEOREM 6.4. *Suppose M is 4-connected mod 2 and $\dim M = n \equiv 11 \pmod{16} > 11$. Then M immerses in \mathbf{R}^{2n-7} if $\text{Sq}^2 H^{n-7}(M; \mathbf{Z}) = \text{Sq}^2 H^{n-7}(M)$ and immerses in \mathbf{R}^{2n-8} if $\text{Sq}^2 \text{Sq}^1 H^{n-8}(M) = \text{Sq}^2 H^{n-7}(M)$.*

Similarly we have by Theorem 5.7:

THEOREM 6.5. *Suppose M satisfies conditions A and B of §3, $w_4(M) = 0$ and $n \equiv 11 \pmod{16} > 11$. If $\text{Sq}^1 H^4(M) = \text{Sq}^2 H^4(M) = 0$ and if either $\chi \text{Sq}^4 H^{n-8}(M) = 0$ or $\phi_3 H^4(M) = 0$ and $\text{Sq}^2 H^{n-7}(M) = 0$ then M immerses in \mathbf{R}^{2n-8} .*

REFERENCES

- [1] J. Adém and S. Gitler, *Secondary characteristic classes and the immersion problem*, Bol. Soc. Mat. Mexicana, **8** (1963), 53–78.
- [2] S. Gitler and M. E. Mahowald, *The geometric dimension of real stable vector bundles*, Bol. Soc. Mat. Mexicana, **8** (1960), 85–106.
- [3] A. Hughes and E. Thomas, *A note on certain secondary cohomology operations*, Bol. Soc. Mat. Mexicana, **71** (1970), 661–667.
- [4] M. E. Mahowald, *The index of a tangent 2-field*, Pacific J. Math., **58** (1975), 539–548.
- [5] M. E. Mahowald and F. P. Peterson, *Secondary cohomology operations on the Thom class*, Topology, **2** (1964), 367–377.
- [6] C. R. F. Maunder, *Cohomology operations of the N -th kind*, Proc. London Math. Soc., (3) (1960), 125–154.
- [7] J. Milgram, *Cartan Formulae*, Illinois J. Math., **75** (1971), 633–647.
- [8] Tze-Beng Ng, *The existence of 7-fields and 8-fields on manifolds*, Quart. J. Math. Oxford, (2) **30** (1979), 197–221.
- [9] ———, *The mod 2 cohomology of $B\hat{\text{S}}\text{O}_n\langle 16 \rangle$* , to appear in Canad. J. Math.
- [10] D. G. Quillen, *The mod 2 cohomology rings of extra-special 2-groups and the spinor groups*, Math. Ann., **194** (1971), 197–212.
- [11] D. Randall, *Tangent frame fields on spin manifolds*, Pacific J. Math., **76** (1978), 157–167.

- [12] E. Thomas, *Postnikov invariants and higher order cohomology operations*, Ann. of Math., (2) **85** (1967), 184–217.
- [13] _____, *Real and complex vector fields on manifolds*, J. Math. and Mechanic, **16** (1967), 1183–1205.
- [14] _____, *The index of a tangent 2-fields*, Comment. Math. Helv., **42** (1967), 86–110.
- [15] _____, *The span of a manifold*, Quart. J. Math., Oxford (2) **19** (1968), 225–244.

Received June 8, 1984.

NATIONAL UNIVERSITY OF SINGAPORE
KENT RIDGE
SINGAPORE 0511

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN (Managing Editor)
University of California
Los Angeles, CA 90024

HEBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

CHARLES R. DEPRIMA
California Institute of Technology
Pasadena, CA 91125

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

H. SAMELSON
Stanford University
Stanford, CA 94305

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906-1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1986 by Pacific Journal of Mathematics

Jorge Almeida , Minimal nonpermutative pseudovarieties of semigroups. I	257
Jorge Almeida , Minimal nonpermutative pseudovarieties of semigroups. II	271
Carlos Andradas Heranz and José Manuel Gamboa Mutuberría , On projections of real algebraic varieties	281
Zeev Ditzian , Inverse theorems for multidimensional Bernstein operators ...	293
M. Furi and Maria Patrizia Pera , A continuation principle for forced oscillations on differentiable manifolds	321
James J. Hebda , The collars of a Riemannian manifold and stable isosystolic inequalities	339
Henryk Hecht and Dragan Miličić , Character identities and asymptotic behavior of matrix coefficients of discrete series	357
Piotr Jakóbczak , The boundary regularity of the solution of the $\bar{\partial}$ -equation in the product of strictly pseudoconvex domains	371
Krzysztof Jarosz , Isometries between injective tensor products of Banach spaces	383
Hans Keller , On valued, complete fields and their automorphisms	397
David Masser and Peter Man-Kit Shiu , On sparsely totient numbers	407
Tze-Beng Ng , Vector bundles over $(8k + 3)$ -dimensional manifolds	427
Thomas Joseph Ransford , The spectrum of an interpolated operator and analytic multivalued functions	445
Akihito Uchiyama , On the radial maximal function of distributions	467
Jang-Mei Gloria Wu , On singularity of harmonic measure in space	485