

Pacific Journal of Mathematics

ASYMPTOTIC EXPANSIONS OF THE LEBESGUE CONSTANTS FOR JACOBI SERIES

C. L. FRENZEN AND RODERICK SUE-CHUEN WONG

ASYMPTOTIC EXPANSIONS OF THE LEBESGUE CONSTANTS FOR JACOBI SERIES

C. L. FRENZEN AND R. WONG

Explicit expressions are obtained for the implied constants in the two O -terms in Lorch's asymptotic expansions of the Lebesgue constants associated with Jacobi series [Amer. J. Math., 81 (1959), 875–888]. In particular, a question of Szegő concerning asymptotic monotonicity of the Lebesgue constants for Laplace series is answered. Our method differs from that of Lorch, and makes use of some recently obtained uniform asymptotic expansions for the Jacobi polynomials and their zeros.

1. Introduction and summary. The n th partial sum of the Fourier series of an arbitrary function can be written in the form of an integral involving the Dirichlet kernel. The integral of the absolute value of this kernel is known as the n th *Lebesgue constant*, and is usually denoted by

$$(1.1) \quad L_n = \frac{1}{\pi} \int_0^\pi \frac{|\sin(n + \frac{1}{2})t|}{\sin(t/2)} dt;$$

see [19, p. 172]. The behavior of the sequence $\{L_n\}$ is closely connected with convergence and divergence properties of Fourier series, and the importance of this sequence has led many mathematicians to be concerned not only with just its asymptotic formula but also with its full asymptotic expansion. First, Fejer [1] showed that

$$(1.2) \quad L_n = \frac{4}{\pi^2} \log n + c_0 + \frac{c_1}{n} + \frac{\alpha(n)}{n^2},$$

where c_0 and c_1 are constants and $\alpha(n) = O(1)$ as $n \rightarrow \infty$. An explicit expression was given for c_0 but not for c_1 . Later, infinite asymptotic expansions were derived by Gronwall [4], Watson [18] and Hardy [7].

In an entirely analogous manner, the n th partial sum of the expansion of an arbitrary function in terms of Jacobi polynomials can be written as an integral involving a kernel; see, e.g., [17, p. 39]. The n th Lebesgue constant in this case has the integral representation

$$(1.3) \quad L_n(\alpha, \beta) = \frac{\Gamma(n + \alpha + \beta + 2)}{\Gamma(\alpha + 1)\Gamma(n + \beta + 1)} \cdot \int_0^\pi \left(\sin \frac{\theta}{2}\right)^{2\alpha+1} \left(\cos \frac{\theta}{2}\right)^{2\beta+1} |P_n^{(\alpha+1, \beta)}(\cos \theta)| d\theta.$$

This result is due to Rau [12]. In view of the identity [17, p. 60]

$$(1.4) \quad P_n^{(1/2, -1/2)}(\cos \theta) = \frac{\Gamma(n + \frac{3}{2})}{\Gamma(n + 1)\Gamma(\frac{3}{2})} \frac{\sin(n + \frac{1}{2})\theta}{(2n + 1)\sin(\theta/2)},$$

Equation (1.3) gives $L_n(-\frac{1}{2}, -\frac{1}{2}) = L_n$, contrary to a statement made in [8, footnote 7].

Rau [12] was the first to show that for $\alpha > -1/2$ and $\beta > -1$,

$$(1.5) \quad L_n(\alpha, \beta) = A_{\alpha\beta}n^{\alpha+1/2} + o(n^{\alpha+1/2}), \quad n \rightarrow \infty,$$

where

$$(1.6) \quad A_{\alpha\beta} = \frac{2}{\pi^{3/2}} \frac{\Gamma(\alpha/2 + 1/4)\Gamma(\beta/2 + 3/4)}{\Gamma(\alpha + 1)\Gamma([\alpha + \beta]/2 + 1)}.$$

Later Szegő [16] had an alternative proof of (1.5), and furthermore showed that

$$(1.7) \quad L_n(-\frac{1}{2}, \beta) = \frac{4}{\pi^2} \log n + o(\log n)$$

for $\beta > -1$, and that for $-1 < \alpha < -\frac{1}{2}$ and $\beta > -1$

$$(1.8) \quad L_n(\alpha, \beta) = \frac{2^{-\alpha}}{\Gamma(\alpha + 1)} \int_0^\infty \theta^\alpha |J_{\alpha+1}(\theta)| d\theta + o(1),$$

where $J_{\alpha+1}(\theta)$ is the Bessel function of first kind.

The above results have been sharpened by Lorch [8, 9], particularly in the cases $\alpha = -\frac{1}{2}$ and $-\frac{1}{2} < \alpha < \frac{1}{2}$. For $-\frac{1}{2} < \alpha < \frac{1}{2}$, $\alpha - \beta < 1$ and $\beta > -1$, Lorch's result can be stated as follows:

$$(1.9) \quad L_n(\alpha, \beta) = A_{\alpha\beta}n^{\alpha+1/2} + B_\alpha + O(n^{\alpha-1/2}) + O(n^{\alpha-\beta-1})$$

where

$$(1.10) \quad B_\alpha = \frac{2^{-\alpha}}{\Gamma(\alpha + 1)} \left\{ -M_1(\alpha) + \int_0^{j_1} x^\alpha J_{\alpha+1}(x) dx \right. \\ \left. + 2\alpha \sum_{k=1}^{\infty} (-1)^k \int_{j_k}^{j_{k+1}} x^{\alpha-1} J_\alpha(x) dx \right. \\ \left. + 2 \sum_{k=1}^{\infty} \left[M_k(\alpha) - \frac{\sqrt{2}}{\pi^{3/2}} \int_{j_{k-1}}^{j_k} x^{\alpha-1/2} dx \right] \right\},$$

both infinite series being absolutely convergent. (Equation (4) in [9] contains two misprints; $M_{k+1}(\alpha)$ should be replaced by $M_k(\alpha)$ and $M_1(\alpha)$ should have a minus sign.) In (1.10), $j_n = j_{\alpha+1, n}$ is the n th positive zero of

$J_{\alpha+1}(x)$, $n = 1, 2, \dots$, $j_0 = 0$, and

$$(1.11) \quad M_k(\alpha) \equiv (-1)^k (j_{\alpha+1, k})^\alpha J_\alpha(j_{\alpha+1, k}) > 0, \quad k = 1, 2, \dots$$

For $\alpha = -\frac{1}{2}$ and $\beta > -1$, Lorch also obtained the result that as $n \rightarrow \infty$,

$$(1.12) \quad L_n(-\tfrac{1}{2}, \beta) = \frac{4}{\pi^2} \log n + C_\beta \\ + O(n^{-1} \log n) + O(n^{-\beta-3/2}),$$

where

$$(1.13) \quad C_\beta = \frac{8}{\pi^2} \log 2 + \frac{2}{\pi} \int_0^1 \theta^{-1} \sin \theta \, d\theta \\ - \frac{4}{\pi^2} \int_0^{\pi/2} \frac{1 - (\cos \theta)^{\beta+1/2}}{\sin \theta} \, d\theta \\ - \frac{2}{\pi} \int_1^\infty \theta^{-1} \left\{ \frac{2}{\pi} - |\sin \theta| \right\} \, d\theta,$$

the last integral being convergent. (There is a typographical error in [9, (8)]; the factor in front of $\log 2$ should be $8/\pi^2$ and not $4/\pi^2$.)

Lorch's investigation [10] was motivated by a question raised by Szegő concerning asymptotic monotonicity of the sequence $\{L_n(0, 0)\}$; see also the editor's comment at the end of [13]. The result in (1.9), however, fails to answer the question of Szegő. Lorch thus posed to us in 1980 the problem of replacing the O -terms in (1.9) and (1.12) by explicitly determined expressions plus terms of lower asymptotic order. The following results provide a solution to his problem, and were announced in [3]. The detailed proofs of these results are the contents of the present paper. The fact that $\{L_n(0, 0)\}$ is an asymptotically increasing sequence is an immediate consequence of the result given in (1.18) below.

First, for the restricted range $-\frac{1}{2} < \alpha < \frac{1}{2}$ and $-\frac{1}{2} < \beta < \frac{1}{2}$, we have

$$(1.14) \quad L_n(\alpha, \beta) = A_{\alpha\beta} n^{\alpha+1/2} + B_\alpha + C_{\alpha\beta} A_{\alpha\beta} n^{\alpha-1/2} \\ + \frac{1}{\Gamma(\alpha+1)} D_\beta n^{\alpha-\beta-1} + O(n^{\alpha-3/2}),$$

where

$$(1.15) \quad C_{\alpha\beta} = \frac{(\alpha + \beta + 2)(\alpha + 1/2)}{2}$$

and

$$(1.16) \quad D_\beta = 2^{-\beta} \sum_{k=1}^{\infty} \left[\hat{M}_k(\beta) - \frac{\sqrt{2}}{\pi^{3/2}} \int_{j'_{k-1}}^{j'_k} x^{\beta+1/2} dx - \frac{\beta + 1/2}{\sqrt{2}\pi} \int_{j'_{k-1}}^{j'_k} x^{\beta-1/2} dx \right].$$

In (1.16), $j'_n \equiv j_{\beta,n}$ is the n th positive zero of $J_\beta(x)$, $n = 1, 2, \dots$, $j'_0 = 0$ and

$$(1.17) \quad \begin{aligned} \hat{M}_k(\beta) &= (-1)^k (j_{\beta,k})^{1+\beta} J_{\beta-1}(j_{\beta,k}) \\ &= (-1)^{k+1} (j_{\beta,k})^{1+\beta} J_{\beta+1}(j_{\beta,k}). \end{aligned}$$

In the important particular case of Laplace series (i.e., the series in terms of Legendre polynomials at the end point $x = 1$), $\alpha = \beta = 0$, and (1.14) becomes

$$(1.18) \quad \begin{aligned} L_n(0,0) &= \frac{2^{3/2}}{\sqrt{\pi}} n^{1/2} \\ &+ \left[1 + 2 \sum_{k=1}^{\infty} \left\{ M_k(0) - \frac{2^{3/2}}{\pi} [k^{1/2} - (k-1)^{1/2}] \right\} \right] \\ &+ \sqrt{\frac{2}{\pi}} n^{-1/2} + n^{-1} \sum_{k=1}^{\infty} \left\{ \hat{M}_k(0) - \frac{2^{3/2}}{3} [k^{3/2} - (k-1)^{3/2}] \right. \\ &\quad \left. - 2^{-3/2} [k^{1/2} - (k-1)^{1/2}] \right\} \\ &+ O(n^{-3/2}). \end{aligned}$$

The principal term in (1.18) was first given by Gronwall [5, 6], and later by Szegö [14, 15] with simpler proofs.

An improved version of (1.12) is

$$(1.19) \quad \begin{aligned} L_n(-\frac{1}{2}, \beta) &= \frac{4}{\pi^2} \log n + C_\beta + E_\beta n^{-1} + \frac{1}{\sqrt{\pi}} D_\beta n^{-\beta-3/2} \\ &+ O(n^{-2} \log n), \end{aligned}$$

valid for $-\frac{1}{2} < \beta < \frac{1}{2}$, where D_β is as given in (1.16) and

$$(1.20) \quad E_\beta = \frac{4}{\pi^2} \left(\frac{\beta + \frac{3}{2}}{2} \right).$$

Finally, we turn to the case $-1 < \alpha < -\frac{1}{2}$. Under the additional restrictions $\alpha - \beta > -1$ and $-\frac{1}{2} < \beta < \frac{1}{2}$, we have the following sharpened form of (1.8):

$$(1.21) \quad L_n(\alpha, \beta) = C_0 + C_1 n^{\alpha+1/2} + C_2 n^{\alpha-1/2} + C_3 n^{\alpha-\beta-1} + O(n^{-2}),$$

where

$$(1.22) \quad C_0 = \frac{2^{-\alpha}}{\Gamma(\alpha + 1)} \int_0^\infty x^\alpha |J_{\alpha+1}(x)| dx$$

$$(1.23) \quad C_1 = \frac{4}{\Gamma(\alpha + 1)\pi^{3/2}} \cdot \left\{ \frac{1}{\alpha + 1/2} \left(\frac{\pi}{2}\right)^{\alpha+1/2} + \int_0^{\pi/2} [(\sin \theta)^{\alpha-1/2}(\cos \theta)^{\beta+1/2} - \theta^{\alpha-1/2}] d\theta \right\}$$

$$(1.24) \quad C_2 = \frac{1}{2}(\alpha + \beta + 2)(\alpha + \frac{1}{2})C_1$$

$$(1.25) \quad C_3 = \frac{1}{\Gamma(\alpha + 1)} D_\beta,$$

D_β again being the same constant given in (1.16).

Lorch’s method essentially consists of replacing the Jacobi polynomial in (1.3) by its asymptotic formula of “Hilb’s type” [17, p. 197], and splitting the interval of integration $(0, \pi)$ at the points $j_{\alpha+1, k}/N$, $k = 1, \dots, [N]$, where $N = n + \frac{1}{2}(\alpha + \beta + 2)$. Our approach differs from that of Lorch. We first split the interval $(0, \pi)$ at the exact zeros of the Jacobi polynomial and then apply recently obtained uniform asymptotic expansions for the Jacobi polynomials and their zeros [2]. Our method may also be extended to give higher order approximations when desired.

2. Sketch of the procedure. For simplicity of presentation, we restrict our attention to the case $-\frac{1}{2} < \alpha < \frac{1}{2}$ and $-\frac{1}{2} < \beta < \frac{1}{2}$. Let θ_k denote the k th zero of $P_n^{(\alpha+1, \beta)}(\cos \theta)$, put $\bar{n} = [n/2]$, and write

$$(2.1) \quad \frac{\Gamma(\alpha + 1)\Gamma(n + \beta + 1)}{\Gamma(n + \alpha + \beta + 2)} L_n(\alpha, \beta) = \int_0^{\theta_{\bar{n}}} + \int_{\theta_{\bar{n}}}^\pi \equiv L_n^{(1)}(\alpha, \beta) + L_n^{(2)}(\alpha, \beta).$$

We shall first be concerned with the constant $L_n^{(1)}(\alpha, \beta)$. The evaluation of $L_n^{(2)}(\alpha, \beta)$ proceeds in a similar manner. For convenience, we set

$\theta_0 \equiv 0$ and define

$$(2.2) \quad I_k = \int_{\theta_k}^{\theta_{k+1}} \left(\sin \frac{\theta}{2}\right)^{2\alpha+1} \left(\cos \frac{\theta}{2}\right)^{2\beta+1} P_n^{(\alpha+1, \beta)}(\cos \theta) d\theta.$$

Since $\cos 0 = 1$ and

$$(2.3) \quad P_n^{(\alpha+1, \beta)}(1) = \binom{n + \alpha + 1}{n} = \frac{\Gamma(n + \alpha + 2)}{\Gamma(n + 1)\Gamma(\alpha + 2)} > 0,$$

we have

$$(2.4) \quad L_n^{(1)}(\alpha, \beta) = \sum_{k=0}^{\bar{n}-1} (-1)^k I_k.$$

To evaluate I_k , we shall use the following results given in [2]; see, in particular, the main theorem, Corollary 2, and the first paragraph in §5 of that reference.

LEMMA 1. For $\alpha + 1 > -\frac{1}{2}$ and $\alpha + \beta + 1 \geq -1$, we have

$$(2.5) \quad \begin{aligned} & \left(\sin \frac{\theta}{2}\right)^{\alpha+1} \left(\cos \frac{\theta}{2}\right)^{\beta} P_n^{(\alpha+1, \beta)}(\cos \theta) \\ &= \frac{\Gamma(n + \alpha + 2)}{n!} \left(\frac{\theta}{\sin \theta}\right)^{1/2} \left[\sum_{l=0}^{m-1} A_l(\theta) \frac{J_{\alpha+l+1}(N\theta)}{N^{\alpha+l+1}} + \sigma_m \right], \end{aligned}$$

where

$$(2.6) \quad N = n + \frac{1}{2}(\alpha + \beta + 2)$$

and

$$(2.7) \quad \sigma_m = \theta^m O(N^{-m-\alpha-1}),$$

the O -term being uniform with respect to $\theta \in [0, \pi - \epsilon]$, $\epsilon > 0$. The coefficients $A_l(\theta)$ are analytic functions in $0 \leq \theta \leq \pi - \epsilon$, and are $O(\theta^l)$ in that interval. In particular, $A_0(\theta) = 1$ and

$$(2.8) \quad A_1(\theta) = \left[(\alpha + 1)^2 - \frac{1}{4} \right] \left(\frac{1 - \theta \cot \theta}{2\theta} \right) - \frac{(\alpha + 1)^2 - \beta^2}{4} \tan \frac{\theta}{2}.$$

LEMMA 2. Let $\alpha + 1 > -\frac{1}{2}$, $\alpha + \beta + 1 \geq -1$ and let $0 < \theta_1 < \theta_2 < \dots < \theta_n < \pi$ be the zeros of $P_n^{(\alpha+1, \beta)}(\cos \theta)$. Then, as $n \rightarrow \infty$,

$$(2.9) \quad \begin{aligned} \theta_l &= \frac{j_{\alpha+1, l}}{N} \\ &+ \frac{1}{N^2} \left\{ \left[(\alpha + 1)^2 - \frac{1}{4} \right] \frac{1 - t \cot t}{2t} - \frac{(\alpha + 1)^2 - \beta^2}{4} \tan \frac{t}{2} \right\} \\ &+ t^2 O(N^{-3}), \end{aligned}$$

where $j_{\alpha+1,l}$ is the l -th positive zero of the Bessel function $J_{\alpha+1}(x)$ and $t = j_{\alpha+1,l}/N$. The O -term is uniformly bounded for all values of $l = 1, 2, \dots, [\gamma n]$, where $\gamma \in (0, 1)$ is a constant.

Taking $m = 3$ in (2.5) and substituting the resulting expression in (2.2) gives

$$(2.10) \quad I_k = \frac{\Gamma(n + \alpha + 2)}{n!N^{\alpha+1}} [I_k^{(1)} + I_k^{(2)} + I_k^{(3)} + I_k^{(4)}],$$

where

$$(2.11) \quad I_k^{(l+1)} = \frac{1}{N^l} \int_{\theta_k}^{\theta_{k+1}} A_l^*(\theta) J_{\alpha+l+1}(N\theta) d\theta, \quad l = 0, 1, 2,$$

and

$$(2.12) \quad I_k^{(4)} = \frac{1}{N^3} O \left\{ \int_{\theta_k}^{\theta_{k+1}} \theta^3 \left(\sin \frac{\theta}{2} \right)^\alpha \left(\cos \frac{\theta}{2} \right)^{\beta+1} \left(\frac{\theta}{\sin \theta} \right)^{1/2} d\theta \right\}$$

with

$$(2.13) \quad A_l^*(\theta) = \left(\frac{\theta}{2} \cot \frac{\theta}{2} \right)^{1/2} \left(\sin \frac{\theta}{2} \right)^\alpha \left(\cos \frac{\theta}{2} \right)^\beta A_l(\theta).$$

Note that the implied constant in the O -symbol in (2.12) is independent of k . It is easily seen that

$$(2.14) \quad \sum_{k=0}^{\bar{n}-1} (-1)^k I_k^{(4)} = O(N^{-3}).$$

Using the identity

$$(2.15) \quad \frac{d}{dz} (z^{-\nu} J_\nu(z)) = -z^{-\nu} J_{\nu+1}(z),$$

we have by integration by parts

$$(2.16) \quad I_k^{(2)} = T_1 + T_2,$$

where

$$(2.17) \quad T_1 = \frac{1}{N^2} [A_1^*(\theta_k) J_{\alpha+1}(N\theta_k) - A_1^*(\theta_{k+1}) J_{\alpha+1}(N\theta_{k+1})]$$

$$(2.18) \quad T_2 = \frac{1}{N^2} \int_{\theta_k}^{\theta_{k+1}} [\theta^{\alpha+1} A_1^*(\theta)]' \theta^{-\alpha-1} J_{\alpha+1}(N\theta) d\theta.$$

For fixed ν , it is well-known that

$$(2.19) \quad j_{\nu,k} = \left(k + \frac{1}{2}\nu - \frac{1}{4} \right) \pi - \frac{4\nu^2 - 1}{8(k + \frac{1}{2}\nu - \frac{1}{4})\pi} + O\left(\frac{1}{k^3} \right).$$

Thus, by Lemma 2, we have

$$(2.20) \quad N\theta_k = j_{\alpha+1,k} + O(k/N^2),$$

where the implied constant in the O -symbol is independent of k for $k = 1, \dots, \bar{n}$. Note that a combination of (2.6), (2.19) and (2.20) shows that for large n , we have $\theta_{\bar{n}} < \pi/2$. From (2.20), it also follows that

$$(2.21) \quad J_{\alpha+1}(N\theta_k) = J_{\alpha+1}(j_{\alpha+1,k}) + J'_{\alpha+1}(\xi)O(k/N^2),$$

where ξ lies between $N\theta_k$ and $j_{\alpha+1,k}$ and hence is $O(k)$. Since $J_{\alpha+1}(j_{\alpha+1,k}) = 0$ and $J'_{\alpha+1}(\xi) = O(\xi^{-1/2})$ as $\xi \rightarrow \infty$, (2.21) yields

$$(2.22) \quad J_{\alpha+1}(N\theta_k) = O(k^{1/2}/N^2).$$

Observe that $A_1^*(\theta) = O(\theta^{\alpha+1})$ and $\theta_k = O(k/N)$. Thus, by coupling (2.17) and (2.22), we obtain

$$(2.23) \quad T_1 = O(k^{\alpha+3/2}/N^{\alpha+5}).$$

Furthermore, since $[\theta^{\alpha+1}A_1^*(\theta)]' = O(\theta^{2\alpha+1})$, it is easily seen that

$$(2.24) \quad T_2 = \frac{1}{N^{\alpha+3}} O\left\{ \int_{N\theta_k}^{N\theta_{k+1}} y^\alpha |J_{\alpha+1}(y)| dy \right\}.$$

The integral inside of the curly bracket is clearly a bounded function of N if $k = 0$, in view of (2.20). For $k = 1, \dots, \bar{n} - 1$, we use (2.19) and the fact $J_{\alpha+1}(y) = O(y^{-1/2})$ to conclude that this integral is $O(k^{\alpha-1/2})$. Now we recall the well-known expansions [11, p. 292]

$$(2.25) \quad \sum_{k=1}^{n-1} k^\alpha - \zeta(-\alpha) \sim \frac{n^{\alpha+1}}{\alpha+1} \sum_{s=0}^{\infty} \binom{\alpha+1}{s} \frac{B_s}{n^s} \quad (\alpha \neq -1),$$

where $\zeta(\mu)$ is the Riemann-Zeta function. A combination of (2.23), (2.24) and (2.25) gives

$$(2.26) \quad \sum_{k=0}^{\bar{n}-1} (-1)^k I_k^{(2)} = O(N^{-5/2}).$$

An entirely similar argument leads to

$$(2.27) \quad \sum_{k=0}^{\bar{n}-1} (-1)^k I_k^{(3)} = O(N^{-5/2}).$$

Here we have used the fact that

$$(2.28) \quad \bar{n} = \frac{n}{2} \left[1 + O\left(\frac{1}{n}\right) \right], \quad \text{as } n \rightarrow \infty.$$

From (2.4) and (2.10), it now follows that as $n \rightarrow \infty$,

$$(2.29) \quad L_n^{(1)}(\alpha, \beta) = \frac{\Gamma(n + \alpha + 2)}{n!N^{\alpha+1}} \left[\sum_{k=0}^{\bar{n}-1} (-1)^k I_k^{(1)} + O(N^{-5/2}) \right].$$

In the next section, it will be shown that

$$\begin{aligned}
 (2.30) \quad & \sum_{k=0}^{\bar{n}-1} (-1)^k I_k^{(1)} \\
 &= \frac{1}{2^\alpha N^{\alpha+1}} \left\{ \int_0^{j_1} y^\alpha J_{\alpha+1}(y) dy + M_1(\alpha) \right. \\
 &\quad \left. + \sqrt{\frac{2}{\pi}} 2^{(\alpha-\beta-1)/2} n^{\alpha-1/2} + 2S_n + 2\alpha R_n \right. \\
 &\quad \left. + O(N^{\alpha-3/2}) \right\},
 \end{aligned}$$

where

$$(2.31) \quad S_n = \sum_{k=2}^{\bar{n}-1} g_{\alpha\beta}(\theta_k) M_k(\alpha),$$

$$(2.32) \quad R_n = \sum_{k=1}^{\bar{n}-1} (-1)^k \int_{j_k}^{j_{k+1}} y^{\alpha-1} J_\alpha(y) dy$$

and

$$(2.33) \quad g_{\alpha\beta}(\theta) = \left(\frac{\theta}{2} \cot \frac{\theta}{2} \right)^{1/2} \left[\frac{\sin \theta/2}{\theta/2} \right]^\alpha \left(\cos \frac{\theta}{2} \right)^\beta.$$

To proceed further, we need the following two lemmas, whose proofs are given in §4.

LEMMA 3. *As $n \rightarrow \infty$, the sum in (2.31) has the asymptotic approximation*

$$\begin{aligned}
 (2.34) \quad S_n \approx & B_\alpha^{(1)} + \frac{\sqrt{2}}{\pi^{3/2}} N^{\alpha+1/2} \left[\int_0^{\pi/2} g_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta \right. \\
 & \left. - \int_0^{j_1/N} g_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta \right. \\
 & \left. - \int_{j_{\bar{n}-1}/N}^{\pi/2} g_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta \right] \\
 & + \frac{n^{\alpha-1/2}}{\sqrt{2\pi}} 2^{(\alpha-\beta-1)/2} + O(n^{\alpha-3/2}),
 \end{aligned}$$

where $B_\alpha^{(1)}$ is a constant given by

$$(2.35) \quad B_\alpha^{(1)} = \sum_{k=2}^{\infty} \left[M_k(\alpha) - \frac{\sqrt{2}}{\pi^{3/2}} \int_{j_{k-1}}^{j_k} x^{\alpha-1/2} dx \right].$$

LEMMA 4. *The asymptotic behavior of the sum in (2.32) is given by*

$$(2.36) \quad R_n = B_\alpha^{(2)} + O(n^{\alpha-3/2}),$$

where

$$(2.37) \quad B_\alpha^{(2)} = \sum_{k=1}^{\infty} (-1)^k \int_{J_k}^{J_{k+1}} x^{\alpha-1} J_\alpha(x) dx.$$

The constants $B_\alpha^{(1)}$ and $B_\alpha^{(2)}$ are related to the constant B_α given in (1.10). The asymptotic approximation of $L_n^{(1)}(\alpha, \beta)$ is obtained by inserting (2.34) and (2.36) in (2.30), and combining the resulting expression with (2.29). It is anticipated that the terms involving the last two integrals in (2.34) will combine with similar terms from $L_n^{(2)}(\alpha, \beta)$.

Evaluation of $L_n^{(2)}(\alpha, \beta)$ proceeds as follows. By definition we have

$$(2.38) \quad L_n^{(2)}(\alpha, \beta) = \int_{\theta_{\bar{n}}}^{\pi} \left(\sin \frac{\theta}{2}\right)^{2\alpha+1} \left(\cos \frac{\theta}{2}\right)^{2\beta+1} |P_n^{(\alpha+1, \beta)}(\cos \theta)| d\theta,$$

In the above integral, we replace θ by $\pi - \theta$. The result is

$$(2.39) \quad L_n^{(2)}(\alpha, \beta) = \int_0^{\pi - \theta_{\bar{n}}} \left(\sin \frac{\theta}{2}\right)^{2\beta+1} \left(\cos \frac{\theta}{2}\right)^{2\alpha+1} |P_n^{(\beta, \alpha+1)}(\cos \theta)| d\theta,$$

on account of the identity [17, p. 59]

$$(2.40) \quad P_n^{(\alpha, \beta)}(\cos \theta) = (-1)^n P_n^{(\beta, \alpha)}(-\cos \theta).$$

From (2.40), it also follows that there is a one-to-one relationship between the zeros of $P_n^{(\alpha, \beta)}$ and $P_n^{(\beta, \alpha)}$. Suppose the zeros of $P_n^{(\alpha, \beta)}(\cos \theta)$ are arranged in the order:

$$0 < \theta_{n,1}^{(\alpha, \beta)} < \theta_{n,2}^{(\alpha, \beta)} < \dots < \theta_{n,n}^{(\alpha, \beta)} < \pi.$$

Since $\theta_{n,1}^{(\alpha, \beta)}$ is the smallest zero of $P_n^{(\alpha, \beta)}(\cos \theta)$, it follows that $\pi - \theta_{n,1}^{(\alpha, \beta)}$ is the largest zero of $P_n^{(\beta, \alpha)}(\cos \theta)$. Thus, $\pi - \theta_{n,1}^{(\alpha, \beta)} = \theta_{n,n}^{(\beta, \alpha)}$. In general, we have $\pi - \theta_{n,p}^{(\alpha, \beta)} = \theta_{n,n-p+1}^{(\beta, \alpha)}$, or equivalently

$$(2.41) \quad \theta_{n,p}^{(\alpha, \beta)} + \theta_{n,n-p+1}^{(\beta, \alpha)} = \pi$$

for $p = 1, \dots, n$. Since $\theta_{\bar{n}} \equiv \theta_{n,\bar{n}}^{(\alpha+1, \beta)}$, (2.41) gives $\pi - \theta_{\bar{n}} = \theta_{n,n-\bar{n}+1}^{(\beta, \alpha+1)}$. Set $\bar{m} = n - \bar{n} + 1$ and $\theta_{\bar{m}} \equiv \theta_{n,\bar{m}}^{(\beta, \alpha+1)}$. Equation (2.39) now becomes

$$(2.42) \quad L_n^{(2)}(\alpha, \beta) = \int_0^{\theta_{\bar{m}}} \left(\sin \frac{\theta}{2}\right)^{2\beta+1} \left(\cos \frac{\theta}{2}\right)^{2\alpha+1} |P_n^{(\beta, \alpha+1)}(\cos \theta)| d\theta.$$

Despite the fact that $\theta_{\bar{m}} > \pi/2$ while $\theta_n < \pi/2$ for large values of n , computation of $L_n^{(2)}(\alpha, \beta)$ proceeds in the same manner as $L_n^{(1)}(\alpha, \beta)$. A brief summary of this calculation is given in §5. Asymptotic expansion (1.14) is obtained by adding the results for $L_n^{(1)}(\alpha, \beta)$ and $L_n^{(2)}(\alpha, \beta)$ together. This is done in §6. Since the derivations of expansions (1.19) and (1.21) are similar to that of (1.14), they will not be presented here.

3. Calculation of $I_k^{(1)}$. From (2.11) and (2.13) we have

$$(3.1) \quad I_k^{(1)} = \int_{\theta_k}^{\theta_{k+1}} A_0^*(\theta) J_{\alpha+1}(N\theta) d\theta$$

where

$$(3.2) \quad A_0^*(\theta) = \left(\frac{\theta}{2} \cot \frac{\theta}{2}\right)^{1/2} \left(\sin \frac{\theta}{2}\right)^\alpha \left(\cos \frac{\theta}{2}\right)^\beta.$$

As in (2.16), integration by parts gives

$$(3.3) \quad I_k^{(1)} = F_1 + F_2,$$

for $k \neq 0$, where

$$(3.4) \quad F_1 = \frac{1}{N} [A_0^*(\theta_k) J_\alpha(N\theta_k) - A_0^*(\theta_{k+1}) J_\alpha(N\theta_{k+1})]$$

and

$$(3.5) \quad F_2 = \frac{1}{N} \int_{\theta_k}^{\theta_{k+1}} [A_0^*(\theta) \theta^\alpha]' \theta^{-\alpha} J_\alpha(N\theta) d\theta.$$

In terms of the function $g_{\alpha\beta}(\theta)$ defined in (2.33) (and also used in [9, (30)]), (3.4) becomes

$$(3.6) \quad F_1 = \frac{1}{2^\alpha N^{\alpha+1}} [g_{\alpha\beta}(\theta_k)(N\theta_k)^\alpha J_\alpha(N\theta_k) - g_{\alpha\beta}(\theta_{k+1})(N\theta_{k+1})^\alpha J_\alpha(N\theta_{k+1})].$$

By (2.20), Taylor's theorem gives

$$(3.7) \quad J_\alpha(N\theta_k) = J_\alpha(j_{\alpha+1,k}) + J'_\alpha(j_{\alpha+1,k}) O\left(\frac{k}{N^2}\right) + J''_\alpha(\xi) O\left(\frac{k^2}{N^4}\right),$$

where ξ lies between $N\theta_k$ and $j_{\alpha+1,k}$. Since $j_{\alpha+1,k} \sim \pi k$ by (2.19), $\xi = O(k)$ and hence $J''_\alpha(\xi) = O(k^{-1/2})$. From the identity $zJ'_\alpha(z) + zJ_{\alpha+1}(z) = \alpha J_\alpha(z)$, we have

$$(3.8) \quad J'_\alpha(j_{\alpha+1,k+1}) = \frac{\alpha}{j_{\alpha+1,k+1}} J_\alpha(j_{\alpha+1,k}) = O(k^{-3/2}).$$

Therefore, (3.7) becomes

$$(3.9) \quad J_\alpha(N\theta_k) = J_\alpha(j_{\alpha+1,k}) + O\left(\frac{1}{k^{1/2}N^2}\right) + O\left(\frac{k^{3/2}}{N^4}\right).$$

Since (2.19) and (2.20) imply

$$(3.10) \quad (N\theta_k)^\alpha = j_{\alpha+1,k}^\alpha + O\left(\frac{k^\alpha}{N^2}\right),$$

it follows from (3.6) that

$$(3.11) \quad F_1 = \frac{(-1)^k}{2^\alpha N^{\alpha+1}} [g_{\alpha\beta}(\theta_k)M_k(\alpha) + g_{\alpha\beta}(\theta_{k+1})M_{k+1}(\alpha)] + O\left(\frac{k^{\alpha-1/2}}{N^{\alpha+3}}\right),$$

where $M_k(\alpha)$ is as given in (1.11). In deriving (3.11), we have also used the fact that $g_{\alpha\beta}(\theta) = 1 + O(\theta^2)$ for $0 \leq \theta \leq \pi/2$.

We now evaluate F_2 given in (3.5). From (3.2), we obtain

$$(3.12) \quad [A_0^*(\theta)\theta^\alpha]' = \frac{\alpha}{2^{\alpha-1}}\theta^{2\alpha-1} + O(\theta^{2\alpha+1}).$$

Inserting (3.12) in (3.5) gives

$$(3.13) \quad F_2 = \frac{\alpha}{2^{\alpha-1}N^{\alpha+1}}Q + R,$$

where

$$(3.14) \quad Q = \int_{N\theta_k}^{N\theta_{k+1}} y^{\alpha-1}J_\alpha(y) dy$$

and

$$(3.15) \quad R = \frac{1}{N^{\alpha+3}}O\left\{\int_{N\theta_k}^{N\theta_{k+1}} y^{\alpha+1}|J_\alpha(y)| dy\right\}.$$

In view of (2.19), (2.20) and the behavior of the Bessel function, it is easily seen that the integral in (3.15) is equal to

$$\left|\int_{j_k}^{j_{k+1}} y^{\alpha+1}J_\alpha(y) dy\right| + O\left(\frac{k^{\alpha+3/2}}{N^2}\right).$$

Since $x^{\alpha+1}J_\alpha(x) = [x^{\alpha+1}J_{\alpha+1}(x)]'$, the last integral is zero. Thus

$$(3.16) \quad R = O(k^{\alpha+3/2}/N^{\alpha+5}).$$

By the same argument, we also have

$$(3.17) \quad Q = \int_{j_k}^{j_{k+1}} y^{\alpha-1}J_\alpha(y) dy + O\left(\frac{k^{\alpha-1/2}}{N^2}\right).$$

Now substitute (3.16) and (3.17) in (3.13), and add the resulting equation to (3.11). From (3.3) it follows that the sum is

$$(3.18) \quad I_k^{(1)} = \frac{(-1)^k}{2^\alpha N^{\alpha+1}} \left[g_{\alpha\beta}(\theta_k) M_k(\alpha) + g_{\alpha\beta}(\theta_{k+1}) M_{k+1}(\alpha) \right] \\ + \frac{\alpha}{2^{\alpha-1} N^{\alpha+1}} \int_{j_k}^{j_{k+1}} y^{\alpha-1} J_\alpha(y) dy + O\left(\frac{k^{\alpha-1/2}}{N^{\alpha+3}}\right)$$

for $k \neq 0$.

Note that (3.2) gives

$$(3.19) \quad A_0^*(\theta) = \left(\frac{\theta}{2}\right)^\alpha [1 + O(\theta^2)].$$

Inserting this in (3.1), we get

$$(3.20) \quad I_0^{(1)} = \frac{1}{2^\alpha} \int_0^{\theta_1} \theta^\alpha J_{\alpha+1}(N\theta) d\theta + O\left\{ \int_0^{\theta_1} \theta^{\alpha+2} J_{\alpha+1}(N\theta) d\theta \right\}.$$

Since (2.20) implies $\theta_1 = O(N^{-1})$, the second integral in (3.20) is $O(N^{-\alpha-3})$. By a similar argument, it can easily be shown that the first integral is equal to

$$\frac{1}{N^{\alpha+1}} \int_0^{j_1} y^\alpha J_{\alpha+1}(y) dy + O\left(\frac{1}{N^{\alpha+3}}\right).$$

Thus we have

$$(3.21) \quad I_0^{(1)} = \frac{1}{2^\alpha N^{\alpha+1}} \int_0^{j_1} y^\alpha J_{\alpha+1}(y) dy + O\left(\frac{1}{N^{\alpha+3}}\right).$$

From (2.25), (3.18) and (3.21), it now follows that

$$(3.22) \quad \sum_{k=0}^{\bar{n}-1} (-1)^k I_k^{(1)} \\ = \frac{1}{2^\alpha N^{\alpha+1}} \sum_{k=1}^{\bar{n}-1} \left[g_{\alpha\beta}(\theta_k) M_k(\alpha) + g_{\alpha\beta}(\theta_{k+1}) M_{k+1}(\alpha) \right] \\ + \frac{1}{2^\alpha N^{\alpha+1}} \int_0^{j_1} y^\alpha J_{\alpha+1}(y) dy + \frac{\alpha}{2^{\alpha-1} N^{\alpha+1}} R_n \\ + O\left(\frac{1}{N^{\alpha+3}}\right) + O(N^{-5/2})$$

where R_n is given in (2.32). The sum on the right-hand side of (3.22) can be written as

$$(3.23) \quad \sum_{k=1}^{\bar{n}-1} \left[g_{\alpha\beta}(\theta_k) M_k(\alpha) + g_{\alpha\beta}(\theta_{k+1}) M_{k+1}(\alpha) \right] \\ = g_{\alpha\beta}(\theta_1) M_1(\alpha) + g_{\alpha\beta}(\theta_{\bar{n}}) M_{\bar{n}}(\alpha) + 2S_n$$

with S_n being given in (2.31). Since $g_{\alpha\beta}(\theta) = 1 + O(\theta^2)$ and $\theta_1 = j_1/N + O(N^{-3})$, we obtain

$$(3.24) \quad g_{\alpha\beta}(\theta_1)M_1(\alpha) = M_1(\alpha) + O(N^{-2}).$$

Inserting (2.19) in the asymptotic expansion of $J_\alpha(x)$ gives

$$(3.25) \quad M_k(\alpha) = \sqrt{\frac{2}{\pi}} \left[j_{\alpha+1,k}^{\alpha-1/2} - \frac{1}{4}(\alpha + \frac{1}{2})(\alpha + \frac{3}{2})j_{\alpha+1,k}^{\alpha-5/2} + O(j_{\alpha+1,k}^{\alpha-7/2}) \right];$$

cf. (1.11). This in particular gives

$$(3.26) \quad M_{\bar{n}}(\alpha) = \left(\frac{\pi}{2}\right)^{\alpha-1} n^{\alpha-1/2} + O(n^{\alpha-3/2}).$$

Since $g_{\alpha\beta}(\theta)$ is analytic in $[0, \pi)$ (see (2.33)), we may expand it at $\theta = \pi/2$. Thus,

$$(3.27) \quad g_{\alpha\beta}(\theta_{\bar{n}}) = g_{\alpha\beta}\left(\frac{\pi}{2} + O(n^{-1})\right) = \left(\frac{\pi}{4}\right)^{1/2-\alpha} 2^{-(\alpha+\beta)/2} + O(n^{-1}),$$

and consequently

$$(3.28) \quad g_{\alpha\beta}(\theta_{\bar{n}})M_{\bar{n}}(\alpha) = \sqrt{\frac{2}{\pi}} 2^{(\alpha-\beta-1)/2} n^{\alpha-1/2} + O(n^{\alpha-3/2}).$$

A combination of (3.22), (3.23), (3.24) and (3.28) yields the desired result (2.30).

4. Proofs of Lemmas 3 and 4. We first give the proof of Lemma 4, which is considerably simpler than that of Lemma 3. Using the identity $[x^{\alpha+1}J_{\alpha+1}(x)]' = x^{\alpha+1}J_\alpha(x)$, we have by integration by parts

$$\int_{j_k}^{j_{k+1}} x^{\alpha-1}J_\alpha(x) dx = 2 \int_{j_k}^{j_{k+1}} x^{\alpha-2}J_{\alpha+1}(x) dx \quad (k \neq 0),$$

the integrated term vanishing since j_k and j_{k+1} are zeros of $J_{\alpha+1}(x)$. In view of the asymptotic behavior of $J_\alpha(x)$, the last integral is $O(k^{\alpha-5/2})$. Thus

$$\sum_{k=\bar{n}}^{\infty} (-1)^k \int_{j_k}^{j_{k+1}} x^{\alpha-1}J_\alpha(x) dx = O(n^{\alpha-3/2}).$$

This establishes Lemma 4.

Now we begin the proof of Lemma 3. By using (2.19) and (3.25), it can be shown that for $k \neq 1$,

$$(4.1) \quad M_k(\alpha) = \frac{2^{1/2}}{\pi^{3/2}} \int_{j_{k-1}}^{j_k} x^{\alpha-1/2} dx + \frac{\alpha - 1/2}{\sqrt{2\pi}} \int_{j_{k-1}}^{j_k} x^{\alpha-3/2} dx + O(k^{\alpha-5/2}).$$

Thus

$$(4.2) \quad \sum_{k=\bar{n}}^{\infty} \left[M_k(\alpha) - \frac{2^{1/2}}{\pi^{3/2}} \int_{j_{k-1}}^{j_k} x^{\alpha-1/2} dx - \frac{\alpha-1/2}{\sqrt{2\pi}} \int_{j_{k-1}}^{j_k} x^{\alpha-3/2} dx \right] = O(n^{\alpha-3/2}).$$

Set

$$(4.3) \quad S_{n,1} = \sum_{k=2}^{\infty} \left[M_k(\alpha) - \frac{2^{1/2}}{\pi^{3/2}} \int_{j_{k-1}}^{j_k} x^{\alpha-1/2} dx - \frac{\alpha-1/2}{\sqrt{2\pi}} \int_{j_{k-1}}^{j_k} x^{\alpha-3/2} dx \right]$$

$$(4.4) \quad S_{n,2} = \frac{2^{1/2}}{\pi^{3/2}} \sum_{k=2}^{\bar{n}-1} \int_{j_{k-1}}^{j_k} g_{\alpha\beta}(\theta_k) x^{\alpha-1/2} dx$$

and

$$(4.5) \quad S_{n,3} = \frac{\alpha-1/2}{(2\pi)^{1/2}} \sum_{k=2}^{\bar{n}-1} \int_{j_{k-1}}^{j_k} g_{\alpha\beta}(\theta_k) x^{\alpha-3/2} dx.$$

Since $g_{\alpha\beta}(\theta_k) = 1 + O(k^2/N^2)$, we can now express S_n as

$$(4.6) \quad S_n = S_{n,1} + S_{n,2} + S_{n,3} + O(n^{\alpha-3/2}).$$

Note that the series $S_{n,1}$ converges absolutely in view of (4.1), and is a constant independent of n .

(A) *Evaluation of $S_{n,2}$.* Make the change of variable $x = \theta N$ in (4.4), and write

$$(4.7) \quad g_{\alpha\beta}(\theta_k) = g_{\alpha\beta}(\theta) + g'_{\alpha\beta}(\theta)(\theta_k - \theta) + \frac{1}{2}g''_{\alpha\beta}(\xi)(\theta_k - \theta)^2,$$

where ξ is between θ_k and θ . Since $\theta \in [j_{k-1}/N, j_k/N]$, by (2.20) we have $\theta_k - \theta = O(1/N)$. Furthermore, since $g_{\alpha\beta}(\theta)$ is analytic in $[0, \pi/2]$, $g''_{\alpha\beta}(\xi)$ is bounded. Thus the remainder term $\frac{1}{2}g''_{\alpha\beta}(\xi)(\theta_k - \theta)^2$ is $O(n^{-2})$, and contributes to $S_{n,2}$ a term of order

$$O\left(n^{-2} \sum_{k=2}^{\bar{n}-1} \int_{j_{k-1}}^{j_k} x^{\alpha-1/2} dx \right) = O\left(n^{-2} \sum_{k=2}^{\bar{n}-1} k^{\alpha-1/2} \right) = O(n^{\alpha-3/2}).$$

Inserting (4.7) in (4.4) then leads to

$$(4.8) \quad S_{n,2} = \frac{2^{1/2}}{\pi^{3/2}} [S_{n,2}^{(1)} + S_{n,2}^{(2)} + S_{n,2}^{(3)}] + O(n^{\alpha-3/2}),$$

where

$$(4.9) \quad S_{n,2}^{(1)} = N^{\alpha+1/2} \sum_{k=2}^{\bar{n}-1} \int_{j_{k-1}/N}^{j_k} g_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta$$

$$(4.10) \quad S_{n,2}^{(2)} = N^{\alpha+1/2} \sum_{n=2}^{n-1} \theta_k \int_{j_{k-1}/N}^{j_k/N} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta$$

and

$$(4.11) \quad S_{n,2}^{(3)} = -N^{\alpha+1/2} \sum_{k=2}^{\bar{n}-1} \int_{j_{k-1}/N}^{j_k/N} g'_{\alpha\beta}(\theta) \theta^{\alpha+1/2} d\theta.$$

Clearly, $S_{n,2}^{(1)}$ can be written as

$$(4.12) \quad S_{n,2}^{(1)} = N^{\alpha+1/2} \left[\int_0^{\pi/2} - \int_0^{j_1/N} - \int_{j_{\bar{n}-1}/N}^{\pi/2} \right] g_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta,$$

which is exactly the second term on the right-hand side of (2.34) (except for the constant factor $\sqrt{2}/\pi^{3/2}$).

In (4.11), we let $g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} = f(\theta)$. With an appropriate change of variable, each integral there can be written as

$$(4.13) \quad \int_{j_{k-1}/N}^{j_k/N} \theta f(\theta) d\theta$$

$$= \frac{1}{2N} (j_k + j_{k-1}) \int_{j_{k-1}/N}^{j_k/N} f(\theta) d\theta$$

$$+ \int_0^{(j_k-j_{k-1})/2N} \theta \left[f\left\{ \frac{1}{2N} (j_k + j_{k-1}) + \theta \right\} \right. \\ \left. - f\left\{ \frac{1}{2N} (j_k + j_{k-1}) - \theta \right\} \right] d\theta.$$

By the mean value theorem the second integral on the right is equal to

$$(4.14) \quad \int_0^{(j_k-j_{k-1})/2N} 2\theta^2 f'(\xi_\theta) d\theta$$

for some ξ_θ satisfying

$$\frac{1}{2N} (j_k + j_{k-1}) - \theta < \xi_\theta < \frac{1}{2N} (j_k + j_{k-1}) + \theta.$$

Since $0 \leq \theta \leq (j_k - j_{k-1})/2N$ in (4.14), we have $j_{k-1}/N \leq \xi_\theta \leq j_k/N$ so that, by (2.19), $\xi_\theta = O(k/N)$. From (2.19) it also follows that $(j_k - j_{k-1})/2N = O(1/N)$. Note that the implied constants in the last two O -terms are independent of k . Since $f'(\theta) = O(\theta^{\alpha-1/2})$, a combination of the above results shows that the integral in (4.14) is $O(k^{\alpha-1/2}/N^{\alpha+5/2})$. By using this estimate and (4.13), the addition of (4.10) and (4.11) yields

$$(4.15) \quad S_{n,2}^{(2)} + S_{n,2}^{(3)}$$

$$= N^{\alpha+1/2} \sum_{k=2}^{\bar{n}-1} \left\{ \left[\theta_k - \frac{1}{2N} (j_k + j_{k-1}) \right] \int_{j_{k-1}/N}^{j_k/N} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta \right\}$$

$$+ O(N^{\alpha-3/2}),$$

where we have also made use of (2.25). From (2.19) and (2.20), it now follows that for $k \geq 1$,

$$(4.16) \quad \theta_k - \frac{1}{2N}(j_k + j_{k-1}) = \frac{\pi}{2N} + O\left(\frac{1}{k^2N}\right) + O\left(\frac{k}{N^3}\right).$$

Since $g_{\alpha\beta}(\theta) = 1 + O(\theta^2)$ for $0 \leq \theta \leq \pi/2$, each integral under the summation sign in (4.15) is $O(k^{\alpha+1/2}/N^{\alpha+3/2})$, where again the implied constant in the last O -term is independent of k . Since $-\frac{1}{2} < \alpha < \frac{1}{2}$, a combination of (4.15) with (4.16), (2.25) and the last estimate yields

$$(4.17) \quad S_{n,2}^{(2)} + S_{n,2}^{(3)} = \frac{\pi}{2} N^{\alpha-1/2} \sum_{k=2}^{\bar{n}-1} \int_{j_{k-1}/N}^{j_k/N} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta + O(N^{\alpha-3/2}).$$

The sum in (4.17) may now be written as

$$(4.18) \quad \int_0^{\pi/2} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta - \int_0^{j_1/N} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta - \int_{j_{\bar{n}-1}/N}^{\pi/2} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta.$$

The second integral in (4.18) is $O(N^{-\alpha-3/2})$ and, since $j_{\bar{n}-1}/N = \pi/2 + O(N^{-1})$, the third integral is $O(N^{-1})$. Coupling (4.17) and (4.18) results in

$$(4.19) \quad S_{n,2}^{(2)} + S_{n,2}^{(3)} = \frac{\pi}{2} N^{\alpha-1/2} \int_0^{\pi/2} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta + O(N^{\alpha-3/2}).$$

Finally, using (4.8), the addition of (4.12) and (4.19) gives

$$(4.20) \quad S_{n,2} = \frac{2^{1/2}}{\pi^{3/2}} N^{\alpha+1/2} \left[\int_0^{\pi/2} - \int_0^{j_1/N} - \int_{j_{\bar{n}-1}/N}^{\pi/2} \right] g_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta + \frac{1}{\sqrt{2\pi}} N^{\alpha-1/2} \int_0^{\pi/2} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta + O(n^{\alpha-3/2}).$$

(B) *Evaluation of $S_{n,3}$.* The analysis here parallels that given for $S_{n,2}$, and is in fact simpler. We first make the change of variable $x = \theta N$ in (4.5), and then substitute (4.7) in (4.5). The result is

$$(4.21) \quad S_{n,3} = \frac{\alpha - 1/2}{(2\pi)^{1/2}} N^{\alpha-1/2} [S_{n,3}^{(1)} + S_{n,3}^{(2)}] + O(n^{-2}),$$

where

$$(4.22) \quad S_{n,3}^{(1)} = \int_{j_1/N}^{j_{\bar{n}-1}/N} g_{\alpha\beta}(\theta) \theta^{\alpha-3/2} d\theta$$

and

$$(4.23) \quad S_{n,3}^{(2)} = \sum_{k=2}^{\bar{n}-1} \int_{j_{k-1}/N}^{j_k/N} g'_{\alpha\beta}(\theta) (\theta_k - \theta) \theta^{\alpha-3/2} d\theta;$$

cf. (4.8). To the integral in (4.22), we apply integration by parts. By using (3.27) and the fact that $g_{\alpha\beta}(\theta) = 1 + O(\theta^2)$ as $\theta \rightarrow 0^+$, it can be shown that the integrated term is equal to

$$\frac{1}{\alpha - 1/2} \left[2^{(\alpha-\beta-1)/2} - (j_1/N)^{\alpha-1/2} + O(n^{-1}) \right].$$

The other term can be replaced by

$$\frac{1}{\alpha - 1/2} \int_0^{\pi/2} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta + O(n^{-1}),$$

as was done in the case of $S_{n,2}$, cf. (4.18). Thus

$$(4.24) \quad S_{n,3}^{(1)} = \frac{1}{\alpha - 1/2} \left[2^{(\alpha-\beta-1)/2} - (j_1/N)^{\alpha-1/2} - \int_0^{\pi/2} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta \right] + O(n^{-1}).$$

Since $g'_{\alpha\beta} = O(\theta)$ and $\theta_k - \theta = O(n^{-1})$ for $j_{k-1}/N \leq \theta \leq j_k/N$, the $(k - 1)$ th integral in (4.23) is $O(k^{\alpha-1/2}/N^{\alpha+3/2})$. This implies

$$(4.25) \quad S_{n,3}^{(2)} = O(N^{-1}).$$

The sum of $S_{n,3}^{(1)}$ and $S_{n,3}^{(2)}$ gives, using (4.21),

$$(4.26) \quad S_{n,3} = -\frac{j_1^{\alpha-1/2}}{\sqrt{2\pi}} + \frac{N^{\alpha-1/2}}{\sqrt{2\pi}} \left[2^{(\alpha-\beta-1)/2} - \int_0^{\pi/2} g'_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta \right] + O(n^{\alpha-3/2}),$$

thus completing the evaluation of $S_{n,3}$.

Observing that the sum $S_{n,1}$ in (4.3) can be written as

$$(4.27) \quad S_{n,1} = B_\alpha^{(1)} + \frac{j_1^{\alpha-1/2}}{\sqrt{2\pi}},$$

where $B_\alpha^{(1)}$ is given in (2.35), the result in (2.34) now follows immediately from (4.6), (4.20) and (4.26). This proves Lemma 3.

5. Evaluation of $L_n^{(2)}(\alpha, \beta)$. In what follows we shall use the same notation as we did in the evaluation of $L_n^{(1)}(\alpha, \beta)$. There should be no confusion resulting from this, when care is taken to distinguish the zeros of $P_n^{(\alpha+1, \beta)}(\cos \theta)$ and $P_n^{(\beta, \alpha+1)}(\cos \theta)$. Thus we again let $\theta_0 \equiv 0$, θ_k denote the k th positive zero of $P_n^{(\beta, \alpha+1)}(\cos \theta)$, and

$$(5.1) \quad I_k = \int_{\theta_k}^{\theta_{k+1}} \left(\sin \frac{\theta}{2} \right)^{2\beta+1} \left(\cos \frac{\theta}{2} \right)^{2\alpha+1} P_n^{(\beta, \alpha+1)}(\cos \theta) d\theta.$$

Since $\cos 0 = 1$ and

$$(5.2) \quad P_n^{(\beta, \alpha+1)}(1) = \frac{\Gamma(n + \beta + 1)}{\Gamma(n + 1)\Gamma(\beta + 1)} > 0,$$

we have from (2.42)

$$(5.3) \quad L_n^{(2)}(\alpha, \beta) = \sum_{k=0}^{\bar{m}-1} (-1)^k I_k.$$

As before, we now use the asymptotic expansion of $P_n^{(\beta, \alpha+1)}(\cos \theta)$ for large n . The result corresponding to (2.29) is

$$(5.4) \quad L_n^{(2)}(\alpha, \beta) = \frac{\Gamma(n + \beta + 1)}{n!N^\beta} \left[\sum_{k=0}^{\bar{m}-1} (-1)^k I_k^{(1)} + O(n^{-5/2}) \right],$$

where

$$(5.5) \quad \sum_{k=0}^{\bar{m}-1} (-1)^k I_k^{(1)} = \frac{1}{2^{\beta+1}N^{\beta+2}} \left\{ \int_0^{j_{\beta,1}} y^{\beta+1} J_\beta(y) dy + \hat{M}_1(\beta) + \sqrt{\frac{2}{\pi}} 2^{(\beta-\alpha+1)/2} n^{\beta+1/2} + 2\hat{S}_n + O(n^{\beta-1/2}) \right\},$$

$$(5.6) \quad \hat{S}_n = \sum_{k=2}^{\bar{m}-1} \hat{g}_{\alpha\beta}(\theta_k) \hat{M}_k(\beta),$$

$\hat{M}_k(\beta)$ being as given in (1.17), and

$$(5.7) \quad \hat{g}_{\alpha\beta}(\theta) = \left(\frac{2}{\theta} \tan \frac{\theta}{2} \right)^{1/2} \left(\frac{2}{\theta} \sin \frac{\theta}{2} \right)^\beta \left(\cos \frac{\theta}{2} \right)^\alpha.$$

Note that \hat{R}_n , the contribution to $L_n^{(2)}(\alpha, \beta)$ corresponding to R_n in (2.32), is absent in (5.5), because

$$\int_{j_{\beta,k}}^{j_{\beta,k+1}} y^\beta J_{\beta-1}(y) dy = \int_{j_{\beta,k}}^{j_{\beta,k+1}} [y^\beta J_\beta(y)]' dy = 0.$$

An analogue of Lemma 3 is the following approximation:

$$(5.8) \quad \hat{S}_n = \frac{\sqrt{2}}{\pi^{3/2}} N^{\beta+3/2} \left(\int_0^{\pi/2} - \int_0^{j_{\beta,1}/N} - \int_{j_{\beta,\bar{m}-1}/N}^{\pi/2} \right) \hat{g}_{\alpha\beta}(\theta) \theta^{\beta+1/2} d\theta + \sqrt{\frac{2}{\pi}} 2^{(\beta-\alpha-1)/2} N^{\beta+1/2} + 2^\beta D_\beta - \hat{M}_1(\beta) + \frac{2^{1/2}}{\pi^{3/2}(\beta + \frac{3}{2})} j_{\beta,1}^{\beta+3/2} + O(N^{\beta-1/2}),$$

where D_β is given in (1.16). Note that the sum in D_β starts with $k = 1$, whereas the sum in $B_\alpha^{(1)}$, the corresponding constant in S_n given in (2.35), starts with $k = 2$. Since $\hat{g}_{\alpha\beta}(\theta) = 1 + O(\theta^2)$, the term involving the second integral on the right of (5.8) cancels with the third from the last term in the same equation. Thus

$$(5.9) \quad \hat{S}_n = \frac{\sqrt{2}}{\pi^{3/2}} N^{\beta+3/2} \left(\int_0^{\pi/2} - \int_{j_\beta, \bar{m}-1/N}^{\pi/2} \right) \hat{g}_{\alpha\beta}(\theta) \theta^{\beta+1/2} d\theta$$

$$+ \sqrt{\frac{2}{\pi}} 2^{(\beta-\alpha-1)/2} N^{\beta+1/2} + 2^\beta D_\beta - \hat{M}_1(\beta)$$

$$+ O(N^{\beta-1/2}).$$

Now we insert (5.9) in (5.5), and observe that

$$(5.10) \quad \int_0^{j_\beta, 1} x^{\beta+1} J_\beta(x) dx = \hat{M}_1(\beta)$$

which follows from (1.17) and the identity $[x^{\beta+1} J_{\beta+1}(x)]' = x^{\beta+1} J_\beta(x)$. The resulting expression from (5.5) coupled with (5.4) gives

$$(5.11) \quad \frac{\Gamma(n + \alpha + \beta + 2)}{\Gamma(\alpha + 1)\Gamma(n + \beta + 1)} L_n^{(2)}(\alpha, \beta)$$

$$= \frac{2^{-1-\beta}}{\Gamma(\alpha + 1)} \left[n^{\alpha-\beta-1} + \frac{1}{2}(\alpha + \beta + 2)(\alpha - \beta - 1)n^{\alpha-\beta-2} \right]$$

$$\cdot \left\{ 2^{\beta+1} D_\beta + 2\sqrt{\frac{2}{\pi}} 2^{(\beta-\alpha+1)/2} n^{\beta+1/2} \right.$$

$$+ \frac{2^{3/2}}{\pi^{3/2}} N^{\beta+3/2} \left(\int_0^{\pi/2} - \int_{j_\beta, \bar{m}-1/N}^{\pi/2} \right)$$

$$\cdot 2^{\beta+1/2} \left(\sin \frac{\theta}{2} \right)^{\beta+1/2} \left(\cos \frac{\theta}{2} \right)^{\alpha-1/2} d\theta \left. \right\}$$

$$+ O(n^{\alpha-3/2}),$$

where use has been made of (5.7) and the asymptotic expansion

$$(5.12) \quad \frac{2^{-1-\beta}\Gamma(n + \alpha + \beta + 2)}{\Gamma(\alpha + 1)\Gamma(n + 1)N^{2\beta+2}}$$

$$= \frac{2^{-1-\beta}}{\Gamma(\alpha + 1)} n^{\alpha-\beta-1} \left[1 + \frac{1}{2n}(\alpha + \beta + 2)(\alpha - \beta - 1) + O(n^{-2}) \right].$$

6. The sum of $L_n^{(1)}(\alpha, \beta)$ and $L_n^{(2)}(\alpha, \beta)$. From (2.29), (2.30) and the fact that

$$(6.1) \quad \frac{\Gamma(n + \alpha + \beta + 2)}{\Gamma(n + \beta + 1)} \frac{\Gamma(n + \alpha + 2)}{n!N^{2\alpha+2}} = 1 + O(n^{-2}),$$

we have

$$(6.2) \quad \begin{aligned} & \frac{\Gamma(n + \alpha + \beta + 2)}{\Gamma(\alpha + 1)\Gamma(n + \beta + 1)} L_n^{(1)}(\alpha, \beta) \\ &= \frac{1}{2^\alpha \Gamma(\alpha + 1)} \left\{ \int_0^{j_1} y^\alpha J_{\alpha+1}(y) dy + M_1(\alpha) \right. \\ & \quad \left. + \sqrt{\frac{2}{\pi}} 2^{(\alpha-\beta-1)/2} n^{\alpha-1/2} + 2S_n + 2\alpha R_n \right\} \\ & \quad + O(N^{\alpha-3/2}). \end{aligned}$$

Since $g_{\alpha\beta}(\theta) = 1 + O(\theta^2)$, the second integral on the right-hand side of (2.34) can be written as

$$(6.3) \quad \int_0^{j_1/N} g_{\alpha\beta}(\theta) \theta^{\alpha-1/2} d\theta = \frac{1}{N^{\alpha+1/2}} \int_0^{j_1} x^{\alpha-1/2} dx + O(N^{-\alpha-5/2}).$$

Now write (2.35) as

$$(6.4) \quad \begin{aligned} B_\alpha^{(1)} &= \sum_{k=1}^\infty \left[M_k(\alpha) - \frac{\sqrt{2}}{\pi^{3/2}} \int_{j_{k-1}}^{j_k} x^{\alpha-1/2} dx \right] \\ & \quad - M_1(\alpha) + \frac{\sqrt{2}}{\pi^{3/2}} \int_0^{j_1} x^{\alpha-1/2} dx, \end{aligned}$$

where we have used the fact that $j_0 = 0$. Combining (6.1), (2.34), (2.36), (6.2) and (6.3) gives

$$(6.5) \quad \begin{aligned} & \frac{\Gamma(n + \alpha + \beta + 2)}{\Gamma(\alpha + 1)\Gamma(n + \beta + 1)} L_n^{(1)}(\alpha, \beta) \\ &= B_\alpha + \frac{1}{2^\alpha \Gamma(\alpha + 1)} \\ & \quad \cdot \left\{ \frac{2^{\alpha+1}}{\pi^{3/2}} N^{\alpha+1/2} \int_0^{\pi/2} \left(\sin \frac{\theta}{2} \right)^{\alpha-1/2} \left(\cos \frac{\theta}{2} \right)^{\beta+1/2} d\theta \right. \\ & \quad + \frac{2^{3/2}}{\pi^{3/2}} 2^{(\alpha-\beta-1)/2} n^{\alpha-1/2} \\ & \quad \left. - \frac{2^{\alpha+1}}{\pi^{3/2}} N^{\alpha+1/2} \int_{j_{\bar{n}-1}/N}^{\pi/2} \left(\sin \frac{\theta}{2} \right)^{\alpha-1/2} \left(\cos \frac{\theta}{2} \right)^{\beta+1/2} d\theta \right\} \\ & \quad + O(n^{\alpha-3/2}), \end{aligned}$$

where B_α is given in (1.10) and we have used (2.33) and (2.6).

We now note that

$$(6.6) \quad \left[n^{\alpha-\beta-1} + \frac{1}{2}(\alpha + \beta + 2)(\alpha - \beta - 1)n^{\alpha-\beta-2} \right] N^{\beta+3/2} \\ = n^{\alpha+1/2} + C_{\alpha\beta}n^{\alpha-1/2} + O(n^{\alpha-3/2}),$$

where $C_{\alpha\beta}$ is given in (1.15). Upon adding (6.5) and (5.11) and using (6.6), we obtain

$$(6.7) \quad L_n(\alpha, \beta) = I_1^* + I_2^* + I_3^* + B_\alpha + \frac{1}{\Gamma(\alpha + 1)} D_\beta \\ + O(n^{\alpha-3/2}),$$

where B_α and D_β are given in (1.10) and (1.16) respectively and where

$$(6.8) \quad I_1^* = \frac{2}{\pi^{3/2}\Gamma(\alpha + 1)} \left(n^{\alpha+1/2} + C_{\alpha\beta}n^{\alpha-1/2} \right) \\ \cdot \left\{ \int_0^{\pi/2} \left(\sin \frac{\theta}{2} \right)^{\alpha-1/2} \left(\cos \frac{\theta}{2} \right)^{\beta+1/2} d\theta \right. \\ \left. + \int_0^{\pi/2} \left(\sin \frac{\theta}{2} \right)^{\beta+1/2} \left(\cos \frac{\theta}{2} \right)^{\alpha-1/2} d\theta \right\} \\ + O(n^{\alpha-3/2}),$$

$$(6.9) \quad I_2^* = -\frac{2n^{\alpha+1/2}}{\Gamma(\alpha + 1)} \left[\int_{j_{n-1}/N}^{\pi/2} \left(\sin \frac{\theta}{2} \right)^{\alpha-1/2} \left(\cos \frac{\theta}{2} \right)^{\beta+1/2} d\theta \right. \\ \left. + \int_{j_{\beta, \bar{m}-1}/N}^{\pi/2} \left(\sin \frac{\theta}{2} \right)^{\beta+1/2} \left(\cos \frac{\theta}{2} \right)^{\alpha-1/2} d\theta \right],$$

and

$$(6.10) \quad I_3^* = \frac{4}{\Gamma(\alpha + 1)} \sqrt{\frac{2}{\pi}} 2^{-(\alpha+\beta+1)/2} n^{\alpha-1/2}.$$

By letting $\theta = \pi - \phi$ in the second integral in (6.8), the two integrals there can be combined into the single integral

$$(6.11) \quad \int_0^\pi \left(\sin \frac{\theta}{2} \right)^{\alpha-1/2} \left(\cos \frac{\theta}{2} \right)^{\beta+1/2} d\theta = \frac{\Gamma(\frac{1}{2}\alpha + \frac{1}{4})\Gamma(\frac{1}{2}\beta + \frac{3}{4})}{\Gamma(\frac{1}{2}(\alpha + \beta) + 1)}.$$

Using the above result in (6.8) yields

$$(6.12) \quad I_1^* = A_{\alpha\beta}n^{\alpha+1/2} + C_{\alpha\beta}A_{\alpha\beta}n^{\alpha-1/2} + O(n^{\alpha-3/2}),$$

where $A_{\alpha\beta}$ is given in (1.6). Making the change of variable $\theta = \pi - \phi$ in the second integral in (6.9), the two integrals there can also be combined

into the single integral

$$(6.13) \quad \int_{j_{\bar{n}-1}/N}^{\pi - j_{\beta, \bar{m}-1}/N} \left(\sin \frac{\theta}{2}\right)^{\alpha-1/2} \left(\cos \frac{\theta}{2}\right)^{\beta+1/2} d\theta.$$

Note that by (2.20) and (2.41), we have

$$\pi - \frac{j_{\beta, \bar{m}-1}}{N} = \theta_{\bar{n}+1} + O(n^{-2}), \quad \frac{j_{\bar{n}-1}}{N} = \theta_{\bar{n}-1} + O(n^{-2}).$$

From (2.19) and (2.20), it also follows that

$$\theta_{\bar{n}+1} = \frac{\pi}{2} + O(n^{-1}), \quad \theta_{\bar{n}-1} = \frac{\pi}{2} + O(n^{-1}),$$

and

$$\theta_{\bar{n}+1} - \theta_{\bar{n}-1} = \frac{2\pi}{n} + O(n^{-2}).$$

Expanding the integrand in (6.13) about $\pi/2$ and using the above results, the integral in (6.13) can be shown to be

$$\frac{2\pi}{n} 2^{-(\alpha+\beta)/2} + O(n^{-2}).$$

This result coupled with (6.9) gives

$$(6.14) \quad I_2^* = -\frac{4}{\Gamma(\alpha+1)} \sqrt{\frac{2}{\pi}} 2^{-(\alpha+\beta+1)/2} n^{\alpha-1/2} + O(n^{\alpha-3/2}).$$

A combination of (6.7), (6.10), (6.12) and (6.14) yields our final result (1.14).

To conclude this paper, we consider the particular case of Laplace series given in (1.18). The constant term and the $O(n^{-1})$ term in (1.18) have a somewhat different appearance from those obtained by putting $\alpha = \beta = 0$ in (1.14). The transition to the form in (1.18) is made by writing the second infinite series in (1.10) as

$$(6.15) \quad \begin{aligned} &\lim_{n \rightarrow \infty} \left[M_1(0) + \dots + M_n(0) - \frac{\sqrt{2}}{\pi^{3/2}} \int_0^{j_{1,n}} x^{-1/2} dx \right] \\ &= \lim_{n \rightarrow \infty} \left[M_1(0) + \dots + M_n(0) - \frac{2^{3/2}}{\pi} n^{1/2} \right] \\ &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \left\{ M_k(0) - \frac{2^{3/2}}{\pi} (k^{1/2} - [k-1]^{1/2}) \right\}, \end{aligned}$$

since from (2.19), $(j_{1,n})^{1/2} = (n\pi)^{1/2} + O(n^{-1/2})$. Now using the fact that, for $\alpha = 0$,

$$(6.16) \quad \int_0^{j_1} x^\alpha J_{\alpha+1}(x) dx = \int_0^{j_{1,1}} J_1(x) dx = J_0(0) - J_0(j_{1,1}) = 1 + M_1(0),$$

the result for the constant term in (1.18) follows. Similarly in (1.16) we write

$$(6.17) \quad D_0 = \lim_{n \rightarrow \infty} \left[\hat{M}_1(0) + \dots + \hat{M}_n(0) - \frac{\sqrt{2}}{3/2} \int_0^{j_{0,n}} x^{1/2} dx - \frac{1}{2^{3/2}\pi^{1/2}} \int_0^{j_{0,n}} x^{-1/2} dx \right] \\ = \lim_{n \rightarrow \infty} \sum_{k=1}^n \left\{ \hat{M}_k(0) - \frac{2^{3/2}}{3} (k^{3/2} - [k-1]^{3/2}) - 2^{-3/2} (k^{1/2} - [k-1]^{1/2}) \right\},$$

where we have used $(j_{0,n})^{3/2} = \pi^{3/2}(n^{3/2} - \frac{3}{8}n^{1/2}) + O(n^{-1/2})$. Grouping these results together yields the form of $L_n(0, 0)$ given in (1.18).

Acknowledgment. We would like to thank Professor Lee Lorch for posing the problem to us as well as providing some historical background.

REFERENCES

- [1] L. Fejer, *Lebesguesche Konstanten und divergente Fourierreihen*, J. Reine Angew. Math., **138** (1910), 22–53.
- [2] C. L. Frenzen and R. Wong, *A uniform asymptotic expansion of the Jacobi polynomials with error bounds*, Canad. J. Math., to appear.
- [3] ———, *On the asymptotic behavior of the Lebesgue constants for Jacobi series*, C. R. Math. Acad. Sci. Canada, **6** (1984), 267–271.
- [4] T. H. Gronwall, *Über die Lebesgueschen Konstanten bei den Fourierschen Reihen*, Math. Ann., **72** (1912), 244–261.
- [5] ———, *Über die Laplacesche Reihe*, Math. Ann., **74** (1913), 213–270.
- [6] ———, *On the degree of convergence of Laplace series*, Trans. Amer. Math. Soc., **15** (1914), 1–30.
- [7] G. H. Hardy, *Note on Lebesgue’s constants in the theory of Fourier series*, J. London Math. Soc., **17** (1942), 4–13.
- [8] L. Lorch, *The Lebesgue constants for Jacobi series*, I, Proc. Amer. Math. Soc., **10** (1959), 756–761.
- [9] ———, *The Lebesgue constants for Jacobi series*, II, Amer. J. Math., **81** (1959), 875–888.
- [10] ———, private communication.

- [11] F. W. J. Olver, *Asymptotics and Special Functions*, Academic Press, New York, 1974.
- [12] H. Rau, *Über die Lebesgueschen Konstanten der Reihenentwicklungen nach Jacobischen Polynomen*, *J. Reine Angew. Math.*, **161** (1929), 237–254.
- [13] G. Szegő, *Über die Lebesgueschen Konstanten bei den Fourierschen Reihen*, in *The Collected Papers of Gabor Szegő*, Vol. 1, pp. 309–313, R. Askey, ed., Birkhauser, Cambridge, MA, 1982.
- [14] _____, *Über einige asymptotische Entwicklungen der Legendreschen Funktionen*, *Proc. London Math. Soc.*, (2), **36** (1932), 427–450.
- [15] _____, *Über eine von Herrn S. Berstein herrührende Abschätzung der Legendreschen Polynome*, *Math. Ann.*, **108** (1933), 360–369.
- [16] _____, *Asymptotische Entwicklungen der Jacobischen Polynome*, *Schr. der König. Gelehr. Gesell. Naturwiss. Kl.*, **10** (1933), 33–112.
- [17] _____, *Orthogonal Polynomials*, Colloquium Publications, Vol. 23, 3rd ed., Amer. Math. Soc., Providence, RI 1967.
- [18] G. N. Watson, *The constants of Landau and Lebesgue*, *Quart. J. Math.*, **1** (1930), 310–318.
- [19] A. Zygmund, *Trigonometric Series*, Warsaw, 1935.

Received September 4, 1984 and in revised form May 31, 1985. The second author is an I. W. Killam Research Fellow, 1982-1984. The work of this author was also partially supported by N.S.E.R.C. Grant No. A7359.

UNIVERSITY OF BRITISH COLUMBIA,
VANCOUVER, B.C., V6T 1Y4, CANADA

AND

UNIVERSITY OF MANITOBA
WINNIPEG, MANITOBA R3T 2N2, CANADA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN (Managing Editor)
University of California
Los Angeles, CA 90024

HEBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

CHARLES R. DEPRIMA
California Institute of Technology
Pasadena, CA 91125

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

H. SAMELSON
Stanford University
Stanford, CA 94305

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1986 by Pacific Journal of Mathematics

Gideon Amit and David Chillag , On a question of Feit concerning character values of finite solvable groups	257
Constantin Gelu Apostol and Frank Larkin Gilfeather , Isomorphisms modulo the compact operators of nest algebras	263
Parviz Azimi and James Neil Hagler , Examples of hereditarily l^1 Banach spaces failing the Schur property	287
Brian Evan Blank , Boundary behavior of limits of discrete series representations of real rank one semisimple groups	299
Jeffrey Carroll , Some undecidability results for lattices in recursion theory	319
Gerald Howard Cliff and Alfred Rheinhold Weiss , Crossed product and hereditary orders	333
Ralph Cohen , Realizing transfer maps for ramified coverings	347
Ronald James Evans , Hermite character sums	357
C. L. Frenzen and Roderick Sue-Chuen Wong , Asymptotic expansions of the Lebesgue constants for Jacobi series	391
Bruno Iochum , Nonassociative L^p -spaces	417
John McDonald , Unimodular approximation in function algebras	435
John Robert Quine, Jr. , Ramification and unintegrated value distribution ...	441
Marc Raphael , Commutants of quasisimilar subnormal operators	449
Parameswaran Sankaran and Peter Zvengrowski , On stable parallelizability of flag manifolds	455
Helga Schirmer , A relative Nielsen number	459
Barry Simon , Schrödinger semigroups on the scale of Sobolev spaces	475
Viakalathur Shankar Sunder , Stochastic integration in Fock space	481
Jan de Vries , A note on the G -space version of Glicksberg's theorem	493