

Pacific Journal of Mathematics

COMMUTANTS OF QUASISIMILAR SUBNORMAL OPERATORS

MARC RAPHAEL

COMMUTANTS OF QUASISIMILAR SUBNORMAL OPERATORS

MARC RAPHAEL

In this paper it is shown if two rationally cyclic subnormal operators are quasisimilar, then they have naturally isomorphic commutants.

1. Introduction. An operator S on a Hilbert space \mathcal{H} is *subnormal* if there is a Hilbert space \mathcal{K} containing \mathcal{H} and a normal operator N on \mathcal{K} such that $N\mathcal{H} \subset \mathcal{H}$ and $S = N|_{\mathcal{H}}$ (the restriction of N to \mathcal{H}). The weak* topology on $\mathcal{B}(\mathcal{H})$ is the topology $\mathcal{B}(\mathcal{H})$ has as the Banach space dual of $\mathcal{B}_1(\mathcal{H})$, the trace class operators [4].

An operator S in $\mathcal{B}(\mathcal{H})$ is *rationally cyclic* if there is an e in \mathcal{H} such that $\{r(S)e: r \in \text{Rat } \sigma(S)\}$ is dense in \mathcal{H} . (Notation: $\text{Rat } \sigma(S)$ is the set of rational functions with poles off $\sigma(S)$, the spectrum of S ; and e is called a *rationally cyclic vector* for S .) The commutant of S is the weak* closed subalgebra of $\mathcal{B}(\mathcal{H})$ defined by: $\{S\}' = \{A \in \mathcal{B}(\mathcal{H}): AS = SA\}$.

A *measure* μ is always a compactly supported, positive, regular Borel measure on the complex plane, \mathbb{C} . If S is a rationally cyclic subnormal operator then there exist a measure μ and a compact set K containing the support of μ such that S is unitarily equivalent to $S(K, \mu)$, the operator of multiplication by z on $R^2(K, \mu)$ = the closure of $\text{Rat } K$ in $L^2(\mu)$ [4]. Yoshino's Theorem [4] states that the map from $R^2(K, \mu) \cap L^\infty(\mu)$ onto $\{S(K, \mu)\}'$ given by $\phi \mapsto \phi(S(K, \mu))$ = multiplication by ϕ is an isometric isomorphism and a weak* homeomorphism. For f in $L^\infty(\mu)$, $\|f\|_\mu$ denotes the μ -essential supremum of f .

If \mathcal{H}_1 and \mathcal{H}_2 are Hilbert spaces, a bounded linear operator $X: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ is said to be *quasi-invertible* if it is injective and has dense range. If $S_j \in \mathcal{B}(\mathcal{H}_j)$ ($j = 1, 2$), then S_1 is *quasisimilar* to S_2 if there are quasi-invertible operators $X_{21}: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ and $X_{12}: \mathcal{H}_2 \rightarrow \mathcal{H}_1$ such that $X_{21}S_1 = S_2X_{21}$ and $X_{12}S_2 = S_1X_{12}$. Unlike similarity, quasisimilar operators need not have equal spectra [5], though their spectra cannot be disjoint [5]. However, quasisimilar subnormal operators must have equal spectra [1, 2, 3], and rationally cyclic ones have the same approximate point spectra [6].

2. The main result. It is proved in [3] that the weak* algebras generated by quasisimilar subnormal operators are isometrically isomorphic and weak* homeomorphic via a natural map. Theorem 1 shows the existence of a similar map between properly larger algebras, but carries the additional hypothesis that the subnormal operators be rationally cyclic. Some restrictive hypothesis on the subnormal operators is necessary since there is an example of an irreducible subnormal operator similar to a reducible one (p. 276, [4]).

THEOREM 1. *If S_1 and S_2 are quasisimilar rationally cyclic subnormal operators, then there exists an isometric isomorphism and weak* homeomorphism $\Lambda: \{S_1\}' \rightarrow \{S_2\}'$ such that $\Lambda(r(S_1)) = r(S_2)$ for all r in $\text{Rat } \sigma(S_1)$.*

Theorem 1 answers an open question posed on page 225 of [4]. Its proof follows from several preliminary results and is postponed until the end of this paper.

The following proposition is essentially the same as Proposition 4.1 of [3]. Its statement and proof for the rationally cyclic case are included here since there will be need for recourse to them later.

PROPOSITION 2 (Conway, [4]). *Suppose S_1 and S_2 are rationally cyclic quasisimilar subnormal operators. Then, for $i = 1, 2$, there exist measures μ_i such that S_i is unitarily equivalent to $S(\sigma(S_i), \mu_i)$, constants c_i , and ϕ in $R^2(\sigma(S_1), \mu_1) \cap L^\infty(\mu_1)$ such that:*

- (a) $\{\phi r: r \in \text{Rat } \sigma(S_1)\}$ is dense in $R^2(\sigma(S_1), \mu_1)$;
- (b) for every r in $\text{Rat } K$,

$$c_2 \int |\phi|^2 |r|^2 d\mu_1 \leq \int |r|^2 d\mu_2 \leq c_1 \int |r|^2 d\mu_1.$$

Proof. Let $Y_{ij}: \mathcal{H}_j \rightarrow \mathcal{H}_i$ be quasi-invertible operators such that $Y_{ij}S_j = S_iY_{ij}$. If e_1 is a rationally cyclic vector for S_1 , then it follows that $e_2 = Y_{21}e_1$ is a rationally cyclic vector for S_2 . Choose measure μ_j and isomorphisms $U_j: \mathcal{H}_j \rightarrow R^2(\sigma(S_j), \mu_j)$ with $U_j e_j = 1$ and $U_j S_j U_j^{-1} = S(\sigma(S_j), \mu_j)$ (p. 146 [4]). Let $X_{ij} = U_i Y_{ij} U_j^{-1}$. So $X_{ij}: R^2(\sigma(S_j), \mu_j) \rightarrow R^2(\sigma(S_i), \mu_i)$ is quasi-invertible. Moreover, it is straightforward to verify that $X_{ij}S(\sigma(S_j), \mu_j) = S(\sigma(S_i), \mu_i)X_{ij}$.

If $r \in \text{Rat } \sigma(S_1)$, then

$$\begin{aligned} X_{21}r &= X_{21}r(S(\sigma(S_1), \mu_1))1 = r(S(\sigma(S_2), \mu_2))X_{21}1 \\ &= r(S(\sigma(S_2), \mu_2))U_2 Y_{21}e_1 = r(S(\sigma(S_2), \mu_2))1 = r. \end{aligned}$$

If $c_1 = \|X_{21}\|^2$, this shows that $\int |r|^2 d\mu_2 \leq c_1 \int |r|^2 d\mu_1$.

To find the constant c_2 , notice that $X_{12}X_{21}$ commutes with $S(\sigma(S_1), \mu_1)$. By Yoshino's Theorem there exists ϕ in $R^2(\sigma(S_1), \mu_1) \cap L^\infty(\mu_1)$ such that $X_{12}X_{21}f = \phi f$ for every f in $R^2(\sigma(S_1), \mu_1)$. Hence for r in $\text{Rat } \sigma(S_2)$, $\phi r = X_{12}X_{21}r = X_{12}r$. Let $c_2 = \|X_{12}\|^{-2}$. \square

The next two results are the keys to establishing Theorem 1.

PROPOSITION 3. *If S_1 and S_2 are similar rationally cyclic subnormal operators, then the conclusion of Theorem 1 is valid.*

Proof. Let $X: \mathcal{H}_1 \rightarrow \mathcal{H}_2$ be an invertible bounded linear operator such that $S_2 = XS_1X^{-1}$. If $A \in \{S_1\}'$ it is easy to see that $XAX^{-1} \in \{S_2\}'$. Define $\Lambda: \{S_1\}' \rightarrow \{S_2\}'$ by $\Lambda A = XAX^{-1}$. It is easy to verify that Λ is onto $\{S_2\}'$ and an algebra isomorphism. By Yoshino's Theorem A and ΛA are subnormal operators. Since A and ΛA are similar, they have the same spectra; hence $\|A\| = \|\Lambda A\|$.

Suppose $\{A_\alpha\} \subset \{S_1\}'$ is a net converging weak* to A in $\{S_1\}'$. Let $T \in \mathcal{B}_1(\mathcal{H}_2)$. Then

$$\begin{aligned} \text{tr}(T\Lambda A_\alpha) &= \text{tr}(TXA_\alpha X^{-1}) = \text{tr}(X^{-1}TXA_\alpha) \\ &\rightarrow \text{tr}(X^{-1}TXA) = \text{tr}(TXAX^{-1}) = \text{tr}(T\Lambda A). \end{aligned}$$

This shows Λ is weak* continuous. Since $\Lambda(r(S_1)) = r(S_2)$ for all r in $\text{Rat } \sigma(S_1)$, the proof is complete. \square

SIMPLIFICATION THEOREM 4. *For $i = 1, 2$ let μ_i be a measure and K be a compact subset of \mathbf{C} such that the support of μ_i is contained in K . Suppose $\phi \in R^2(K, \mu_1) \cap L^\infty(\mu_1)$ and $\{\phi r: r \in \text{Rat } K\}$ is dense in $R^2(K, \mu_1)$.*

If the bounded linear operators $X_{21}^1: R^2(K, \mu_1) \rightarrow R^2(K, \mu_2)$ and $X_{12}^\phi: R^2(K, \mu_2) \rightarrow R^2(K, \mu_1)$ are quasi-invertible and satisfy

- (a) $X_{21}^1 r = r$ for all r in $\text{Rat } K$
- (b) $X_{12}^\phi r = \phi r$ for all r in $\text{Rat } K$,

then $\Lambda: R^2(K, \mu_1) \cap L^\infty(\mu_1) \rightarrow R^2(K, \mu_2) \cap L^\infty(\mu_2)$, defined by

$$\Lambda f = X_{21}^1 f,$$

is an isometric isomorphism and a weak homeomorphism.*

The author does not know if the map in Theorem 1 is uniquely determined by the condition that $r(S_1) \mapsto r(S_2)$ for all r in $\text{Rat } \sigma(S_1)$. Couched in terms of the Simplification Theorem, it is easy to see that the aforementioned map is unique if the following query has an affirmative answer.

If $\Lambda: R^2(K, \mu) \cap L^\infty(\mu) \rightarrow R^2(K, \mu) \cap L^\infty(\mu)$ is an isometric isomorphism and a weak* homeomorphism that is the identity map when restricted to $\text{Rat } K$, then must Λ be the identity map on $R^2(K, \mu) \cap L^\infty(\mu)$?

Two lemmas are now stated and proved. They are used in the proof of the Simplification Theorem.

LEMMA 5. *Assume the hypothesis of Simplification Theorem 4 with the exception that X_{21}^1 need not be bounded on $\text{Rat } K$. Then for all f in $R^2(K, \mu_2) \cap L^\infty(\mu_2)$ and all non-negative integers n ,*

$$(6) \quad X_{12}^\phi f^n = (X_{12}^\phi f)^n \cdot \phi^{1-n} \mu_1 - a.e.$$

Consequently, X_{12}^ϕ maps $R^2(K, \mu_2) \cap L^\infty(\mu_2)$ into $R^2(K, \mu_1) \cap L^\infty(\mu_1)$ and $\|X_{12}^\phi f\|_{\mu_1} \leq \|\phi\|_{\mu_1} \|f\|_{\mu_2}$.

Proof. Equation (6) will be shown by induction on n . If $n = 1$ then (6) certainly holds. Suppose (6) is true for $1 \leq k \leq n$. To show that (6) holds for $k = n + 1$, let $f \in R^2(K, \mu_2) \cap L^\infty(\mu_2)$ and let $\{r_s\}$ and $\{q_m\}$ be sequences in $\text{Rat } K$ such that $r_s \rightarrow f$ in $L^2(\mu_2)$ and $q_m \rightarrow f^n$ in $L^2(\mu_2)$. Then

$$\begin{aligned} X_{12}^\phi f^{n+1} &= \lim_{m \rightarrow \infty} X_{12}^\phi(fq_m) = \lim_{m \rightarrow \infty} \left\{ X_{12}^\phi \left(\lim_{s \rightarrow \infty} r_s q_m \right) \right\} \\ &= \lim_{m \rightarrow \infty} \left\{ q_m \left(\lim_{s \rightarrow \infty} X_{12}^\phi r_s \right) \right\} = \phi^{-1} \left(\lim_{m \rightarrow \infty} X_{12}^\phi q_m \right) (X_{12}^\phi f) \\ &= \phi^{-1} (X_{12}^\phi f^n) (X_{12}^\phi f) = (X_{12}^\phi f)^{n+1} \cdot \phi^{-n}, \end{aligned}$$

where all limits are taken in the appropriate L^2 -space and the induction hypothesis is used to obtain the next to last equality. This substantiates (6).

In order to obtain the norm estimate involving X_{12}^ϕ let $f \in R^2(K, \mu_2) \cap L^\infty(\mu_2)$ such that $\|f\|_{\mu_2} < 1$. Since $f^n \rightarrow 0$ in $L^2(\mu_2)$ and X_{12}^ϕ is bounded, it follows from (6) that

$$\|X_{12}^\phi f^n\|_2^2 = \int |X_{12}^\phi f|^{2n} |\phi|^{2-2n} d\mu_1 \rightarrow 0$$

as $n \rightarrow \infty$. Hence $|X_{12}^\phi f| \leq |\phi|$ μ_1 -a.e.; so $\|X_{12}^\phi f\|_{\mu_1} \leq \|\phi\|_{\mu_1}$. Since $\|f\|_{\mu_2} < 1$, it follows easily that $\|X_{12}^\phi f\|_{\mu_1} \leq \|\phi\|_{\mu_1} \|f\|_{\mu_2}$ for all f in $R^2(K, \mu_2) \cap L^\infty(\mu_2)$. \square

LEMMA 7. Assume the hypothesis and notation of Simplification Theorem 4. Then Λ is an isometry on $R^2(K, \mu_1) \cap L^\infty(\mu_1)$, and $X_{21}^1(fg) = (X_{21}^1 f)(X_{21}^1 g)$ whenever $f \in R^2(K, \mu_1) \cap L^\infty(\mu_1)$ and $g \in R^2(K, \mu_1)$.

Proof. Let $\{g_n\}$ be a sequence in $\text{Rat } K$ such that $g_n \rightarrow g$ in $L^2(\mu_1)$. Then

$$\begin{aligned} X_{21}^1(fg_n) &= X_{21}^1 g_n(S(K, \mu_1))f = g_n(S(K, \mu_2))X_{21}^1 f \\ &= (X_{21}^1 g_n)(X_{21}^1 f). \end{aligned}$$

Letting $n \rightarrow \infty$ shows that $X_{21}^1(fg) = (X_{21}^1 f)(X_{21}^1 g)$.

It follows from Lemma 5 with $\phi = 1$ that $\|X_{21}^1 f\|_{\mu_2} \leq \|f\|_{\mu_1}$ when $f \in R^2(K, \mu_1) \cap L^\infty(\mu_1)$. To show the desired reverse inequality, it may be assumed that $\|X_{21}^1 f\|_{\mu_2} < 1$ and $\|\phi\|_{\mu_1} = 1$.

Let $\{r_s\}$ be a sequence in $\text{Rat } K$ such that $r_s \rightarrow f$ in $L^2(\mu_1)$. Then

$$(X_{12}^\phi X_{21}^1)(f) = \lim_{s \rightarrow \infty} (X_{12}^\phi X_{21}^1)(r_s) = \lim_{s \rightarrow \infty} \phi r_s = \phi f,$$

where all limits are in $L^2(\mu_1)$. So, for $n \geq 1$, it follows easily from (6) that

$$X_{12}^\phi (X_{21}^1 f)^n = \phi f^n \quad \mu_1\text{-a.e.}$$

Since, by the last part of Lemma 5, $\|X_{12}^\phi (X_{21}^1 f)^n\|_{\mu_1} \leq \|(X_{21}^1 f)^n\|_{\mu_2} \rightarrow 0$ as $n \rightarrow \infty$, it must be that $\|f\|_{\mu_1} \leq 1$. This completes the proof of Lemma 7. \square

Proof of Simplification Theorem 4. By Lemma 7 and the Krein-Smulian Theorem, it suffices to show Λ is onto $R^2(K, \mu_2) \cap L^\infty(\mu_2)$ and weak* continuous. By the proof of Proposition 2 it is possible to do the following: choose measures ν_i ($i = 1, 2$) so that if $T_i = S(K, \nu_i)$ and $S_i = S(K, \mu_i)$, then S_i and T_i are unitarily equivalent; and the operators defined by

$$\begin{aligned} Y_{12}^1: R^2(K, \nu_2) &\rightarrow R^2(K, \nu_1) \\ r &\mapsto r, \quad \text{and} \\ Y_{21}^\lambda: R^2(K, \nu_1) &\rightarrow R^2(K, \nu_2) \\ r &\mapsto \lambda r \end{aligned}$$

are quasi-invertible. (Here λ is a fixed member of $R^2(K, \nu_2) \cap L^\infty(\nu_2)$ and $r \in \text{Rat } K$.)

In order to prove that Λ is onto $R^2(K, \mu_2) \cap L^\infty(\mu_2)$ it is shown that $X_{21}^1 Y f = f$ for all f in $R^2(K, \mu_2) \cap L^\infty(\mu_2)$, where $Y = I_1 Y_{12}^1 I_2$ and $I_1: R^2(K, \nu_1) \cap L^\infty(\nu_1) \rightarrow R^2(K, \mu_1) \cap L^\infty(\mu_1)$, $I_2: R^2(K, \mu_2) \cap L^\infty(\mu_2) \rightarrow R^2(K, \nu_2) \cap L^\infty(\nu_2)$ are the "identity" maps in the proof of Proposition 3.

Since S_1 and T_1 are unitarily equivalent operators there exist an isomorphism $V: R^2(K, \nu_1) \rightarrow R^2(K, \mu_1)$ and $\Psi \in R^2(K, \mu_1)$ such that

$Vr = \Psi r$ for all r in $\text{Rat } K$. Given f in $R^2(K, \mu_2) \cap L^\infty(\mu_2)$ let $\{r_s\}$ be a sequence in $\text{Rat } K$ such that

$$(8) \quad \begin{aligned} r_s &\rightarrow f = I_2 f \text{ in } L^2(\nu_2) \quad \text{and} \quad \nu_2\text{-a.e.}, \quad \text{and} \\ r_s &\rightarrow Y_{12}^1 I_2 f \text{ in } L^2(\nu_1) \quad \text{and} \quad \nu_1\text{-a.e.} \end{aligned}$$

Then,

$$\begin{aligned} [X_{21}^1 \Psi] [X_{21}^1 (I_1 Y_{12}^1 I_2 f)] &= X_{21}^1 [\Psi (I_1 Y_{12}^1 I_2 f)] \\ &= \lim_{s \rightarrow \infty} [X_{21}^1 V r_s] = \lim_{s \rightarrow \infty} X_{21}^1 (\Psi r_s) \\ &= \lim_{s \rightarrow \infty} (X_{21}^1 \Psi) (X_{21}^1 r_s) = (X_{21}^1 \Psi) f. \end{aligned}$$

In the above equation, Lemma 7 is used in the first and fourth equality. All limits are in $L^2(\mu_2)$ and are justifiable by (8). Since $X_{21}^1 \Psi \neq 0$ μ_2 -a.e., it follows that $X_{21}^1 Y f = f$.

To finish the proof note that S_1 and $T = S(K, \mu_1 + \mu_2)$ are similar via W : $R^2(K, \mu_1) \rightarrow R^2(K, \mu_1 + \mu_2)$, where $Wr = r$ for all r in $\text{Rat } K$. Since Z : $R^2(K, \mu_1 + \mu_2) \cap L^\infty(\mu_1 + \mu_2) \rightarrow R^2(K, \mu_2) \cap L^\infty(\mu_2)$ defined by $Zf = f$ is weak* continuous, it follows from Proposition 3 that $\Lambda = ZW|R^2(K, \mu_1) \cap L^\infty(\mu_1)$ is also weak* continuous. \square

The proof of Theorem 1 can now be given. By Proposition 2 and the Simplification Theorem, there exist measures μ_i ($i = 1, 2$) such that S_i and $S(\sigma(S_i), \mu_i)$ are unitarily equivalent and $\{S(\sigma(S_i), \mu_i)\}'$ satisfy the conclusion of the Simplification Theorem. An application of Proposition 3 completes the proof. \square

REFERENCES

- [1] W. S. Clary, *Equality of spectra of quasi-similar hyponormal operators*, Proc. Amer. Math. Soc., **53** (1975), 88–90.
- [2] ———, *Quasi-similarity and subnormal operators*, Ph.D. thesis, University of Michigan, 1973.
- [3] J. B. Conway, *On quasisimilarity for subnormal operators*, Illinois J. Math., **24** (1980), 689–702.
- [4] ———, *Subnormal Operators*, Pitman Publishing Inc., Marshfield, Mass., 1981.
- [5] T. B. Hoover, *Quasisimilarity of operators*, Illinois J. Math., **16** (1972), 678–686.
- [6] M. Raphael, *Quasisimilarity and essential spectra for subnormal operators*, Indiana Univ. Math. J., **31**, No. 2 (1982), 243–246.

Received June 1, 1984.

UNIVERSITY OF MISSOURI
ROLLA, MO 65401

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN (Managing Editor)
University of California
Los Angeles, CA 90024

HEBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

CHARLES R. DEPRIMA
California Institute of Technology
Pasadena, CA 91125

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

H. SAMELSON
Stanford University
Stanford, CA 94305

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: Send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1986 by Pacific Journal of Mathematics

Gideon Amit and David Chillag , On a question of Feit concerning character values of finite solvable groups	257
Constantin Gelu Apostol and Frank Larkin Gilfeather , Isomorphisms modulo the compact operators of nest algebras	263
Parviz Azimi and James Neil Hagler , Examples of hereditarily l^1 Banach spaces failing the Schur property	287
Brian Evan Blank , Boundary behavior of limits of discrete series representations of real rank one semisimple groups	299
Jeffrey Carroll , Some undecidability results for lattices in recursion theory	319
Gerald Howard Cliff and Alfred Rheinhold Weiss , Crossed product and hereditary orders	333
Ralph Cohen , Realizing transfer maps for ramified coverings	347
Ronald James Evans , Hermite character sums	357
C. L. Frenzen and Roderick Sue-Chuen Wong , Asymptotic expansions of the Lebesgue constants for Jacobi series	391
Bruno Iochum , Nonassociative L^p -spaces	417
John McDonald , Unimodular approximation in function algebras	435
John Robert Quine, Jr. , Ramification and unintegrated value distribution ...	441
Marc Raphael , Commutants of quasisimilar subnormal operators	449
Parameswaran Sankaran and Peter Zvengrowski , On stable parallelizability of flag manifolds	455
Helga Schirmer , A relative Nielsen number	459
Barry Simon , Schrödinger semigroups on the scale of Sobolev spaces	475
Viakalathur Shankar Sunder , Stochastic integration in Fock space	481
Jan de Vries , A note on the G -space version of Glicksberg's theorem	493