

Pacific Journal of Mathematics

**WEAK*-CLOSED COMPLEMENTED INVARIANT SUBSPACES
OF $L_\infty(G)$ AND AMENABLE LOCALLY COMPACT GROUPS**

ANTHONY TO-MING LAU AND VIKTOR LOSERT

WEAK*-CLOSED COMPLEMENTED INVARIANT SUBSPACES OF $L_\infty(G)$ AND AMENABLE LOCALLY COMPACT GROUPS

ANTHONY TO-MING LAU AND VIKTOR LOSERT

One of the main results of this paper implies that a locally compact group G is amenable if and only if whenever X is a weak*-closed left translation invariant complemented subspace of $L_\infty(G)$, X is the range of a projection on $L_\infty(G)$ commuting with left translations. We also prove that if G is a locally compact group and M is an invariant W^* -subalgebra of the von Neumann algebra $VN(G)$ generated by the left translation operators l_g , $g \in G$, on $L_2(G)$, and $\Sigma(M) = \{g \in G; l_g \in M\}$ is a normal subgroup of G , then M is the range of a projection on $VN(G)$ commuting with the action of the Fourier algebra $A(G)$ on $VN(G)$.

1. Introduction. Let G be a locally compact group and $L_\infty(G)$ be the algebra of essentially bounded measurable complex-valued functions on G with pointwise operations and essential sup norm. Let X be a weak*-closed left translation invariant subspace of $L_\infty(G)$. Then X is *invariantly complemented* in $L_\infty(G)$ if X admits a left translation invariant closed complement, or equivalently, X is the range of a continuous projection on $L_\infty(G)$ commuting with left translations.

H. Rosenthal proved in [13] that if G is an abelian locally compact group and X is a weak*-closed translation invariant complemented subspace of $L_\infty(G)$, then X is invariantly complemented in $L_\infty(G)$. Recently Lau [11, Theorem 3.3] proved that a locally compact group G is left amenable if and only if every left translation invariant weak*-closed subalgebra of $L_\infty(G)$ which is closed under conjugation is invariantly complemented. Note that if T is the circle group, then the Hardy space H_∞ is a weak*-closed translation invariant subalgebra of $L_\infty(T)$ and *not* (invariantly) complemented (see [15] and Corollary 4).

In [20, Lemma 4], Y. Takahashi proved that if G is a compact group, then any weak*-closed complemented left translation invariant subspace of $L_\infty(G)$ is invariantly complemented. However, there is a gap in Takahashi's adaptation of Rosenthal's argument (see Zentralblatt für Mathematik 1982: 483.43002). It should be observed that Rosenthal's original argument in [13, Theorem 1.1] is valid only for locally compact

groups G which is amenable as discrete (for example when G is solvable). Indeed it follows from [21, Theorem 16] that under Martin's Axiom, if P is a bounded projection of $L_\infty(G)$ onto \mathbf{C} (which is a weak*-closed and left translation invariant subspace of $L_\infty(G)$), the functions $x \rightarrow \langle l_{x^{-1}}Pl_x f, h \rangle = \langle Pl_x f, h \rangle$, where $f \in L_\infty(G)$ and $h \in L_1(G)$, is in general bounded but not measurable even when G is compact.

In §3 of this paper, we generalize Rosenthal's result to all amenable locally compact groups (and thus giving a correct proof of Takahashi's Lemma 4 in [20] for all compact groups). More precisely, our Theorem 1 implies that a locally compact group G is amenable if and only if whenever X is a weak*-closed translation invariant complemented subspace of $L_\infty(G)$, X is invariantly complemented. Furthermore (Corollary 4), if G is compact, then X is even the range of a weak*-weak* continuous projection which commutes with left translations. Also in this case, $L_\infty(G)$ has a unique left invariant mean (for example when $G = \text{SO}(n, \mathbf{R})$, $n \geq 5$) if and only if every bounded projection of $L_\infty(G)$ into $L_\infty(G)$ which commutes with left translations is weak*-weak* continuous.

Our proof of Theorem 1 depends heavily on a recent result of Losert and Rindler [12] on the existence of an asymptotically central unit in $L_1(G)$ of an amenable locally compact group.

Finally in §4 we give a non-commutative analogue of Lau's result [11, Theorem 3.3]. We prove that (Theorem 4) if M is an invariant W^* -subalgebra of the von Neuman algebra $\text{VN}(G)$ generated by the left translation operators $\{l_g; g \in G\}$ on $L_2(G)$ of a locally compact group G and $\Sigma(M) = \{g \in G; l_g \in M\}$ is a normal subgroup of G , then M is invariantly complement. However, we do not know if the normality condition on $\Sigma(M)$ may be dropped or not unless $\Sigma(M)$ is compact or open.

2. Preliminaries. If E is a Banach space, then E^* denotes its continuous dual. Also if $\phi \in E^*$ and $x \in E$, then the value of ϕ at x will be written as $\phi(x)$ or $\langle \phi, x \rangle$.

Throughout this paper, G denotes a locally compact group with a fixed left Haar measure. Let $C(G)$ denote the Banach algebra of bounded continuous complex-valued functions on G with the supremum norm, and let $C_0(G)$ be the closed subspace of $C(G)$ consisting of all functions in $C(G)$ which vanish at infinity. The Banach spaces $L_p(G)$, $1 \leq p \leq \infty$, are as defined in [7]. If f is a complex-valued function defined locally almost everywhere on G , and if $a, t \in G$, then $(l_a f)(t) = f(a^{-1}t)$ and $(r_a f)(t) = f(ta)$ whenever this is defined. We say that G is *amenable* if there exists

$m \in L_\infty(G)^*$ such that $m \geq 0$, $\|m\| = 1$ and $m(l_a f) = m(f)$ for which $f \in L_\infty(G)$ and $a \in G$ (m is called a *left invariant mean*). Amenable locally compact groups include all compact groups and all solvable groups. However, the free group on two generators is not amenable (see [4]).

For $g \in G$, the corresponding inner automorphism induces a map τ'_g on $L_\infty(G)$ by $\tau'_g f(x) = f(gxg^{-1})$. The adjoint map τ_g on $L_1(G)$ is given by $\tau_g \phi(x) = \phi(g^{-1}xg)\Delta(g)$, where Δ is the Haar modulus function of G . This can also be written as $\tau_g \phi = \delta_g * \phi * \delta_{g^{-1}}$, where δ_g stands for the Dirac measure concentrated at $g \in G$ (convolution as defined in [7]). A net $\{u_\alpha\}$ in $L_1(G)$ is called an *approximate unit* if $\lim_\alpha \|u_\alpha * \phi - \phi\|_1 = \lim_\alpha \|\phi * u_\alpha - \phi\|_1 = 0$ for all $\phi \in L_1(G)$. The net $\{u_\alpha\}$ is said to be *asymptotically central* if $\lim_\alpha \|u_\alpha\|^{-1} \|\tau_g u_\alpha - u_\alpha\| = 0$ for all $g \in G$. The following result of Losert and Rindler is the key to the proof of one of our main results:

LEMMA 1 ([12, Theorem 3]). *Let G be an amenable locally compact group, then $L_1(G)$ has an asymptotically central approximate unit $\{u_\alpha\}$ with $\|u_\alpha\| \leq 1$.*

3. Subspaces of $L_\infty(G)$. A *left Banach G -module* X is a Banach space X which is left G -module such that

- (i) $\|s \cdot x\| \leq \|x\|$ for all $x \in X, s \in G$.
- (ii) for all $x \in X$, the map $s \rightarrow s \cdot x$ is continuous from G into X .

In this case, we define for each $f \in X^*, s \in G, x \in X$

$$\langle f \cdot s, x \rangle = \langle f, s \cdot x \rangle.$$

Define also $\langle f \cdot \mu, x \rangle = \int \langle f, s \cdot x \rangle d\mu(s)$, $\mu \in M(G), f \in X^*, x \in X$, where $M(G)$ is the space of (complex, bounded) Radon measures on G . Then $f \cdot \mu \in X^*, f \cdot \mu = f \cdot s$ if $\mu = \delta_s$ and $(f \cdot \mu_1) \cdot \mu_2 = f \cdot (\mu_1 * \mu_2)$ for $\mu_1, \mu_2 \in M(G)$.

A subspace $L \subseteq X^*$ is called *G -invariant* if $L \cdot s \subseteq L$ for all $s \in G$.

LEMMA 2. *Let L be a weak*-closed subspace of X^* . Then L is G -invariant if and only if $L \cdot \phi \subseteq L$ for each $\phi \in L_1(G)$.*

Proof. Suppose that L is G -invariant and $\phi \in L_1(G), \phi \geq 0$ and $\|\phi\|_1 = 1$. Define $\Phi(f) = \int f(t)\phi(t) dt, f \in C(G)$. Then Φ is a positive functional on $C(G)$ with norm one. Hence there exists a net $\{m_\alpha\}$ in $C(G)^*$ such that each m_α is a convex combination of point evaluations

and m_α converges to Φ in the weak* topology of $C(G)^*$. If $m_\alpha = \sum_{i=1}^n \lambda_i p_{s_i}$, where $p_s(h) = h(s)$, $h \in C(G)$, $s \in S$, and $f \in L$, then $f \cdot m_\alpha = \sum_{i=1}^n \lambda_i f \cdot s_i$ converges to $f \cdot \phi$ in the weak*-topology of X^* . Hence $f \cdot \phi \in L$.

Conversely, if $L \cdot \phi \subseteq L$ for each $\phi \in L_1(G)$ and $s \in G$, let $m \in L_\infty(G)^*$ such that m extends $p_s \in C(G)^*$ and $\|m\| = \|p_s\| = 1$. Then $m \geq 0$. Hence there exists a net $\{\phi_\alpha\} \subseteq L_1(G)$, $\phi_\alpha \geq 0$, $\|\phi_\alpha\|_1 = 1$, such that $\{\phi_\alpha\}$ converges to m in the weak* topology of $L_\infty(G)^*$. Consequently, if $f \in L$, then $f \cdot \phi_\alpha$ converges in the weak* topology of X^* to $f \cdot s$.

A left Banach G -module X is called *non-degenerate* if the closed linear span of $\{g \cdot x; g \in G, x \in X\}$ is X .

THEOREM 1. *Let G be a locally compact group. Then G is amenable if and only if whenever X is a non-degenerate left Banach G -module and L is a weak*-closed G -invariant subspace of X which is complemented in X , then there exists a projection Q of X^* onto L such that $Q(f \cdot s) = Q(f) \cdot s$ for all $s \in G, f \in X^*$.*

Proof. If G is amenable, there exists an asymptotically central approximate unit $\{u_\alpha\}$ in $L_1(G)$, $\|u_\alpha\| \leq 1$ (Lemma 1). Let m be an invariant mean on $L_\infty(G)$. For each $s \in G, f \in X^*$, put $P_{\alpha,s}(f) = O(f \cdot (u_\alpha * \delta_s)) \cdot (\delta_{s^{-1}} * u_\alpha)$. By Lemma 2, $P_{\alpha,s}: X^* \rightarrow L$ and $\|P_{\alpha,s}\| \leq \|P\|$. For each fixed $\alpha, f \in X^*, x \in X$, the function $s \rightarrow \langle x, P_{\alpha,s}(f) \rangle$ is bounded and continuous. Hence we may define the mean P_α of the family $\{P_{\alpha,s}\}_{s \in G}$ by

$$\langle x, P_\alpha f \rangle = m \{ s \rightarrow \langle x, P_{\alpha,s}(f) \rangle \}.$$

Then $P_\alpha: X^* \rightarrow L$ (since L is weak*-closed and if $x \in X$ is annihilated by L , then $\langle x, P_\alpha f \rangle = 0$ by Lemma 2), and $\|P_\alpha f\| \leq \|P\|$. Finally define $Q(f) = \text{weak*} \lim_\alpha P_\alpha(f)$. Again $Q: X^* \rightarrow L, \|Q\| \leq \|P\|$. For $f \in L, f \cdot (u_\alpha * \delta_s) \in L$. Hence $(P_{\alpha,s})(f) = f \cdot (u_\alpha * u_\alpha)$. Now $\{u_\alpha * u_\alpha\}$ is also an approximate unit in $L_1(G)$. Since X is non-degenerate, Cohen's factorization theorem [8, 32.26] implies that each y in X has the form $\phi \cdot x, x \in X, \phi \in L_1(G)$. Hence

$$\langle f \cdot u_\alpha * u_\alpha - f, y \rangle = \langle f, (u_\alpha * u_\alpha) \cdot (\phi \cdot x) - \phi \cdot x \rangle \rightarrow 0$$

i.e. $P_{\alpha,s}(f) = f$.

Now for each $t \in G$

$$\begin{aligned} P_{\alpha,s}(f \cdot t) - P_{\alpha,ts}(f) \cdot t &= P(f \cdot t \cdot (u_\alpha * \delta_{t^{-1}} * \delta_{ts})) \cdot (\delta_{s^{-1}} * u_\alpha) \\ &\quad - P(f \cdot (u_\alpha * \delta_{ts})) \cdot (\delta_{s^{-1}} * u_\alpha) \\ &\quad + P(f \cdot (u_\alpha * \delta_{ts})) \cdot (\delta_{(ts)^{-1}} * \delta_t * u_\alpha * \delta_{t^{-1}} * \delta_t) \\ &\quad - P(f \cdot (u_\alpha * \delta_{ts})) (\delta_{(ts)^{-1}} * u_\alpha * \delta_t). \end{aligned}$$

Hence

$$\|P_{\alpha,s}(f \cdot t) - P_{\alpha,s}(f) \cdot t\| \leq 2\|P\| \|f\| \|\delta_t * u_\alpha * \delta_{t^{-1}} - u_\alpha\|$$

and this estimate carries over to $\|P_\alpha(f \cdot t) - P_\alpha(f) \cdot t\|$ by invariance of m . Since we assume $\|\delta_t * u_\alpha * \delta_{t^{-1}} - u_\alpha\| \rightarrow 0$, we get $Q(f \cdot t) = Q(f) \cdot t$.

The converse follows as in the proof of Theorem 3.3 in [11] by considering $X = L_1(G)$ and $(s \cdot \phi)(t) = \phi(s^{-1}t)$, $s \in G$, $t \in G$, $\phi \in L_1(G)$. Then if $f \in L_\infty(G)$, $(f \cdot s)(t) = f(st) = (l_{s^{-1}}f)(t)$.

Let Z be a locally compact Hausdorff space. Consider a jointly continuous action $G \times Z \rightarrow Z$. Assume that Z has a quasi-invariant measure ν . For each $s \in G$, define $\chi_s(E) = \nu(s^{-1}E)$. Then $\nu_s \ll \nu$. Hence there is a locally ν -integrable Radon Nikodym derivative $(d\nu_s/d\nu)$ such that $\nu_s = (d\nu_s/d\nu) \cdot \nu$. Also $L_1(Z, \nu)$ is a non-degenerate Banach left G -module (see [5, Lemma 2.3]): $s \cdot \phi = \delta_s * \phi$, $s \in G$, $\phi \in L_1(Z, \nu)$ where $(\delta_s * \phi)(\xi) = (d\nu_s/d\nu)(\xi)(s^{-1}\xi)$ defined ν -a.e. on Z . Hence Theorem 1 implies:

COROLLARY 1. *Let G be a locally compact group. Then G is amenable if and only if for any locally compact Hausdorff space Z and jointly continuous action $G \times Z \rightarrow Z$ such that Z has a quasi-invariant measure, then any weak*-closed G -invariant complemented subspace of $L_\infty(Z, \nu)$ is invariantly complemented.*

REMARK. Theorem 1 also implies Lemma 3.1 of [13] for $L_p(G)$, $1 < p < \infty$, and Theorem 4.1 of [11].

If H is a closed subgroup of a locally compact group, then there exists a non-trivial quasi-invariant measure ν on the coset space $G/H = \{xH; x \in G\}$ which is essentially unique. Write $L_\infty(G/H) = L_\infty(G/H, \nu)$.

COROLLARY 2. *Let G be a locally compact group. Then G is amenable if and only if every weak*-closed complemented invariant subspace of*

$L_\infty(G/H)$, H a closed subgroup of G , is the range of a projection on $L_\infty(G/H)$ which commutes with translation.

COROLLARY 3. *Let G be an amenable locally compact group and X be a weak*-closed left translation invariant subspace of $L_\infty(G)$. Then X is the range of a weak*-weak* continuous projection on $L_\infty(G)$ commuting with left translation if and only if $X \cap C_0(G)$ is weak*-dense in X .*

Proof. This follows from Corollary 2 and Lemma 5.2 of [11].

COROLLARY 4. *Let G be a locally compact group. Then G is compact if and only if G has the following property:*

(*) Whenever X is a weak*-closed complemented left translation invariant subspace of $L_\infty(G)$, there exists a weak*-weak* continuous projection from $L_\infty(G)$ onto X commuting with left translations.

Proof. If G is compact, property (*) follows from Corollary 2, and Lemma 2.1, Lemma 5.2 of [11]. Conversely, if (*) holds, then apply the property to the one-dimensional subspace $X = \mathbb{C}$. It follows that there exists $\phi \in L_1(G)$, $\phi \geq 0$, $\phi(1) = 1$ such that $\phi(l_s f) = \phi(f)$ for all $f \in L_\infty(G)$, $s \in G$. In particular, G is compact.

A bounded linear operator T from $L_\infty(G)$ into $L_\infty(G)$ is said to commute with convolution from the left if $T(\phi * f) = \phi * T(f)$ for all $\phi \in L_1(G)$ and $f \in L_\infty(G)$. In this case, T also commutes with left translations i.e. $T(l_s f) = l_s T(f)$ for all $s \in G$ (see [10, Lemma 2]).

LEMMA 3. *If T is a weak*-weak* continuous linear operator from $L_\infty(G)$ into $L_\infty(G)$ and T commutes with left translations, then T also commutes with convolutions from the left.*

Proof. Let $\phi \in L_1(G)$, $\phi \geq 0$ and $\|\phi\|_1 = 1$. Let $\phi_\alpha = \sum_{i=1}^n \lambda_i \delta_{s_i}$ be a net of convex combinations of point measures on G such that $\int f(t) d\phi_\alpha(t)$ converges to $\int f(t) d\phi(t)$ for each $f \in C(G)$. Hence if $h \in L_\infty(G)$, then the net

$$\langle \phi_\alpha * h, k \rangle = \langle k * \tilde{h}, \phi_\alpha \rangle \rightarrow \langle k * \tilde{h}, \phi \rangle = \langle \phi * h, k \rangle$$

for each $k \in L_1(G)$ ($\tilde{h}(t) = h(t^{-1})$). Consequently,

$$T(\phi * h) = \lim_\alpha T(\phi_\alpha * h) = \lim_\alpha \phi_\alpha * T(h) = \phi * T(h).$$

LEMMA 4 [10]. *If G is compact, then any bounded linear operator T from $L_\infty(G)$ into $L_\infty(G)$ which commutes with convolution from the left is weak*-weak* continuous.*

Proof. This is proved in [10, Theorem 2]¹. We give here a different proof. Indeed if $\phi \in L_1(G)$, then $\phi = \phi_1 * \phi_2$, $\phi_1, \phi_2 \in L_1(G)$ by Cohen's factorization theorem. Hence if $f \in L_\infty(G)$, then

$$\begin{aligned} \langle T^*(\phi), f \rangle &= \langle \phi_1 * \phi_2, T(f) \rangle = \langle \phi_2, \tilde{\phi}_1 * T(f) \rangle \\ &= \langle \phi_2, T(\tilde{\phi}_1 * f) \rangle = \langle T^*(\phi_2), \tilde{\phi}_1 * f \rangle = \langle \phi_1 \odot T^*(\phi_2), f \rangle \end{aligned}$$

i.e. $T^*(\phi) = \alpha_1 \odot T^*(\phi_2)$, where \odot is the Arens product defined on the second conjugate algebra $L_\infty(G)^* = L_1(G)^{**}$. Since G is compact, $L_1(G)$ is an ideal in $L_\infty(G)^*$ (see [6]). Hence $T^*(\phi) \in L_1(G)$, i.e. T is weak*-weak* continuous.

PROPOSITION 1. *Let G be a compact group. The following are equivalent:*

- (a) $L_\infty(G)$ has a unique left invariant mean.
- (b) *If E is a finite dimensional G -invariant subspace of $L_\infty(G)^*$ (i.e. $l_s^*E \subseteq E$ for all $s \in G$) such that the map $s \rightarrow l_s^*\psi$ of G into E is continuous, then $E \subseteq L_1(G)$.*
- (c) *Any bounded (projection) linear operator T from $L_\infty(G)$ into $L_\infty(G)$ which commutes with left translations is weak*-weak* continuous.*
- (d) *Any bounded (projection) linear operator T from $L_\infty(G)$ into $L_\infty(G)$ which commutes with left translation also commutes with convolution from the left.*

Proof. (a) \Rightarrow (b). Consider a continuous representation π of G on E defined by $\pi(s)(m) = l_{s^{-1}}^*m$, $s \in G$, $m \in E$. Since E is finite dimensional, there exists an inner product $\langle \cdot, \cdot \rangle$ on E such that π is unitary. We may further assume that π is irreducible. Let $\{\psi_1, \dots, \psi_n\}$ be an orthonormal basis of E . Write $e_{ij}(s) = \langle \pi(s)\psi_j, \psi_i \rangle$ for the coefficients of π . For $g \in L_\infty(G)$, $\psi \in L_\infty(G)^*$, define $\psi \cdot g \in L_\infty(G)^*$ by $\langle \psi \cdot g, f \rangle = \langle \psi, gf \rangle$, $f \in L_\infty(G)$. Then for any $f, g \in L_\infty(G)$, $\psi \in L_\infty(G)^*$, we have

$$\begin{aligned} \langle f, l_s^*(\psi \cdot g) \rangle &= \langle l_s f, \psi \cdot g \rangle = \langle g \cdot (l_s f), \psi \rangle \\ &= \langle l_s((l_{s^{-1}}g) \cdot f), \psi \rangle = \langle f, (l_s^*\psi) \cdot (l_{s^{-1}}g) \rangle. \end{aligned}$$

¹ The converse to Theorem 2 in [10] was omitted in print. It is stated on page 352.

Consequently $l_s^*(\psi \cdot g) = (l_s^*\psi) \cdot (l_{s^{-1}}g)$. Furthermore, observe that

$$l_s^*\psi_i = \sum_{j=1}^n e_{ji}(s^{-1})\psi_j, \quad l_{s^{-1}}e_{lk} = \sum_{l=1}^n e_{li}(s^{-1})e_{lk}.$$

Since π is unitary, $\sum_i e_{ji}(x)\overline{e_{li}(x)} = \delta_{jl}$. Put $\phi_k = \sum_{i=1}^n \psi_i \cdot \bar{e}_{ik}$ ($\bar{}$ denotes the complex conjugate). Then

$$\begin{aligned} l_s^*\phi_k &= \sum_i (l_s^*\psi_i) \cdot (l_{s^{-1}}(\bar{e}_{ik})) \\ &= \sum_{i,j,l} e_{ji}(s^{-1})\psi_j \overline{e_{li}(s^{-1})} \cdot \bar{e}_{lk} = \sum_j \psi_j \bar{e}_{jk} = \phi_k \end{aligned}$$

for all $s \in G$. By assumption, $\phi_k \in L_1(G)$. Finally

$$\sum_k \phi_k \cdot e_{lk} = \sum_i \psi_i \left(\sum_k \bar{e}_{ik} \cdot e_{lk} \right) = \psi_l$$

and $\phi \cdot f \in L_1(G)$ whenever $\phi \in L_1(G)$, $f \in L_\infty(G)$. Hence $\psi_l \in L_1(G)$ for all $l = 1, 2, \dots, n$.

(b) \Rightarrow (c). Since G is compact, it follows that $T(\phi * f) = \phi * T(f)$ for all $\phi \in L_1(G)$, $f \in C(G)$. If $\phi \in L_1(G)$ such that $\{l_s^*\phi; s \in G\}$ belongs to a finite-dimensional G -invariant subspace of $L_\infty(G)^*$, then the same is true for $T^*\phi$. Hence $T^*\phi \in L_1(G)$ by (b). Since elements of this type are dense in $L_1(G)$, $T^*(L_1(G)) \subseteq L_1(G)$ i.e. T is weak*-weak* continuous.

That (c) \Rightarrow (d) follows from Lemma 3.

(d) \Rightarrow (a). If $L_\infty(G)$ has more than one left invariant mean, then there exists a left invariant mean m such that $m \notin L_1(G)$. Now define $T(f) = m(f) \cdot 1$, $f \in L_\infty(G)$. Then T is a projection of $L_\infty(G)$ into $L_\infty(G)$ commuting with left translations. But T does not commute with convolution by Lemma 4.

REMARK. As known (see [3], [15] and [16]) if G is a nondiscrete compact abelian group (or more generally, G is amenable as discrete), then $L_\infty(G)$ has more than one left invariant mean. However, if $n \geq 5$, and $G = \text{SO}(n, \mathbf{R})$, then $L_\infty(G)$ has a unique left invariant mean (see [14] and [17] for more details).

4. Subspaces of $\text{VN}(G)$. Let $P(G)$ be the continuous positive definite functions on G (see [6]). If H is a closed subgroup of G , let

$$P_H = \{ \phi \in P(G); \phi(g) = 1 \text{ for all } g \in H \}$$

Then P_H is a subsemigroup of $P(G)$.

LEMMA 5. *If H is a closed normal subgroup of G , $g \notin H$, there exists $\phi \in P_H$ such that $\phi(g) = 0$.*

Proof. Consider the quotient group G/H and let $\psi \in P(G/H)$ such that $\psi(gH) = 0$ and $\psi(H) = 1$. Define $\phi = \psi \circ \pi$, where π is the canonical mapping of G onto G/H . Then $\phi \in P(G)$, $\phi(h) = 1$, for all $h \in H$ and $\phi(g) = 0$ (see [2, p. 199]).

Let $VN(G)$ denote the von Neumann algebra generated by the left translation operators l_g , $g \in G$, on $L_2(G)$. Then the predual of $VN(G)$ may be identified with $A(G)$, a subalgebra of $C_0(G)$ with pointwise multiplication, consisting of all functions ϕ of the form $\phi(g) = \int h(g^{-1}t)\overline{k(t)}dt$, $h, k \in L_2(G)$. Furthermore, $A(G)$ with the predual norm is a semi-simple commutative Banach algebra and a closed two sided ideal of $B(G)$, the linear span of $P(G)$. There is a natural action of $A(G)$ on $VN(G)$ defined by $\langle \phi \cdot x, \psi \rangle = \langle x, \phi\psi \rangle$, $x \in VN(G)$. When G is commutative, then $A(G)$ and $VN(G)$ are isometrically isomorphic to $L_1(\hat{G})$ and $L_\infty(\hat{G})$ respectively (where \hat{G} is the dual group of G) and the action of $A(G)$ on $VN(G)$ corresponds to convolution of functions in $L_1(\hat{G})$ and $L_\infty(\hat{G})$. (see [2] for more details.)

A subspace M of $VN(G)$ is called *invariant* if $\phi \cdot x \in M$ for all $\phi \in A(G)$, $x \in M$. Define

$$\Sigma(M) = \{g \in G; l_g \in M\}.$$

If M is an invariant W^* -subalgebra of $VN(G)$, then $\Sigma(M) = H$ is a non-empty closed subgroup of G and $M = N_H$, the ultraweak closure of the linear span of $\{l_g; g \in H\}$ in $VN(G)$ (see [18, Theorems 6 and 8]).

LEMMA 6. *Let M be an invariant W^* -subalgebra of $VN(G)$ such that $\Sigma(M) = H$ is a normal subgroup of G . Then $M = \{x \in VN(G); \phi \cdot x = x \text{ for all } \phi \in P_H\}$.*

Proof. Let $N = \{x \in VN(G); \phi \cdot x = x \text{ for all } \phi \in P_H\}$. Then N is weak*-closed, invariant and $N \supseteq N_H = M$ (since $\phi \cdot l_g = \phi(g)l_g = l_g$ for $\phi \in P_H$, $g \in H$). Now if $g \in G$ and $l_g \in N$, then $\phi(g) = 1$ for all $\phi \in P_H$. In particular $\Sigma(M) \subseteq H$ by Lemma 4. Hence if $x \in N$, then $\text{supp}(x) \subseteq \Sigma(N) \subseteq H$ by Proposition 4.4 [2]. Consequently, $x \in N_H$ by Theorem 3 [19].

The following implies one direction of Theorem 3.3 [11] when G is abelian:

THEOREM 2. *Let M be an invariant W^* -subalgebra of $VN(G)$ such that $\Sigma(M) = H$ is a normal subgroup of G . Then there exists a continuous projection P of $VN(G)$ onto M such that $P(\phi \cdot x) = \phi \cdot P(x)$ for all*

$\phi \in A(G)$ and $x \in \text{VN}(G)$. In particular, M admits a closed complement which is also invariant.

Proof. By Lemma 6, $M = \{x \in \text{VN}(G); \phi \cdot x = x \text{ for all } \phi \in P_H\}$. For each $x \in \text{VN}(G)$, let K_x denote the weak*-closed convex hull of $\{\phi \cdot x; \phi \in P_1(G)\}$, where $P_1(G) = \{\phi \in P(G); \phi(e) = 1\}$, and $\langle \phi \cdot x, \psi \rangle = \langle x, \phi\psi \rangle$, $\psi \in A(G)$. Then K_x is a weak*-closed subset of $\text{VN}(G)$. For each $\psi \in P_H$, let $T_\psi: K_x \rightarrow K_x$ be defined by $T_\psi(y) = \psi \cdot y$, $y \in K_x$. Then T_ψ is weak*-weak* continuous and affine. Since P_H is a commutative semigroup, an application of the Markov-Kakutani fixed point theorem ([1, p. 456]) shows that $M \cap K_x$ is nonempty for each $x \in \text{VN}(G)$. By Theorem 2.1 in [9], there exists a projection P from $\text{VN}(G)$ onto M and P commutes with any weak*-weak* continuous operator from M into M which commutes with $\{T_\psi; \psi \in P_H\}$. Hence $P(\phi \cdot x) = \phi \cdot P(x)$ for each $\phi \in A(G)$, $x \in \text{VN}(G)$.

REMARK. Lemma 5 (hence Lemma 6 and Theorem 2) holds for any compact subgroup (see Eymard [2, Lemma 3.2]) and any open subgroup H of G (see Hewitt and Ross [8, 32.43]) without normality.

REFERENCES

- [1] N. Dunford and J. T. Schwartz, *Linear Operators 1*, Interscience Publishers, Inc., New York, 1957.
- [2] P. Eymard, *L'algèbre de Fourier d'un groupe localement compact*, Bull. Soc. Math. France, **92** (1964), 181–236.
- [3] E. Granirer, *Criteria for compactness and for discreteness of locally compact amenable groups*, Proc. Amer. Math. Soc., **40** (1973), 615–624.
- [4] F. P. Greenleaf, *Invariant Means on Topological Groups and Their Applications*, Van Nostrand (1969).
- [5] ———, *Amenable actions of locally compact groups*, J. Funct. Anal., **4** (1969), 295–315.
- [6] M. Grosser, *$L_1(G)$ as an ideal in its second dual space*, Proc. Amer. Math. Soc., **73** (1979), 363–364.
- [7] E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis I*, Springer-Verlag, 1963.
- [8] ———, *Abstract Harmonic Analysis II*, Springer-Verlag, 1970.
- [9] A. T. Lau, *Semigroup of operators on dual Banach spaces*, Proc. Amer. Math. Soc., **54** (1976), 393–396.
- [10] ———, *Operators which commute with convolutions on subspaces of $L_\infty(G)$* , Colloquium Mathematicum, **39** (1978), 351–359.
- [11] ———, *Invariantly complemented subspaces of $L_\infty(G)$ and amenable locally compact groups*, Illinois J. Math., **26** (1982), 226–235.
- [12] V. Losert and H. Rindler, *Asymptotically Central Functions and Invariant Extensions of Dirac Measure*, Probability Measures on Groups VII, Proceedings Oberwolfach 1983, Springer Lecture Notes Math. **1069** (1984), 368–378.

- [13] H. P. Rosenthal, *Projections onto translation invariant subspace of $L^p(G)$* , *Memoirs Amer. Math. Soc.*, **63** (1966).
- [14] J. Rosenblatt, *Uniqueness of invariant means for measure-preserving transformations*, *Trans. Amer. Math. Soc.*, **265** (1981), 623–636.
- [15] W. Rudin, *Invariant means on L^∞* , *Studia Mathematica*, **44** (1972), 219–227.
- [16] J. D. Stafney, *Arens multiplication and convolution*, *Pacific J. Math.*, **14** (1964), 1423–1447.
- [17] D. Sullivan, *For $n > 3$ there is only one finitely additive rotationally invariant measure on the n -sphere defined on all Lebesgue measurable subsets*, *Bull. Amer. Math. Soc.*, **4** (1981), 121–123.
- [18] M. Takesaki and N. Tatsuuma, *Duality and subgroups*, *Ann. of Math.*, **93** (1971), 344–364.
- [19] ———, *Duality and subgroups II*, *J. Funct. Anal.*, **11** (1972), 184–190.
- [20] Y. Takahashi, *A characterization of certain weak*-closed subalgebras of $L_\infty(G)$* , *Hokkaido Math. J.*, **11** (1982), 116–124.
- [21] M. Talgrand, *Closed convex hull of set of measurable functions, Riemann-measurable functions and measurability of translations*, *Ann. Inst. Fourier, Grenoble*, **32** (1982), 39–69.

Received January 10, 1985. The first author is supported by NSERC Grant A-7679.

UNIVERSITY OF ALBERTA
EDMONTON, ALBERTA
CANADA T6G-2G1

AND

UNIVERSITÄT WIEN
STRUDLHOFGASSE 4
A-1090 WIEN
AUSTRIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720
H. SAMELSON
Stanford University
Stanford, CA 94305
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Maria Emilia Alonso García , <i>A note on orderings on algebraic varieties</i>	1
F. S. De Blasi and Józef Myjak , <i>On continuous approximations for multifunctions</i>	9
Frank Albert Farris , <i>An intrinsic construction of Fefferman's CR metric</i>	33
Antonio Giambruno, P. Misso and Francisco César Polcino Milies , <i>Derivations with invertible values in rings with involution</i>	47
Dan Haran and Moshe Jarden , <i>The absolute Galois group of a pseudo real closed algebraic field</i>	55
Telemachos E. Hatziafratis , <i>Integral representation formulas on analytic varieties</i>	71
Douglas Austin Hensley , <i>Dirichlet's theorem for the ring of polynomials over GF(2)</i>	93
Sofia Kalpazidou , <i>On a problem of Gauss-Kuzmin type for continued fraction with odd partial quotients</i>	103
Harvey Bayard Keynes and Mahesh Nerurkar , <i>Ergodicity in affine skew-product toral extensions</i>	115
Thomas Landes , <i>Normal structure and the sum-property</i>	127
Anthony To-Ming Lau and Viktor Losert , <i>Weak*-closed complemented invariant subspaces of $L_\infty(G)$ and amenable locally compact groups</i>	149
Andrew Lelek , <i>Continua of constant distances in span theory</i>	161
Dominikus Noll , <i>Sums and products of B_r spaces</i>	173
Lucimar Nova , <i>Fixed point theorems for some discontinuous operators</i>	189
A. A. S. Perera and Donald Rayl Wilken , <i>On extreme points and support points of the family of starlike functions of order α</i>	197
Massimo A. Picardello , <i>Positive definite functions and L^p convolution operators on amalgams</i>	209
Friedrich Roesler , <i>Squarefree integers in nonlinear sequences</i>	223
Theodore Shifrin , <i>The osculatory behavior of surfaces in \mathbf{P}^5</i>	227