

Pacific Journal of Mathematics

FIXED POINT THEOREMS FOR SOME DISCONTINUOUS OPERATORS

LUCIMAR NOVA

FIXED POINT THEOREMS FOR SOME DISCONTINUOUS OPERATORS

LUCIMAR NOVA G.

The purpose of this paper is to show the existence of fixed points for operators T defined on a subset K of a Banach space X and belonging to a class that the author calls $D(a, b)$ with $0 \leq a, b \leq 1$.

1. Introduction. Let T be a mapping of a set K into itself. An immediate question is whether some point is mapped onto itself; that is, does the equation

$$(1) \quad Tx = x$$

have a solution? If so, x is called a *fixed point* of T . This question generates a theory which began in 1912 with the work of L. E. J. Brouwer, who proved that any continuous mapping T of an n -ball into itself has a fixed point, and was followed in 1922 by S. Banach's Contraction Principle, which states that any mapping T of a complete metric space X into itself that satisfies, for some $0 < k < 1$, the inequality

$$(2) \quad d(Tx, Ty) \leq kd(x, y)$$

for all x and y in X , has a unique fixed point. Here d denotes the metric on X . J. Schauder [13], Tychonoff [16], S. Lefschetz [10], F. Browder [2], W. A. Kirk [7], and many others have added to and generalized these basic results.

In 1969 and 1971, R. Kannan [5], [6], proved some fixed point theorems for operators T mapping a Banach space X into itself which, instead of the contraction property in (2), satisfy the condition:

$$(3) \quad \|Tx - Ty\| \leq \alpha[\|x - Tx\| + \|y - Ty\|],$$

for all x, y in X ; where $0 < \alpha < 1/2$. G. Hardy and T. Rogers [4] generalized this result to continuous mappings T of a complete metric space X into itself that satisfy:

$$(4) \quad d(Tx, Ty) \leq a_1d(x, y) + a_2d(x, Tx) + a_3d(y, Ty) \\ + a_4d(x, Ty) + a_5d(y, Tx),$$

for all x and y in X , where $a_i \geq 0$ and $a_1 + a_2 + a_3 + a_4 + a_5 < 1$. K. Goebel, W. A. Kirk, and T. N. Shimi [3], extended the last result to

continuous mappings of a nonempty bounded, closed and convex subset K of a uniformly convex Banach space into itself, where $a_i \geq 0$ and $a_1 + a_2 + a_3 + a_4 + a_5 \leq 1$.

In this article we will prove some fixed point theorems for operators T defined on a subset K of a Banach space X that satisfy the inequality.

$$(5) \quad \|Tx - Ty\| \leq a\|x - y\| + b[\|x - Tx\| + \|y - Ty\|]$$

for all x and y in K , where $0 \leq a, b \leq 1$. Any operator T satisfying condition (5) will be said to belong to class $D(a, b)$. A contraction operator is in class $D(k, 0)$ with $0 < k < 1$.

Note that although condition (2) implies the continuity of the operator T , condition (3) to (5) may hold even if the operator is not continuous. Indeed, any operator T is in $D(1, 1)$, since by the triangle inequality:

$$\|Tx - Ty\| \leq \|Tx - x\| + \|x - y\| + \|y - Ty\|.$$

Furthermore, inequality (5) is a direct consequence of (4) and the triangle inequality, provided we forego the upper bounds required in [3].

2. Discussion. If we carefully examine the statement and proof of Banach's Contraction Principle (see for example Kreyszig [9, pp. 300–302]) we observe that the main conclusions are

- (i) There exists a unique fixed point.
- (ii) A contraction mapping is an asymptotically regular operator for any point, that is, $\|T^{n-1}x - T^n x\| \rightarrow 0$ as $n \rightarrow \infty$.
- (iii) The sequence $x_n = T^n x_0$, of Picard iterates converges to the unique fixed point.

Which of these conclusions hold for operators T in the class $D(a, b)$ with $0 \leq a, b < 1$?

First, observe that these classes are not empty: consider the discontinuous operator

$$Tx = \begin{cases} \gamma x, & 0 \leq x < 1/2, \\ \rho x, & 1/2 \leq x \leq 1, \end{cases}$$

with $0 < \gamma, \rho < 1$, $\gamma \neq \rho$. Then T is in $D(0, \mu/(1 - \mu))$ where $\mu = \max\{\gamma, \rho\}$ because

$$\frac{\gamma}{1 - \gamma}(x_i - Tx_i) = \gamma x_i, \quad \text{for } x_i \in [0, 1/2),$$

so that

$$\begin{aligned} |Tx_1 - Tx_2| &\leq \gamma(x_1 + x_2) = \frac{\gamma}{1 - \gamma} \{|x_1 - Tx_1| + |x_2 - Tx_2|\} \\ &\leq \frac{\mu}{1 - \mu} \{|x_1 - Tx_1| + |x_2 - Tx_2|\}. \end{aligned}$$

The same inequality holds if $x_i \in [1/2, 1]$. Now if $x_1 < 1/2 \leq x_2$; then

$$\frac{\gamma}{1-\gamma}(x_1 - Tx_1) = \gamma x_1; \quad \frac{\rho}{1-\rho}(x_2 - Tx_2) = \rho x_2$$

and

$$|Tx_1 - Tx_2| \leq \gamma x_1 + \rho x_2 \leq \frac{\mu}{1-\mu} \{ |x_1 - Tx_1| + |x_2 - Tx_2| \}.$$

Moreover, (i) as is obvious from its graph, T has a unique fixed point at $x = 0$,

$$(ii) \quad \frac{\gamma}{1-\gamma}(T^n x - T^{n+1} x) = \gamma T^n x \leq \mu^{n+1} x$$

for n sufficiently large, arbitrary x in $[0, 1]$, and $0 < \mu < 1$, so that T is asymptotically regular at any point, and (iii) $\{x_n - T^n x\}_n$ converges to 0.

How many fixed points can an operator in the class $D(a, b)$, $0 \leq a, b < 1$, have? We shall show that the behaviour of the classes $D(a, b)$ is identical whether or not $b = 0$.

LEMMA 1. *Let T be in the class $D(a, b)$, $a, b \geq 0$, $a < 1$. If $F_T = \{x \in K | Tx = x\}$ is not empty, then F_T consists of a single point.*

Proof. Assume that $x_i, i = 1, 2$, are fixed point of T , and T satisfies $D(a, b)$. Then

$$\|x_1 - x_2\| = \|Tx_1 - Tx_2\| \leq a\|x_1 - x_2\|$$

which only holds if $x_1 = x_2$. □

When does T have a fixed point? Before answering this question we will need the following three interesting facts.

LEMMA 2. *If $T \in D(a, b)$, $a + 2b < 1$, then $\text{Inf}_{x \in K} \|x - Tx\| = 0$.*

Proof. Define $x_n = T^n x_0$, with x_0 an arbitrary point. Then

$$\begin{aligned} \|x_n - x_{n+1}\| &= \|Tx_{n-1} - Tx_n\| \\ &\leq a\|x_{n-1} - x_n\| + b[\|x_{n-1} - x_n\| + \|Tx_{n-1} - Tx_n\|] \end{aligned}$$

so that

$$(1 - b)\|(I - T)x_n\| \leq (a + b)\|(I - T)x_{n-1}\|.$$

Hence

$$\|(I - T)x_n\| \leq \left(\frac{a + b}{1 - b}\right)^n \|(I - T)x_0\|$$

and as $(a + b)/(1 - b) < 1$ it follows that $\|x_n - Tx_n\| \rightarrow 0$ as $n \rightarrow \infty$; that is $\text{Inf}_{x \in K} \|x - Tx\| = 0$. \square

Let X be a Banach space, T be a mapping of X into itself, and x be a point in X . The mapping T is *asymptotically regular in x* if $\|T^{n+1}x - T^n x\| \rightarrow 0$ as $n \rightarrow \infty$.

Because of the definition of x_n in the proof of Lemma 2 we observe that $T^n x_0 - T^{n+1} x_0 = x_n - x_{n+1} = (I - T)x_n \rightarrow 0$ as $n \rightarrow \infty$, so that we have

LEMMA 3. *If $T \in D(a, b)$, $a + 2b < 1$, then T is asymptotically regular at any point.*

From Banach's Contraction Principle, we know that for T in $D(a, 0)$, the Picard iterates converge to the unique fixed point. Although we have not yet proved the existence of a fixed point, is there any property that a sequence of points $\{x_n\}$ in K could satisfy that would imply the existence of a fixed point and the convergence of that sequence to that point? The following result answers this question.

THEOREM 1. *Let K be a closed subset of a Banach space X and let $T \in D(a, b)$ with $0 \leq a, b < 1$. Then the sequence $\{x_n\}_n$ contained in the set K , satisfies*

$$(6) \quad \lim_{n \rightarrow \infty} (x_n - Tx_n) = 0,$$

if and only if the sequence converges to the unique fixed point of T .

Proof. The condition is necessary because

$$\|Tx_n - Tx_m\| \leq a\|x_n - x_m\| + B\{\|x_n - Tx_n\| + \|x_m - Tx_m\|\}$$

and applying the triangle inequality we have

$$\|Tx_n - Tx_m\| \leq \frac{a + b}{1 - a} \{\|x_n - Tx_n\| + \|x_m - Tx_m\|\}.$$

Thus, it follows, from the hypothesis that $\{Tx_n\}_n$ is a Cauchy sequence. Since X is complete and K is closed there exists $z \in K$ such that

$$\lim_{n \rightarrow \infty} Tx_n = z$$

and since $x_n - Tx_n \rightarrow 0$ as $n \rightarrow \infty$, then $x_n \rightarrow z$ as $n \rightarrow \infty$. Using the triangle inequality and the fact that $T \in D(a, b)$ with $b < 1$, we have:

$$\|z - Tz\| \leq \frac{1 + a}{1 - b} \|z - x_n\| + \frac{1 + b}{1 - b} \|x_n - Tx_n\|$$

and since $x_n \rightarrow z$, and $x_n - Tx_n \rightarrow 0$ as $n \rightarrow \infty$ it follows that z is a fixed point under T . By Lemma 1 it is unique.

For the sufficiency part of the theorem we assume that there exists $z \in K$ such that $z = Tz$ and

$$\lim_{n \rightarrow \infty} x_n = z.$$

Since $T \in D(a, b)$ with $b < 1$, using the triangle inequality, we have:

$$\|Tx_n - x_n\| - \|x_n - z\| \leq \|Tx_n - z\| \leq a\|x_n - z\| + b\|x_n - Tx_n\|.$$

Thus,

$$(1 - b)\|Tx_n - x_n\| \leq (1 + a)\|x_n - z\|,$$

by hypothesis, $Tx_n - x_n \rightarrow 0$ as $n \rightarrow \infty$. □

We have already seen that the Picard iterates of any point x in K satisfy equation (6). Hence we have proved

THEOREM 2. *Let K be a closed subset of a Banach space X , let $T \in D(a, b)$ with $a + 2b < 1$. Then T has a unique fixed point z in K .*

Moreover, the Picard iterates of any point x in K converge to z .

When $a + 2b < 1$ we can estimate the rate of convergence of the Picard iterates:

$$\begin{aligned} \|Tx - z\| &= \|Tx - Tz\| \leq a\|x - z\| + b\|x - Tx\| \\ &\leq (a + b)\|x - z\| + b\|z - Tx\| \end{aligned}$$

or

$$(7) \quad \|Tx - z\| \leq \left(\frac{a + b}{1 - b}\right)\|x - z\|.$$

Hence

$$\|T^n x - z\| \leq \left(\frac{a + b}{1 - b}\right)^n \|x - z\|$$

and $a + b < 1 - b$.

Here is an example of an operator T in $D(0, 1)$ which does not have a fixed point:

Consider the function

$$Tx = \begin{cases} x/4 + 19/50, & \text{if } 0 \leq x < 1/2, \\ x/5 + 19/50, & \text{if } 1/2 \leq x \leq 1. \end{cases}$$

It is enough to see the case $x \in [0, 1/2)$, $y \in [1/2, 1]$ and to compare

$$|Tx - Ty| = \frac{|5x - 4y|}{20}$$

with

$$|x - Tx| + |y - Ty| = \frac{16y - 15x}{20}.$$

Analyzing the cases $5x - 4y \leq 0$ or $5x - 4y > 0$, we observe that $T \in D(0, 1)$ but T does not have a fixed point in $[0, 1]$.

The fixed point of the following operator T solves a differential equation which is not covered by the usual Picard Theorem, although the solution is found by the same iterative process.

Let

$$Tx(t) = \begin{cases} \gamma x(t) + \int_0^t k(s, t, x(s)) ds, & 0 \leq x(t) \leq A, \\ \rho x(t) + \int_0^t k(s, t, x(s)) ds, & x(t) > A, \end{cases}$$

where $k(s, t, x(s)) = ce^{-a(t-s)}x(s)$, $a, c > 0$, and $1 > \gamma > \rho > 0$. Let $0 \leq x(t) \leq A < y(t)$ such that $2x(s) \leq (1 + (1 - \gamma)/(1 - \rho))y(s)$ for all $0 \leq s \leq t$. Then

$$x(t) - Tx(t) = (1 - \gamma)x(t) - \int_0^t k(s, t, x(s)) ds$$

so that

$$\begin{aligned} |Tx(t) - Ty(t)| = & \left| \frac{\gamma}{1 - \gamma} \left[(x(t) - Tx(t)) + \frac{1}{\gamma} \int_0^t k(s, t, x(s)) ds \right] \right. \\ & \left. - \frac{\rho}{1 - \rho} \left[(y(t) - Ty(t)) + \frac{1}{\rho} \int_0^t k(s, t, y(s)) ds \right] \right| \end{aligned}$$

and

$$\begin{aligned} (9) \quad & |Tx(t) - Ty(t)| \\ & \leq \frac{\gamma}{1 - \gamma} \|x - Tx\| + \frac{\rho}{1 - \rho} \|y - Ty\| \\ & \quad + \left| \frac{1}{1 - \gamma} \int_0^t k(s, t, x(s)) ds - \frac{1}{1 - \rho} \int_0^t k(s, t, y(s)) ds \right| \\ & \leq \frac{\gamma}{1 - \gamma} [\|x - Tx\| + \|y - Ty\|] \\ & \quad + \frac{1}{1 - \gamma} \left| \int_0^t ce^{-a(t-s)} ds \right| \|x - y\| \\ & \leq \frac{c}{a(1 - \gamma)} \|x - y\| + \frac{\gamma}{1 - \gamma} [\|x - Tx\| + \|y - Ty\|]. \end{aligned}$$

Inequality (9) also holds for any other positive choices of $x(t)$ and $y(t)$.

Thus

$$T \in D\left(\frac{c}{a(1-\gamma)}, \frac{\gamma}{1-\gamma}\right),$$

and if $0 < \gamma < \min(1/2, 1/3(1 - c/a))$, the operator T will have a unique fixed point and any Picard iterates will converge to that fixed point.

Acknowledgments. The author would like to express her deep gratitude to Dr. William Derrick for suggesting the problems on which this paper is based and for many helpful conversations during its preparation, and to the referee for suggestions that lead to a better presentation of this paper.

REFERENCES

- [1] F. E. Browder and W. V. Petryshyn, *The solution by iteration of non-linear functional equations in Banach spaces*, Bull. Amer. Math. Soc., Number 3 (1966), 571–575.
- [2] F. E. Browder, *On a generalization of the Schauder fixed point theorem*, Duke Math. J., **26** (1959), 291–303.
- [3] K. Goebel, W. A. Kirk, T. N. Shimi, *A fixed point theorem in uniformly convex spaces*, Boll. Un. Mat. Ital., (4) **7** (1973), 63–75.
- [4] G. Hardy and T. Rogers, *A generalization of a fixed point theorem of Reich*, Canad. Math. Bull., **16** (1973), 201–206.
- [5] R. Kannan, *Some results on fixed points II*, Amer. Math. Monthly, **76** (1969), 405–408.
- [6] ———, *Some results on fixed points III*, Fund. Math., **70** (1971), 169–177.
- [7] W. A. Kirk, *A fixed point for mappings which do not increase distance*, Amer. Math. Monthly, **72** (1965), 1004–1006.
- [8] ———, *On successive approximations for non-expansive mappings in Banach spaces*, Glasgow Math. J., **12** (1971), 6–9.
- [9] E. Kreyszig, *Introductory Functional Analysis with Applications*, John Wiley & Sons, 1978.
- [10] S. Lefschetz, *Algebraic Topology*, Amer. Math. Soc., New York, 1942.
- [11] W. V. Petryshyn and T. E. Williamson, Jr. *A necessary and sufficient condition for the convergence of a sequence of iterates for quasi-non-expansive mappings*, Bull. Amer. Math. Soc., **78**, #6, (1972), 1027–1031.
- [12] B. E. Rhoades, *Some fixed point theorems for generalized Non-expansive mappings*, Lecture Notes in pure and applied Mathematics Non-Linear analysis and applications, **80** (1982), 223–228.
- [13] J. Schauder, *Der Fixpunktstz in Funktionalraumen*, Studia Math., **2** (1930), 171–180.
- [14] T. N. Shimi, *Approximation of fixed points of Certain Non-Linear mappings*, J. Math. Anal. Appl., **65** (1978), 565–571.

- [15] P. Soardi, *Su un problema di punto Unito di S. Reich*, Boll. Un. Mat. Ital. (4), **4** (1971), 841–845.
- [16] A. Tychonoff, *Ein Fixpunktsatz*, Math. Ann., **111** (1935), 767–776.

Received February 22, 1985. Research supported by COLCIENCIAS, grant #97024-1-02-83.

UNIVERSIDAD NACIONAL DE COLOMBIA
BOGOTA, COLOMBIA, S.A.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720
H. SAMELSON
Stanford University
Stanford, CA 94305
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Maria Emilia Alonso García , <i>A note on orderings on algebraic varieties</i>	1
F. S. De Blasi and Józef Myjak , <i>On continuous approximations for multifunctions</i>	9
Frank Albert Farris , <i>An intrinsic construction of Fefferman's CR metric</i>	33
Antonio Giambruno, P. Misso and Francisco César Polcino Milies , <i>Derivations with invertible values in rings with involution</i>	47
Dan Haran and Moshe Jarden , <i>The absolute Galois group of a pseudo real closed algebraic field</i>	55
Telemachos E. Hatziafratis , <i>Integral representation formulas on analytic varieties</i>	71
Douglas Austin Hensley , <i>Dirichlet's theorem for the ring of polynomials over GF(2)</i>	93
Sofia Kalpazidou , <i>On a problem of Gauss-Kuzmin type for continued fraction with odd partial quotients</i>	103
Harvey Bayard Keynes and Mahesh Nerurkar , <i>Ergodicity in affine skew-product toral extensions</i>	115
Thomas Landes , <i>Normal structure and the sum-property</i>	127
Anthony To-Ming Lau and Viktor Losert , <i>Weak*-closed complemented invariant subspaces of $L_\infty(G)$ and amenable locally compact groups</i>	149
Andrew Lelek , <i>Continua of constant distances in span theory</i>	161
Dominikus Noll , <i>Sums and products of B_r spaces</i>	173
Lucimar Nova , <i>Fixed point theorems for some discontinuous operators</i>	189
A. A. S. Perera and Donald Rayl Wilken , <i>On extreme points and support points of the family of starlike functions of order α</i>	197
Massimo A. Picardello , <i>Positive definite functions and L^p convolution operators on amalgams</i>	209
Friedrich Roesler , <i>Squarefree integers in nonlinear sequences</i>	223
Theodore Shifrin , <i>The osculatory behavior of surfaces in \mathbf{P}^5</i>	227