

Pacific Journal of Mathematics

SQUAREFREE INTEGERS IN NONLINEAR SEQUENCES

FRIEDRICH ROESLER

SQUAREFREE INTEGERS IN NON-LINEAR SEQUENCES

FRIEDRICH ROESLER

The number of squarefree integers in sequences $[f(n+x)]$, $n = 1, 2, 3, \dots, N$ is $(6/\pi^2 + o(1))N$ for almost all $x \geq 0$, when f is a polynomial function of degree $k \geq 2$ or an exponential function.

Following a paper of Stux [4] Rieger has shown in [2] that for all real c with $1 < c < 3/2$ the equation

$$\sum_{\substack{1 \leq n \leq N \\ [n^c] \text{ squarefree}}} 1 = \frac{6}{\pi^2} N + O_c(N^{(2c+1)/4})$$

is an immediate consequence from estimates of Deshouillers [1] concerning the distribution of the integer-sequences $[n^c]$ modulo k .

Using the same method and results of [3] we prove for functions f belonging to one of the following classes

$$\mathfrak{F}_{\text{pol}} = \left\{ ay^k + \sum_{i=1}^m a_i y^{k_i}; a > 0, k > k_1 > \dots > k_m \geq 0 \right\},$$

$$\mathfrak{F}_{\text{exp}} = \{ e^{ky+l} + f(y); k > 0, f \in \mathfrak{F}_{\text{pol}} \cup \{0\} \}:$$

THEOREM. For $f \in \mathfrak{F}_{\text{pol}}$ of degree $k \geq 2$ or $f \in \mathfrak{F}_{\text{exp}}$,

$$\lim_{N \rightarrow \infty} \frac{1}{N} \sum_{\substack{1 \leq n \leq N \\ [f(n+x)] \text{ squarefree}}} 1 = \frac{6}{\pi^2}$$

holds for almost all real $x \geq 0$.

Proof. Let \mathfrak{Q} denote the set of squarefree integers, $0 \leq x < 1$, and

$$S_N(\mathfrak{Q}; x) = \sum_{\substack{1 \leq n \leq N \\ [f(n+x)] \in \mathfrak{Q}}} 1,$$

$$S_N(d\mathbf{N}; x) = \sum_{\substack{1 \leq n \leq N \\ [f(n+x)] \in d\mathbf{N}}} 1.$$

Then

$$|\mu(n)| = \sum_{d > 0, d^2 | n} \mu(d)$$

gives

$$S_N(\mathfrak{Q}; x) = \sum_{d>0} \mu(d) S_N(d^2\mathbf{N}; x)$$

and therefore

$$\begin{aligned} (1) \quad \int_0^1 \left| S_N(\mathfrak{Q}; x) - \frac{6}{\pi^2} N \right| dx &= \int_0^1 \left| \sum_{d>0} \mu(d) \left(S_N(d^2\mathbf{N}; x) - \frac{N}{d^2} \right) \right| dx \\ &= O\left(\sum_{d>0} \int_0^1 \left| S_N(d^2\mathbf{N}; x) - \frac{N}{d^2} \right| dx \right). \end{aligned}$$

To estimate the sum on the right side of (1) we use

$$\int_0^1 \left| S_N(d\mathbf{N}; x) - \frac{N}{d} \right| dx = d^{1/2} O_f(N^{3/4} \log N)$$

which for functions $f \in \mathfrak{F}_{\text{pol}}$ of degree $k \geq 2$ is Satz 9 in [3] and for $f \in \mathfrak{F}_{\text{exp}}$ follows easily from the proof of Satz 6(ii) in [3]. Thus we have for small values of d

$$(2) \quad \sum_{0 < d \leq M} \int_0^1 \left| S_N(d^2\mathbf{N}; x) - \frac{N}{d^2} \right| dx = O_f(M^2 N^{3/4} \log N).$$

The rest of the sum in (1) is estimated by

$$\begin{aligned} (3) \quad \sum_{d>M} \int_0^1 \left| S_N(d^2\mathbf{N}; x) - \frac{N}{d^2} \right| dx &\leq \sum_{d>M} \int_0^1 S_N(d^2\mathbf{N}; x) dx + N \sum_{d>M} \frac{1}{d^2} \\ &\leq \sum_{d>M} \int_0^1 S_N(d^2\mathbf{N}; x) dx + \frac{N}{M}. \end{aligned}$$

The explicit formula for the value of the integral $\int_0^1 S_N^{\text{nat}}(\mathfrak{P}; x) dx$ as a finite sum of interval-lengths

$$\sum_{1 \leq n \leq N} \sum_{p \in \mathfrak{P}_n(0,1)} \mu(I_{p,n}^{0,1})$$

as given at the end of part 1 in [3] shows in particular

$$(4) \quad \int_0^1 S_N(d^2\mathbf{N}; x) dx = O_f \left(\sum_{\substack{m \in d^2\mathbf{N} \\ f(0) \leq m \leq f(N+1)}} (f^{-1})'(m) \right).$$

For $f \in \mathfrak{F}_{\text{pol}}$, $\deg f = k \geq 2$, we have

$$(f^{-1})'(y) = O_f(y^{1/k-1})$$

and thus get from (4)

$$\begin{aligned}
 (5) \quad \int_0^1 S_N(d^2\mathbf{N}; x) dx &= O_f \left(\sum_{1 \leq n \leq f(N+1)/d^2} (d^2n)^{1/k-1} \right) \\
 &= O_f \left(d^{2/k-2} \sum_{1 \leq n \leq N^k/d^2} n^{1/k-1} \right) = O_f \left(\frac{N}{d^2} \right).
 \end{aligned}$$

For $f \in \mathfrak{F}_{\text{exp}}$, we even have

$$(f^{-1})'(y) = O_f(y^{-1})$$

and obtain (5) again.

Finally we choose $M = N^{1/12}$ and (1), (2), (3), (5) give

$$\int_0^1 \left| \frac{1}{N} S_N(\mathfrak{Q}; x) - \frac{6}{\pi^2} \right| dx = O_f(N^{-1/12} \log N).$$

This implies first

$$\lim_{N \rightarrow \infty} \frac{1}{N^{13}} S_{N^{13}}(\mathfrak{Q}; x) = \frac{6}{\pi^2}$$

for almost all x in the interval $[0, 1[$ (with respect to the Lebesgue-measure on \mathbf{R}) and then by the principle of Hilfssatz 1 in [3]

$$\lim_{N \rightarrow \infty} \frac{1}{N} S_N(\mathfrak{Q}; x) = \frac{6}{\pi^2}$$

for almost all $x \geq 0$.

REFERENCES

[1] J.-M. Deshouillers, *Sur la répartition des nombres $[n^c]$ dans les progressions arithmétiques*, C. R. Acad. Sci. Paris, **277** Ser. A., (1973), 647–650.
 [2] G. J. Rieger, *Remark on a paper of Stux concerning squarefree numbers in non-linear sequences*, Pacific J. Math., **78** (1978), 241–242.
 [3] F. Roesler, *Über die Verteilung der Primzahlen in Folgen der Form $[f(n+x)]$* , Acta Arith., **35** (1979), 117–174.
 [4] I. E. Stux, *Distribution of squarefree integers in non-linear sequences*, Pacific J. Math., **59** (1975), 577–584.

Received March 9, 1979.

TECHNISCHE UNIVERSITÄT
MÜNCHEN, GERMANY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720
H. SAMELSON
Stanford University
Stanford, CA 94305
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Maria Emilia Alonso García , A note on orderings on algebraic varieties	1
F. S. De Blasi and Józef Myjak , On continuous approximations for multifunctions	9
Frank Albert Farris , An intrinsic construction of Fefferman's CR metric	33
Antonio Giambruno, P. Misso and Francisco César Polcino Milies , Derivations with invertible values in rings with involution	47
Dan Haran and Moshe Jarden , The absolute Galois group of a pseudo real closed algebraic field	55
Telemachos E. Hatziafratis , Integral representation formulas on analytic varieties	71
Douglas Austin Hensley , Dirichlet's theorem for the ring of polynomials over $GF(2)$	93
Sofia Kalpazidou , On a problem of Gauss-Kuzmin type for continued fraction with odd partial quotients	103
Harvey Bayard Keynes and Mahesh Nerurkar , Ergodicity in affine skew-product toral extensions	115
Thomas Landes , Normal structure and the sum-property	127
Anthony To-Ming Lau and Viktor Losert , Weak*-closed complemented invariant subspaces of $L_\infty(G)$ and amenable locally compact groups	149
Andrew Lelek , Continua of constant distances in span theory	161
Dominikus Noll , Sums and products of B_r spaces	173
Lucimar Nova , Fixed point theorems for some discontinuous operators	189
A. A. S. Perera and Donald Rayl Wilken , On extreme points and support points of the family of starlike functions of order α	197
Massimo A. Picardello , Positive definite functions and L^p convolution operators on amalgams	209
Friedrich Roesler , Squarefree integers in nonlinear sequences	223
Theodore Shifrin , The osculatory behavior of surfaces in \mathbf{P}^5	227