

Pacific Journal of Mathematics

ON RESTRICTION THEOREMS OF MAXIMAL-TYPE

CHRISTOPHER DONALD SOGGE

ON RESTRICTION THEOREMS OF MAXIMAL-TYPE

CHRISTOPHER D. SOGGE

In this paper we prove a vector-valued restriction theorem for the Fourier transform. This result allows us to establish certain sharp “maximal-type” restriction theorems and a partial generalization of the Riemann-Lebesgue lemma for certain $p > 1$.

1. Introduction. Vector-valued inequalities for Bochner-Riesz multipliers have been known for some time in two dimensions and for certain exponents in higher dimensions (cf. [2], [3], [4]). In this note our main result is the corresponding vector-valued restriction theorem for the Fourier transform which is valid everywhere the restriction problem is now known to be true. In two dimensions the result is proved by using the techniques of oscillatory integrals while in higher dimensions our inequalities follow trivially from known ones. Finally, using these results we then obtain a lacunary restriction theorem of maximal-type for the Fourier transform.

2. Vector-valued inequalities. Our main results for this section will be the following sharp vector-valued restriction theorems. By $d\sigma$ we will mean Lebesgue measure on the unit sphere $S^{n-1} \subset \mathbf{R}^n$.

THEOREM 1. *If $n = 2$, $1 \leq p < 4/3$, $q = p'/3$, $R_j > 0$, $j = 1, 2, 3, \dots$*

$$\left\| \left(\sum |\hat{f}_j(R_j \cdot) R_j^{2/p'}|^2 \right)^{1/2} \right\|_{L^q(S^1)} \leq C_p \left\| \left(\sum |f_j|^2 \right)^{1/2} \right\|_{L^p(\mathbf{R}^2)},$$

whenever $f_j \in \mathcal{S}$.

THEOREM 2. *If*

$$n \geq 3, \quad 1 \leq p \leq 2(n+1)/(n+3),$$

$$q = (n-1)p'/(n-1), \quad R_j > 0,$$

$$\left\| \left(\sum |\hat{f}_j(R_j \cdot) R_j^{n/p'}|^2 \right)^{1/2} \right\|_{L^q(S^{n-1})} \leq C_p \left\| \left(\sum |f_j|^2 \right)^{1/2} \right\|_{L^p(\mathbf{R}^n)},$$

whenever $f_j \in \mathcal{S}$.

We start out by proving Theorem 1. For this purpose we define for $x \in \mathbf{R}^2, R_j > 0$,

$$T_j g(x) = \int_{S^1} \exp[iR_j(x \cdot y)] g(y) d\sigma(y).$$

It then follows from duality that Theorem 1 is equivalent to the following:

THEOREM 1'. *If $1 \leq p < 4, q = 3p'$*

$$\left\| \left(\sum |T_j g_j|^2 \right)^{1/2} \right\|_{L^q(\mathbf{R}^2)} \leq C_p \left\| \left(\sum |g_j(\cdot) R_j^{-2/q}|^2 \right)^{1/2} \right\|_{L^p(S^1)}.$$

To prove this we will make use of the following lemma which follows trivially from Minkowski's integral inequality (cf. [8], p. 19).

LEMMA. *If (X, dx) is a measure space and $1 \leq r \leq 2$ then*

$$(1) \quad \left(\sum \|f_j\|_r^2 \right)^{1/2} \leq \left\| \left(\sum |f_j|^2 \right)^{1/2} \right\|_r$$

and,

$$(2) \quad \left\| \left(\sum |f_j|^2 \right)^{1/2} \right\|_{r'} \leq \left(\sum \|f_j\|_r^2 \right)^{1/2}.$$

We now come to the proof of Theorem 1'. It is based on a modification of the arguments of Zygmund [9] using inequalities (1) and (2).

Proof of Theorem 1'. Fix p and q as in the statement and put $r = q/2$. Then $r > 2$ and so by (2) we have

$$(3) \quad \left\| \left(\sum |T_j g_j|^2 \right)^{1/2} \right\|_q^2 \leq \left(\sum \sum \|T_j g_j T_k g_k\|_r^2 \right)^{1/2}.$$

However,

$$\begin{aligned} & \|T_j g_j T_k g_k\|_r \\ &= \left(\int_{\mathbf{R}^2} \left| \int_0^{2\pi} \int_0^{2\pi} \exp[ix \cdot (R_j \cos s - R_k \cos t, R_j \sin s - R_k \sin t)] \right. \right. \\ & \quad \left. \left. \times g_j(s) g_k(t) ds dt \right|^r dx \right)^{1/r}. \end{aligned}$$

Therefore, we put for a fixed pair j, k

$$u = (R_j \cos s - R_k \cos t, R_j \sin s - R_k \sin t)$$

and since $|\partial u/\partial(s, t)| = R_j R_k |\sin(s - t)|$ we see that the last integral equals

$$\begin{aligned}
 (4) \quad & \left(\int_{R^2} \left| \int_{R^2} e^{iu \cdot x} g_j(s) g_k(t) |R_j R_k \sin(s - t)|^{-1} du \right|^r dx \right)^{1/r} \\
 & \leq \left(\int |g_j(s) g_k(t) (R_j R_k \sin(s - t))^{-1}|^{r'} du \right)^{1/r'} \\
 & = \left(\iint |R_j^{-2/q} g_j(s)|^{r'} |R_k^{-2/q} g_k(t)|^{r'} |\sin(s - t)|^{-r'+1} ds dt \right)^{1/r'}.
 \end{aligned}$$

The last inequality comes from the Hausdorff-Young Theorem. Since $r' < 2$ we now see from inequalities (1), (3), (4) that

$$\begin{aligned}
 & \left\| \left(\sum |T_j g_j|^2 \right)^{1/2} \right\|_q^2 \\
 & \leq \left[\sum \sum \left(\iint |R_j^{-2/q} g_j(s)|^{r'} |R_k^{-2/q} g_k(t)|^{r'} \right. \right. \\
 & \qquad \qquad \qquad \left. \left. \times |\sin(s - t)|^{-r'+1} ds dt \right)^{2/r'} \right]^{1/2} \\
 & \leq \left[\iint \left(\sum |R_j^{-2/q} g_j(s)|^2 \right)^{r'/2} \left(\sum |R_k^{-2/q} g_k(t)|^2 \right)^{r'/2} \right. \\
 & \qquad \qquad \qquad \left. \times |\sin(s - t)|^{-r'+1} ds dt \right]^{1/r'}.
 \end{aligned}$$

If now as in [9] we use the classical inequality or Hardy-Littlewood for fractional integrals we find that this last expression is dominated by

$$\left(\int \left(\sum |R_j^{-2/q} g_j(s)|^2 \right)^{p/2} ds \right)^{2/p} = \left\| \left(\sum |R_j^{-2/q} g_j|^2 \right)^{1/2} \right\|_p^2.$$

This completes the proof of Theorem 1'.

Theorem 2 follows easily from the L^2 -restriction theorem of Stein-Tomas [7] and our lemma. In fact if, as before, we define

$$T_j g(x) = \int_{S^{n-1}} \exp[iR_j(x \cdot y)] g(y) d\sigma(y)$$

then the restriction theorem of Stein-Tomas implies that for each $1 \leq p \leq 2$, $q = p'(n + 1)/(n - 1)$,

$$\|T_j g\|_{L^q(R^n)} \leq C \|R_j^{-n/q} g_j\|_{L^p(S^{n-1})}.$$

Therefore, since $q > 2$ and $p < 2$ this inequality and the lemma show that

$$\begin{aligned} \left\| \left(\sum |T_j g_j|^2 \right)^{1/2} \right\|_q &\leq \left(\sum \|T_j g_j\|_q^2 \right)^{1/2} \leq C \left(\sum \|R_j^{-n/q} g_j\|_p^2 \right)^{1/2} \\ &\leq C \left\| \left(\sum |R_j^{-n/q} g_j|^2 \right)^{1/2} \right\|_p. \end{aligned}$$

As before, Theorem 2 follows by duality from this last inequality.

3. Restriction theorems of maximal-type. In this section we indicate how the above results imply the following sharp lacunary restriction theorems of maximal-type.

We define the lacunary maximal restriction operators:

$$M_p \hat{f}(y) = \sup_{j \in \mathbf{Z}} |\hat{f}(2^j y) 2^{nj/p'}|, \quad y \in S^{n-1}, f \in \mathcal{S}.$$

THEOREM 3. *Let $n = 2, 1 \leq p < 4/3, q = p'/3$. Then*

$$\|M_p \hat{f}\|_{L^q(S^{n-1})} \leq C_p \|f\|_{L^p(\mathbf{R}^n)}, \quad f \in \mathcal{S}.$$

THEOREM 4. *Let $n \geq 3,$*

$$1 \leq p \leq 2(n+1)/(n+3), \quad q = p'(n-1)/(n+1).$$

Then

$$\|M_p \hat{f}\|_{L^p(S^{n-1})} \leq C_p \|f\|_{L^p(\mathbf{R}^n)}, \quad f \in \mathcal{S}.$$

Proofs. Fix $n \geq 2$ and p and q as above. Clearly, to prove these results it suffices to show that whenever $\{E_j\}, j \in \mathbf{Z}$, is a collection of disjoint measurable subsets of S^{n-1} there is an absolute constant C for which

$$(5) \quad \left\| \sum \chi_{E_j}(\cdot) \hat{f}(2^j \cdot) 2^{nj/p'} \right\|_{L^q(S^{n-1})} \leq C \|f\|_p.$$

Now let $b \in C_0^\infty$ be a bump function with the properties that $b(1) = 1, \text{supp}(b) \subset (1/4, 2)$ and $\sum b(2^j t) = 1, t \in \mathbf{R}_+$. If we define f_j by $\hat{f}_j = b(2^{-j}|y|) \hat{f}(y)$ then the left hand side of (5) is majorized by

$$\left\| \left(\sum |\hat{f}_j(2^j \cdot) 2^{nj/p'}|^2 \right)^{1/2} \right\|_{L^q(S^{n-1})} \leq C \left\| \left(\sum |f_j|^2 \right)^{1/2} \right\|_{L^p(\mathbf{R}^n)}.$$

The inequality follows from Theorems 1 and 2. Our desired result now follows via Littlewood-Paley theory from the inequality $\|(\sum |f_j|^2)^{1/2}\|_p \leq C \|f\|_p$ if $p > 1$. The case $p = 1$ is the Riemann-Lebesgue lemma.

This finishes the proofs.

If now $n \geq 2$ and p is as in Theorem 1 or 2 it follows that $\hat{f}(2^j\xi)$ can be defined as an element of $L^q(S^{n-1})$, $q = p'(n - 1)/(n + 1)$. Elements of $L^q(S^{n-1})$ of course are only defined almost everywhere so in what follows to avoid ambiguity we assume that for each $f \in L^p(\mathbf{R}^n)$ and $j \in \mathbf{Z}$ we have fixed a function $\hat{f}(2^j\xi)$ which represents this element. If we now define $M_p\hat{f}$ as before, it follows from a simple limiting argument that the a priori inequalities of Theorems 3 and 4 now hold for general $f \in L^p(\mathbf{R}^n)$. With these things in mind we now come to the following partial generalizations of the classical Riemann-Lebesgue lemma.

COROLLARY 1. *If $1 < p < 4/3$, $f \in L^p(\mathbf{R}^2)$ then for almost every direction $\xi \in S^1$ we have:*

$$2^{2j/p'}\hat{f}(2^j\xi) \rightarrow 0 \quad \text{as } |j| \rightarrow \infty.$$

COROLLARY 2. *If $n \geq 3$, $1 < p \leq (2n + 2)/(n + 3)$ and $f \in L^p(\mathbf{R}^n)$ then for almost every every direction $\xi \in S^{n-1}$ we have:*

$$2^{nj/p'}\hat{f}(2^j\xi) \rightarrow 0 \quad \text{as } |j| \rightarrow \infty.$$

Proofs. Fix $n \geq 2$ and p and q as above. Define

$$\Omega\hat{f}(\xi) = \limsup_{j \rightarrow \infty} \left[|2^{nj/p'}\hat{f}(2^j\xi)| + |2^{-nj/p'}\hat{f}(2^{-j}\xi)| \right].$$

We must show that $\Omega\hat{f} = 0$ for almost every direction $\xi \in S^{n-1}$. Clearly, $\Omega\hat{f}(\xi) \leq 2M_p\hat{f}(\xi)$ and so by Theorems 3 and 4

$$m\{\xi \in S^{n-1}: \Omega\hat{f}(\xi) > \varepsilon\} \leq C/\varepsilon^q \|f\|_p^q.$$

It is also clear that if $h \in \mathcal{S}$ then $\Omega\hat{h} = 0$ almost everywhere. Furthermore, $f \in L^p(\mathbf{R}^n)$ can be written as $f = g + h$ with $h \in \mathcal{S}$ and $\|g\|_p$ arbitrarily small. Consequently, as $\Omega\hat{f} \leq \Omega\hat{g} + \Omega\hat{h}$ we get that

$$m\{\xi \in S^{n-1}: \Omega\hat{f}(\xi) > \varepsilon\} \leq C/\varepsilon^q \|g\|_p^q.$$

Finally, since the size of $\|g\|_p$ is at our disposal, we see that $\Omega\hat{f} = 0$ almost everywhere as desired.

REMARKS. We now show that how the results of this section are of the best possible nature in the sense that in \mathbf{R}^n , $n \geq 2$, no inequality of the kind

$$\int_{S^{n-1}} \sup_{1 < r < 2} |\hat{f}(r\xi)| d\sigma \leq A_p \|f\|_{L^p(\mathbf{R}^n)}$$

can hold for any $p > 1$.

In fact consider $f = f_1(x')f_2(x_n)$ where $f_1(x') = e^{-|x'|^2}$, and

$$\hat{f}_2(\xi_n) = \begin{cases} |\xi_n - 1|^{-1/2p'}, & |\xi_n| < 2 \\ 0, & \text{otherwise.} \end{cases}$$

Then $\hat{f}(\xi', \xi_n) = c_n |\xi_n - 1|^{-1/2p'} e^{-|\pi\xi'|^2}$ for (ξ', ξ_n) near $(0, \dots, 0, 1)$, and hence $\sup_{1 < r < 2} |\hat{f}(r\xi)| = \infty$ $\xi \in S^{n-1}$ in a neighborhood of $(0, \dots, 0, 1)$ even though $f \in L^p(\mathbf{R}^n)$.

This counterexample was kindly pointed out to the author by E. M. Stein.

Finally, we also remark that one would expect that Theorems 2 and 4 should also hold for $(2n + 2)/(n + 3) < p < 2n/(n + 1)$, since this is the conjectured range for the restriction problem for the Fourier transform (cf. [5]).

Acknowledgments. I would like to thank E. M. Stein for many helpful conversations. I would also like to thank Alfred P. Sloan Foundation for the generous support which they provided through the Doctoral Dissertation Fellowship Program.

REFERENCES

- [1] A. Carbery, *The boundedness of the maximal Bochner-Riesz operator on $L^4(\mathbf{R}^2)$* , Duke Math. J., **50** (1983), 409–416.
- [2] M. Christ, *On almost everywhere convergence of Bochner-Riesz means in higher dimensions*, preprint.
- [3] A. Córdoba and B. Lopez-Melero, *Spherical summation: a Problem of E. M. Stein*, Ann. Inst. Fourier, **31** (1982), 147–152.
- [4] S. Igari, *Decomposition theorem and lacunary convergence of Bochner-Riesz means of Fourier transforms of two variables*, Tohoku Math. J., **33** (1981), 413–419.
- [5] E. M. Stein, *Some problems in harmonic analysis*, Proc. Sym. Pure Math., Amer. Math. Soc., XXXV Part I (1979), 3–20.
- [6] E. M. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton Univ. Press, 1971.
- [7] P. Tomas, *Restriction theorems for the Fourier transform*, Proc. Sym. Pure Math., Amer. Math. Soc., XXXV Part (1979), 111–114.
- [8] A. Zygmund, *Trigonometric Series*, Cambridge Univ. Press, Cambridge England, 1979.
- [9] ———, *On Fourier coefficients and transforms of functions of two variables*, Studia Math., **50** (1974), 189–202.

Received February 25, 1985.

PRINCETON UNIVERSITY
PRINCETON, NJ 08544

Current address: University of Chicago
Chicago, IL 60637

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720
H. SAMELSON
Stanford University
Stanford, CA 94305
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA	UNIVERSITY OF OREGON
UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA	UNIVERSITY OF HAWAII
MONTANA STATE UNIVERSITY	UNIVERSITY OF TOKYO
UNIVERSITY OF NEVADA, RENO	UNIVERSITY OF UTAH
NEW MEXICO STATE UNIVERSITY	WASHINGTON STATE UNIVERSITY
OREGON STATE UNIVERSITY	UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1986 by Pacific Journal of Mathematics

David Jay Anick , A loop space whose homology has torsion of all orders . . .	257
Steven P. Diaz , Space curves that intersect often	263
Thierry Fack and Hideki Kosaki , Generalized s -numbers of τ -measurable operators	269
Karl Heinrich Hofmann and Karl Strambach , Lie's fundamental theorems for local analytical loops	301
James Secord Howland , On the Kato-Rosenblum theorem	329
Frieder Knüppel and Edzard Salow , Plane elliptic geometry over rings . . .	337
Alan Noell , Peak points in boundaries not of finite type	385
William J. Ralph , An extension of singular homology to Banach algebras	391
Wade C. Ramey , Averaging properties of pluriharmonic boundary values . .	407
Thomas Joseph Ransford , On the range of an analytic multivalued function	421
Christopher Donald Sogge , On restriction theorems of maximal-type	441
Edwin Spanier , Cohomology with supports	447
Emil J. Straube , Orthogonal projections onto subspaces of the harmonic Bergman space	465
Thomas Vogel , Asymptotic behavior of two semilinear elliptic free boundary problems	477