

Pacific Journal of Mathematics

ON ACCRETIVE OPERATORS ON l_n^∞

HUBERT BERENS AND LOTHAR HETZELT

ON ACCRETIVE OPERATORS ON l_n^∞

H. BERENS AND L. HETZELT

To Professor H. G. Tillmann on the occasion of his 60th birthday

It is the object of the paper to discuss the result of Crandall and Liggett on m -accretive operators in l_n^p in greater detail.

1. Introduction. By Zorn's Lemma any accretive operator has a maximally accretive extension. In this respect G. J. Minty [12] proved in 1962 that if H is a Hilbert space the accretive (= monotone) operator $A \subset H \times H$ is maximally accretive exactly when there exists a $\lambda \in \mathbf{R}^+$ such that (and consequently for all $\lambda \in \mathbf{R}^+$)

$$(1.1) \quad \forall y \in H \quad x + \lambda A = y \quad \text{has a (unique) solution in } A,$$

i.e., $I + \lambda A$ defines a bijection of A onto H . In this case, A is said to be m -accretive. See the following section for the relevant definitions and notations.

In contrast to Minty's result, M. G. Crandall and T. W. Liggett [6] showed in 1971 that for l_n^p , $n \in \mathbf{N}$, ≥ 2 and $1 < p \leq \infty$, the class of m -accretive operators coincides with the class of maximally accretive ones exactly when $p = 2$, or ∞ . In particular, we want to reprove their

(1.2) **THEOREM (Crandall-Liggett).** *In l_n^∞ , $n \in \mathbf{N}$, the class of maximally accretive operators coincides with the class of m -accretive ones.*

In the following section we do the necessary preliminary work, while §3 is devoted to the proof of the theorem. In §4 we comment on the theorem. Section 5, finally, deals with the domain and range of m -accretive operators.

We would like to thank Professor K. Donner for the many valuable discussions we had on this and related subjects.

2. Definitions, Notations, and Preliminaries. Let X be a finite dimensional, real, normed vector space with $\|\cdot\|$. Its elements are denoted by x, y, z, \dots . For a subset K of X , \bar{K} , $\overset{\circ}{K}$, and ∂K denote its closure, interior, and boundary, respectively. The open ball centered at x with radius $r \in \mathbf{R}^+$ is denoted by $b_r(x)$.

The *semi-inner product* $\langle \cdot, \cdot \rangle_s: X \times X \rightarrow \mathbf{R}$ is defined by

$$\langle y, x \rangle_s := \lim_{t \rightarrow 0+} \frac{\|x + ty\|^2 - \|x\|^2}{2t}, \quad (y, x) \in X \times X.$$

If F denotes the duality map on X into its dual— F is the subdifferential of $\|\cdot\|^2/2: X \rightarrow \mathbf{R}$ —then

$$\langle y, x \rangle_s = \max\{\langle y, w \rangle: w \in F(x)\}.$$

It follows that $\langle \cdot, \cdot \rangle_s$ is upper semi-continuous on $X \times X$. Furthermore, if X is strictly convex, then $F(x) \cap F(x') = \emptyset$ whenever $x \neq x'$. If X is smooth, then F is single-valued and $\langle y, x \rangle_s$ is just the derivative of $\|\cdot\|^2/2$ at x in the direction y . If, in particular, X is an inner product space, the semi-inner product reduces to the inner product on $X \times X$. For l_n^p , $1 < p \leq \infty$,

$$\langle y, x \rangle_s = \begin{cases} \sum_{i=1}^n y_i \operatorname{sgn} x_i |x_i|^{p-1} / \|x\|_p^{p-2}, & 1 < p < \infty, \\ \max\{y_i x_i: 1 \leq i \leq n, |x_i| = \|x\|_\infty\}, & p = \infty, \end{cases}$$

$$x, y \in l_n^p, x \neq 0.$$

For $x, y \in X$, $x \neq y$,

$$C(y, x) := \bigcup_{\lambda > 0} b_{\lambda\|x-y\|}(y + \lambda(x - y))$$

denotes the *cone of decrease* of x with vertex at y . $C(y, x)$ is the open tangential cone of $b_{\|x-y\|}(x)$ at y . With use of the semi-inner product the cone of decrease can be rewritten as

$$C(y, x) = \{z \in X: \langle y - z, x - y \rangle_s < 0\}.$$

If $x = y$ $C(y, x)$ is defined to be the empty set.

The set of intermediate points between x and y in X is defined by

$$Z(x, y) := \{z \in X: \|x - z\| + \|z - y\| = \|x - y\|\}.$$

A $\|\cdot\|$ -segment in X is a curve whose length equals the distance of its endpoints.

We say that $K \subset X$ is $\|\cdot\|$ -convex, if

$$\forall k', k'' \in K \exists k \in K, k \neq k', k'' \ni k \in Z(k', k'').$$

If K is a closed subset of X there is a useful characterization of $\|\cdot\|$ -convexity, see [5, p. 29].

(2.1) LEMMA. *Let K be a closed set in X . K is $\|\cdot\|$ -convex exactly when every two points in K can be joint by a $\|\cdot\|$ -segment completely contained in K .*

If X is strictly convex, then $\|\cdot\|$ -convexity reduces to the classical notion of convexity. For l_n^1 —this is the case we are particularly interested in—the intermediate points of x and y are given by the parallelepiped spanned by x and y . And $[0, \|y - x\|_1] \ni t \mapsto z(t)$ defines a $\|\cdot\|_1$ -segment joining x and y , if it is a continuous curve with x and y as endpoints which is monotone in each component $z_i(t)$, $0 \leq t \leq \|y - x\|_1$.

A set-valued map A from X into itself is called an operator on X . It is convenient to identify A with its graph in $X \times X$. $D(A) := \{x \in X: (x, a) \in A\}$ and $R(A) := \{a \in X: (x, a) \in A\}$ denote the domain and range of A , respectively. $A^{-1} := \{(a, x) \in X \times X: (x, a) \in A\}$. If A^1 and A^2 are two operators on X , $A^1 + A^2 := \{(x, a^1 + a^2) \in X \times X: (x, a^1) \in A^1 \text{ and } (x, a^2) \in A^2\}$, and for $\lambda \in \mathbf{R}^+$ $\lambda A := \{(x, \lambda a) \in X \times X: (x, a) \in A\}$.

The operator A is said to be *accretive* if

$$\forall (x, a), (x', a') \in A \quad 0 \leq \langle a - a', x - x' \rangle_s.$$

This is equivalent to the fact that

$$\forall \lambda \in \mathbf{R}^+ \quad (I + \lambda A)^{-1} \text{ defines a contraction from } X \text{ to itself,}$$

i.e.,

$$\forall (x, a), (x', a') \in A$$

$$\|x - x'\| \leq \|(x + \lambda a) - (x' + \lambda a')\|, \quad \forall \lambda \in \mathbf{R}^+.$$

A is said to be *m-accretive* if A is accretive and if for all $\lambda \in \mathbf{R}^+$ $I + \lambda A$ is surjective. (It follows trivially from the accretiveness of A that $I + \lambda A$ is injective, thus if A is *m-accretive* for each $\lambda \in \mathbf{R}^+$ $I + \lambda A$ is bijective.) An *m-accretive* operator on X is a maximal element within the class of accretive operators on X ordered by inclusion.

If X is an inner product space it is common to speak of monotone operators instead of accretive ones.

3. Proof of the Theorem. The key to our proof is

(3.1) THEOREM. *Let A be a finite accretive set in $l_n^\infty \times l_n^\infty$, then*

$$\forall x \in l_n^\infty \quad B_A(x) := \{a \in l_n^\infty: \langle a' - a, x' - x \rangle_s \geq 0 \quad \forall (x', a') \in A\}$$

is nonempty, closed, and $\|\cdot\|_1$ -convex. If Q_A denotes the smallest closed parallelepiped¹ in l_n^∞ containing $R(A)$, then even $B_A^Q(x) := Q_A \cap B_A(x)$ is nonempty, compact, and $\|\cdot\|_1$ -convex.

¹The faces of Q_A are assumed to be parallel to the main axes.

Let $A := \{(x^1, a^1), \dots, (x^m, a^m)\}$, $m \in \mathbf{N}$, and let $x \in l_n^\infty$. Setting

$$C^j := C(a^j, a^j + x^j - x), \quad 1 \leq j \leq m,$$

Theorem 3.1 claims that

$$B_A(x) = \bigcap_{j=1}^m \mathbf{C}C^j \text{ is } \neq \emptyset, \text{ closed and } \|\cdot\|_1\text{-convex.}$$

Let us introduce some further notation

$$I_\pm^j := \{1 \leq i \leq n: x_i^j - x_i = \pm \|x^j - x\|_\infty\} \quad \text{and} \quad I^j := I_+^j \cup I_-^j.$$

To prove Theorem 3.1 we start with

(3.2) LEMMA.

$$\forall 1 \leq j, l \leq m \quad C^j \cap C^l \text{ is a cone.}$$

Proof. It is an easy exercise to verify that

$$\begin{aligned} C^j \cap C^l &= \{z \in l_n^\infty: z_i > a_i^j, i \in I_+^j \setminus I^l\} \\ &\quad \cap \{z \in l_n^\infty: z_i < a_i^j, i \in I_-^j \setminus I^l\} \\ &\quad \cap \{z \in l_n^\infty: z_i > a_i^l, i \in I_+^l \setminus I^j\} \\ &\quad \cap \{z \in l_n^\infty: z_i < a_i^l, i \in I_-^l \setminus I^j\} \\ &\quad \cap \{z \in l_n^\infty: z_i > \max(a_i^j, a_i^l), i \in I_+^j \cap I_+^l\} \\ &\quad \cap \{z \in l_n^\infty: z_i < \min(a_i^j, a_i^l), i \in I_-^j \cap I_-^l\} \\ &\quad \cap \{z \in l_n^\infty: a_i^j < z_i < a_i^l, i \in I_+^j \cap I_-^l\} \\ &\quad \cap \{z \in l_n^\infty: a_i^l < z_i < a_i^j, i \in I_-^j \cap I_+^l\}. \end{aligned}$$

Hence, if $C^j \cap C^l \neq \emptyset$

$$C^j \cap C^l \text{ is a cone iff } I_+^j \cap I_-^l = \emptyset \text{ and } I_-^j \cap I_+^l = \emptyset.$$

But if $i \in I_+^j \cap I_-^l$, then

$$\begin{aligned} x_i^j - x_i^l &= (x_i^j - x_i) - (x_i^l - x_i) \\ &= \|x^j - x\|_\infty + \|x^l - x\|_\infty = \|x^j - x^l\|_\infty, \end{aligned}$$

and consequently,

$$(3.3) \quad (a_i^j - a_i^l)(x_i^j - x_i^l) = \langle a^j - a^l, x^j - x^l \rangle_s \geq 0$$

implies $a_i^l \leq a_i^j$, or $C^j \cap C^l = \emptyset$. Similarly, if $i \in I_-^j \cap I_+^l$, then $x_i^j - x_i^l = -\|x^j - x^l\|_\infty$, by (3.3) $a_i^j \leq a_i^l$ which again implies $C^j \cap C^l = \emptyset$. \square

Proof of Theorem 3.1. We prove the theorem by induction on $m = \text{card}(A)$. For $m = 1$ there is nothing to prove. Let us assume that for all accretive sets of cardinality $m - 1$ the theorem holds true.

If $x = x^j$ for some $1 \leq j \leq m$, we are done. Let us therefore assume that $x \neq x^j$, $1 \leq j \leq m$.

By the induction hypothesis

$$(3.4) \quad \bigcap_{j=1}^{m-1} \mathbf{C}C^j \neq \emptyset, \text{ closed, and } \|\cdot\|_1\text{-convex.}$$

Obviously, $B_A(x)$ is closed. First, we prove that it is nonempty. In contrast to our claim, let us assume that $B_A(x) = \emptyset$. Consequently,

$$(3.5) \quad \mathbf{C}C^m \subseteq \bigcup_{j=1}^{m-1} C^j.$$

W.l.o.g. let $I^m = \{1, 2, \dots, s\}$, $1 \leq s \leq n$. We define

$$J_0 := \{1 \leq j \leq m-1 : C^j \cap S^m \neq \emptyset\},$$

where $S^m = \{b \in I_n^\infty : b_i = a_i^m, i \in I^m\}$ is the vertex set of C^m . Since $S^m \subset \mathbf{C}C^m$, by (3.5) $J_0 \neq \emptyset$. We claim,

$$(3.6) \quad \exists j_0 \in J_0 \ni I^{j_0} \subset I^m.$$

If not, then $J_0 = \bigcup_{i=s+1}^n D^i$, where $D^i = D_+^i \cup D_-^i$ and $D_\pm^i = \{j \in J_0 : i \in I_\pm^j\}$. We select a sub-manifold $S_0^m \subset S^m$ subject to the following restrictions. For $s+1 \leq i \leq n$ and $D^i \neq \emptyset$ we set $b_i^0 = \min\{a_i^j : j \in D_+^i\}$, in case $D_+^i \neq \emptyset$, otherwise $= \max\{a_i^j : j \in D_-^i\}$. In the first case, if $D_-^i \neq \emptyset$ too, for all $l \in D_-^i$ $a_i^l \leq b_i^0$, for, if for some $j \in D_+^i$ and some $l \in D_-^i$ $a_i^j < a_i^l$, then $C^j \cap C^l \neq \emptyset$ while $I_+^j \cap I_-^l \neq \emptyset$, contradicting Lemma 3.2.

We define $S_0^m := \{b \in S^m : b_i = b_i^0, s+1 \leq i \leq n, D_i \neq \emptyset\}$. By construction $S_0^m \subset \mathbf{C}C^j$ for all $j \in D^i$, $s+1 \leq i \leq n$, hence for all $j \in J_0$. Trivially, $S_0^m \subset \mathbf{C}C^m$ and $S_0^m \subset \mathbf{C}C^j$ for all $1 \leq j \leq m-1$ not belonging to J_0 . Thus $S_0^m \subset B_A(x)$, which contradicts the assumption that $B_A(x) = \emptyset$.

Hence there exists an index $j_0 \in J_0$ such that $I^{j_0} \subseteq I^m$. Since $C^{j_0} \cap C^m \neq \emptyset$, by use of Lemma 3.2 $I_+^{j_0} \cap I_-^m$ and $I_-^{j_0} \cap I_+^m$ are empty, i.e., $I_+^{j_0} \subseteq I_+^m$ and $I_-^{j_0} \subseteq I_-^m$. But this implies that $C^m \subset C^{j_0}$, and consequently, $B_A(x) = \bigcap_{j=1}^{m-1} \mathbf{C}C^j$ which is nonempty by the induction hypothesis (3.4), again in contradiction to our assumption that $B_A(x) = \emptyset$.

To prove that $B_A(x)$ is $\|\cdot\|_1$ -convex, let us assume there exist two points in $B_A(x)$ which have no proper intermediate points. W.l.o.g. we may assume that 0 and a are these points where $a_i > 0$, $1 \leq i \leq s$ and $a_i = 0$, $s+1 \leq i \leq n$, for some $1 \leq s \leq n$. Hence

$$Z(0, a) \cap B_A(x) = (Z(0, a) \cap \mathbb{C}C^m) \cap \left(Z(0, a) \cap \bigcap_{j=1}^{m-1} \mathbb{C}C^j \right) = \{0, a\}.$$

Since by the induction hypothesis (3.4) $\bigcap_{j=1}^{m-1} \mathbb{C}C^j$ is $\|\cdot\|_1$ -convex, it follows that

$$\{0, a\} \subset \partial C^m \quad \text{and} \quad Z(0, a) \cap C^m \neq \emptyset$$

—indeed, any $\|\cdot\|_1$ -segment in $\bigcap_{j=1}^{m-1} \mathbb{C}C^j$ connecting 0 and a belongs to C^m except for the two endpoints. Consequently, there are two indices i_0 and i_a , $1 \leq i_0, i_a \leq s$, $i_0 \neq i_a$ (by use of Lemma 3.2) such that $i_0 \in I_+^m$ and $i_a \in I_-^m$. W.l.o.g. we may further assume that $i_0 = 1$.

Consider the point

$$b_\varepsilon := (\varepsilon, a_2, \dots, a_n) \text{ in } Z(0, a), \quad 0 < \varepsilon < a_1.$$

It is a proper intermediate point of 0 and a located on ∂C^m . Since by assumption $Z(0, a) \cap B_A(x) = \{0, a\}$, there exists an index j , $1 \leq j \leq m-1$, such that $b_\varepsilon \in C^j$. Obviously, $C^m \cap C^j \neq \emptyset$. By Lemma 3.2 $I_+^m \cap I_-^j$ and $I_-^m \cap I_+^j$ are empty.

If $1 \notin I^j$, then $a \in C^j$, contradicting $a \in B_A(x)$. Hence $1 \in I^j$. If $1 \in I_+^j$, then again $a \in C^j$, contradicting $a \in B_A(x)$. Hence $1 \in I_-^j$. Since $1 \in I_+^m$, this is in contradiction to $I_+^m \cap I_-^j = \emptyset$. Thus any given two points in $B_A(x)$ have proper intermediate points in $B_A(x)$ with respect to the $\|\cdot\|_1$ -norm.

Let Q_A be the smallest closed parallelepiped containing $R(A)$. To prove that even

$$Q_A \cap B_A(x) \text{ is nonempty, compact, and } \|\cdot\|_1\text{-convex,}$$

we extend A via

$$A_{\text{ext}} := A \cup \bigcup_{i=1}^n \{(x^{i,\pm}, a^{i,\pm}) \in l_n^\infty \times l_n^\infty\},$$

where for each $1 \leq i \leq n$

$$x^{i,\pm} := x \pm r e_i,$$

e_i being the i th unit vector in l_n^∞ and $r = 2 \max\{\|x - x^j\|_\infty : 1 \leq j \leq m\}$, and

$$a^{i,\pm} \in Q_A, \quad a_i^{i,+} := \max_{1 \leq j \leq m} a_i^j \quad \text{and} \quad a_i^{i,-} := \min_{1 \leq j \leq m} a_i^j.$$

By definition, A_{ext} is accretive and $B_{A_{\text{ext}}}(x) = Q_A \cap B_A(x)$. \square

(3.7) PROPOSITION. *Let A be a maximally accretive operator on l_n^∞ . $\forall x \in D(A)$ $A(x)$ is closed and $\|\cdot\|_1$ -convex.*

Proof. By the maximality of A and by the u. semi-continuity of the semi-inner product for each $x \in D(A)$ $A(x)$ is closed.

To see that $A(x)$ is $\|\cdot\|_1$ -convex, let Γ be the net of all finite subsets $A_\gamma = \{(x^1, a^1), \dots, (x^{m_\gamma}, a^{m_\gamma})\}$, $m_\gamma \in \mathbb{N}$, of A subject to the restriction that $x^j \neq x$, $1 \leq j \leq m_\gamma$; $\gamma < \gamma'$ if $A_\gamma \subseteq A_{\gamma'}$. Clearly,

$$\begin{aligned} A(x) \subseteq B_A(x) &:= \bigcap_{\substack{(x', a') \in A \\ x' \neq x}} \{b \in l_n^\infty : \langle a' - b, x' - x \rangle_s \geq 0\} \\ &= \lim_{\gamma \in \Gamma} B_{A_\gamma}(x). \end{aligned}$$

We show that $B_A(x)$ is $\|\cdot\|_1$ -convex. Indeed, take b' and b'' in $B_A(x)$, and for each $\gamma \in \Gamma$ let $[0, \|b'' - b'\|_1] \ni t \mapsto b_\gamma(t)$ be a $\|\cdot\|_1$ -segment in $B_{A_\gamma}(x)$ connecting b' and b'' . By definition, b_γ is Lipschitz-continuous, i.e.,

$$\forall 0 \leq t' < t'' \leq \|b'' - b'\|_1, \quad \|b_\gamma(t'') - b_\gamma(t')\|_1 = t'' - t'.$$

By the theorem of Arzela-Ascoli, any accumulation point of the net $\{b_\gamma : \gamma \in \Gamma\}$ defines a $\|\cdot\|_1$ -segment in $B_A(x)$ connecting b' and b'' .

By the maximality of A , however, $A(x) = B_A(x)$. \square

(3.8) PROPOSITION. *Let A be an accretive operator with $R(A) \subseteq Q$, a compact parallelepiped.*

If A is maximal with respect to all accretive operators with range in Q , then $\forall x \in l_n^\infty$ $A(x)$ is nonempty, compact, and $\|\cdot\|_1$ -convex.

Proof. By use of the notation introduced in the proofs of Theorem 3.1 and Proposition 3.7, for each $x \in l_n^\infty$

$$\begin{aligned} A(x) \subseteq B_A^Q(x) &:= \bigcap_{\substack{(x', a') \in A \\ x' \neq x}} \{b \in Q : \langle a' - b, x' - x \rangle_s \geq 0\} \\ &= \lim_{\gamma \in \Gamma} B_{A_\gamma}^Q(x). \end{aligned}$$

By Theorem 3.1 for each $\gamma \in \Gamma$ $B_{A_\gamma}^Q(x)$ is nonempty, compact and $\|\cdot\|_1$ -convex, and so is $B_A^Q(x)$ by a compactness argument. The maximality of A again implies that for each x $A(x) = B_A^Q(x)$. \square

Next we want to prove that under the assumptions of Proposition 3.8 an accretive operator is m -accretive. The proof rests upon a theorem of surjectivity for set-valued mappings (J. M. Lasry and R. Robert [11, Corollaire 1.18, p. 98]): Let $\Gamma: \mathbf{R}^n \rightarrow \mathbf{R}^n$ be an upper semi-continuous, compact-valued, acyclic mapping. If Γ is coercive, then it is surjective.—For each $x \in \mathbf{R}^n$, let $\gamma(x) = \min\{\langle y, x \rangle: y \in \Gamma(x)\}$. Γ is said to be coercive if $\gamma(x)/\|x\| \rightarrow \infty$ when $\|x\| \rightarrow \infty$.

Let A be an accretive operator on l_n^∞ satisfying the assumptions of Proposition 3.8, and let $\lambda \in \mathbf{R}^+$ be fixed. By maximality $I + \lambda A$ is upper semi-continuous, and Proposition 3.8 assures that

$$\forall x \in l_n^\infty \quad A(x) \neq \emptyset, \text{ compact, and } \|\cdot\|_1\text{-convex.}$$

In [1] the authors proved

(3.9) LEMMA. *A nonempty, compact, $\|\cdot\|_1$ -convex set in l_n^∞ is a R_δ -set, i.e., it is the intersection of a decreasing sequence of compact sets which are contractible in itself.*

By J. M. Lasry and R. Robert [11, Proposition 2.1, p. 110] R_δ -sets in \mathbf{R}^n are acyclic (in the sense of Čech-cohomology).

Since $R(A)$ is contained in the compact parallelepiped Q , $I + \lambda A$ is trivially coercive. Thus the conditions of the theorem of surjectivity are fulfilled, giving

(3.10) PROPOSITION. *Let A be an accretive operator on l_n^∞ with $R(A) \subseteq Q$, a compact parallelepiped.*

If A is maximal with respect to all accretive operators with range in Q , then A is m -accretive.

REMARK. To prove Proposition 3.10, instead of the theorem of surjectivity due to Lasry-Robert we may use a set-valued version of the theorem of the invariance of domain (A. Granas and J. W. Jaworowski [8]): Let U be an open subset of \mathbf{R}^n and let $\phi: U \rightarrow \mathbf{R}^n$ be an upper semi-continuous, compact-valued, acyclic mapping such that $\Phi(x) \cap \Phi(x') = \emptyset$ whenever $x \neq x'$, then $\Phi(U)$ is open. Indeed, for each $\lambda \in \mathbf{R}^+$ $I + \lambda A$ satisfies the assumptions of the theorem.

Now we are ready to prove the Theorem.

Let A be a maximally accretive operator on l_n^∞ . W.l.o.g. let $(0, 0) \in A$.

For each $m \in \mathbf{N}$, we define the restriction $A^m = \{(x, a) \in A: \|a\|_\infty \leq m\}$ of A . Let B^m be a maximal extension of A^m subject to the restriction that for all $(x, b) \in B^m$ $\|b\|_\infty \leq m$. By Proposition 3.10 B^m is m -accretive.

Let $\lambda \in \mathbf{R}^+$ be fixed. Assume that $I + \lambda A$ is not surjective. We take a point in l_n^∞ , say y , contained in the complement of $(I + \lambda A)(l_n^\infty)$. Since for each $m \in \mathbf{N}$ B^m is m -accretive

$$y = x^m + \lambda b^m \quad \text{for some } (x^m, b^m) \in B^m.$$

Since $\|x^m\|_\infty \leq \|y\|_\infty$, there exists a convergent subsequence, say $\lim_j x^{m_j} = x$. Consequently, $\lim_j b^{m_j} = (y - x)/\lambda =: b$. Take a pair $(x', a') \in A$. For $m_j \geq \|a'\|_\infty$ $(x', a') \in B^{m_j}$ and by the u. semi-continuity of the semi-inner product

$$0 \leq \overline{\lim}_j \langle b^{m_j} - a', x^{m_j} - x' \rangle_s \leq \langle b - a', x - x' \rangle_s.$$

By construction, $(x, b) \notin A$. Hence $A \cup (x, b)$ properly extends A , which contradicts its maximality. \square

REMARK. The proof of Crandall and Liggett runs as follows:

Let A be a maximally accretive operator on l_n^∞ .

Fix an element $y \in l_n^\infty$. They claim that $y \in R(I + A)$ (λ is set to be equal to 1). To prove their claim they define

$$\forall (x', a') \in A \quad V(x', a') = \{z \in l_n^\infty: \langle y - z - a', z - x' \rangle_s \geq 0\}.$$

If $\bigcap \{V(x', a'): (x', a') \in A\}$ is not empty, say z^0 belongs to the intersection, then by the maximality of A $(z^0, y - z^0) \in A$, giving $y = z^0 + (y - z^0)$. For each $(x', a') \in A$ $V(x', a')$ is nonempty and compact. Thus it remains to verify that $\{V(x', a'): (x', a') \in A\}$ has the finite intersection property.

To do this, let $B = \{(x^1, b^1), \dots, (x^m, b^m)\}$, $m \in \mathbf{N}$ be a finite accretive operator and let D_B be the smallest closed parallelepiped which contains $y - b^j$, $1 \leq j \leq m$. Define $T: D_B \rightarrow D_B$ by

$$D_B \ni x \mapsto T(x) = \{z \in D_B: \langle (y - z) - b^j, x - x^j \rangle_s \geq 0, 1 \leq j \leq m\}.$$

The crucial part of their proof is the verification of the following

(3.11) **LEMMA.**

$\forall x \in D_B \quad T(x) \neq \emptyset$, compact, and contractible in itself.

Obviously, T is upper semi-continuous. By the fixed point theorem of Eilenberg and Montgomery T has a fixed point in D_B . This proves that $\{V(x', a') : (x', a') \in A\}$ has the finite intersection property.

4. Remarks about the Theorem. The counterpart to Theorem 3.1 in \mathbf{R}^n was formulated and proved by G. J. Minty [13] in 1962.

(4.1) **THEOREM.** *Let A be a finite monotone set in $\mathbf{R}^n \times \mathbf{R}^n$. Then $\forall x \in \mathbf{R}^n$ $B_A(x) := \{a \in \mathbf{R}^n : \langle a' - a, x' - x \rangle \geq 0 \forall (x', a') \in A\}$ is nonempty, closed and convex.*

If $Q_A = \text{co}\{a' \in \mathbf{R}^n : (x', a') \in A\}$, then even the intersection $Q_A \cap B_A(x)$ is nonempty, compact, and convex.

To be more precise than above, the first statement of the theorem is due to Minty, its extension was proved by H. Debrunner and P. Flor [7] in 1964, and the proof of Minty's maximality theorem given in H. Brezis [3, Theorem 2.1, p. 23f] is based on their extension.

For l_p^2 , $1 < p < \infty$, $p \neq 2$, Crandall and Liggett considered the following operator A : Let $\{e_1, e_2\}$ be the natural basis in l_p^2 , and let

$$A := \{(0, 0), (e_1, e_2), (e_2, -e_1)\}.$$

They pointed out that no maximally accretive extension of A on l_p^2 is defined on the triangle $\{(\xi_1, \xi_2) \in l_p^2 : 0 < \xi_1 < \xi_2, \xi_1 + \xi_2 < 1\}$ if $1 < p < 2$, respectively on the triangle $\{(\xi_1, \xi_2) \in l_p^2 : 0 < \xi_2 < \xi_1, \xi_1 + \xi_2 < 1\}$ if $2 < p < \infty$, in contrast to the fact that the closure of an m -accretive operator on l_p^2 is convex, see Theorem 5.1 below.

In [2] the authors extended their result as follows:

(4.2) *For the plane endowed with a strictly convex and smooth norm the class of maximally accretive operators coincides with the class of m -accretive ones exactly when the norm generates an inner product.*

The stronger statement: In the normed plane the two classes coincide exactly when the unit ball is either an ellipse or a parallelogram, as well as the obvious extension for the n -space, are still open, see also the comments of Crandall and Liggett, loc. cit., on this subject.

We want to conclude the section with two statements on maximally accretive operators on the normed plane, which are in the vein of our paper.

Via the quadratic and skew-symmetric form

$$S(x, y) := \eta_1 \xi_2 - \xi_1 \eta_2 \quad x = (\xi_1, \xi_2) \text{ and } y = (\eta_1, \eta_2)$$

we define the so-called dual $*$ -norm

$$\forall x \quad \|x\|^* := \sup\{S(x, y) : \|y\| \leq 1\}.$$

The unit ball of the plane w.r.t. the *dual $*$ -norm* is just the unit ball w.r.t. the dual norm rotated by 90° , and $\|\cdot\|^{**} = \|\cdot\|$. The following lemma, due to H. Busemann, is of interest in connection with accretiveness on the plane.

(4.3) LEMMA.

$$\forall x \quad \langle y, x \rangle_s \geq 0 \Leftrightarrow \langle x, y \rangle_{s^*} \geq 0.$$

We have

(4.4) *Let A be a maximally accretive operator on the plane.*

$$\forall x \in D(A) \quad A(x) \text{ is closed and } \|\cdot\|^* \text{-convex.}$$

(4.5) *Let A be a maximally accretive operator on the plane.*

If A is defined on the whole plane then A is m -accretive.

We do not want to give formal proofs of the two propositions. The first one is not difficult to verify, while the second one follows from the fact that under the assumptions of the proposition A is upper semi-continuous and

$$\forall x \quad A(x) \text{ is nonempty, compact, and contractible in itself.}$$

5. On the domain and the range of an m -accretive operator. In the following A is assumed to be m -accretive on X . Hence for each λ the Yosida-resolvent $J_\lambda := (I + \lambda A)^{-1}$ defines a contraction on X .

Following H. Brezis [3, Theorem 2.2] and R. C. Bruck [4] it is not difficult to prove

(5.1) THEOREM. *Let A be m -accretive on X .*

$$(5.2) \quad \forall y \in X \quad \exists x \in \overline{D(A)} \ni 0 \leq \langle y - x, x - x' \rangle_s \quad \forall x' \in \overline{D(A)}.$$

Moreover, $\overline{D(A)}$ is $\|\cdot\|$ -convex.

If X is strictly convex, $\overline{D(A)}$ is just convex. If X is smooth, then $\overline{D(A)}$ is the range of a uniquely defined contractive projection, say $P_{\overline{D(A)}}$, and

$$\forall y \in X \quad \lim_{\lambda \rightarrow 0^+} J_\lambda y = P_{\overline{D(A)}} y = x.$$

Proof. Take an element $y \in X$.

$$\forall \lambda \in \mathbf{R}^+ \quad \exists (x_\lambda, a_\lambda) \in A \ni x_\lambda + \lambda a_\lambda = y.$$

By the accretiveness of A

$$0 \leq \langle a_\lambda - a', x_\lambda - x' \rangle_s \quad \forall (x', a') \in A,$$

or, multiplying the inequality by λ and replacing λa_λ by $y - x_\lambda$,

$$0 \leq \langle y - x_\lambda - \lambda a', x_\lambda - x' \rangle_s \quad \forall (x', a') \in A.$$

On the other hand, $\{x_\lambda\}_{\lambda > 0}$ is bounded for $\lambda \rightarrow 0 +$ —indeed, for each $(x', a') \in A$ $\|x_\lambda - x'\| \leq \|y - (x' + \lambda a')\|$. Since the semi-inner product is upper semi-continuous, for any accumulation point x of $\{x_\lambda\}_{\lambda > 0}$ for $\lambda \rightarrow 0 +$,

$$0 \leq \langle y - x, x - x' \rangle_s \quad \forall x' \in \overline{D(A)}.$$

Since $\{x_\lambda\}_{\lambda > 0} \subset D(A)$, $x \in \overline{D(A)}$.

The inequality (5.2) implies that

$$(5.3) \quad \|x - x'\|^2 \leq \langle y - x', x - x' \rangle_s$$

$$\leq \|y - x'\| \|x - x'\|, \quad \forall x' \in \overline{D(A)},$$

from which we easily derive that $\overline{D(A)}$ is $\|\cdot\|$ -convex. Indeed, let x' and x'' be elements of $\overline{D(A)}$. If $y = (x' + x'')/2 \in \overline{D(A)}$, we are done. If not, let $x \in \overline{D(A)}$ be such that (5.2) holds. By (5.3),

$$\|x - x'\| \leq \|y - x'\| \quad \text{and} \quad \|x - x''\| \leq \|y - x''\|,$$

but $\|y - x'\| = \|y - x''\| = \|x' - x''\|/2$ which implies that $x \neq x'$, x'' and that

$$\|x' - x''\| \leq \|x' - x\| + \|x'' - x\| \leq \|x' - x''\|,$$

x is consequently a proper intermediate point between x' and x'' in $\overline{D(A)}$.

If x is smooth then the semi-inner product is linear in its first variable. Let $y \in X$ and $x_1, x_2 \in D(A)$ be such that

$$0 \leq \langle y - x_i, x_i - x' \rangle_s$$

for all $x' \in \overline{D(A)}$, $i = 1, 2$. It follows that

$$\begin{aligned} \|x_1 - x_2\|^2 &= \langle x_1 - x_2, x_1 - x_2 \rangle_s \\ &= -\langle y - x_1, x_1 - x_2 \rangle_s - \langle y - x_2, x_2 - x_1 \rangle_s \leq 0. \end{aligned}$$

Thus for each $y \in X$ there exists at most one element $x \in \overline{D(A)}$ satisfying the inequality in (5.2). By (5.2) there exists such an element, namely,

$$\lim_{\lambda \rightarrow 0+} J_\lambda y = x.$$

□

Let K be a proper closed nonempty subset of X . In connection with his study of the fixed point set of contractive mappings, F. E. Browder introduced the so-called *approximation region* $A(y; K)$ between y ($\in X$) and K :

$$A(y; K) = \{z \in X: 0 \leq \langle y - z, z - k \rangle, \forall k \in K\}.$$

If for each $y \in X$, $A(y; K) \cap K \neq \emptyset$ then K is said to be a *co-sun*, a notion which was introduced by P. L. Papini and I. Singer in connection with problems within the theory of best approximation, see L. Hetzelt [10] and U. Westphal [14] for details. With use of this notion, for an m -accretive operator on X the closure of its domain is a co-sun.

We want to state a few facts about co-suns which seem to be of relevance in connection with accretive operators, see [9] for proofs.

(5.4) *A subset in \mathbf{R}^n is a co-sun exactly when it is closed and convex, and the metric projection onto it is the uniquely defined contractive retraction of \mathbf{R}^n onto it (F. O. L. Klore).*

(5.5) *A subset in the normed plane is a co-sun exactly when it is closed and $\|\cdot\|$ -convex which in turn is the range of a contractive ray retraction (L. A. Karlovitz, P. Gruber, L. Hetzelt).*

In 1941 F. Bohnenblust characterized those subspaces in l_n^p , $1 < p < \infty$, $p \neq 2$ which are the ranges of contractive linear projectinos. He proved, a hyper-subspace is the range of a contractive linear projection exactly when its normal vector contains at most two nonzero coefficients, and concluded that a subspace has this property when and only when it can be written as the intersection of such hyper-subspaces. The second named author extended Bohnenblust's characterization as follows.

(5.6) *Let U_n denote the set of unit vectors in \mathbf{R}^n which have at most two nonzero coefficients. A subset in l_n^p , $1 < p < \infty$, $p \neq 2$, is a co-sun exactly when it is the intersection of a family of closed half spaces the normal vectors of which belong to U_n .*

For l_n^1 and likewise for l_n^∞ those subspaces which are the ranges of linear contractions have been characterized, but as far as we know there are no descriptions of co-suns for these spaces.

Also there is not much known about the ranges of m -accretive operators.

If X is an inner product space and A a maximally monotone operator on X , so is A^{-1} and, consequently, the closure of $R(A)$ is convex. This result has its counterpart for the normed plane.

(5.7) *If A is accretive on the normed plane, then, by Lemma 4.2, A^{-1} is accretive with respect to the dual $\|\cdot\|$ -norm. Consequently, if A is m -accretive then $\overline{R(A)}$ is $\|\cdot\|$ -convex.*

Note added in proof. Professor S. Reich kindly pointed out to the authors that A. Cernès [Israel J. Math. **19** (1974), 335–48] already proved (4.2) even for n -spaces. In the plane he further verified that the two notions of accretiveness coincide exactly when the unit ball is either an ellipse or a parallelogram. In [J. Funct. Anal. **26** (1977), 378–95] S. Reich among others extended Cernès first statement to smooth spaces.

Finally, following Reich's ideas on approximating zeros it is not difficult to prove: *Let A be m -accretive on X ,*

$$\forall y \in X \exists a \in \overline{R(A)} \ni 0 \leq \langle a - a', y - a \rangle_s \quad \forall a' \in \overline{R(A)},$$

i.e., $\overline{R(A)}$ is a sun in the setting of best approximation.

REFERENCES

- [1] H. Berens and L. Hetzelt, *Die metrische Struktur der Sonnen in $l_\infty(n)$* , Aequationes Math., **27** (1984), 274–287.
- [2] ———, *On maximally accretive operators in the plane*, in *Anniversary Volume on Approximation Theory and Functional Analysis*, ed. by P. L. Butzer, R. L. Stens, and B. Sz. Nagy, ISNM **65**. Birkhäuser Verlag, Basel 1984, 109–116.
- [3] H. Brezis, *Opérateurs Maximaux Monotones*, Notas de Matemática (50). North-Holland Publ. Co., Amsterdam, London, 1973.
- [4] R. C. Bruck, *Nonexpansive projections on subsets of Banach spaces*, Pacific J. Math., **47** (1973), 341–355.
- [5] H. Busemann, *The Geometry of Geodesics*, Academic Press, New York 1955.
- [6] M. G. Crandall and T. M. Liggett, *A theorem and a counterexample in the theory of semigroups of nonlinear transformations*, Trans. Amer. Math. Soc., **160** (1971), 263–278.
- [7] H. Debrunner and P. Flor, *Ein Erweiterungssatz für monotone Mengen*, Arch. Math., **15** (1964), 445–447.
- [8] A. Granas and J. W. Jaworowski, *Some theorems on multi-valued mappings of subsets of the Euclidean space*, Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys., **7**, No. 5 (1959), 227–283.
- [9] L. Hetzelt, *Über die beste Coapproximation im \mathbb{R}^n* , Dissertation U Erlangen-Nürnberg 1981.
- [10] ———, *On suns and cosuns in finite dimensional normed vector spaces*, Acta Math. Acad. Sci. Hung., **45** (1–2) (1985), 53–68.

- [11] J. M. Lasry and R. Robert, *Analyse nonlinéaire multivoque*, Cahiers de Mathématiques de la Décision No. 7611, Université Paris IX, Dauphine 1980.
- [12] G. J. Minty, *Monotone (nonlinear) operators in Hilbert space*, Duke Math. J., **29** (1962), 341–346.
- [13] ———, *On the simultaneous solution of a certain system of linear inequalities*, Proc. Amer. Math. Soc., **13** (1962), 11–12.
- [14] U. Westphal, *Über die Existenz- und Eindeutigkeitsmengen bei der besten Ko-Approximation*, in *Linear Spaces and Approximation*, ed. by P. L. Butzer, E. Görlich and B. Sz.-Nagy, ISNM **60**, Birkhäuser Verlag Basel 1981, 225–264.

Received April 30, 1985.

UNIVERSITÄT ERLANGEN-NÜRNBERG
BISMARCKSTRASSE 1½
D-8520 ERLANGEN, W. GERMANY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720
H. SAMELSON
Stanford University
Stanford, CA 94305
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA	UNIVERSITY OF OREGON
UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA	UNIVERSITY OF HAWAII
MONTANA STATE UNIVERSITY	UNIVERSITY OF TOKYO
UNIVERSITY OF NEVADA, RENO	UNIVERSITY OF UTAH
NEW MEXICO STATE UNIVERSITY	WASHINGTON STATE UNIVERSITY
OREGON STATE UNIVERSITY	UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1986 by Pacific Journal of Mathematics

Dale Edward Alspach, On $\mathcal{L}_{p,\lambda}$ spaces for small λ	257
Jong Sook Bae and Sangsuk Yie, Range of Gateaux differentiable operators and local expansions	289
Hubert Berens and Lothar Hetzelt, On accretive operators on l_n^∞	301
Marie-Françoise Bidaut-Véron, Global existence and uniqueness results for singular solutions of the capillarity equation	317
Donald M. Davis and Mark Mahowald, Classification of the stable homotopy types of stunted real projective spaces	335
Aad Dijkema, Heinz K. Langer and Hendrik S. V. de Snoo, Unitary colligations in Π_κ -spaces, characteristic functions and Štraus extensions	347
Michel Enock and Jean-Marie Schwartz, Algèbres de Kac moyennables	363
Seppo Granlund, Peter Lindqvist and Olli Martio, Note on the PWB-method in the nonlinear case	381
Palle E. T. Jorgensen, Analytic continuation of local representations of Lie groups	397
Robert P. Kaufman, Plane curves and removable sets	409
José M. Montesinos and Wilbur Carrington Whitten, Constructions of two-fold branched covering spaces	415
Benedict Seifert, Highly transitive group actions on trees and normalizing Tits systems	447
Charles Stuart Stanton, Counting functions and majorization for Jensen measures	459
Luen-Fai Tam, On existence criteria for capillary free surfaces without gravity	469
Zhuocheng Yang, Exposed points of left invariant means	487