

# Pacific Journal of Mathematics

## **CLASSIFICATION OF THE STABLE HOMOTOPY TYPES OF STUNTED REAL PROJECTIVE SPACES**

DONALD M. DAVIS AND MARK MAHOWALD

# CLASSIFICATION OF THE STABLE HOMOTOPY TYPES OF STUNTED REAL PROJECTIVE SPACES

DONALD M. DAVIS AND MARK MAHOWALD

Let  $P_n^{n+k}$  denote the stunted real projective space  $RP^{n+k}/RP^{n-1}$ . In this paper we complete the classification of stable homotopy type of stunted real projective spaces begun by Feder, Gitler, and Mahowald in 1977.

Let  $\phi(m, n)$  denote the number of integers  $j$  satisfying (a)  $j \equiv 0, 1, 2$ , or  $4 \pmod{8}$  and (b)  $m \leq j \leq n$  if  $m \not\equiv 0 \pmod{4}$ , or  $m < j \leq n$  if  $m \equiv 0 \pmod{4}$ . Let  $\phi(n) = \phi(1, n)$ , and let  $\nu(n)$  denote the exponent of 2 in  $n$ .

**THEOREM 1.**  $P_n^{n+k}$  and  $P_m^{m+k}$  have the same stable homotopy type ( $\sim$ ) if and only if  $\nu(n - m)$

$$\geq \begin{cases} \phi(k) & \text{if } k \leq 8 \text{ or } \nu(n) \geq \phi(k) - 1 \text{ or } \nu(n + k + 1) \geq \phi(k) - 1 \\ \max\{\phi(n, n + k) - 1, \phi(-n - k - 1, -n - 1) - 1\} & \text{otherwise.} \end{cases}$$

This max will equal either  $\phi(k)$  or  $\phi(k) - 1$ . A mindless restatement of Theorem 1 is given by the following result.

**THEOREM 1'.**  $P_m^{m+k} \sim P_n^{n+k}$  iff  $\nu(n - m) \geq \phi(k) - \varepsilon(m, k)$ , where  $\varepsilon(m, k)$  is either 0 or 1. In the following cases  $\varepsilon(m, k)$  is 0; otherwise it is 1.

- (i)  $k \leq 8$
- (ii)  $\nu(m) \geq \phi(k) - 1$
- (iii)  $\nu(m + k + 1) \geq \phi(k) - 1$ .
- (iv) The congruence classes of  $m \pmod{4}$  and of  $k \pmod{8}$  are such that  $a *$  occurs in the following table.

$m \equiv$	$k \equiv$	0	1	2	3	4	5	6	7
0									
1		*	*		*				*
2		*						*	*
3					*		*	*	*

Readers familiar with the spectrum  $J(= \text{fibre}(bo \rightarrow \Sigma^4 bsp))$  will appreciate the following corollary.

**COROLLARY 2.** *Stunted real projective spaces have the same stable homotopy type if and only if their  $J$ -homology and  $J$ -cohomology groups are isomorphic as graded abelian groups (with dimension shift).*

This can be established by comparing Theorem 1 with the calculations in [11] of  $J_*(P_n^{n+k})$ . Here we use Whitehead's theorem ([13]) that  $h_*(X) \approx h^*(DX)$ , where  $DX$  is the  $S$ -dual of  $X$ , to express  $J_*(P_{-n-k-1}^{-n-1})$  as  $J^*(P_n^{n+k})$ .

[7] had conjectured Theorem 1 and had established it in certain congruences. (Their table has a misprint when  $n$  is odd and  $k \equiv 2$ .) Our proof follows the approach of [7].

That  $\nu(n-m) \geq \phi(k)$  is sufficient for  $P_m^{m+k} \sim P_n^{n+k}$  is clear from [3] since the spaces are then Thom complexes of  $J$ -equivalent vector bundles. That  $\nu(n-m) \geq \phi(k)$  when  $k \leq 8$  is necessary is easily deduced from the action of the Steenrod algebra. That  $\nu(n-m) \geq \max$  is necessary was proved using Adams operations in [9] and [7; 2.1]. That  $\nu(n-m) \geq \phi(k)$  when  $\nu(n) \geq \phi(k) - 1$  or  $\nu(n+k+1) \geq \phi(k) - 1$  is necessary was stated in [7; 1.1] and proved using Adams operations in [8].

We will prove that

- (i)  $P_{8a}^{8a+8b+3} \sim P_{8a'}^{8a'+8b+3}$  if  $\nu(8a) < 4b+1 \leq \nu(8a-8a')$ , and
- (ii)  $P_{4a}^{4a+8b+7} \sim P_{4a'}^{4a'+8b+7}$  if  $\max\{\nu(4a), \nu(4a+8b+8)\} < 4b+2 \leq \nu(4a-4a')$ .

Of course the equivalence of (ii) also implies equivalences if cells are removed from the bottom or top. This, and an application of  $S$ -duality to (i), then handles all cases unresolved in [7].

We begin with (i). Write  $8a' = 8a + c \cdot 2^{4b+1}$  and let  $L$  be such that  $\nu(L) \geq 4b+2$  and  $L > 8a' + 8b + 4$ . The multiple  $(L - 8a - 8b - 4)\xi_{8b+3}$  of the Hopf bundle is classified by the composite

$$P^{8b+3} \xrightarrow{\Delta} P^{8b+3} \times P^{8b+3} \xrightarrow{c \cdot 2^{4b+1}\xi \times d\xi} BO_{c \cdot 2^{4b+1}} \times BO_d \rightarrow BO_{L-8a-8b-4},$$

where  $d = L - 8a' - 8b - 4$ . Since  $2^{4b+1}\xi_{8b+1}$  is trivial and  $\pi_{8b+3}(BO) = 0$ , the first factor of the middle map can be factored as

$$P^{8b+3} \xrightarrow{c} P_{8b+2}^{8b+3} = S^{8b+2} \vee S^{8b+3} \xrightarrow{c} S^{8b+2} \xrightarrow{\theta} BO_{c \cdot 2^{4b+1}}$$

with  $\theta$  the nonzero element of  $\pi_{8b+2}(BO_{c \cdot 2^{4b+1}})$ . The induced map of Thom spaces is

$$P_{L-8a-8b-4}^{L-8a-1} \rightarrow \left( S^{c \cdot 2^{4b+1}} \cup_{\beta} e^{c \cdot 2^{4b+1} + 8b+2} \right) \wedge P_{L-8a'-8b-4}^{L-8a'-1},$$

where  $\beta$  generates the image of  $J$  in the  $(8b + 1)$ -stem. If the map is made skeletal, we obtain

$$(*) \quad P_{L-8a-8b-4}^{L-8a-1} \rightarrow \Sigma^{c \cdot 2^{4b+1}} \left( P_{L-8a'-8b-4}^{L-8a'-1} \cup_{\beta \vee \beta} (e^{L-8a'-2} \vee e^{L-8a'-1}) \right),$$

where the attaching map is

$$\begin{aligned} S^{L-8a'-3} \vee S^{L-8a'-2} &\xrightarrow{\beta \vee \beta} S^{L-8a'-8b-4} \vee S^{L-8a'-8b-3} \\ &= P_{L-8a'-8b-4}^{L-8a'-3} \hookrightarrow P_{L-8a'-8b-4}^{L-8a'-1}. \end{aligned}$$

We will prove later

**PROPOSITION 3.** *If  $\nu(4A + 8b + 4) < 4b + 1$ , then the composite*

$$S^{4A+8b+2} \xrightarrow{\beta} S^{4A+1} \hookrightarrow S^{4A} \vee S^{4A+1} = P_{4A}^{4A+1} \hookrightarrow P_{4A}^{4A+8b+2}$$

*is null-homotopic, where  $\beta$  generates the image of  $J$ .*

The hypotheses of (i) imply  $\nu(L - 8a') < 4b + 1$ , so that Proposition 3 implies that the  $e^{L-8a'-1}$  of the target in  $(*)$  splits off, giving us a map

$$P_{L-8a-8b-4}^{L-8a-1} \xrightarrow{g} \Sigma^{c \cdot 2^{4b+1}} \left( P_{L-8a'-8b-4}^{L-8a'-1} \cup_{\beta} e^{L-8a'-2} \right).$$

That this map is injective in  $H^*(; \mathbf{Z})$  on the projective space is clear for all except the top cell; that it is also true on the top cell follows from the argument of [7; 3.1]. This is proved by using the diagram

$$\begin{array}{ccc} \Sigma^{c \cdot 2^{4b+1}} \left( P_d^{L-8a'-2} \cup e^{L-8a'-2} \cup e^{L-8a'-1} \right) & \xrightarrow{r} & \Sigma^{c \cdot 2^{4b+1}} \left( P_d^{L-8a'-2} \cup e^{L-8a'-2} \right) \\ \downarrow & & \downarrow \\ P_{L-8a-8b-4}^{L-8a-1} & \xrightarrow{g'} \Sigma^{c \cdot 2^{4b+1}} \left( P_d^{L-8a'-1} \cup e^{L-8a'-2} \cup e^{L-8a'-1} \right) & \xrightarrow{\hat{r}} \Sigma^{c \cdot 2^{4b+1}} \left( P_d^{L-8a'-1} \cup e^{L-8a'-2} \right), \\ & \underbrace{\hspace{15em}}_{g} & \end{array}$$

where  $r$  and  $\hat{r}$  are retractions for the splitting, and  $r$  exists because the dimension of the target in Proposition 3 is one less than one might have naively required.  $g'^*$  sends both classes in  $H^{L-8a-1} (; \mathbf{Z})$  nontrivially, but  $\hat{r}^*( )$  hits only the first.

The  $S$ -dual of  $g$  is

$$(**) \quad \left( S^{8a'+1} \vee P_{8a'}^{8a'+8b+2} \right) \cup_{\beta \vee \alpha} e^{8a'+8b+3} \xrightarrow{f} \Sigma^{c \cdot 2^{4b+1}} P_{8a}^{8a+8b+3},$$

where  $\beta$  as before generates the image of  $J$  and  $\alpha$  is the attaching map for the top cell of  $P_{8a'}^{8a'+8b+3}$ . Now

$$\pi_{8a'+1} \left( \Sigma^{c \cdot 2^{4b+1}} P_{8a}^{8a+8b+3} \right) \approx \pi_{8a+1} \left( P_{8a}^{8a+8b+3} \right) \approx \mathbf{Z}_2 \oplus \mathbf{Z}_2,$$

with one generator  $\eta$  on the bottom cell and the other of degree 1 on the next-to-bottom cell. By Proposition 3,  $\beta$  composed with the latter class is 0, while  $\beta$  composed with  $\eta$  is 0. Thus the restriction of the map  $f$  of  $(**)$  to  $S^{8a'+1}$  preceded by  $\beta$  is trivial. The existence of  $f$  says that

$$(f | S^{8a'+1} \vee P_{8a'}^{8a'+8b+2}) \circ (\beta \vee \alpha) = 0.$$

Subtracting, we deduce that  $f | P_{8a'}^{8a'+8b+2}$  extends over  $P_{8a'}^{8a'+8b+3}$  to give the desired map, which is certainly an equivalence below the top cell and at the odd primes, and has odd degree on the top cell by  $Sq^4$ .

The argument for (ii) is similar, but slightly more elaborate. Let  $4a' = 4a + c \cdot 2^{4b+2}$  and  $L > 4a' + 8b + 8$  with  $\nu(L) \geq 4b + 3$ . A map

$$P_{L-4a-8b-8}^{L-4a-1} \rightarrow \left( \Sigma^{c \cdot 2^{4b+2}} \cup_{\beta} e^{c \cdot 2^{4b+2}+8b+4} \right) \wedge P_{L-4a'-8b-8}^{L-4a'-1},$$

with  $\beta$  a generator of the image of  $J$  in the  $(8b + 3)$ -stem, is constructed as before. This yields

$$\begin{aligned} & P_{L-4a-8b-8}^{L-4a-1} \\ & \xrightarrow{\Psi} \Sigma^{c \cdot 2^{4b+2}} \left( P_{L-4a'-8b-8}^{L-4a'-1} \cup_{\beta \vee \beta \vee \beta} \left( e^{L-4a'-4} \vee CM \vee e^{L-4a'-1} \right) \right) \end{aligned}$$

where  $CM$  is the cone on  $M = S^{L-4a'-4} \cup_2 e^{L-4a'-3}$ . We prove later

**PROPOSITION 4.** *If  $\beta$  generates the image of  $J$  in the  $(8b + 3)$ -stem, the composite*

$$S^{4A+8b+4} \cup_2 e^{4A+8b+5} \xrightarrow{\beta \wedge 1} P_{4A+1}^{4A+2} \rightarrow S^{4A} \vee P_{4A+1}^{4A+2} = P_{4A}^{4A+2} \hookrightarrow P_{4A}^{4A+8b+6}$$

*is null-homotopic.*

**PROPOSITION 5.** *If  $\nu(4A + 8b + 8) < 4b + 2$ , the composite*

$$S^{4A+8b+6} \xrightarrow{\beta} S^{4A+3} \hookrightarrow S^{4A} \vee P_{4A+1}^{4A+2} \vee S^{4A+3} = P_{4A}^{4A+3} \hookrightarrow P_{4A}^{4A+8b+6}$$

*is null-homotopic.*

These propositions imply that in the map  $\Psi$  above the top 2 parts of the 3-part wedge split off, and as before the resulting map is injective in  $H^*(\ ; \mathbf{Z})$  on the projective space. The dual of the resulting map is analogous to (\*\*):

$$\left( S^{4a'+3} \vee P_{4a'}^{4a'+8b+6} \right) \cup_{\beta \vee \alpha} e^{4a'+8b+7} \xrightarrow{f} \Sigma^{c \cdot 2^{4b+1}} P_{4a}^{4a+8b+7}.$$

An argument similar to that applied to (\*\*), using Proposition 5, shows that  $f|P_{4a'}^{4a'+8b+6}$  extends over  $P_{4a'}^{4a'+8b+7}$ , as desired. This extension induces an isomorphism in the top cohomology group because  $f^*$  does, while the extension

$$S^{4a'+3} \cup_{\beta} e^{4a'+8b+7} \rightarrow \Sigma^{c \cdot 2^{4b+2}} P_{4a}^{4a+8b+7} \quad \text{of } f|S^{4a'+3}$$

does not since by Proposition 5 the null-homotopy can be factored through  $\Sigma^{c \cdot 2^{4b+2}} P_{4a}^{4a+8b+6}$ .

*Proof of Proposition 3.* We let  $N = 4A + 8b + 4$ . If  $A = 2\bar{A}$  is even, then the argument is straightforward:  $\beta$  has filtration  $4b$  by [11; 8.2] and is annihilated by  $\eta$ . But  $\pi_{8\bar{A}+8b+2}(P_{8\bar{A}}^{8\bar{A}+8b+2})$  in filtration  $\geq 4b$  is acted on injectively by  $\eta$ . To see this, we use charts of [12] to get


$$\begin{array}{c} \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \quad \diagup \quad \diagdown \\ \bullet \quad \bullet \quad \bullet \quad \bullet \\ \diagup \quad \diagdown \quad \diagup \quad \diagdown \quad \diagup \quad \diagdown \\ \bullet \quad \bullet \quad \bullet \quad \bullet \quad \bullet \end{array} \\ \uparrow \\ t - s = 8\bar{A} + 8b + 2 \end{array} \quad \leftarrow s = 4b$$

and follow into  $J_*(\ )$  to see that none of these classes is hit by a differential.

Thus we may assume  $N \equiv 0 \pmod{8}$ . In the diagram

$$\begin{array}{ccc} S^{N-8b-5} \cup_{\eta} e^{N-8b-3} & \xrightarrow{i_1} & P_{N-8b-5}^{N-2} \\ c \downarrow & & c \downarrow \\ S^{N-8b-3} & \xrightarrow{i_2} & P_{N-8b-4}^{N-2} \end{array}$$

the charts for  $\text{Ext}_{\mathcal{A}}^{s,t}(H^*(\ ), \mathbf{Z}_2)$  in  $s \geq 4b - 1$ ,  $N - 3 \leq t - s \leq N$ , are



Here the maps denoted  $c$  are collapse maps, and the horizontal inclusions maps are of cells near the bottom of the stunted projective spaces. The Ext charts are taken from the tables of [12]. That they are as claimed can be proved by the Adams periodicity theorem ([1]). The chart in the upper left is obtained from  $\text{Ext}_A^{*,*}(\mathbb{Z}_2, \mathbb{Z}_2) + \text{Ext}_A^{*,*+2}(\mathbb{Z}_2, \mathbb{Z}_2)$  by inserting differentials between classes related by  $h_1$ .

In these charts, all homomorphisms in  $(t-s, s) = (N-2, 4b)$  are isomorphisms. This can be deduced from [12] or by low-level calculation and periodicity. It is proved in [11; 8.2] that the generator of the image of  $J$  in  $\pi_{N-2}(S^{N-8b-3})$  is represented by the element in  $(N-2, 4b)$  in the chart in the lower left. Thus the composite of Proposition 3 is represented by the element in  $(N-2, 4b)$  in the chart in the lower right. The following lemma implies that this class must be 0 in  $\pi_*(P_{N-8b-4}^{N-2})$ , proving Proposition 3. (The desired class cannot be nonzero in higher filtration since it is annihilated by  $\eta$ , but the higher-filtration classes are not.)

**LEMMA 6.** *The image-of- $J$  generator of  $\pi_{N-2}(S^{N-8b-3})$  coextends to a filtration- $4b$  element of  $\pi_{N-2}(S^{N-8b-5} \cup_{\eta} e^{N-8b-3})$ , and if  $3 \leq v(N) \leq 4b$ , then  $i_{1*}$  sends this class to 0 in  $\pi_{N-2}(P_{N-8b-5}^{N-2})$ .*

*Proof.* The first part is equivalent to the dual statement—that there is a map  $S^{8b+1} \cup_{\eta} e^{8b+3} \rightarrow S^0$  of filtration  $4b$  whose restriction to  $S^{8b+1}$  generates the image of  $J$ . By [11; 8.2]  $\beta_{8b+1}$  lifts to  $E_{4b}$ , the  $4b$ th stage of an Adams resolution of  $S^0$ . Since  $\pi_{8b+2}(SO) = 0$ ,  $\eta\beta_{8b+1} = 0$ . The only

way that the composite

$$S^{8b+2} \xrightarrow{\eta} S^{8b+1} \rightarrow E_{4b}$$

might fail to be 0 is if in the Adams spectral sequence (ASS) for  $\pi_*(S^0)$  there was a nonzero differential  $E_r^{s,s+8b+3} \rightarrow E_r^{s+r,s+r+8b+2}$  with  $s < 4b < s + r$ . Since the only possible target element ( $s + r = 4b + 2$ ) survives to a homotopy class, the first half of the lemma is established.

We will prove

**LEMMA 7.** *If  $3 \leq 4c - 3 \leq \nu(N) \leq 4c$ , then*

$$\pi_{N-2}(P_{N-8c-5}^{N-2}) \xrightarrow{\eta_*} \pi_{N-1}(P_{N-8c-5}^{N-2})$$

*is injective on elements of filtration  $\geq 4c$ , or equivalently (by the previous ASS chart (C1)) the element in  $(N - 2, 4c)$  is hit by a differential in the ASS.*

From this we now easily deduce the second half of Lemma 6, and hence Proposition 3. The hypothesis of Lemma 6 says that  $\nu(N)$  can be written as in Lemma 7 for some positive  $c \leq b$ . Let

$$P_{N-8c-5}^{N-2} \rightarrow P_{N-8b-5}^{N-2}$$

be the composite of  $b - c$  of the filtration 4 maps considered in [10], [4], or [5]. As these induce an isomorphism in Ext near the upper edge, we use Lemma 7 to deduce that the element in  $(N - 2, 4b)$  of  $\text{ASS}(P_{N-8b-5}^{N-2})$  is hit by a differential, implying the second half of Lemma 6.

The proof of Lemma 7 has two main parts. The first is to prove that if  $\nu(N) = 4c - \varepsilon$ , then there is a commutative diagram as below with  $\beta$  a generator of the image of  $J$  and  $i\tilde{\beta} \simeq *$ . The case  $\varepsilon = 2$  is representative,

$  \begin{array}{ccccc}  \varepsilon=0 & & & & \\  & P_{N-8c-3}^{N-8c-2} & \xrightarrow{i} & P_{N-8c-3} & \\  \tilde{\beta} \nearrow & \downarrow & & \downarrow & \\  S^{N-2} \xrightarrow{\beta} & S^{N-8c-2} & \rightarrow & P_{N-8c-2} & \\  \\  \varepsilon=2 & & & & \\  & P_{N-8c-1}^{N-8c+3} & \xrightarrow{i} & P_{N-8c-1} & \\  \tilde{\beta} \nearrow & \downarrow & & \downarrow & \\  S^{N-2} \xrightarrow{\beta} & S^{N-8c+3} & \rightarrow & P_{N-8c+3} &  \end{array}  $	$  \begin{array}{ccccc}  \varepsilon=1 & & & & \\  & P_{N-8c-2}^{N-8c-1} & \xrightarrow{i} & P_{N-8c-2} & \\  \tilde{\beta} \nearrow & \downarrow & & \downarrow & \\  S^{N-2} \xrightarrow{\beta} & S^{N-8c-1} & \rightarrow & P_{N-8c-1} & \\  \\  \varepsilon=0 & & & & \\  & P_{N-8c+3}^{N-8c+5} & \xrightarrow{i} & P_{N-8c+3} & \\  \tilde{\beta} \nearrow & \downarrow & & \downarrow & \\  S^{N-2} \xrightarrow{\beta} & S^{N-8c+5} & \rightarrow & P_{N-8c+5} &  \end{array}  $
--	--






preceding  $\varepsilon = 2$  diagram, and show that  $\tilde{\beta}$  is detected in filtration  $4c - 3$ , from which Ext charts show that the same is true of  $b$ .

This analysis of the filtration of  $\tilde{\beta}$  is similar to the proof of the first half of Lemma 6. We prove the dual result—if  $E_{4c-3}$  is the indicated stage of a minimal Adams resolution of  $S^0$ , then the lifting to  $E_{4c-3}$  of the generator of the image of  $J$  extends to  $\gamma$  as indicated

$$\begin{array}{ccc} \Sigma^{-1}P_{8c-4}^{8c} & \xrightarrow{\gamma} & E_{4c-3} \\ \uparrow & \nearrow & \downarrow \\ S^{8c-5} & \xrightarrow{\beta} & S^0. \end{array}$$

This is proved by noting that a chart for  $\pi_*(E_{4c-3})$  begins



and so the composite

$$S^{8c-3} \cup_2 e^{8c-2} \xrightarrow{c} S^{8c-2} \xrightarrow{v} S^{8c-5} \rightarrow E_{4c-3}$$

is trivial.  $\square$

The proof of Proposition 4 is much easier. We just need to note that  $\beta \wedge 1$  has filtration  $\geq 4b + 1$  by [11; 8.2] and that  $\text{Ext}_{\mathcal{A}}^{s,t}(H * P_{4A}, \mathbf{Z}_2) = 0$  if  $4A + 8b + 4 \leq t - s \leq 4A + 8b + 5$  and  $s \geq 4b$  by the tables of [12] and Adams periodicity.

We complete the paper by giving the proof of Proposition 5, which is very similar to that of Proposition 3. The following result, which implies Proposition 5, is analogous to Lemma 6. As before,  $N \equiv 0 \pmod{8}$ .

**LEMMA 8.** *The image-of- $J$  generator of  $\pi_{N-2}(S^{N-8b-5})$  coextends to a filtration  $(4b + 1)$  element of  $\pi_{N-2}(P_{N-8b-9}^{N-8b-5})$ , and if  $v(N) \leq 4b + 1$  then this class goes to 0 in  $\pi_{N-2}(P_{N-8b-9}^{N-2})$ .*

The first half of Lemma 8 was proved above, in the consideration of  $\tilde{\beta}$  when  $\varepsilon = 2$ . The second half of Lemma 8 is proved as before by applying filtration 4 maps to the following analogue of Lemma 7.

LEMMA 9. If  $4c - 2 \leq \nu(N) \leq 4c + 1$ , then the elements in  $\pi_{N-2}(P_{N-8c-9}^{N-2})$  of filtration  $\geq 4c + 1$  are hit by differentials in the ASS.

The first step of the proof of Lemma 9 is to show that if  $\nu(N) = 4c + \varepsilon$ , then there is a commutative diagram as below with  $\beta$  a generator of the image of  $J$  and  $i\tilde{\beta} \simeq *$ . The proof mimics that of the analogous previous result.

$$\begin{array}{ccccc}
 & P_{N-8c-l}^{N-8c-k} & \xrightarrow{i_1} & P_{N-8c-l} & \\
 \tilde{\beta} \nearrow & \downarrow & & \downarrow & \\
 S^{N-2} & \xrightarrow{\beta} & S^{N-8c-k} & \rightarrow & P_{N-8c-k}
 \end{array}
 \quad
 \begin{array}{c|cc}
 \varepsilon & k & l \\
 \hline
 1 & 3 & 7 \\
 0 & 2 & 6 \\
 -1 & 1 & 5 \\
 -2 & -3 & 1
 \end{array}$$

The second half of the proof of Lemma 9 mimics that of Lemma 7. We shall consider the case  $\varepsilon = -1$ , which contains one subtle point that should have been addressed in the case  $\varepsilon = 1$  of the proof of Lemma 7. The filtration of  $\beta_{8c-1}$  is better behaved in the Moore space than in the sphere. Therefore we modify the above diagram in this case to

$$\begin{array}{ccccc}
 & P_{N-8c-5}^{N-8c} & \xrightarrow{i_1} & P_{N-8c-5} & \\
 \tilde{\beta} \nearrow & \downarrow & & \downarrow & \\
 S^{N-2} & \xrightarrow{\beta} & S_{N-8c-1}^{N-8c} & \hookrightarrow & P_{N-8c-1}
 \end{array}$$

Once we have shown that  $\tilde{\beta}$  is nontrivial in filtration  $4c - 2$ , then the analogue of (\*\*\*) can be used to complete the proof. It follows from [11; 8.3] that  $\beta$  has filtration exactly  $4c - 2$ ; to show the same is true of  $\tilde{\beta}$ , we dualize, so that we must show  $\Sigma^{-1}P_{8c-1}^{8c} \rightarrow E_{4c-2}$  extends over  $\Sigma^{-1}P_{8c-1}^{8c+4}$ , or equivalently that  $\Sigma^{-2}P_{8c+1}^{8c+4} \xrightarrow{\alpha} \Sigma^{-1}P_{8c-1}^{8c} \rightarrow E_{4c-2}$  is trivial. We calculate that all elements of  $[\Sigma^{-2}P_{8c+1}^{8c+4}, E_{4c-2}]$  of positive filtration map nontrivially to  $[\Sigma^{-2}P_{8c+1}^{8c+4}, S^0]$ , but here our map is certainly trivial since it is the obstruction to existence of an existing map, namely the dual of  $\tilde{\beta}$ .

## REFERENCES

- [1] J. F. Adams, *A periodicity theorem in homological algebra*, Proc. Camb. Phil. Soc., **62** (1966), 365–377.
- [2] ———, *On the groups  $J(X)$  – IV*, Topology, **5** (1966), 21–71.
- [3] M. F. Atiyah, *Thom complexes*, Proc. London Math. Soc. **11**, (1961), 291–310.
- [4] D. M. Davis, S. Gitler, and M. Mahowald, *The stable geometric dimension of vector bundles over real projective spaces*, Trans. Amer. Math. Soc., **268** (1981), 39–62.
- [5] D. M. Davis, M. Mahowald, and H. Miller, *Mapping telescopes and  $K_*$ -localizations*, to appear in Proc. John Moore Conf.

- [6] S. Feder and S. Gitler, *Stable homotopy type of Thom complexes*, Quart. J. Math., Oxford, **25** (1974), 143–149.
- [7] S. Feder, S. Gitler, and M. Mahowald, *On the stable homotopy type of stunted projective spaces*, Bol. Soc. Mat. Mex., **22** (1977), 1–5.
- [8] T. Kobayashi, *Stable homotopy types of stunted real projective spaces*, Proc. Amer. Math. Soc., **87** (1983), 555–556.
- [9] T. Kobayashi and M. Sugawara, *KT-rings of lens spaces  $L^n(4)$* , Hiroshima Math. J., **1** (1971), 253–271.
- [10] K. Y. Lam, *KO-equivalences and existence of nonsingular bilinear maps*, Pacific J. Math., **82** (1979), 145–153.
- [11] M. Mahowald, *The image of  $J$  in the EHP sequence*, Annals of Math., **116** (1982), 65–112.
- [12] ———, *The metastable homotopy of  $S^n$* , Memoirs of Amer. Math. Soc., **72** (1967).
- [13] G. W. Whitehead, *Generalized homology theories*, Trans. Amer. Math. Soc., **102** (1962), 227–283.

Received July 22, 1985. The authors were supported by the National Science Foundation at the University of Washington while this work was performed.

LEHIGH UNIVERSITY  
BETHLEHEM, PA 18015

AND

NORTHWESTERN UNIVERSITY  
EVANSTON, IL 60201



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024  
HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112  
R. FINN  
Stanford University  
Stanford, CA 94305

HERMANN FLASCHKA  
University of Arizona  
Tucson, AZ 85721  
RAMESH A. GANGOLLI  
University of Washington  
Seattle, WA 98195  
VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720  
ROBION KIRBY  
University of California  
Berkeley, CA 94720

C. C. MOORE  
University of California  
Berkeley, CA 94720  
H. SAMELSON  
Stanford University  
Stanford, CA 94305  
HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS      E. F. BECKENBACH      B. H. NEUMANN      F. WOLF      K. YOSHIDA  
(1906–1982)

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA	UNIVERSITY OF OREGON
UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA	UNIVERSITY OF HAWAII
MONTANA STATE UNIVERSITY	UNIVERSITY OF TOKYO
UNIVERSITY OF NEVADA, RENO	UNIVERSITY OF UTAH
NEW MEXICO STATE UNIVERSITY	WASHINGTON STATE UNIVERSITY
OREGON STATE UNIVERSITY	UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1986 by Pacific Journal of Mathematics

Dale Edward Alspach, On $\mathcal{L}_{p,\lambda}$ spaces for small $\lambda$	257
Jong Sook Bae and Sangsuk Yie, Range of Gateaux differentiable operators and local expansions	289
Hubert Berens and Lothar Hetzelt, On accretive operators on $l_n^\infty$	301
Marie-Françoise Bidaut-Véron, Global existence and uniqueness results for singular solutions of the capillarity equation	317
Donald M. Davis and Mark Mahowald, Classification of the stable homotopy types of stunted real projective spaces	335
Aad Dijkzma, Heinz K. Langer and Hendrik S. V. de Snoo, Unitary colligations in $\Pi_\kappa$ -spaces, characteristic functions and Štraus extensions	347
Michel Enock and Jean-Marie Schwartz, Algèbres de Kac moyennables	363
Seppo Granlund, Peter Lindqvist and Olli Martio, Note on the PWB-method in the nonlinear case	381
Palle E. T. Jorgensen, Analytic continuation of local representations of Lie groups	397
Robert P. Kaufman, Plane curves and removable sets	409
José M. Montesinos and Wilbur Carrington Whitten, Constructions of two-fold branched covering spaces	415
Benedict Seifert, Highly transitive group actions on trees and normalizing Tits systems	447
Charles Stuart Stanton, Counting functions and majorization for Jensen measures	459
Luen-Fai Tam, On existence criteria for capillary free surfaces without gravity	469
Zhuocheng Yang, Exposed points of left invariant means	487