

Pacific Journal of Mathematics

**REALIZING CERTAIN POLYNOMIAL ALGEBRAS AS
COHOMOLOGY RINGS OF SPACES OF FINITE TYPE
FIBERED OVER $\times BU(d)$**

LARRY SMITH

REALIZING CERTAIN POLYNOMIAL ALGEBRAS AS COHOMOLOGY RINGS OF SPACES OF FINITE TYPE FIBERED OVER $\times BU(d)$

LARRY SMITH

The problem of constructing topological spaces whose cohomology ring with coefficients in the field of p elements is a polynomial algebra has attracted the attention of algebraic topologists for many decades. Apart from the naturally occurring examples, classifying spaces of Lie groups away from their torsion primes, rather little progress was made until the construction of Clark and Ewing of a vast number of new non-modular examples. The completeness of their construction in the non-modular case was shown by Adams and Wilkerson (see Smith and Switzer for a compact-proof). One interest in the construction of spaces with polynomial cohomology is that they are related to the study of finite H -spaces, which should appear as their loop spaces; "should" because the construction of Clark and Ewing does not yield a simply connected CW complex of finite type. On the contrary the construction of Clark and Ewing yields non-simply connected spaces that are p -adically complete. By forming their finite completion they can be made simply connected. But considerably more effort would be required to show that they have the homotopy type of the p -completion of a simply connected CW complex of finite type.

We will avoid these drawbacks by constructing for certain of the examples of Clark and Ewing a simply connected space of finite type with the requisite cohomology.

Recall that the construction of [6] depends on a group $G < GL(V)$, $V = \bigoplus_n \mathbb{F}_p$, where \mathbb{F}_p is the field of p -elements, and which satisfies: G is generated by pseudo reflections and $|G| \not\equiv O(p)$. A theorem of Chevalley [2; V §5 no. 5, 3, Thm. 3] [24] shows that:

$$P(V^*)^G \simeq P[\rho_1, \dots, \rho_n]$$

where $P(V^*)$ denotes the ring of polynomials on the dual vector space V^* of V upon which G acts, and $P(V^*)^G \leq P(V^*)$ is the ring of invariant polynomials. We will say that $P(V^*)^G$ satisfies the *weak splitting principle* iff we can find polynomial generators $\rho_1, \dots, \rho_n \in P(V^*)^G$, and polynomials $f_1(X), \dots, f_b(X) \in P(V^*)^G[X]$, where X is an indeterminate of degree 2, such that:

- (1) ρ_1, \dots, ρ_n are among the coefficients of $f_1(X), \dots, f_b(X)$, and

(2) $f_i(X)$ splits into a product of linear factors in $P(V^*)[X]$ for $i = 1, \dots, b$. If we can choose $b = 1$ then we say $P(V^*)^G$ satisfies the *splitting principle*. (N.B. The polynomials $f_i(X)$ are homogeneous and their roots lie in V^* .)

THEOREM. *Suppose $G < \mathrm{GL}(V)$ is generated by pseudo reflections. $|G| \not\equiv O(p)$, and $P(V^*)^G$ satisfies the weak splitting principle. Then there exist integers d_1, \dots, d_a depending on $G < \mathrm{GL}(V)$ and a convergent tower of fibrations*

$$\cdots \rightarrow X_n \rightarrow X_{n-1} \rightarrow \cdots \rightarrow X_1 \rightarrow \bigtimes_{i=1}^a BU(d_i)$$

such that: if W is the space at the top of the tower then W is simply connected, of the homotopy type of a CW complex of finite type, and further

$$H^*(W, \mathbf{F}_p) \simeq P(V^*)^G$$

and the induced map

$$H^*\left(\bigtimes_{i=1}^a BU(d_i); \mathbf{F}_p\right) \rightarrow P(V^*)^G$$

is surjective.

There are two aspects to the construction of the tower to which we wish to draw attention. The first is the simple idea of “trapping” $P(V^*)^G$, which we wish to realize as a cohomology ring, as the image of an induced map inside $H^*(BV; \mathbf{F}_p)$; where BV is the classifying space of the elementary abelian p group V . (See Lemma 5 and the discussion around it.) This allows us to solve the sort of extension problems that occur in [8], [9], [14], [15], etc.

The second aspect of the construction worthwhile remarking is the use of the Carlsson [4], Miller [10] result that $H^*(BV; \mathbf{F}_p)$ is an injective object in the category of unstable modules over the Steenrod algebra (see also [23]) to continue trapping $P(V^*)^G < H^*(BV; \mathbf{F}_p)$ as the image of an induced map as we move up the tower.

The method employed should be of use in other situations, and is perhaps more important than the actual results obtained here. A number of further applications will be appearing in a joint paper with Paul Goerss.

We postpone a discussion of examples of groups satisfying the splitting principles to the end of this paper and [17], where, among other things, we show that, when p is large compared to $|G|$, $P(V^*)^G$ satisfies weak splitting.

The remainder of this paper is organized as follows. After some preliminary maneuvers we take up the construction of the tower required for the proof of the main theorem. We then complete the proof of the main theorem, deferring to the end the proofs of the technical results, so as to not interrupt the flow of the argument.

The tower construction. We denote by BV the classifying space of the abelian p -group $V = \bigoplus_n \mathbf{F}_p$, and identify

$$H^2(BV, \mathbf{Z}) \simeq V^* \simeq \text{Ext}_{\mathbf{Z}}(V, \mathbf{Z}).$$

We may then write

$$H^*(BV; \mathbf{F}_p) \simeq E(\beta^{-1}V^*) \otimes P(V^*)$$

where β is the mod p Bockstein operator and $E(\)$ the exterior algebra functor. If $G < \text{GL}(V)$ then there is an induced action of G on BV (which we may assume to be free) and hence on $H^*(BV; \mathbf{F}_p)$. In this way we obtain an inclusion

$$P(V^*)^G \hookrightarrow H^*(BV; \mathbf{F}_p)^G$$

and an isomorphism (see [1] for more information)

$$P(V^*)^G \simeq \frac{H^*(BV; \mathbf{F}_p)^G}{\sqrt{O}},$$

where \sqrt{O} denotes the ideal of nilpotents.

PROPOSITION 1. *Suppose $G < \text{GL}(V)$, $V = \bigoplus_n \mathbf{F}_p$, and $P(V^*)^G$ satisfies the weak splitting principle. Then there exist G vector bundles $\xi_i \downarrow BV$, $i = 1, \dots, a$, such that*

$$c_i(\xi_j) \in P(V^*)^G; \quad i = 1, \dots, \dim \xi_j; \quad j = 1, \dots, a$$

and moreover $\{c_i(\xi_j)\}$ generate $P(V^)^G$ as an algebra.*

Proof. By hypothesis there are polynomials

$$f_i(X)S \in P(V^*)^G[X]$$

such that:

$$f_i(X) = \prod_{v \in A_i} (X + v) \in P(V^*)[X]; \quad A_i \subset V$$

and $P(V^*)^G$ is generated as an algebra by the coefficients of $f_1(X), \dots, f_a(X)$. Since $f_1(X), \dots, f_a(X) \in P(V^*)^G[X]$, it follows that A_1, \dots, A_a are G invariant subsets of $V^* \simeq H^2(BV; \mathbf{Z})$. Corresponding to

each element $v \in A_j$ we may find a (unique) line bundle $\lambda(v) \downarrow BV$ with $c_1(\lambda(v)) = v$. The vector bundle $\xi_i = \bigoplus_{v \in A_i} \lambda(v) \downarrow BV$ is then a G bundle and satisfies

$$c(\xi_i) = f(1) \in P(V^*)^G,$$

where $c(\)$ denotes the total Chern class as required. \square

COROLLARY 2. *Suppose $G < \mathrm{GL}(V)$, $V = \bigoplus_n \mathbf{F}_p$ and $P(V^*)^G$ satisfies the weak splitting principle. Then there is an epimorphism of algebras over the Steenrod algebra*

$$\bigotimes_{i=1}^a H^*(BU(d_i); \mathbf{F}_p) \rightarrow P(V^*)^G. \quad \square$$

The preceding corollary suggests a rather obvious way to attempt to construct a space W with $H^*(W; \mathbf{F}_p) \simeq P(V^*)^G$. Namely, we let

$$f: \bigtimes_{i=1}^a BU(d_i) \rightarrow K = \bigtimes_{i=1}^c K(\mathbf{H}\mathbf{Z}, m_j)$$

be a map such that the classes

$$f^*(i_{m_j}) \in H^*\left(\bigtimes_{i=1}^a BU(d_i); \mathbf{F}_p\right): \quad j = 1, \dots, c$$

are a minimal set of ideal generators for the kernel of an epimorphism

$$\bigotimes_{i=1}^a H^*(BU(d_i); \mathbf{F}_p) \rightarrow P(V^*)^G.$$

Introduce the fibre square

$$(\mathcal{F}) \quad \begin{array}{ccc} X & \rightarrow & L \\ \downarrow \tau & & \downarrow \\ \bigtimes_{i=1}^a BU(D_i) & \xrightarrow{f} & K \end{array}$$

where $L \downarrow K$ is the usual pathspace fibration. It is of course too much to hope that

$$H^*(X; \mathbf{F}_p) \simeq P(V^*)^G \simeq H^*\left(\bigtimes_{i=1}^a BU(d_i); \mathbf{F}_p\right) // f^*.$$

It is however not too much to hope that $H^*(X; \mathbf{F}_p)$ is a good approximation to $P(V^*)^G$, and in fact this is the case as we now proceed to show.

LEMMA 3. *In the preceding diagram set $S^* = \mathrm{Im} f^*$. Then S^* is a polynomial algebra and $H^*(\bigtimes_{i=1}^s BU(d_i); \mathbf{F}_p)$ is a free S^* -module.*

LEMMA 4. $\text{Ker } f^* < H^*(K; \mathbb{F}_p)$ is a Borel ideal.

We may now apply [14; II.2] to compute $H^*(X; \mathbb{F}_p)$. We obtain an isomorphism of algebras over the Steenrod algebra \mathcal{A}^*

$$H^*(X; \mathbb{F}_p) \simeq \bigcup_{R^*} (F^{-1}),$$

a coexact sequence of \mathcal{A}^* algebras (for a discussion of coexactness see [14, II] or [15])

$$1 \rightarrow R^* \rightarrow H^*(X; \mathbb{F}_p) \rightarrow \bigcup(M) \rightarrow 1,$$

and an exact sequence of \mathcal{A}^* modules

$$1 \rightarrow R^* \rightarrow F^{-1} \rightarrow E_{\infty}^{-1,*} \rightarrow 0$$

where:

(a) $R^* =: H^*(\times_{i=1}^a BU(d_i); hF_p) // f^* \simeq P(V^*)^G$;

(b) $F^{-1} < H^*(X; hF_p)$ is the submodule of elements of filtration -1 with respect to the Eilenberg-Moore spectral sequence;

(c) $M = \mathbb{F}_p \otimes_{R^*} E_{\infty}^{-1,*}$;

(d) $\bigcup(M) \simeq \text{Im}\{i^*: H^*(X; \mathbb{F}_p) \rightarrow H^*(\Omega K; \mathbb{F}_p)\}$.

(For a discussion of the functors $\bigcup(\)$ and $\bigcup_{R^*}(\)$ see [8] and [9].) All this is standard. What we would now like to do is to choose a submodule $N < H^*(X; \mathbb{F}_p)$ mapping isomorphically onto M ; represent the elements of N by a map

$$X \xrightarrow{f'} \bigtimes_{i=1}^{\infty} K(N^i, i) = K',$$

form the pullback diagram

$$\begin{array}{ccc} X' & \rightarrow & L' \\ \downarrow \tau' & & \downarrow \\ X & \xrightarrow{f'} & K' \end{array}$$

etc. The problem is that we do not a priori know that N can be chosen to be an \mathcal{A}^* submodule, and so we cannot assure that τ'^* maps $P(V^*)^G \simeq R^* < H^*(X; \mathbb{F}_p)$ monomorphically. To deal with this problem let us decorate the fibre square (\mathcal{F}) to a diagram

$$(\mathfrak{F}) \quad \begin{array}{ccccc} & & & X & \rightarrow L \\ & & \nearrow \sigma & \downarrow \tau & \downarrow \\ BV & \xrightarrow{g} & \bigtimes_{i=1}^a BU(d_i) & \xrightarrow{f} & K \end{array}$$

where

$$g_j\colon BV \xrightarrow{g} \bigtimes_{i=1}^a BU(d_i) \xrightarrow{\text{pr}_j} BU(d_j)$$

classifies the bundle $\xi_j \downarrow BV$ of Proposition 1. By construction $f \circ g$ is null homotopic, so there is a lift σ as indicated.

LEMMA 5. *There is a choice of σ such that*

$$\text{Im } \sigma^* \simeq P(V^*)^G.$$

We obtain therefore a diagram

$$\begin{array}{ccccc} & & H^*(X, \mathbb{F}_p) & & \\ & \swarrow \sigma^* & \uparrow \tau^* & \nwarrow & \\ H^*(BV; \mathbb{F}_p) & \xrightarrow{g^*} & \bigotimes_{i=1}^a H^*(BU(d_i); \mathbb{F}_p) & \xrightarrow{\quad} & R^* \simeq P(V^*)^G \end{array}$$

the composite

$$R^* \hookrightarrow H^*(X; \mathbb{F}_p) \xrightarrow{\sigma^*} P(V^*)^G$$

being the identity. The map σ^* therefore provides a splitting over $R \odot \mathcal{A}^*$ of both of the sequences

$$\begin{aligned} 0 \rightarrow R^* \rightarrow F^{-1}H^*(X; \mathbb{F}_p) \rightarrow E_{\infty}^{-1,*} \rightarrow 0, \\ 1 \rightarrow R^* \rightarrow H^*(X; \mathbb{F}_p) \rightarrow \cup(M) \rightarrow 0. \end{aligned}$$

If we set

$$N = \text{Ker}\big\{\sigma^*\colon F^{-1}H^*(X; \mathbb{F}_p) \rightarrow P(V^*)^G\big\},$$

we then obtain:

PROPOSITION 6. *With the notations preceding there is an isomorphism*

$$H^*(X; \mathbb{F}_p) \simeq R^* \otimes \cup(N)$$

such that

$$\begin{array}{ccc} H^*(\Omega K; \mathbb{F}_p) & \hookleftarrow & \cup(N) \\ \uparrow i^* & & \uparrow \varepsilon \otimes 1 \\ H^*(X; \mathbb{F}_p) & \simeq & R^* \otimes \cup(N) \\ \downarrow \sigma^* & & \downarrow 1 \otimes \varepsilon \\ H^*(BV; \mathbb{F}_p) & \hookleftarrow & P(V^*)^G \simeq R^* \end{array}$$

where ε is the augmentation. The module N is at least 3 connected. If $x \in N$ is a non-zero element of minimal degree, then some higher order Bockstein applied to x is non-zero.

REMARK. For a discussion of higher order Bockstein operators β_r , see [3]. If $u \in H^d(X, \mathbb{F}_p)$ and $\beta_r u = 0$, $r < s$, $\beta_s u \neq 0$, then

$$u \in \text{Im}\{H^*(X; \mathbb{Z}/p^s) \rightarrow H^*(X, \mathbb{Z}/p)\},$$

but

$$u \notin \text{Im}\{H^*(X; \mathbb{Z}/p^{s+1}) \rightarrow H^*(X; \mathbb{Z}/p)\}.$$

This implies that we can represent u by a map

$$\varphi: X \rightarrow K(\mathbb{Z}/p^s, d)$$

such that $\varphi^*(i) = u$, and hence $\ker \varphi^* = 0$ in degrees $< d + 2$.

PROPOSITION 7. With the notations preceding we may choose f , such that

$$H^*(X; \mathbb{Q}) \simeq H^*\left(\bigtimes_{i=1}^a BU(d_i), \mathbb{Q}\right) // f^*$$

is a polynomial algebra of the same type as $P(V^*)^G$. In fact there is a subalgebra $R^* < H^*(X; \mathbb{Z})$ such that $R^* \otimes_{\mathbb{Z}} \mathbb{Q} \simeq H^*(X; \mathbb{Q})$ and $R^* \otimes_{\mathbb{Z}} \mathbb{Z}/p \simeq R^* \simeq P(V^*)^G$.

Proof of the main theorem. We are going to construct inductively a convergent tower of fibrations and lifts

$$\begin{array}{ccc} & & \downarrow \\ & & X_n \\ & \nearrow \sigma_n & \downarrow \tau_n \\ & & \vdots \\ & \nearrow \sigma_1 & \downarrow \tau_1 \\ & & X_1 \\ & \nearrow \bar{g} & \downarrow \tau_1 \\ g: BV \xrightarrow{\pi} B & \xrightarrow{\bar{g}} & \bigtimes_{i=1}^a BU(d_i) \end{array}$$

where $B = BV/G$ and \bar{g} factors g over the orbit map $\pi: BV \downarrow B$, and

(a) $\text{Im } g^* = \text{Im } \pi^* \sigma_1^* = \dots = \text{Im } \pi^* \sigma_n^* = P(V^*)^G$;

(b) $H^*(X_n; \mathbb{F}_p) \simeq P(V^*)^G \otimes \cup(N(n))$ as algebra over the Steenrod algebra; and

(c) the diagram

$$\begin{array}{ccc} H^*(X_n; \mathbf{F}_p) & \simeq & P(V^*)^G \otimes \cup(N(n)) \\ \downarrow \pi^* \sigma_n^* & & \downarrow 1 \otimes \varepsilon \\ H^*(BV; \mathbf{F}_p) & \leftarrow & P(V^*)^G \end{array}$$

commutes.

For $n = 1$ this is just Proposition 6, so the induction starts and we turn to the inductive step. We will first describe the construction of X_{n+1} from X_n . We then locate inside $H^*(X_{n+1}; \mathbf{F}_p)$ a well-placed copy of $P(V^*)^G$ (condition (a)). Next we show how to split this copy of $P(V^*)^G$ off (condition (b)), and finally we adjust the lift σ_{n+1} (condition (c)) so that we may repeat the splitting argument at the next stage.

Construction of X_{n+1} . Let

$$f_{n+1}: X_n \rightarrow K_{n+1} := \times K(N(n)^j; j)$$

be defined by requiring that

$$f_{n+1}^*: H^*(K_{n+1}; \mathbf{F}_p) \rightarrow H^*(X_n; \mathbf{F}_p)$$

map $N(n)$ isomorphically. There is then the fibre square

$$\begin{array}{ccc} X_{n+1} & \rightarrow & L \\ \tau_{n+1} \downarrow & & \downarrow \\ X_n & \xrightarrow{f_{n+1}} & K_{n+1} \end{array}$$

(\mathcal{F}_n)

defining X_{n+1} where $L_{n+1} \downarrow K_{n+1}$ is the path space fibration.

Locating $P(V^)^G$ inside $H^*(X_{n+1}; \mathbf{F}_p)$.* An Eilenberg-Moore spectral sequence argument (of the type to be found in [15; §4] say) applied to (\mathcal{F}_n) yields an isomorphism of algebras over the Steenrod algebra

$$H^*(X_{n+1}; \mathbf{F}_p) \simeq \bigcup_{R^*} (F^{-1}),$$

a coexact sequence of \mathcal{A}^* algebras

$$1 \rightarrow R^* \rightarrow H^*(X_{n+1}; \mathbf{F}_p) \rightarrow \cup(M(n+1)) \rightarrow 1$$

and an exact sequence of \mathcal{A}^* modules

$$1 \rightarrow R^* \rightarrow F^{-1} \rightarrow E_{\infty}^{-1,*} \rightarrow 0,$$

where

$$R^* = H^*(X_n; \mathbf{F}_p) \simeq P(V^*)^G,$$

and

$$\cup(M(n+1)) \simeq \text{Im}\{H^*(X_n; \mathbb{F}_p) \rightarrow H^*(\Omega K_{n+1}; \mathbb{F}_p)\}.$$

Splitting $H^(X_{n+1}; \mathbb{F}_p)$.* By property (c) of the inductive hypothesis we may decorate (\mathcal{F}_n) to a diagram

$$\begin{array}{ccccc} & & X_{n+1} & \xrightarrow{\quad} & L_{n+1} \\ & \nearrow \sigma_{n+1} & \downarrow \tau_{n+1} & & \downarrow \\ BV \xrightarrow{\pi} B & \xrightarrow{\sigma_n} & X_n & \xrightarrow{f_{n+1}} & K_{n+1} \end{array}$$

because the composite $f_{n+1} \cdot \sigma_n$ is null homotopic by construction. Since $|G| \not\equiv O(p)$ a transfer argument shows

$$H^*(B; \mathbb{F}_p) \simeq H^*(BV; \mathbb{F}_p)^G.$$

From [1] it follows

$$H^*(BV; \mathbb{F}_p)^G / \sqrt{O} \simeq P(V^*)^G.$$

The composite

$$H^*(X_{n+1}; \mathbb{F}_p) \xrightarrow{\sigma_{n+1}^*} H^*(B; \mathbb{F}_p) \rightarrow P(V^*)^G$$

splits the above exact sequences to give an isomorphism of $R^* \odot \mathcal{A}^*$ modules

$$F^{-1} \simeq R^* \oplus E_{\infty}^{-1,*}.$$

Since the functor $\cup_{R^*}(\)$ sends direct sums into tensor products and $E_{\infty}^{-1,*}$ is a free R^* module, we obtain

$$H^*(X_{n+1}; \mathbb{F}_p) \simeq R^* \otimes \cup(M(n+1))$$

where

$$M(n+1) = \mathbb{F}_p \otimes_{R^*} E_{\infty}^{-1,*}.$$

Adjusting the lift σ_{n+1} . It remains to show that we can choose the lift σ_{n+1} so that $\sigma_{n+1}^*(N(n+1)) = 0$. To this end consider the diagram (P denotes primitives)

$$\begin{array}{ccccc} & & PH^*(K_{n+1}; \mathbb{F}_p) & & \\ & \nwarrow \alpha & & \uparrow i^* & \\ H^*(B; \mathbb{F}_p) & \xleftarrow{\sigma_{n+1}^*} & H^*(X_{n+1}; \mathbb{F}_p) & \xleftarrow{\quad} & N(n) \end{array}$$

By the dual of a theorem of Carlsson [4] and Miller [10], [23] $H^*(BV; \mathbb{F}_p)$ is an injective object in the category of unstable modules over the Steenrod algebra. By a transfer argument $H^*(B; \mathbb{F}_p)$ is an \mathcal{A}^* direct summand in $H^*(BV; \mathbb{F}_p)$. Therefore $H^*(B; \mathbb{F}_p)$ is also an injective in the category of unstable modules over the Steenrod algebra, and therefore we obtain a dotted map α . Cutting α down to fundamental classes defines a map $\varphi: B \rightarrow K_{n+1}$ such that

$$\sigma_{n+1}^* = \varphi^* i^*: N(n) \rightarrow H^*(B; \mathbb{F}_p).$$

Since the fibration

$$\Omega K_{n+1} \rightarrow X_{n+1} \rightarrow X_n$$

is a principal bundle, we can form the twisted lift

$$\sigma'_{n+1}: B \xrightarrow{(\varphi, \sigma_{n+1})} \Omega K_{n+1} \times X_{n+1} \xrightarrow{\mu} X_{n+1}$$

where μ is the principal action. The adjusted lift σ'_{n+1} satisfies

$$\sigma_{n+1}'^*(N(n+1)) = 0$$

by construction.

This completes the inductive construction of the tower. It remains to verify that the tower converges. To this end, note that as a consequence of Propositions 6 and 7 the space X_{n+1} arises from X_n by using an \mathbb{F}_p vector space to annihilate p -torsion classes in $H_*(X_n; \mathbb{Z})$ of finite order. Thus the connectivity of $N(n)$ goes to infinity with n and the tower converges. \square

It remains to prove (3)–(7).

Proof of (3). We first show that $H^*(\times_{i=1}^a BU(d_i); \mathbb{F}_p)$ is a free S^* -module. Note by construction we have a coexact sequence of commutative graded algebras

$$1 \rightarrow S^* \rightarrow H^* \rightarrow R^* \rightarrow 1,$$

where we have abbreviated $H^*(\times_{i=1}^a BU(d_i); \mathbb{F}_p)$ to H^* , and $P(V^*)^G$ to R^* . Polynomial algebras being free commutative, this sequence splits to yield an isomorphism of algebras $H^* \simeq S^* \otimes R^*$. Hence H^* is a free S^* -module.

To verify that S^* is a polynomial algebra we employ Serre's converse to Hilbert's syzygy theorem [12] (see also [20; Thm. 8]). According to this result a graded connected algebra over a field is a polynomial algebra iff it

has finite projective dimension. So it will suffice to show that the projective dimension of S^* is finite. To this end suppose M is a graded S^* -module. Set $d = d_1 + \cdots + d_a$. Let

$$(\mathcal{P}) \quad 0 \rightarrow K_{d+1} \rightarrow P_d \rightarrow \cdots \rightarrow P_0 \rightarrow M \rightarrow 0$$

be a partial projective resolution of M , that is, P_0, \dots, P_d are projective S^* -modules and the sequence is exact. Since H^* is a free S^* -module the functor $\otimes_{S^*} H^*$ is exact and sends projective S^* -modules into projective H^* -modules. Thus

$$0 \rightarrow K_{d+1} \otimes_{S^*} H^* \rightarrow P_d \otimes_{S^*} H^* \rightarrow \cdots \rightarrow M \otimes_{S^*} H^* \rightarrow 0$$

is an exact sequence of H^* -modules, where all but perhaps the last term are projective. By Hilbert's syzygy theorem H^* has projective dimension d and hence the last term $K_{d+1} \otimes_{S^*} H^*$ is also H^* projective [5; VI.2.1]. Since H^* is free over S^* , projective H^* -modules are also projective S^* -modules. Note

$$S^* \hookrightarrow H^* \simeq S^* \otimes R^* \xrightarrow{1 \otimes \varepsilon} S^*$$

where ε is the augmentation, presents S^* as an S^* -module direct summand in H^* . Thus

$$K_{d+1} \simeq K_{d+1} \otimes_{S^*} S^* \rightleftarrows K_{d+1} \otimes_{S^*} H^*$$

represents K_{d+1} as an S^* direct summand in the projective S^* -module $K_{d+1} \otimes_{S^*} H^*$. Hence K_{d+1} is a projective S^* -module, and thus (\mathcal{P}) is a projective resolution of M , so the projective dimension of S^* is at most d . \square

Proof of (4). From the epimorphism

$$f^*: H^*(K; \mathbb{F}_p) \rightarrow S^*$$

we obtain by passing to the indecomposable quotients an epimorphism

$$Qf^*: QH^*(K; \mathbb{F}_p) \rightarrow QS^*.$$

Since polynomial algebras are free commutative algebras there is an algebra splitting of f^*

$$s: S^* \rightarrow H^*(K; \mathbb{F}_p)$$

by Lemma 3. Let T be the kernel of Qf^* . By choosing representatives in $H^*(K; \mathbb{F}_p)$ for a basis of T and using the splitting s , we can define

$$\varphi: P(T) \otimes S^* \rightarrow H^*(K; \mathbb{F}_p),$$

which is an isomorphism on the module of indecomposables. Since the domain and range of φ are polynomial algebras φ is an isomorphism. Thus $\varphi(T)$ generates the kernel of f^* . The iso φ shows that $H^*(K; \mathbb{F}_p)$ is a free $P(\varphi T)$ -module, so the result follows from [14; II.1.1]. \square

Proof of (5) (after Zabrodsky). The action of G on V induces an action on BV , which, without loss of generality we may assume to be free. Let $\pi: BV \downarrow B$ be the orbit map. By Proposition 1 the bundles ξ_1, \dots, ξ_a descend to B so we may form the diagram

$$\begin{array}{ccccc}
 & & X & \rightarrow & L \\
 & \nearrow \bar{\sigma} & \downarrow \tau & & \downarrow \\
 BV & \xrightarrow{\pi} B & \xrightarrow{\bar{g}} \bigtimes_{i=1}^a BU(d_i) & \xrightarrow{f} & K
 \end{array}$$

where \bar{g} classifies the descended bundles. By replacing f by $|G| \cdot f$ if need be we may assume that

$$\bar{g}^* \cdot f^*: \tilde{H}^*(K; \mathbb{Z}/q) \rightarrow \tilde{H}^*(B; \mathbb{Z}/q)$$

is the zero map for all primes $q \neq p$. This composite also vanishes for $q = p$ by construction and the fact that

$$H^*(B; \mathbb{F}_p) \xrightarrow[\cong]{\pi^*} H^*(BV; \mathbb{F}_p)^G.$$

Since also $\tilde{H}^*(B; \mathbb{Q}) = 0$, it follows that

$$\bar{g}^* \cdot f^*: \tilde{H}^*(K; \mathbb{Z}); \mathbb{Z} \rightarrow \tilde{H}^*(B; \mathbb{Z})$$

is the zero map. Thus $\bar{g} \cdot f$ is null homotopic, and there exists a lift $\bar{\sigma}$ as indicated.

Consider the diagram

$$\begin{array}{ccccc}
 & & PH^*(\Omega K; \mathbb{F}_p) & & \\
 & \nwarrow & \uparrow & & \\
 H^*(BV; \mathbb{F}_p) & \xleftarrow{\pi^*} & H^*(B; \mathbb{F}_p) & \xleftarrow{\bar{\sigma}^*} & M
 \end{array}$$

By the dual of a theorem of Carlsson [4] and Miller [10] [23] $H^*(BV; \mathbb{F}_p)$ is an injective object in the category of unstable modules over the Steenrod algebra. Since $|G| \not\equiv 0(p)$ we may form the averaging operator

$$A := \frac{1}{|G|} \sum_{g \in G} g: H^*(Bv; \mathbb{F}_p) \rightarrow H^*(BV; \mathbb{F}_p)$$

whose image is $H^*(BV; \mathbb{F}_p)^G \simeq H^*(B; \mathbb{F}_p)$. Since A is \mathcal{A}^* linear it follows that $H^*(B; \mathbb{F}_p)$ is an \mathcal{A}^* direct summand in $H^*(BV; \mathbb{F}_p)$, and therefore $H^*(B; \mathbb{F}_p)$ is an injective object in the category of unstable modules over the Steenrod algebra. Therefore there is a dotted morphism as in the preceding diagram. Now recall that by construction

$$\Omega K = \times K(\mathbb{Z}, 2j - 1)$$

for suitable integers j . Therefore, $PH^*(\Omega K; \mathbb{F}_p)$ being generated as an \mathcal{A}^* -module by its fundamental classes, is generated by odd dimensional classes with vanishing Bocksteins. However, by [1; §3].

$$H^*(BV; \mathbb{F}_p)^G \simeq E[\tau_1, \dots, \tau_n] \otimes P[\rho_1, \dots, \rho_n]$$

where $\beta\tau_i = \tau_i$, $i = 1, \dots, n$, so

$$H^*(B; \mathbb{F}_p) \simeq H^*(BV; \mathbb{F}_p)^G$$

contains no odd dimensional classes with vanishing Bocksteins. Therefore the dotted map is zero. Thus $\pi^*\bar{\sigma}^*(m) = 0$. The fibration

$$\Omega K \rightarrow X \rightarrow \bigtimes_{i=1}^a BU(d_i)$$

is a principal ΩK bundle. Thus any two lifts σ', σ'' of g over τ differ by a map $BV \rightarrow \Omega K$. But the preceding considerations with Bockstein shows any such map induces the zero map in \mathbb{F}_p cohomology. Thus $\sigma^* = \pi^*\bar{\sigma}^*$ and so $\sigma^*(M) = 0$. \square

Proof of (7). By Lemma 3 and [14; II.1.1]

$$f^*(i_{m_j}) \in H^*\left(\bigtimes_{i=1}^a BU(d_i); \mathbb{F}_p\right) \quad j = 1, \dots, c$$

is an ESP-sequence. Therefore

$$p, f^*(i_{m_1}), \dots, f^*(i_{m_c}) \in H^*\left(\bigtimes_{i=1}^a BU(d_i); \mathbb{Z}\left[\frac{1}{p'}\right]\right)$$

is an ESP-sequence. A standard Eilenberg-Moore spectral sequence argument now completes the proof. \square

Proof of (6). It only remains to verify the statement about Bocksteins. But this follows because Proposition 7 assures us that X contains no unwanted rational cohomology classes. \square

We turn now to a discussion of examples of groups G satisfying the weak splitting principle.

EXAMPLE 1. Let $G = \text{GL}(V)$. Then [7] [18]

$$P(V^*)^G = D^*(n) = P[y_1, \dots, y_n]$$

where

$$X^{p^n-1} + y_1 X^{p^{n-1}-1} + \dots + y_n = \prod_{v \neq 0 \in V^*} (X + v).$$

This is perhaps the most basic example, but of course for $n > 1$ does not satisfy the non-modularity condition $|G| \not\equiv O(p)$. As previously noted, the groups $G < \text{GL}(V)$ generated by pseudo reflections and satisfying $|G| \not\equiv O(p)$ are products of irreducible examples that have been listed by Clark and Ewing [6]. In this list there are three infinite families.

EXAMPLE 2. Let $G = \Sigma_n$ be the symmetric group on n letters acting on $V = \bigoplus_n \mathbb{F}_p$ by permutation of the basis vectors. Then

$$P(V^*)^{\Sigma_n} = P[\sigma_1, \dots, \sigma_n]$$

where

$$X^n + \sigma_1 X^{n-1} + \dots + \sigma_n = \prod_{i=1}^n (X + t_i),$$

$\{t_1, \dots, t_n\}$ being the basis of V permuted by Σ_n . Thus Σ_n satisfies the splitting principle.

EXAMPLE 3. $G = D_m = \mathbb{Z}/m \ltimes \mathbb{Z}/2$, the dihedral group of order $|D_m| = 2m$, where m is a divisor of $p - 1$, acting on $V = \mathbb{F}_p \oplus \mathbb{F}_p$ via the matrices

$$\begin{pmatrix} \theta & 0 \\ 0 & \theta^{-1} \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

where $\theta \in \mathbb{F}_p^*$ is a primitive m th root of unity. If we write $\{u, v\}$ for the dual basis in V^* , then

$$P(V^*)^{D_m} \simeq P[\rho_1, \rho_2]$$

where we may choose

$$\rho_1 = muv, \quad \rho_2 = -(u^m + v^m),$$

and

$$X^m + \rho_1 X^{m-2} + \dots + \rho_2 = \prod_{i=1}^{m-1} (X + (\theta^i u + \theta^{-i} v)).$$

Thus D_m satisfies the weak splitting principle.

EXAMPLE 4. $G(n, m, r)$. These are the groups of type 2b in the Clark-Ewing list. Here $p \equiv 1 \pmod m$, $p \nmid n!$, and $r|m$. If $\theta \in \mathbb{F}_p^*$ is a primitive m th root of unity, then $G(n, m, r)$ may be taken to be the subgroup of $\text{GL}(V)$ generated by the permutation matrices and the matrices

$$\begin{pmatrix} \theta^{v_1} & & & 0 \\ & \theta^{v_2} & & \\ & & \ddots & \\ 0 & & & \theta^{v_n} \end{pmatrix}$$

with respect to some fixed basis of V , where $v_1 + \cdots + v_n \equiv 0 \pmod r$. One has

$$P(V)^G = P[\rho_1, \dots, \rho_n]$$

where

$$\rho_i = \begin{cases} \sigma_i(t_1^m, \dots, t_n^m): & 1 \leq i \leq n-1, \\ \sigma_n(t_1^{m/r}, \dots, t_n^{m/r}): & i = n. \end{cases}$$

Moreover

$$\prod_{\substack{1 \leq i \leq m \\ 1 \leq j \leq n}} (X + \theta^i t_j) = X^{nm} + \rho_1 X^{(n-1)m} + \cdots + \rho_n^r.$$

Let $w = t_1 + \cdots + t_n$. The orbit of w “sees” r ; to wit

$$[w] = \{ \theta^{v_1} t_1 + \cdots + \theta^{v_n} t_n \mid v_1 + \cdots + v_n \equiv 0 \pmod r \}.$$

This orbit has $m^n r$ elements. Let

$$s = \sum_{v \in [w]} v^{nm/r} = \sum_{v_1 + \cdots + v_n \equiv 0(r)} (\theta^{v_1} t_1 + \cdots + \theta^{v_n} t_n)^{nm/r}.$$

Since s is a polynomial in the elementary symmetric functions, it lies in the subalgebra generated by the coefficients of $\varphi_{[w]}(X)$. Let us look at the coefficient of $\rho_n = t_1^{m/r} \cdots t_n^{m/r}$ in s . The sum defining s has nm/r terms, namely

$$\theta^{(v_1 + \cdots + v_n)m/r} \frac{(nm/r)!}{[(m/r)!]^n} = \frac{(nm/r)!}{[(m/r)!]^n} \not\equiv O(p)$$

as $p > nm/r$ and $p \nmid n!$. Therefore the coefficient of $\rho_n = t_1^{m/r} \cdots t_n^{m/r}$ in s is

$$\frac{(nm/r)(nm/r)!}{[(m/r)!]^n} \equiv O(p),$$

so we conclude that

$$s \equiv \rho_n \in P(V)^{G(n,m,r)} \text{ mod decomposable.}$$

Thus for $p > nm/r$, $p \nmid n!$, $P(V)^{G(n,m,r)}$ satisfies the weak splitting principle.

Acknowledgment. I am grateful to R. E. Stong for assistance with the preceding example.

For a thorough discussion of splitting principles we refer to [17].

A Consequence of the Main Theorem

THEOREM. *Suppose $G < \text{GL}(V)$ is generated by pseudo reflections, $|G| \neq 0 \text{ mod } p$, and $P(V)^G$ satisfies the weak splitting principle. Let $B \subset V$ be an orbit with orbit Chern class $c(B) \in P(V^*)^G$. Then there exists a simply connected space X of finite type and a vector bundle $\xi_B \downarrow X$ such that*

$$H^*(X; \mathbb{F}_p) \simeq P(V^*)^G$$

and under this identification

$$c(B) = c(\xi_B)$$

where $c(\xi_B)$ is the total Chern class of the vector bundle ξ_B .

Proof. We modify ever so slightly the proof of the main theorem. Let

$$\mathbf{B} = \{B \subset V \mid B \text{ is an orbit of } G \text{ in } V\}.$$

For $B \in \mathbf{B}$ let $|B|$ be the cardinality of B . The orbit Chern classes then provide a map

$$H^*(BU(|B|); \mathbb{F}_p) \rightarrow P(V^*)^G$$

and the induced map

$$H^*\left(\bigtimes_{b \in \mathbf{B}} BU(|B|); \mathbb{F}_p\right)$$

is an epimorphism. We now start the construction of the tower in the main theorem with this product $\bigtimes_{b \in \mathbf{B}} BU(|B|)$ to construct the desired space X . The desired vector bundle ξ_B is then classified by the composite

$$X \rightarrow \bigtimes_{b \in \mathbf{B}} BU(|B|) \rightarrow BU(|B|)$$

where the last map is projection onto the factor corresponding to the orbit B . □

REFERENCES

- [1] J. Aguadé and L. Smith, *Modular cohomology algebras*, Amer. J. Math., (1985), 507–530.
- [2] N. Bourbaki, *Groupes et Algèbres de Lie*, Ch. 4, 5, 6 Masson Paris 1981.
- [3] W. Browder, *Torsion in H-spaces*, Annals of Math., **74** (1961), 24–51.
- [4] G. Carlsson, *G. B. Segal's Burnside ring conjecture for $(\mathbb{Z}/2)^k$* , Topology, **22** (1983), 83–103.
- [5] H. Cartan and S. Eilenberg, *Homological Algebra*, Princeton University Press 1956.
- [6] A. Clark and J. Ewing, *The realization of polynomial algebras as cohomology rings*, Pacific J. Math., **50** (1974), 425–434.
- [7] L. E. Dickson, *A fundamental system of invariants of the general modular linear group*, Trans. Amer. Math. Soc., **12** (1911), 75–98.
- [8] W. S. Massey and F. P. Peterson, *The cohomology structure of certain fibre spaces*, Topology, **4** (1965), 47–65.
- [9] W. S. Massey and F. P. Peterson, *The mod 2 Cohomology Structure of Certain Fibre Spaces*, Mem. Amer. Math. Soc., **74** (1967),
- [10] H. Miller, *The Sullivan Conjecture on Maps from Classifying Spaces* (in Algebraic Topology Aarhus 1982 (Ed. Oliver and Madsen SLNM 1051) and Univ. of Washington Preprint August 1983).
- [11] J. Milnor and J. C. Moore, *The structure of Hopf algebras*, Ann. of Math., **81** (1965), 211–264.
- [12] J. P. Serre, *Sur la dimension homologique des anneaux et des modules noetheriens*, Tokyo Symposium 1955.
- [13] G. D. Shephard and J. A. Todd, *Finite unitary reflection groups*, Canad. J. Math., **6** (1954), 274–304.
- [14] L. Smith, *Homological Algebra and the Eilenberg-Moore Spectral Sequence*, Trans. Amer. Math. Soc., **129** (1967), 58–93.
- [15] ———, *Hopf Fibration Towers and the Unstable Adams Spectral Sequence*, Proc. Conf. Categorical Algebra, 1968.
- [16] ———, *Remarks on realizing Dickson covariants as cohomology rings*, Quart. J. Math. Oxford (2), **36** (1985), 113–115.
- [17] L. Smith and R. E. Stong, *Chern classes, orbit polynomials and rings of invariants*, J. Algebra, (to appear).
- [18] L. Smith and R. M. Switzer, *On the realization and non-realization of Dickson algebras as cohomology rings*, Proc. Amer. Math. Soc., **89** (1983), 303–313.
- [19] ———, *Polynomial algebras over the Steenrod algebra*, Proc. Edinburgh Math. Soc., **27** (1984), 11–19.
- [20] J. Tate, *Homology of Noetherian and local rings*, Illinois J. Math., (1957), 14–28.
- [21] C. W. Wilkerson, *Integral closure of unstable Steenrod algebra actions*, J of P. and P. Algebra, **13** (1978), 49–55.
- [22] A. Zabrodsky, *Maps Between Classifying Spaces*, Univ. of Jerusalem Preprint 1984.
- [23] J. Lannes and S. Zarati, *Sur les U-Injectifs*, Ann. École. Norm. Sup. 4^{ème}, **19** (1986), 1–31.
- [24] L. Smith, *On the invariant theory of finite pseudo reflection groups*, Arch. Math., **44** (1985), 222–228.

Received April 16, 1985.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720
H. SAMELSON
Stanford University
Stanford, CA 94305
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906–1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA	UNIVERSITY OF OREGON
UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA	UNIVERSITY OF HAWAII
MONTANA STATE UNIVERSITY	UNIVERSITY OF TOKYO
UNIVERSITY OF NEVADA, RENO	UNIVERSITY OF UTAH
NEW MEXICO STATE UNIVERSITY	WASHINGTON STATE UNIVERSITY
OREGON STATE UNIVERSITY	UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1987 by Pacific Journal of Mathematics

Charles James Keith Batty, Derivations on the line and flows along orbits	209
Paul Erdős, Adolf J. Hildebrand, Andrew Odlyzko, Paul Pudaite and Bruce Reznick, The asymptotic behavior of a family of sequences	227
Gregory James Galloway, A note on the fundamental group of a compact minimal hypersurface	243
Nigel Higson, A characterization of KK -theory	253
Anthony To-Ming Lau and Wataru Takahashi, Weak convergence and nonlinear ergodic theorems for reversible semigroups of nonexpansive mappings	277
Pere Menal and Juame Moncasi, Lifting units in self-injective rings and an index theory for Rickart C^* -algebras	295
Yoshimi Saito, Schrödinger operators with a nonspherical radiation condition	331
Larry Smith, Realizing certain polynomial algebras as cohomology rings of spaces of finite type fibered over $\times BU(d)$	361
Carl E. Swenson and Calvin T. Long, Necessary and sufficient conditions for simple A -bases	379
Kenneth Giovanni Valente, The p -primes of a commutative ring	385