

Pacific Journal of Mathematics

EXTENSIONS OF GENERALIZED HOMOLOGY THEORIES

FRIEDRICH-WILHELM BAUER

EXTENSIONS OF GENERALIZED HOMOLOGY THEORIES

FRIEDRICH W. BAUER

James Dugundji in memoriam

Relations between different kinds of homology theories on the category $\overline{\text{Com}}$ (of compacta) resp. the related strong shape category $\overline{\text{Com}}$ are studied. In particular homology theories satisfying a *clusteraxiom* (as for example the strong shape homology \overline{E}_* with coefficients in a spectrum \underline{E} , for a restricted class of spectra being defined on the category of finite dimensional compacta) allow interesting characterizations. As an application this provides new proofs of classical theorems concerning Steenrod-Sitnikov homology theories, including a result on the Brown-Douglas-Fillmore homology ε_* .

0. Let \underline{E} be any CW-spectrum and \underline{E}_* the corresponding homology theory [1] on the category \underline{P}_0 of based, finite CW-spaces.

We are concerned with the following questions:

(1) Can \underline{E}_* be extended over the categories $\overline{\text{Com}}$ (= based compact metric spaces) or alternatively over \overline{K} , the strong shape category?

(2) Under what circumstances is such an extension uniquely determined?

These two problems are both extension problems. The first one deals with the extension of \underline{E}_* itself, while the second one requires the extension of a given isomorphism between two homology theories.

The solution of both problems deserves interest because it turns out that numerous problems in topology and analysis can be reduced to (1) or (2).

There are for example two "natural" (not in the sense of "canonical") extensions of complex K -homology $\overline{\text{BU}}_*$ over a category containing \underline{P}_0 : (1) The homology theory $\overline{\text{BU}}_*$ defined on $\overline{\text{Com}}$ (the subcategory of \overline{K} , determined by compact metric spaces) and (2) the Brown-Douglas-Fillmore homology theory ε_* defined on $\overline{\text{Com}}$ ([6], [8], [9]) by purely functional analytic methods. Theorem 6.6 confirms that ε_* and \overline{E}_* are naturally isomorphic for finite dimensional spaces in $\overline{\text{Com}}$.

D. Edwards and H. Hastings [7] as well as the authors of [8] also dealt with the problem of establishing generalized, so called *Steenrod homology theories*. The relation between the homology theory ${}^s\overline{E}_*$ of [8] and our \overline{E}_*

is the subject of [3], while Theorem 5.7 of the present paper confirms the existence of an isomorphism between ${}^s\bar{E}_*$ and the Edwards-Hastings kind of Steenrod extensions of a homology theory (cf. remark at the end of §7). The strong shape category \bar{K} (resp. the homotopy category \bar{K}_h (which is used in this paper (cf. §7) differs formally from that of [7]. However Ju. Lisica [11] established in the meantime the equivalence of \bar{K}_h with the strong shape category s-Sh of [7]. This clarifies the relations between different concepts of Steenrod homology theories, shape homology theories resp. the related strong shape categories, which were introduced independently but approximately at the same time.

J. Milnor characterized in [12] ordinary homology theories (i.e. those with $\bar{E} = K(G)$, an Eilenberg-Mac Lane spectrum) on Com axiomatically. We get back this result in the following form (Theorem 6.4.): $\bar{K}(G)_*$ on Com , is up to an isomorphism, uniquely determined by the Eilenberg-Steenrod axioms with a strong excision axiom (§5 A2)) and the cluster-axiom (§5 A1)). This is a corollary of a more general result (Corollary 5.7) which deals with arbitrary CW-spectra \bar{E} having the property that a clusteraxiom for \bar{E}_* on compact metric spaces is valid. Because the spectrum BU (the classifying spectrum of complex K -theory [1]) turns out to be, at least for finite dimensional compact metric spaces (Proposition 4.4), of this kind, the above mentioned result on ϵ_* is another corollary of this assertion.

The extension problem can be approached from a somewhat different point of view leading to the concept of *shape singular homology theory* with coefficients in a CW spectrum \bar{E} (§§1–3): Let \bar{K} be the strong shape category with arbitrary based spaces. Does there exist an extension of \bar{E}_* over \bar{K}_h (the corresponding homotopy category) having the following two properties:

- (1) Every weak homotopy equivalence in \bar{K}_h (Definition 1.2) induces an isomorphism of the homology groups.
- (2) The extended homology theory has compact carrier (Definition 1.7)?

Theorem 1.6 provides us with an affirmative answer to this question. Moreover the extension (denoted by \tilde{E}_*) is unique up to an isomorphism (Theorem 3.1).

Even for compact metric X we can in general not expect to obtain an isomorphism $\tilde{E}_*(X) \approx \bar{E}_*(X)$. However one has for all X an isomorphism (Theorem 6.1):

$$\tilde{E}_*(X) \approx \bar{E}_*(|\bar{S}(X)|),$$

where $\bar{S}(X)$ denotes the shape-singular complex of the space X . So \tilde{E}_* reveals itself as shape singular homology. The question under what circumstances \tilde{E}_* is isomorphic to \bar{E}_* is settled in Theorem 6.1. This result in combination with Theorem 6.2 (asserting that Steenrod-Sitnikov homology groups $H_*^S(X; G)$ being for all compact metric X and all abelian groups G naturally isomorphic to $\overline{K(G)}_*(X)$) yields another proof of Theorem 7.7 in [2], claiming that $H_*^S(X; S)$ is naturally isomorphic to $H_*(|\bar{S}(X)|; \mathbf{Z})$ for shape connected compacta X . It turns out that for s -continua, \tilde{E}_* can be regarded as the appropriate generalization of Borel-Moore homology theory (cf. [10]) while \bar{E}_* corresponds to Steenrod-Sitnikov homology theory. In a final section (§7) we indicate the construction of the shape category and of the shape homology in this category with coefficients in a CW-spectrum. This is done without going into the details, referring to the relevant expositions (cf. [4]). The present paper is very closely related (but independent of) [3], which has already appeared. Therefore [3] refers occasionally to the present paper (concerning some details of proofs).

1. The existence theorem. Unless stated otherwise, we denote in the first three sections by \underline{K} the category Top_0 . We are working within the strong shape category \bar{K} resp. the associated homotopy category \bar{K}_h , whose construction is briefly recorded in §7. For more details see [3], [4].

1.1. DEFINITION. A morphism $\bar{f} \in \bar{K}(X, Y) = \bar{K}((X, x_0), (Y, y_0))$ is called a *weak homotopy equivalence* whenever

$$|\bar{S}(\bar{f})| \in \text{Top}_0(|\bar{S}(X)|, |\bar{S}(Y)|)$$

is a homotopy equivalence, $\bar{S}(\)$ denoting the shape singular complex, cf. §7. Let $\mathfrak{S}_* = \{\mathfrak{S}_n\}: \bar{K}_h \rightarrow \text{Ab}^{\mathbf{Z}}$ ($=$ category of \mathbf{Z} -graded abelian groups) be a functor, then we introduce the following *Whitehead axiom*:

1.2. DEFINITION. (W) Let $\bar{f} \in \bar{K}(X, Y)$ be a weak homotopy equivalence, then $\mathfrak{S}_*(\bar{f})$ is an isomorphism.

1.3. DEFINITION. A functor $\mathfrak{S}_*: \bar{K}_h \rightarrow \text{Ab}^{\mathbf{Z}}$ together with a natural transformation $\sigma: \mathfrak{S}_*(\) \rightarrow \mathfrak{S}_{*+1}(\Sigma \)$ ($\Sigma =$ reduced suspension) is called a *shape singular homology theory* $\mathfrak{S}_* = \{\mathfrak{S}_n, \sigma\}$ on \bar{K}_h whenever \mathfrak{S}_* satisfies the Whitehead axiom (W). We will during this and the two ensuing sections simply talk about a *homology theory* \mathfrak{S}_* . A natural transformation $\varphi: \mathfrak{S}_* \rightarrow \mathfrak{S}'_*$ between the functors $\mathfrak{S}_*, \mathfrak{S}'_*$ is called a

transformation of homology theories whenever φ commutes with the corresponding transformations $\sigma: \mathfrak{H}_*() \rightarrow \mathfrak{H}_{*+1}(\Sigma)$ resp. $\sigma': \mathfrak{H}'_*() \rightarrow \mathfrak{H}'_{*+1}(\Sigma)$.

REMARK. Here we do not require that the natural transformation σ is an isomorphism. This changes in §5.

Let $\underline{P}_1(\underline{P}_0)$ be the category of (compact) CW-spaces, understood as a subcategory of \underline{K} . Because of 7.1 \underline{P}_{1h} is a full subcategory of \underline{K}_h . Our first aim is to extend a given homology theory $\mathfrak{H}_*: \underline{P}_{1h} \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ over \underline{K}_h . We need ([2] Theorem 5.1.c).

1.4. LEMMA. *Let $X \in \underline{\text{Top}}_0$ be any based space, then the natural transformation*

$$\bar{\omega}_X \in \underline{K}(|\bar{S}(X)|, X)$$

(cf. §7) *is a weak homotopy equivalence. Every weak homotopy equivalence induces isomorphisms*

$$\bar{\pi}_*(\bar{f}) \quad (\text{with } \bar{\pi}_n(X, x_0) = \underline{K}_h((S^n, *), (X, x_0))).$$

Suppose that $X, Y \in \underline{\text{Top}}_0$ are s-connected (= shape connected, i.e. $\bar{\pi}_0(X) = \bar{\pi}_0(Y) = 0$); then also the converse holds.

Proof. The first part follows because $|\bar{S}(\bar{\omega}_X)|: |\bar{S}(|\bar{S}(X)|)| \rightarrow |\bar{S}(X)|$ is evidently (because everything happens in \underline{P}_1) a homotopy equivalence. Let $\bar{f} \in \underline{K}(X, Y)$ be a weak homotopy equivalence. Then we have the commutative diagram

$$\begin{array}{ccc} X & \xrightarrow{\bar{f}} & Y \\ \bar{\omega}_X \uparrow & & \downarrow \bar{\omega}_Y \\ |\bar{S}(X)| & \xrightarrow{f_{\#}} & |\bar{S}(Y)| \end{array} \quad |\bar{S}(\bar{f})| = f_{\#},$$

$\bar{\pi}_*(\bar{\omega}_X)$, $\bar{\pi}_*(\bar{\omega}_Y)$ being isomorphisms. We have $\bar{\pi}_*(|\bar{S}(X)|) \approx \pi_*(|\bar{S}(X)|)$ (resp. for Y) so that together with $\pi_*(f_{\#})$ also $\bar{\pi}_*(f_{\#})$ and therefore $\bar{\pi}_*(\bar{f})$ is an isomorphism. This proves the second part. Assume X, Y being s-connected; then $|\bar{S}(X)|, |\bar{S}(Y)|$ are connected. Suppose that $\bar{\pi}_*(\bar{f})$ is an isomorphism; then $\pi_*(f_{\#})$ is an isomorphism and the conclusion follows from the classical Whitehead theorem.

This enables us to define:

$$(1) \quad \tilde{\mathfrak{H}}_*(X) = \mathfrak{H}_*(|\bar{S}(X)|)$$

and for any $\bar{f} \in \bar{K}(X, Y)$

$$(2) \quad \tilde{\mathfrak{S}}_*(\bar{f}) = \mathfrak{S}_*(|\bar{S}(\bar{f})|).$$

We get a natural transformation $\tilde{\sigma}: \tilde{\mathfrak{S}}_*() \rightarrow \tilde{\mathfrak{S}}_{*+1}(\Sigma)$ as follows: We have a natural mapping

$$\lambda: S^1 \wedge |\bar{S}(X)| \rightarrow |\bar{S}(S^1 \wedge X)| = |\bar{S}(\Sigma X)|$$

for any space X (cf. Theorem 4.1 in [4] and Theorem 2.1 below). So we can define

$$\tilde{\sigma}: \tilde{\mathfrak{S}}_*(X) \rightarrow \tilde{\mathfrak{S}}_{*+1}(\Sigma X)$$

by composing $\tilde{\sigma}$ with $\mathfrak{S}_{*+1}(\lambda)$. Lemma 1.4 implies:

1.5. LEMMA. Let $\mathfrak{S}_*: \underline{P}_{1h} \rightarrow \text{Ab}^{\mathbb{Z}}$ be a given homology theory, then $\mathfrak{S}_*: \bar{K}_h \rightarrow \text{Ab}^{\mathbb{Z}}$ is a homology theory (i.e. one, satisfying the Whitehead axiom \underline{W}). Moreover the natural transformation $\bar{\omega}: |\bar{S}()| \rightarrow 1_{\bar{K}}$ induces a natural isomorphism

$$\tilde{\mathfrak{S}}_*|_{\underline{P}_1} \approx \mathfrak{S}_*.$$

We summarize:

1.6. THEOREM. To each homology theory $\mathfrak{S}_*: \underline{P}_{1h} \rightarrow \text{Ab}^{\mathbb{Z}}$ there exists a homology theory $\tilde{\mathfrak{S}}_*: \bar{K}_h \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ and a natural isomorphism $\tilde{\mathfrak{S}}_*|_{\underline{P}_1} \approx \mathfrak{S}_*$.

The importance of the concept of *compact support* is well-known from classical algebraic topology. In shape theory we have to define this notion in the following way:

1.7. DEFINITION. A homology theory $\mathfrak{S}_*: \bar{K}_h \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ is defined to have *compact support* whenever the following holds:

(C) (a) To any $\zeta \in \mathfrak{S}_n(X)$, $X \in \bar{K}$ there exists a compact metric K , a $\bar{f} \in \bar{K}(K, X)$ and a $\zeta' \in \mathfrak{S}_n(K)$ such that

$$\mathfrak{S}_n(\bar{f})(\zeta') = \bar{f}_*(\zeta') = \zeta.$$

(b) Let ζ' , K , \bar{f} be as in (a) and assume that $\bar{f}_*(\zeta') = 0$. Then there exists a homotopy commutative diagram in \bar{K}

$$\begin{array}{ccc} K & \xrightarrow{\bar{f}} & X \\ \bar{r} \searrow & & \nearrow i \\ & L & \end{array}$$

with compact metric L , such that $\bar{r}_*(\zeta') = 0$. □

On the other hand let $\mathfrak{S}_*: \underline{P}_{1h} \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ be a homology theory, then we say that \mathfrak{S}_* has compact support, whenever the following holds:

(PC) (a) Let $\zeta \in \mathfrak{S}_*(X)$ be any element, then there exists a compact subspace $K \subset X$, $K \in \underline{P}_1$ and a $\zeta' \in \mathfrak{S}_*(K)$ such that $\mathfrak{S}_*(i)(\zeta') = \zeta$, $i: K \subset X$ denoting the inclusion.

(b) Let ζ' , K , be as in (a) such that now $i_*(\zeta') = \mathfrak{S}_*(i)(\zeta') = 0$, then there exists a $L \in \underline{P}_0$, $K \subset L \subset X$, such that $\mathfrak{S}_*(j)(\zeta') = 0$ ($j: K \subset L$ being the inclusion).

REMARK. The difference between (C) and (PC) is caused by the fact that a continuous mapping $f: K \rightarrow X \in \underline{P}_1$, \bar{K} compact, has *always* a compact image. This is no longer true for shape mappings: Even for a point $*$ and a shape mapping $\bar{f} \in \bar{K}(*, X)$ into a non-compact space X , there does in general not exist a compact $K \subset X$ over which \bar{f} factors.

The relation between (C) and (PC) is embodied in the following:

1.8. PROPOSITION. Let $\mathfrak{S}_*: \underline{P}_{1h} \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ be a homology theory satisfying (PC). Then the extension $\tilde{\mathfrak{S}}_*() = \mathfrak{S}_*(|\bar{S}()|): \bar{K}_h \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ of \mathfrak{S}_* over \bar{K}_h satisfies (C).

Proof. Let $\zeta \in \mathfrak{S}_*(|\bar{S}(X)|)$ be a given element, then there exists a finite subcomplex $K \subset |\bar{S}(X)|$ and a $\zeta'' \in \mathfrak{S}_*(K)$ such that $\mathfrak{S}_*(i)(\zeta'') = i_*(\zeta'') = \zeta$.

Moreover we have the following commutative diagram

$$\begin{array}{ccccc}
 K & \xrightarrow[\subset]{} & |\bar{S}(X)| & \xrightarrow{\bar{\omega}_X} & X \\
 \bar{\omega}_K \uparrow & & \uparrow \bar{\omega}_{|\bar{S}(X)|} & & \uparrow \bar{\omega}_X \\
 |\bar{S}(K)| & \xrightarrow{|\bar{S}(i)|} & |\bar{S}(|\bar{S}(X)|)| & \xrightarrow{|\bar{S}(\bar{\omega}_X)|} & |\bar{S}(X)|
 \end{array}$$

where $\bar{\omega}_K$ is a homotopy equivalence in \bar{K} between CW spaces, hence homotopic to a continuous homotopy equivalence ω in \underline{K} (cf. 7.1). The mapping $\tilde{\mathfrak{S}}_*(\bar{\omega}_X) = \mathfrak{S}_*(|\bar{S}(\bar{\omega}_X)|)$ is an isomorphism (cf. Lemma 1.5). Setting $\zeta' = \mathfrak{S}_*(\omega)^{-1}(\zeta'')$ and $\bar{f} = \bar{\omega}_X i$, we conclude

$$\mathfrak{S}_*(|\bar{S}(\bar{\omega}_X)|) \mathfrak{S}_*(|\bar{S}(\bar{f})|)(\zeta') = \mathfrak{S}_*(|\bar{S}(\bar{\omega}_X)|) \mathfrak{S}_*(i) \mathfrak{S}_*(\omega)(\zeta''),$$

therefore

$$\tilde{\mathfrak{S}}_*(\bar{f})(\zeta') = \mathfrak{S}_*(i)(\zeta'') = \zeta.$$

This proves (C) (a). The proof of (C) (b) is similar.

2. The homology theory \bar{E}_* . A spectrum $\underline{E} = \{E_n, \sigma_n: E_n \rightarrow E_{n+1}, n \in \mathbb{Z}\}$ is a sequence of based spaces and continuous maps σ_n . A CW spectrum has the additional property that all E_n are CW complexes and that all σ_n are supposed to be cellular. The category $\underline{\text{Spec}}$ has these CW spectra as objects and so-called functions of spectra $\underline{f} = \{f_n: E_n \rightarrow F_n\}$ as morphisms, where we assume that the f_n are compatible with the corresponding $\sigma_n: \Sigma E_n \rightarrow E_{n+1}$ resp. $\Sigma F_n \rightarrow f_{n+1}$.

The following full subcategory $\underline{\text{CSpec}} \subset \underline{\text{Spec}}$ deserves particular interest: Its objects are those spectra $\underline{E} = \{E_n\}$, having the property that (1) all m -skeletons $(E_n)^m$ are for all m and all $n \in \mathbb{Z}$ compact and that (2) there exists a n_0 such that all E_n , $n \geq n_0$ are simply connected. We call such a spectrum also a *cs-spectrum*. It can be verified that an Eilenberg-Mac Lane spectrum $\underline{K(G)}$ for finitely generated abelian group G has this property.

Let $\underline{\text{Com}} \subset \bar{K}$ be the full subcategory of based compacta, then we have defined in [4] the homology theory

$$(1) \quad \bar{E}_n(X) = \varinjlim_k \bar{\pi}_{n+k}(E_k \bar{\wedge} X), \quad X \in \underline{\text{Com}}.$$

We have that $\bar{E}_*: \underline{\text{Com}}_h \rightarrow \text{Ab}^{\mathbb{Z}}$ is a functor. Moreover there are natural isomorphisms $\sigma = \sigma_n: \bar{E}_n \approx \bar{E}_{n+1} \cdot \Sigma$. However the Whitehead axiom (W) is not necessarily fulfilled so $\bar{E}_* = \{\bar{E}_n, \sigma\}$ is not always a shape singular homology theory (cf. Definition 1.3).

A space $X \in \underline{\text{Com}}$ is called *s-continuum*, whenever $\bar{\pi}_0(X) = 0$.

So the solenoid for example is not *s-connected*, while it can be proved that the double suspension of the solenoid is a *s-continuum*. Let E be a CW complex, X a compactum, then we have a homotopy equivalence

$$E \wedge |\bar{S}(X)| \simeq |\bar{S}(E)| \wedge |\bar{S}(X)|$$

and a continuous mapping $\lambda: |\bar{S}(E)| \wedge |\bar{S}(X)| \rightarrow |\bar{S}(E \bar{\wedge} X)|$. As a consequence of Theorem 4.1 in [4] we have:

2.1. THEOREM. *Let X be a *s-continuum*, E a simply connected CW complex of finite type (i.e. all m -skeletons are compact), then λ induces a natural homotopy equivalence*

$$|\bar{S}(E \bar{\wedge} X)| \stackrel{\simeq}{\leftarrow} |\bar{S}(E)| \wedge |\bar{S}(X)| \simeq E \wedge |\bar{S}(X)|,$$

rendering the following diagram homotopy commutative:

$$\begin{array}{ccc}
 |\bar{S}(E)| \wedge |\bar{S}(X)| & \xrightarrow{\bar{\omega}_E \wedge 1} & E \wedge |\bar{S}(X)| \\
 \downarrow \parallel & & \downarrow 1 \wedge \bar{\omega}_X \\
 |\bar{S}(E \bar{\wedge} X)| & \xrightarrow{\bar{\omega}_{E \bar{\wedge} X}} & E \bar{\wedge} X
 \end{array}$$

We define $\underline{E}_n(X) = \varinjlim_k \pi_{n+k}(E_k \bar{\wedge} X)$ (all mappings and homotopies being continuous). As a consequence we have:

2.2. THEOREM. (Cf. [4] Theorem 4.2.) *Let the s -continuum X and the spectrum $\underline{E} \in \underline{\text{CSpec}}$ be given, then there exists a natural isomorphism*

$$\bar{E}_*(X) \approx \underline{E}_*(|\bar{S}(X)|).$$

Proof. We have the following series of isomorphisms

$$\begin{aligned}
 \bar{E}_n(X) &= \varinjlim_k \bar{\pi}_{n+k}(E_k \bar{\wedge} X) \approx \varinjlim_k \pi_{n+k}(|\bar{S}(E_k \bar{\wedge} X)|) \\
 &\approx \varinjlim_k \pi_{n+k}(E_k \wedge |\bar{S}(X)|) = \underline{E}_n(|\bar{S}(X)|),
 \end{aligned}$$

which are clearly compatible with the natural transformations σ .

In accordance with §1(1), (2) we denote the homology theory

$$\underline{E}_*(|\bar{S}(\)|): \bar{K}_h \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$$

by \tilde{E}_* .

2.3. PROPOSITION. *Let $f: \underline{E} \rightarrow \underline{F}$ be any morphism in $\underline{\text{Spec}}$, then we have (1) an induced natural transformation*

$$f_{\#}: \bar{E}_* \rightarrow \bar{F}_*$$

and (2) a natural transformation

$$\tilde{f}_{\#}: \tilde{E}_* \rightarrow \tilde{F}_*.$$

Proof. This is in both cases an immediate consequence of the definitions of \bar{E}_* resp. of \tilde{E}_* : For example in the first case we have

$$E_k \bar{\wedge} X \xrightarrow{f_k \bar{\wedge} 1_X} F_k \bar{\wedge} X$$

which is well-defined (cf. [4] §2). This provides us easily with the required $f_{\#}$. \square

2.4. PROPOSITION. *Let $\underline{E} \in \underline{\text{Spec}}$, $X \in \underline{K}$ and $\zeta \in \tilde{\underline{E}}_n(X)$ be given, then there exists (1) a $\underline{\text{cs-spectrum}}$ \underline{E}' (2) a $\underline{\varphi} \in \underline{\text{Spec}}(\underline{E}', \underline{E})$ and (3) a $\zeta' \in \tilde{\underline{E}}'_n(X)$ such that $\tilde{\varphi}_{\#}(\zeta') = \zeta$. Moreover assume ζ' , \underline{E}' , $\underline{\varphi}$ be given such that $\tilde{\varphi}_{\#}(\zeta') = 0$, then there exists a $\underline{E}'' \in \underline{\text{CSpec}}$ as well as a factorization*

$$\begin{array}{ccc} \underline{E}' & \xrightarrow{\underline{\varphi}} & \underline{E} \\ \psi \searrow & & \nearrow \eta \\ & \underline{E}'' & \end{array}$$

such that $\tilde{\psi}_{\#}(\zeta') = 0$.

Proof. We abbreviate $|\bar{S}(X)|$ by Y and call a $\underline{E} \in \underline{\text{Spec}}$ *simple* whenever there exists a $n_0 \in \mathbb{Z}$ such that all E_n , $n \geq n_0$ are simply connected. Let us in a first step deal with a simple spectrum \underline{E} . We construct \underline{E}' , $\underline{\varphi}$ in the following way:

Let $(f: S^{n+k} \rightarrow E_k \wedge Y)$ be a representative of ζ (observe that Y is a CW space!). Then there exists a compact $E'_k \subset E_k$ such that f factors over $E'_k \wedge Y$. By eventually suspending f we obtain a E'_k which is simply connected. Now we set $E'_l = *$ for $l < k$. In order to establish E'_{k+1} we take $\sigma_k(\Sigma E'_k) \subset E_{k+1}$ and kill the fundamental group of this space in E_{k+1} (which is by assumption simply connected). This provides us with a $E'_{k+1} \subset E_{k+1}$ and a $\sigma'_k: \Sigma E'_k \rightarrow E'_{k+1}$. Proceeding inductively we obtain a $\underline{\varphi}: \underline{E}' \rightarrow \underline{E}$ such that $\underline{E}' \in \underline{\text{CSpec}}$ and a $\zeta' \in \underline{E}_*(Y)$ with $\underline{\varphi}_{\#}(\zeta') = \zeta$. The second statement is proved similarly.

This completes the proof of the assertion for simple spectra. In order to get rid of this last assumption, we construct to any $\underline{E} \in \underline{\text{Spec}}$ the following spectrum $\hat{\underline{E}} = \{\hat{E}_n\}$:

$$\hat{E}_n = \Sigma^2 E_{n-2},$$

$$\Sigma^2 \sigma_{n-2} = \hat{\sigma}_n: \Sigma \hat{E}_n = \Sigma^3 E_{n-2} \rightarrow \Sigma^2 \hat{E}_{n-1} = \hat{E}_{n+1}.$$

This spectrum is clearly simple. There exists a morphism $\tau_{\underline{E}} = \tau: \hat{\underline{E}} \rightarrow \underline{E}$, $\tau_n = \sigma_{n-1} \cdot \Sigma \sigma_{n-2}: \hat{E}_n = \Sigma^2 E_{n-2} \rightarrow E_n$, $\tau \in \underline{\text{Spec}}(\hat{\underline{E}}, \underline{E})$. On \underline{P}_1 this transformation clearly induces a natural isomorphism

$$\tau_{\#}: \hat{E}_n(\) \approx E_n(\),$$

hence a natural isomorphism

$$\tau_{\#}: \tilde{\hat{\underline{E}}}_n(\) \approx \tilde{\underline{E}}_n(\).$$

Now we can apply the previous construction of \underline{E}' to $\hat{\underline{E}}$ (instead of \underline{E}) thereby completing the proof of 2.4.

REMARK. There exist a s -continuum X and a not finitely generated abelian group G such that

$$\overline{K(G)}_n(X) \neq \overline{K(G)}_n(X).$$

(Cf. remark following 6.5.)

In particular this confirms that 2.4 does not hold for $\overline{K(G)}_*$ (even not for s -connected compacta).

2.5. THEOREM. *The homology theory \tilde{E}_* : $\bar{K}_h \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ has compact support (i.e. satisfies \underline{C}) in Definition 1.7.*

Proof. The fact that \tilde{E}_* is a homology theory follows from Lemma 1.5. Because we have a natural isomorphism $\underline{E}_*() \approx \tilde{E}_*()$ on \underline{P}_1 and since \underline{E}_* on \underline{P}_1 has compact support, the assertion follows from 1.8. \square

REMARK. In general we cannot expect that

$$\sigma: \tilde{E}_n(X) \rightarrow \tilde{E}_{n+1}(\Sigma X)$$

is becoming an isomorphism, unless X is a shape simply connected compactum (i.e. one has $\pi_i(X) = 0$ for $i = 0, 1$). In the latter case we have due to Theorem 4.1 in [4] a homotopy equivalence

$$|\bar{S}(S^1 \wedge X)| \simeq S^1 \wedge |\bar{S}(X)|$$

furnishing an isomorphism

$$\tilde{E}_n(X) = \underline{E}_n(|\bar{S}(X)|) \approx \underline{E}_{n+1}(S^1 \wedge |\bar{S}(X)|) = \tilde{E}_{n+1}(\Sigma X).$$

3. **The uniqueness theorem.** Let $\mathfrak{S}_*, \mathfrak{S}'_*: \bar{K}_h \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ be two homology theories (i.e. those satisfying (W)). Suppose we have a natural transformation of homology theories $\alpha: \mathfrak{S}_*|_{\underline{P}_{1h}} \rightarrow \mathfrak{S}'_*|_{\underline{P}_{1h}}$.

3.1. THEOREM. *There exists a unique extension $\tilde{\alpha}: \mathfrak{S}_* \rightarrow \mathfrak{S}'_*$ of α over \bar{K}_h .*

Proof. The natural transformation $\bar{\omega}_X: |\bar{S}(X)| \rightarrow X$ is a weak homotopy equivalence in \bar{K}_h , hence we are allowed to define

$$\tilde{\alpha}_X: \mathfrak{S}_*(X) \rightarrow \mathfrak{S}'_*(X)$$

by

$$(1) \quad \tilde{\alpha}_X = \mathfrak{S}'_*(\bar{\omega}_X) \alpha_{|\bar{S}(X)|} \mathfrak{S}_*(\bar{\omega}_X)^{-1}.$$

The following assertions are all more or less immediate.

(1) $\tilde{\alpha}$ is natural with respect to X .

Proof. This is obvious because $\bar{\omega}: |\bar{S}(\cdot)| \rightarrow 1_{\bar{K}}$ as well as α are natural.

(2) $\tilde{\alpha}$ is a homomorphism.

This is trivial.

(3) $\tilde{\alpha} \mid \underline{P}_{1h} = \alpha$.

Proof. Let $X \in \underline{P}_1$ be a CW space, then we have a commutative diagram

$$(2) \quad \begin{array}{ccc} \mathfrak{S}_*(X) & \xleftarrow{\mathfrak{S}_*(\bar{\omega}_X)} & \mathfrak{S}_*(|\bar{S}(X)|) \\ \alpha_X \downarrow & & \downarrow \alpha_{|\bar{S}(X)|} \\ \mathfrak{S}'_*(X) & \xleftarrow{\mathfrak{S}'_*(\bar{\omega}_X)} & \mathfrak{S}'_*(|S(X)|) \end{array}$$

and therefore

$$\tilde{\alpha}_X = \mathfrak{S}'_*(\bar{\omega}_X) \alpha_{|\bar{S}(X)|} \mathfrak{S}_*(\bar{\omega}_X)^{-1} = \alpha_X.$$

(4) $\tilde{\alpha}$ is compatible with the bonding maps σ, σ' .

Proof. This follows immediately because α is supposed to have this property on the category P_1 .

(5) Let $\beta: \mathfrak{S}_* \rightarrow \mathfrak{S}'_*$ be any natural transformation extending α , then we have $\beta = \tilde{\alpha}$.

Proof. This is trivial in view of (2) (now for any $X \in \underline{K}$ and after replacing α_X by β_X).

This completes the proof of Theorem 3.1.

We have the following corollaries:

3.2. COROLLARY. *Let $\alpha: \mathfrak{S}_*|_{\underline{P}_{1h}} \approx \mathfrak{S}'_*|_{\underline{P}_{1h}}: \underline{P}_{1h} \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ be an isomorphism of homology theories, then there exists a unique isomorphism $\tilde{\alpha}: \mathfrak{S}_* \approx \mathfrak{S}'_*$ on \underline{K}_h , extending α .*

3.3. COROLLARY. *Let $\mathfrak{S}_*: \underline{\bar{K}}_h \rightarrow \text{Ab}^{\mathbb{Z}}$ be any homology theory and $\alpha: \mathfrak{S}_* \approx \underline{E}_*$ an isomorphism of homology theories on the category \underline{P}_{1h} . Then there exists a unique extension $\tilde{\alpha}: \mathfrak{S}_* \approx \tilde{E}_*$ of α to an isomorphism on the category $\underline{\bar{K}}_h$. Furthermore \mathfrak{S}_* is of compact support (Definition 1.7).*

Proof. The first part follows simply by applying Theorem 3.1 to $\mathfrak{S}'_* = \tilde{E}_*$. The second part follows from the first because 2.5 makes sure that $\tilde{E}_* \approx \mathfrak{S}_*$ is of compact support.

4. \bar{E}_* and the clusteraxiom. In [12] J. Milnor introduced the *clusteraxiom* (or strong wedge axiom) for a homology theory H_* defined on the category of compact metric spaces (with base-points).

(C1) Let $(X_i, x_{0i}) = X_i, i = 1, 2, \dots$ be a sequence of compact metric spaces and

$$\text{Cl}_{i=1}^{\infty} X_i = \varprojlim_m (X_1 \vee \dots \vee X_m)$$

the cluster (or strong wedge), then the natural homomorphism

$$H_*\left(\text{Cl}_{i=1}^{\infty} X_i\right) \rightarrow \prod_{i=1}^{\infty} H_*(X_i)$$

becomes an isomorphism.

He was able to show that the Steenrod-Sitnikov homology $H_*^S(X, Y; G)$ groups with arbitrary coefficient group G can be characterized (now in the category of compact metric pairs) by the ordinary Eilenberg-Steenrod axioms (with a strong excision axiom $H_*^S(X, Y; G) \approx H_*^S(X/Y, *; G)$) together with the clusteraxiom.

In this section we are going to find out under what circumstances $\bar{E}_*h: \underline{\text{Com}} \rightarrow \text{Ab}^{\mathbb{Z}}$ for a given CW spectrum \underline{E} fulfills the cluster axiom.

We formulate the following property of a CW-spectrum:

(S) There exists a $p \in \mathbb{Z}$ having the following property:

The mapping $\sigma: \Sigma E_i \rightarrow E_{i+1}$ induces an isomorphism of the $(2i + p)$ -skeleton for all i .

We are able to prove:

4.1. PROPOSITION. *Let $\underline{E} \in \underline{\text{Spec}}$ be such that (S) is satisfied, then the cluster axiom (C1) holds for \bar{E}_* .*

Proof. Set $X = \text{Cl}_{i=1}^{\infty} X_i$ and let

$$p_n^{n+1}: \bigvee_{i=1}^{n+1} X_i \rightarrow \bigvee_{i=1}^n X_i$$

resp.

$$q_n: \prod_{i=1}^{\infty} X_i \rightarrow X_n$$

be the projections. We have the natural homomorphism

$$\varphi: \underline{E}_*(X) \rightarrow \prod_{i=1}^{\infty} \underline{E}_*(X_i)$$

defined by

$$\varphi(\zeta) = \{q_{i*}(\zeta)\}.$$

In order to construct an inverse to φ , let $\{\zeta_i\} \in \prod_{i=1}^{\infty} \underline{E}_n(X_i)$ be any element in the direct product and $\zeta_i \ni \tilde{f}_i: S^{n+1(i)} \rightarrow E_{1(i)} \overline{\wedge} X_i$ be representatives. We fix a continuous $g_i: X_i \rightarrow P_i \in \underline{P}$ for each i and define

$$f_i = \tilde{f}_i(g_i): S^{n+1(i)} \rightarrow E_{1(i)} \wedge P_i.$$

We have to distinguish two cases:

1. $l(i) > n - p + 2$: We put $q = p - 2$ and observe that f_i factors over $(E_{l(i)} \wedge P_i)^{n+1(i)} \subset (E_{l(i)} \wedge P_i)^{2l(i)+q}$. However for cellular reasons we have:

$$(E_{l(i)} \wedge P_i)^{2l(i)+q} \subset (E_{l(i)})^{2l(i)+q} \wedge P_i$$

and, by our assumption (S), an isomorphism

$$(E_{l(i)})^{2l(i)+q} \wedge P_i \approx (\Sigma E_{l(i)-1})^{2l(i)+q} \wedge P_i.$$

Hence f_i turns out to be stably homotopic to a

$$f'_i: S^{n+1(i)} \rightarrow \Sigma(E_{l(i)-1} \wedge P_i)$$

because we have

$$(\Sigma E_{l(i)-1})^k \approx \Sigma(E_{l(i)-1})^{k-1}$$

for all $k \in N$, this procedure can be iterated until we reach a

$$f''_i: S^{n+1(i)} \rightarrow \Sigma^{l(i)-(n-q)}(E_{n-q} \wedge P_i).$$

Putting $k = l(i) - n + 1$ we have $n + l(i) = 2n - q + k$ and therefore we find

$$f''_i \in \pi_{2n-1+k}(\Sigma^k(E_{n-q} \wedge P_i)).$$

The stability theorem ensures the existence of an index m such that the suspension homomorphism

$$\Sigma_*: \pi_{2n-1+k}(\Sigma^k(E_{n-q} \wedge P_i)) \rightarrow \pi_{2n-1+k+1}(\Sigma^{k+1}(E_{n-q} \wedge P_i))$$

becomes an isomorphism for $k \geq m$. Therefore the number

$$l = n - q + m$$

has the property that to each f_i there always exists a representative $\hat{f}_i: S^{n+l} \rightarrow E_l \wedge P_i$. Observe that l is independent of i , $l(i)$ and g_i . As a result we can henceforth assume without loss of generality that $\bar{f}_i \in \zeta_i$ is of the form:

$$\bar{f}_i \in \bar{K}_h(S^{n+l}, E_l \bar{\wedge} X_i), \quad n + l \geq 1,$$

with a universal l (not depending upon i).

The second case:

2. $l(i) \leq n - p + 2$: can be settled by eventually suspending until we reach the same universal level l for all i .

Now it turns out to be rather simple to compose all these \bar{f}_i , yielding a $\psi\{\bar{f}_i\}: S^{n+l} \rightarrow E_l \bar{\wedge} X$.

To this end we must apply the explicit construction of a shape mapping, laid down, for example, in [4] appendix: Let $g: X \rightarrow P \in \underline{P}$ be any object in \underline{P}_X , then we factorize over a suitable

$$g^m: X \rightarrow \bigvee_{j=1}^m P_j, \quad g^m = \left(\bigvee_{j=1}^m g_j \right) p^m,$$

where

$$p^m: \bigcap_{i=1}^{\infty} X_i \rightarrow \bigvee_{i=1}^m X_i$$

denotes the projection and $g_j: X_j \rightarrow P_j$ is a given object in \underline{P}_{X_j} . Hence we merely have to evaluate $\psi\{\bar{f}_i\}$ on mappings of the form g^m . This can obviously be accomplished by

$$(1) \quad \psi\{\bar{f}_i\}(g^m) = (f_1 \vee \cdots \vee f_m) \kappa$$

where $\kappa: S^{n+l} \rightarrow S^{n+l} \vee \cdots \vee S^{n+l}$ denotes the m -fold comultiplication and where we put: $f_i = \bar{f}_i(g_i)$ as before. The verification of the fact that (1) establishes a

$$\psi\{\bar{f}_i\}: S^{n+l} \rightarrow E_l \bar{\wedge} X$$

is in view of the techniques developed in [4] appendix merely routine. This gives rise to a mapping $\psi: \prod_{i=1}^{\infty} \bar{E}_n(X_i) \rightarrow \bar{E}_n(X)$ having the property

$$(2) \quad \varphi\psi = 1.$$

We still have to verify that φ is also a monomorphism. To this end let

$$(\bar{f}: S^{n+r} \rightarrow E_r \overline{\wedge} X) \in \zeta \in \bar{E}_n(X)$$

be given and consider

$$\bar{f}_i = (1_{E_r} \overline{\wedge} q_i) \bar{f}: S^{n+r} \rightarrow E_r \overline{\wedge} X_i.$$

Assume the existence of indexes $l(i) \geq r$ (eventually depending upon i) such that the corresponding representatives $\bar{f}'_i: S^{n+l(i)} \rightarrow E_{l(i)} \overline{\wedge} X_i$ are null-homotopic.

The same kind of reasoning as above provides us with a universal index l and representatives

$$\bar{f}''_i: S^{n+l} \rightarrow E_l \overline{\wedge} X_i$$

such that $\bar{f}''_i \simeq 0$. Now these homotopies can be composed to a unified homotopy of the original \bar{f} to the constant map (in the same way in which we constructed $\psi\{\bar{f}_i\}$ before). This assures us that φ is a monomorphism and completes the proof of the proposition. \square

There are two basic applications of this result, which we are going to develop:

Recall that a spectrum \underline{E} is called *connective*, whenever

(1) \underline{E} is an Ω -spectrum (i.e. one has $E_i \simeq \Omega E_{i+1}$) and

(2) E_i is $(i-1)$ connected (a condition which is of course meaningless for $i < 0$).

We have the following general assertion:

4.2. LEMMA. *Let \underline{E} be connective, then there exists a spectrum $\underline{E}' = \{E'_i\} \in \underline{\text{Spec}}$ such that (S) holds as well as a function of spectra $\underline{f}: \underline{E} \rightarrow \underline{E}'$ with f_n being a homotopy equivalence for all n . In particular \underline{f} induces a natural isomorphism $\bar{E}_* \approx \bar{E}'_*$.*

Proof. This is a quite elementary fact which can be proved inductively: Setting $E'_i = E_i$ for $i \leq 0$, we assume that E'_m and f_m have already been constructed for $m \leq n$. According to the stability theorem we obtain a suspension isomorphism

$$\pi_{i+1}(E_{n+1}) \approx \pi_i(E'_n) \approx \pi_{i+1}(E'_n)$$

for $i \leq 2(n-1)$.

This isomorphism is easily recognized to be induced by $\tau_n: \Sigma E_n^2 \rightarrow E_{n+1}$ (with τ_n being defined by $\sigma_n = (\Sigma f_n)\tau_n$): Let σ'_n resp. p be the adjoints to σ_n resp. the identity $1: \Sigma E'_n \rightarrow \Sigma E'_n$, then we have

$$(\Omega \tau_n)p = \tau'_n.$$

As a result we have a commutative diagram

$$\begin{array}{ccc}
 \pi_{i+1}(E_{n+1}) & \approx & \pi_i(\Omega E_{n+1}) \\
 \sigma_{n\#} \uparrow & & \uparrow \Omega \sigma_{n\#} \\
 \pi_{i+1}(\Sigma E'_n) & \approx & \pi_i(\Omega \Sigma E'_n)
 \end{array}
 \begin{array}{c}
 \swarrow \sigma'_{n\#} \\
 \leftarrow p_{\#} \pi_i(E'_n)
 \end{array}$$

Since all induced homomorphisms on the right-hand side are isomorphism (for $i \leq 2(n-1)$), we conclude that $\sigma_{n\#}$ is also an isomorphism.

In defining E'_{n+1} , we set $(E'_{n+1})^{2n-3} = (\Sigma E'_n)^{2n-3}$ and recall that $\tau_n: \Sigma E'_n \rightarrow E_{n+1}$ induces an isomorphism of the homotopy groups π_i for $i \leq 2n-3$. We attach $2n-1$ -cells to $\Sigma E'_n$ killing the kernel of $\pi_{2n-2}(\tau_n)$ and $2n-2$ -cells representing the cokernel of $\pi_{2n-2}(\tau_n)$. As a result we find a new space $X_1 \supset \Sigma E'_n$ and a continuous $f_1: X \rightarrow E_{n+1}$, extending τ_n such that $\pi_i(f_1)$ is isomorphism for $i \leq 2n-2$. Proceeding inductively, we find f_2, X_2, \dots and finally a CW-complex $X \supset \Sigma E'_n$ as well as a continuous $f: X \rightarrow E'_n$ extending σ_n such that $\pi_i(f)$ becomes an isomorphism for all i . We define $X = E'_{n+1}$ and conclude that $(E'_{n+1})^{2n-3} = (\Sigma E'_n)^{2n-3}$, and that there exists a homotopy equivalence $f_{n+1}: E'_{n+1} \xrightarrow{\sigma'_n} E_{n+1}$ which is compatible with the resp. bonding maps. $\Sigma E'_n \xrightarrow{\sigma'_n} E'_{n+1}$ (being the inclusion $\Sigma E'_n \xrightarrow{\subset} X = E'_{n+1}$).

As a corollary we obtain our first application:

4.3. PROPOSITION. *Let G be any abelian group, then for $\overline{K(G)}_{\star}$ the clusteraxiom holds.*

Proof. Because $K(G)$, the Eilenberg-Mac Lane spectrum is connective, Lemma 4.2 and finally Proposition 4.1 applies.

Our second application deals with complex K -theory \underline{BU}_{\star} which is determined by the spectrum $\underline{BU} = \{E_n\}$ with

$$E_{2n} = \mathbf{Z} \times BU, \quad E_{2n+1} = U.$$

We obtain a connective spectrum $\underline{bu} = \underline{L} = \{L_n\}$ by killing the first $(i-1)$ bottom homotopy groups of \overline{E}_i in a well-known way (e.g. by taking the fibers of certain Postnikov-decompositions of E_i).

Consider the category $\underline{\text{Com}}^f$ of all finite dimensional compact metric spaces, then we have:

(1) An isomorphism

$$\underline{BU}^n(X) \approx \underline{L}^n(X)$$

for all CW-spaces X and $n < 0$.

(2) Let $X \in \underline{\text{Com}}^f$ be embedded in some S^{n+1} , then we have an isomorphism ([4] Theorem 7.1)

$$\underline{E}^{n-k}(S^{n+1} \setminus X) \approx \underline{E}_k(X).$$

(3) $\underline{\text{BU}}_*$ is periodic of period 2, i.e.

$$\underline{\text{BU}}_n(X) \approx \underline{\text{BU}}_{n+2}(X), \quad n \in \mathbf{Z}.$$

The first statement follows immediately by definition. The second one is simply Alexander duality. The third assertion can be achieved in the same way as in the classical case as an immediate consequence of Bott-periodicity.

4.4. PROPOSITION. *On the category $\underline{\text{Com}}^f$ the homology theory $\underline{\text{BU}}_*$ fulfills a clusteraxiom.*

Proof. Let $X = \text{Cl}_{i=1}^\infty X_i$, $X \in \underline{\text{Com}}^f$ be embedded in some S^N . In establishing the isomorphism

$$(3) \quad \underline{\text{BU}}_n(X) \approx \prod_{i=1}^\infty \underline{\text{BU}}_n(X_i)$$

we can in view of (3) without loss of generality assume that $N - n < 0$. Under these circumstances we obtain

$$(4) \quad \underline{\text{BU}}_n(X) \approx \underline{\text{BU}}^{N-n}(S^N \setminus X) \approx \underline{L}^{N-n}(S^N \setminus X) \approx \underline{L}_n(X).$$

Application of Lemma 4.1, 4.2 yields the clusteraxiom for $\underline{L}_n(X)$. However the isomorphism (4) is compatible with the inclusions $X_i \rightarrow \text{Cl}_{i=1}^\infty X_i$. On the other hand the desired isomorphism (3) can also be regarded as being induced by these inclusions (rather than by the projections $\text{Cl}_{i=1}^\infty X_i \rightarrow X_i$). This completes the proof of the assertion.

EXAMPLES. (1) Every suspension spectrum $\underline{E} = \{\Sigma^n E \mid E \in \underline{P}_1\}$ has property (**S**).

(2) The same kind of argument which leads to Proposition 4.4 can of course be applied to $\underline{\text{BO}}_*$, the homology theory of real K -theory.

(3) Let $\underline{E} = \{E_n\}$ be a spectrum having the following property:

(a) $\sigma: \Sigma E_n \rightarrow E_{n+1}$ is cellular embedding

(b) $E_{n+1} = S^{n+1} \vee \Sigma E_n$.

It is a rather trivial task, to construct a spectrum of this kind. Put $X_i = S^0 = \{x_i, *\}$, then we have

$$E_k \overline{\wedge} X_i = E_k \wedge X_i = E_k.$$

Let $\xi_i \in \underline{E}_0(X_i)$ be defined by the inclusion f_i of $S^{0+i} = S^i$ into the copy of $S^i \subset E_i \setminus \Sigma E_{i-1}$. There is certainly no $\zeta \in \overline{E}_0(X)$ available such that $\varphi\zeta = \{\xi_i\}$: Assume to the contrary the existence of such a ζ , then there must necessarily exist a universal l such that f_i is stably equivalent to $f'_i: S^l \rightarrow E_i$.

However this is just not true by construction.

This assures us that the clusteraxiom does not hold for every spectrum $\underline{E} \in \underline{\text{Spec}}$.

More generally we have proved:

4.5. LEMMA. *Let \underline{E} be a CW-spectrum such that the clusteraxiom holds for \underline{E}_* . Let $X = \text{Cl}_{i=1}^\infty X_i$ and a family*

$$\tilde{f}_i: S^{n+l(i)} \rightarrow E_{l(i)} \overline{\wedge} X_i$$

be given. Then there exists a universal l , independent of i such that \tilde{f}_i is stably homotopic to a

$$\tilde{f}'_i: S^{n+l} \rightarrow E_l \overline{\wedge} X_i.$$

Proof. Use the same argument that led to the last conclusion in the preceding third example.

REMARK. Proposition 4.4 allows an extension to arbitrary Ω -spectra $\underline{E} \in \underline{\text{Spec}}$ (cf. Theorem 3.1 in [3]). The proof of this assertion follows by Proposition 4.1 and a considerable refinement of the arguments that led to Proposition 4.2. In [3] this is needed as the main tool for comparing \underline{E}_* and sE_* (cf. [8]) on the category $\underline{\text{Com}}^f$. Proposition 4.3 however is, as far as Eilenberg-Mac Lane spectra are concerned, better than Theorem 3.1 in [3] because it works for all (rather, than for finite dimensional) compacta.

5. The Milnor axioms. As we already mentioned J. Milnor characterized an ordinary homology theory on the category \mathfrak{A}_{CM} of compact pairs (this is J. Milnor's original terminology) by means of the following axioms:

(A1) *The clusteraxiom* for the subcategory $\underline{\text{Com}} \subset \mathfrak{A}_{\text{CM}}$ (the category of based compact metric spaces).

The preceding section was entirely devoted to this axiom.

(A2) *The strong excision axiom* which requires that for a homology theory $\{H_n(_, \partial)\}$ the projection $p: (X, A) \rightarrow (X/A, *)$ induces an isomorphism

$$(1) \quad H_*(X, A) \rightarrow H_*(X/A, *).$$

(A3) *The exactness axiom.*

(A4) *The homotopy axiom.*

Contrary to J. Milnor's treatment, we do not have a dimension axiom.

5.1. DEFINITION. A functor $H_*: \mathfrak{A}_{\text{CM}} \rightarrow \text{Ab}^{\mathbb{Z}}$ together with the corresponding boundary operators is called a *homology theory* on \mathfrak{A}_{CM} $H_* = \{H_n, \partial\}$ whenever (A1)–(A4) are satisfied.

It is well-known how to transform such a homology theory into a reduced homology theory on Com and vice-versa. We will freely make use of this correspondence and call a functor $H_*: \text{Com} \rightarrow \text{Ab}^{\mathbb{Z}}$ (together with natural transformations $\sigma: H_n \rightarrow H_{n+1}\Sigma$) a *homology theory on Com*, whenever the related functor on \mathfrak{A}_{CM} fulfills 5.1.

To this corresponds the concept of a natural transformation $\varphi: H_* \rightarrow H'_*$ between homology theories (cf. Definition 1.3).

This terminology should not be mixed up with that of 1.3 where we dealt with singular homology theories on a shape category.

Let $\underline{E} \in \underline{\text{Spec}}$ be any spectrum, then the functor $h: \text{Com} \rightarrow \overline{\text{Com}}$ gives rise to a homology theory $\underline{E}_*h: \text{Com} \rightarrow \text{Ab}^{\mathbb{Z}}$. According to our custom not to write down the functor h explicitly, we will henceforth write \underline{E}_* instead of \underline{E}_*h .

In [4] §3 we have proved for \underline{E}_* , $\underline{E} \in \underline{\text{Spec}}$, all the axioms (A2)–(A4). Concerning strong excision, recall that $\underline{E}_n(X, A)$ is defined as $\underline{E}_n(X \cup CA, *)$, where $*$ denotes the vertex of the cone CA . However since every inclusion in Com is a cofibration ([4] Proposition (A9) in $\overline{\text{Com}}$, we have a homotopy equivalence

$$X \cup CA \simeq X/A \quad \text{in } \overline{\text{Com}}.$$

This proves (1) for \underline{E}_* .

The preceding section was devoted to a treatment of the cluster axiom, so that we are able to summarize:

5.2. THEOREM. *On the category Com there exists to each CW spectrum \underline{E} a functor $\underline{E}_*: \text{Com} \rightarrow \text{Ab}^{\mathbb{Z}}$ (together with a natural transformation $\sigma: \underline{E}_n(X) \rightarrow \underline{E}_{n+1}(\Sigma X)$) such that the axioms (A2)–(A4) of a homology theory on $\overline{\text{Com}}$ are satisfied.*

The main objective of this section is to establish the following

5.3. THEOREM. *Let $H_*, H'_*: \text{Com} \rightarrow \text{Ab}^{\mathbb{Z}}$ be two homology theories on Com and $\alpha: H_* \approx H'_*$ be an isomorphism on the subcategory $\underline{P}_0 \subset \overline{\text{Com}}$ of compact (based) CW-spaces. Then there exists a unique extension of α to an isomorphism $\tilde{\alpha}: H_* \approx H'_*$ over $\overline{\text{Com}}$.*

The proof is preceded by some remarks and two lemmas:

(1) We write H_* also for the corresponding homology theory on \mathfrak{A}_{CM} . The isomorphism α can be extended to an isomorphism (also denoted by the same letter) between H_* and H'_* defined on the category of polyhedral pairs.

(2) Instead of working with actual CW-spaces (resp. CW-pairs or polyhedral pairs), we will use ANR subspaces of a Hilbert cube Q in which any space $X \in \underline{\text{Com}}$ is supposed to be embedded. This does clearly not cause any difficulties.

(3) Let $\bar{\zeta} \in H_n(X)$ be any element. A simple exactness argument guarantees that $H_{n+1}(Q, X) \overset{\partial}{\approx} H_n(X, x_0)$, hence there exists a uniquely determined $\zeta \in H_{n+1}(Q, X)$ such that $\partial\zeta = \bar{\zeta}$. It is actually this ζ with which we are going to operate.

(4) Let $(X, A), (X_1, A_1), (X_2, A_2) \in \mathfrak{A}_{\text{CM}}$ be three pairs such that $X = X_1 \cup X_2, X_1 \cap X_2 = A_1 \cap A_2$.

Denote by

$$i: (X, A) \subset (X, A_1 \cup A_2 \cup A), \quad i_j: (X_j, A_j) \subset (X, A_1 \cup A_2 \cup A)$$

the inclusions. Let $\zeta_i \in H_m(X_i, A_i)$ be two fixed elements, then we denote by $\zeta = \zeta_1 \oplus \zeta_2 \in H_m(X, A)$ any element, having the property

$$i_*\zeta = i_{1*}\zeta_1 + i_{2*}\zeta_2.$$

This ζ may not exist nor is it necessarily uniquely determined. We can of course extend this terminology to finitely many summands.

(5) The cluster axiom enables us to extend this construction to countably many summands:

Let to this end Q be the Hilbert cube, $X \subset Q$ a given compact space and let $\bar{\zeta} \in H_n(X), \zeta \in H_{n+1}(Q, X)$ with $\partial\zeta = \bar{\zeta}$ be fixed elements. We take a decreasing sequence $P_1 = Q, P_2 \supset \dots P_k \supset \dots$ of compact ANRs with $\bigcap P_i = X$. Moreover we establish an increasing sequence of compact ANRs $\emptyset = U_1 \subset U_2 \subset \dots$ in Q having the following properties:

(1) $U_i \cap P_{i+k} = \emptyset$ for $k \geq 1$, (2) $\overline{P_j \setminus P_i} \subset \text{Int } U_i, i > j$. All this can be obviously achieved.

We set:

$$R_i = P_i \cap U_i, \quad X_i = P_i \cap U_{i+1}, \quad A_i = R_i \cup R_{i+1} \quad (\text{Fig. 1}).$$

$$\bar{R}_i = R_i \cup X, \quad \bar{X}_i = X_i \cup X, \quad \bar{A}_i = A_i \cup X.$$




FIGURE 1

Let $\zeta_i \in H_m(X_i, A_i)$ for all i be given elements and $\bar{\zeta}_i$ the images of ζ_i under the inclusion $(X_i, A_i) \subset (\bar{X}_i, \bar{A}_i)$. We assume:

(*) The existence of elements $\eta_r \in H_m(X_1 \cup \dots \cup X_r, R_{r+1})$ which are inductively determined

$$\zeta_1 = \eta_1, \eta_2 = \eta_1 \oplus \zeta_2, \dots, \eta_r = \eta_{r-1} \oplus \zeta_r,$$

and denote by $\bar{\eta}_r \in H_m(\bar{X}_1 \cup \dots \cup \bar{X}_r, \bar{R}_{r+1})$ the image of η_r under the corresponding inclusion.

Since we have

$$R_i \cap R_{i+1} = \emptyset,$$

the definition of (X_i, A_i) implies that there exists $\rho_i^{(1)}, \rho_i^{(2)} \in H_{m-1}(R_i)$ such that

$$\rho_i^{(1)} - \rho_{i+1}^{(2)} = \partial \zeta_i,$$

where we omitted the inclusions $R_i \subset A_i$ from our notation. Because $R_1 = Q \cap U_1 = \emptyset$, we have $\rho_1^{(1)} = \rho_1^{(2)} = 0$.

The existence of η_r implies that we have

$$(2) \quad \rho_i^{(1)} = \rho_i^{(2)}.$$

The following lemma asserts that we are able to find a $\zeta \in H_m(Q, X) = H_m(\bigcup_{i=1}^{\infty} \bar{X}_i, X)$ which can be considered as some kind of η_{∞} :

5.4. LEMMA. *Assume that (*) is satisfied, then (1) there exists a $\zeta \in H_m(Q, X)$ such that*

$$(3) \quad k_r * \zeta = l_r * \bar{\eta}_r$$

with inclusions

$$k_r: (Q, X) \subset \left(Q, \bigcup_{i=r+1}^{\infty} \bar{X}_i \right) \\ l_r: \left(\bigcup_{i=1}^r \bar{X}_i, \bar{R}_{r+1} \right) \subset \left(Q, \bigcup_{i=r+1}^{\infty} \bar{X}_i \right);$$

(2) *any other elements ζ' satisfying (2) (with possibly different family $\{\bar{\zeta}_r\}$ and same $\{\zeta_i\}$) is equal to ζ .*

Proof. Let $Z_i = \bar{X}_i / \bar{A}_i$; form the cluster of different copies of the spaces \bar{X}_i and define

$$Z = \text{Cl}(\bar{X}_i / X).$$

Observe that $\bar{X}_i / X = X_i \cup X / X = X_i^+$ and $\bar{A}_i / X = A_i \cup X / X = A_i^+$.

Due to (A2) we have an isomorphism

$$(4) \quad H_m(\text{Cl } Z_i, *) \approx H_m(Z, \text{Cl } \bar{A}_i / X),$$

while (A1) provides us with a $\eta' \in H_m(\text{Cl } Z_i, *)$, satisfying $\eta' | \bar{X}_i / \bar{A}_i = \zeta'_i \in H_m(\bar{X}_i / \bar{A}_i) \approx H_m(\bar{X}_i, \bar{A}_i)$ where ζ'_i corresponds to ζ_i under this isomorphism.

Here we denote by $\eta' | \bar{X}_i / \bar{A}_i$ the i th factor in the element corresponding to η' by the isomorphism (A1):

$$H_m\left(\bigcup_1^{\infty} Z_i, *\right) \approx \prod_{i=1}^{\infty} H_m(Z_i, *).$$

To η' there exists under the isomorphism (4) a $\eta \in H_m(Z, \text{Cl}_1^{\infty} A_i^+)$. We have an infinite folding map:

$$f: \left(Z, \bigcup_1^{\infty} A_i^+ \right) \rightarrow (Q/X, \cup A_i^+)$$

(identifying back the different copies of the X_i as subsets of Q resp. for the quotient spaces Q/X) and obtain $\eta'' = f_* \eta \in H_m(Q/X, \bigcup_{i=1}^{\infty} A_i^+)$.

(**) There exists an element $\zeta \in H_m(Q, X) \approx H_m(Q/X, *)$ such that $j_* \zeta = \eta''$, $j: (Q/X, *) \xrightarrow{\subset} (Q/X, \cup A_i^+)$.

Proof. Because of the exact (reduced) homology sequence of the pair $(Q/X, \bigcup_{i=1}^{\infty} A_i^+)$ it suffices to prove that $\partial\eta'' = 0$. To this end we have the commutative diagram

$$(5) \quad \begin{array}{ccc} \prod_{i=1}^{\infty} H_m(\bar{X}_i/\bar{A}_i) & \xrightarrow{\Delta} & \prod_{i=1}^{\infty} H_{m-1}(A_i^+) \\ \Downarrow \varphi & & \Downarrow \varphi \\ H_m\left(\bigcup_{i=1}^{\infty} X_i^+, \bigcup_{i=1}^{\infty} A_i^+\right) & \xrightarrow{\partial} & H_{m-1}\left(\bigcup_{i=1}^{\infty} A_i^+\right) \\ \downarrow f_* & & \downarrow f_* \\ H_m\left(Q/X, \bigcup_{i=1}^{\infty} A_i^+\right) & \xrightarrow{\partial} & H_{m-1}\left(\bigcup_{i=1}^{\infty} A_i^+\right) \end{array}$$

where the vertical isomorphisms stem from (A1) (or from (A1) in combination with (A2)) and the vertical arrows f_* are induced by the folding map f .

According to (5) we have proved $\partial\eta'' = 0$ whenever we are able to confirm that $f_*\varphi\{\Delta\xi'_i\} = 0 \in H_{m-1}(\bigcup_{i=1}^{\infty} A_i^+)$, where $\Delta: H_m(\bar{X}_i/\bar{A}_i) \rightarrow H_{m-1}(A_i^+)$ is the composite of $\partial: H_m(X_i^+, A_i^+) \rightarrow H_{m-1}(A_i^+)$ with the natural isomorphism

$$H_m(\bar{X}_i/\bar{A}_i) \approx H_m(X_i^+, A_i^+).$$

On the other hand we have $A_i = R_i \cup R_{i+1}$. Hence the space R_i appears in the cluster $\bigcup_{i=1}^{\infty} A_i^+$ twice, namely as a subset $R_i^{(1)} \subset A_i$ and secondly as a subset $R_i^{(2)}$ of A_{i-1} . The folding map f identifies both copies. We have already detected the elements $\rho_i^{(k)} \in H_{m-1}(R_i^{(k)})$, $k = 1, 2$ such that in $Q(2)$ (now for A_i^+ instead of A_i) is satisfied.

We can rearrange the cluster

$$\bigcup_{i=1}^{\infty} A_i^+ = \bigcup_{i=1}^{\infty} (R_i^{(1)} \cup R_{i+1}^{(2)})^+$$

getting a new cluster $C = \bigcup_{i=1}^{\infty} (R_i^{(1)} \cup R_i^{(2)})^+$ with folding map $g: C \rightarrow \bigcup_{i=1}^{\infty} A_i^+$.

Moreover there exist isomorphisms in a commutative diagram

$$\begin{array}{ccc} \prod_{i=1}^{\infty} H_{m-1}(R_i^{(1)} \cup R_i^{(2)}) & \approx & \prod_{i=1}^{\infty} H_{m-1}(A_i^+) \\ \Downarrow \varphi & & \Downarrow \varphi \\ H_{m-1}(C) & \approx & H_{m-1}\left(\bigcup_{i=1}^{\infty} A_i^+\right) \\ g_* \searrow & & \swarrow f_* \\ & H_{m-1}\left(\bigcup_{i=1}^{\infty} A_i^+\right) & \end{array}$$

As a result we have

$$0 = g_*\psi \{ \rho_i^{(1)} - \rho_i^{(2)} \} = f_*\varphi \{ \rho_i^{(1)} - \rho_{i+1}^{(2)} \} = \partial\eta''.$$

This completes the construction of $\zeta \in H_m(Q, X)$.

Condition (3) asserts that $\bar{\eta}_r = l_r^{-1} * k_{r*} \zeta$ (since l_r is easily recognized to be an excision) and therefore simply states that by replacing infinite clusters by finite (r -fold) wedges we get back η_r . The verification of this fact is straightforward.

This completes the proof of the assertion (1) in 5.4.

We accomplish a proof of 5.4 (2) by collecting the following remarks:

(1) the inclusion

$$j: \left(\bigcup_{i=1}^{\infty} \bar{A}_i, x_0 \right) \subset \left(\bigcup_{i=1}^{\infty} \bar{A}_i, X \right) \quad \left(\text{resp. } j_k: \left(\bigcup_1^k \bar{A}_i, x_0 \right) \subset \left(\bigcup_1^k \bar{A}_i, X \right) \right)$$

induce an epimorphism in homology.

Proof. We have $X \cap A_i = \emptyset$ and therefore a commutative diagram

$$\begin{array}{ccc} H_m \left(\bigcup_1^{\infty} \bar{A}_i, x_0 \right) & \xrightarrow{j_*} & H_m \left(\bigcup_1^{\infty} \bar{A}_i, X \right) \\ & \rho_* \nwarrow & \parallel \\ & & H_m \left(\bigcup_1^{\infty} A_i^+, * \right) \end{array}$$

with inclusion ρ where the vertical arrow is an excision.

(2) The sequence $\{\zeta_i\}$ determines η'' uniquely. Let $\zeta, \zeta' \in H_m(Q/X, *) = H_m(Q, \bar{X})$ be two elements which map into η'' in

$$H_m(Q/X, *) \rightarrow H_m \left(Q/X, \bigcup_1^{\infty} A_i^+ \right)$$

then by exactness, $\zeta - \zeta' = q_*\alpha$, with $\alpha \in H_m(\bigcup_1^{\infty} A_i^+)$, $q: (\bigcup_1^{\infty} A_i^+, x_0) \subset (Q/X, *)$.

Consider the diagram

$$\begin{array}{ccccc} & & H_m \left(\bigcup_1^{\infty} A_i^+, * \right) & & \\ & & \parallel & & \\ H_m(\bigcup \bar{A}_i, x_0) & \xleftarrow{j_*} & H_m \left(\bigcup_1^{\infty} \bar{A}_i, X \right) & \rightarrow & H_m(Q, X) \\ \partial'' \downarrow & & \partial' \downarrow & & \partial \downarrow \\ H_{m-1}(x_0) & \leftarrow & H_{m-1}(X, x_0) & \approx & H_{m-1}(X, x_0) \end{array}$$

whose commutativity ensures that $\partial\alpha = \partial\zeta - \partial\zeta' = 0$. This proves that two ζ, ζ' which are achieved by our construction satisfy $\partial\zeta = \partial\zeta'$.

(3) Let $\zeta \in H_m(Q, X)$ be given such that for fixed $\{\zeta_i\}$ and some selection of $\{\bar{\eta}_r\}$ (3) holds. Observe that l_r is an excision, hence l_{r*} an isomorphism. We have a commutative diagram for a $\zeta \in H_m(X, Q)$ such that (3) (for all r) holds:

$$\begin{array}{ccc}
 \zeta \in H_m(Q, X) & & \\
 \downarrow \approx & & \\
 \bar{\zeta} \in H_m(Q/X) & \rightarrow & H_m\left(Q/X, \bigcup_{i=1}^{\infty} A_i^+\right) \ni \eta'' \\
 & \uparrow & \uparrow \\
 & \prod_{i=1}^{\infty} H_m(\bar{X}_i, \bar{A}_i) \ni \{\bar{\zeta}_i\} &
 \end{array}$$

which guarantees that ζ is in fact obtained by the construction process which we established during the proof of (1). These remarks ensure that we have $\partial\zeta = \partial\zeta'$ for any other ζ' which satisfies (3) (using an eventually different $\{\bar{\eta}_r\}$ related to the same family $\{\zeta_i\}$). Now $\zeta = \zeta'$ follows by the fact that $\partial: H_{n+1}(Q, X) \approx H_n(X, x_0)$ is an isomorphism (cf. Remark (3) at the beginning of this proof).

We can of course get back the η_r and even the ζ_i from ζ (which in analogy to the finite case will be denoted by $\zeta = \bigoplus_{i=1}^{\infty} \zeta_i$).

5.5. LEMMA. *Let $(Q, X), (X_i, A_i)$ be as before and fix a $\zeta \in H_m(Q, X)$. Then there exists a $\zeta_i \in H_m(X_i, A_i)$ for all i , such that*

$$\zeta = \bigoplus_{i=1}^{\infty} \zeta_i.$$

Proof. We have $\zeta_j \in H_m(Q, \bigcup_{i \neq j} X_i)$ for any j and obtain a $\zeta_j \in H_m(X_j, A_j)$ by excision. Also the elements η_r are obtained by excision from $\eta'_r \in H_m(Q, \bigcup_{i=r}^{\infty} X_i \cup X)$. The verification of (2) is now immediate.

Proof of Theorem 5.3. The basic idea of the proof of Theorem 5.3 is this: Starting with a $\bar{\zeta} \in H_{m-1}(X, x_0)$ we find a $\zeta \in H_m(Q, X)$ which can be cut into pieces $\{\zeta_i\}$ by means of Lemma 5.5. On the other hand Lemma 5.4 allows us to paste together these pieces $\{\zeta_i\}$ (which are required to satisfy certain compatibility conditions) to the effect that we get a $\zeta \in H_m(Q, x)$ and finally a $\partial\zeta = \bar{\zeta} \in H_{m-1}(X, x_0)$. While $\bar{\zeta} \in H_{m-1}(X, x_0)$ is eventually defined outside \underline{P}_0 , the pieces ζ_i are all defined for pairs in \underline{P}_0^2 and therefore accessible to the transformation α .

Let $\bar{\zeta} \in H_n(X)$, $\zeta \in H_{n+1}(Q, X)$ be given, such that $\partial\zeta = \bar{\zeta}$. We write for simplicity $H_*(X)$ ($H'_*(X)$) instead of $H_*(X, x_0)$ ($H'_*(X, x_0)$); all homology groups in this proof are understood to be *reduced*.

Lemma 5.5 yields a sequence $\{\zeta_i \in H_{n+1}(X_i, A_i)\}$ such that $\zeta = \bigoplus_{i=1}^{\infty} \zeta_i$. This enables us to extend $\alpha: H_* \approx H'_*$ in the following way: Let $\bar{\zeta} \in H_n(X)$ be given, then we go over to $H_{n+1}(Q, X)$ and the family $\{\zeta_i\}$. Because all ζ_i are defined for spaces in a subcategory where α has been already established, we have the family $\{\zeta_i\}$ (see Remarks (1), (2) at the beginning of this proof) which in turn establishes (due to Lemma 5.4) again an element $\tilde{\alpha}\bar{\zeta} \in H'_n(X)$. By Remark (3) and the last conclusion of Lemma 5.4, this element $\tilde{\alpha}\bar{\zeta}$ is uniquely determined by $\bar{\zeta}$ (because every intermediate step has this property, with fixed P_i, U_i).

Observe that we can instead of a family $\{\zeta_i\}$ equally well deal with a related family $\{\tilde{\eta}_r \in H_{n+1}(Q, P_r)\}$ which determines (1) the family $\{\eta_r \in H_{n+1}(\bigcup_{i=1}^r \bar{X}_i, \bar{R}_{r+1})\}$ by excision and finally (2) a family $\{\zeta_i \in H_{n+1}(\bar{X}_i, \bar{A}_i)\}$. Since $\{\zeta_i\}$ determines $\bar{\zeta} = \partial\zeta$ uniquely, we are naturally led to the question under what conditions two different families $\{\tilde{\eta}_r \in H_{n+1}(Q, P_r)\}$, $\{\tilde{\eta}'_r \in H_{n+1}(Q, P'_r)\}$ (with a different approximating $\{P'_i, U'_i\}$) determine the same ζ : This problem is settled by means of the following observation:

(F1) Let $\{l(r)\}$ be an infinite subsequence of the integers, then $\{\tilde{\eta}_r\}$ and $\{\tilde{\eta}_{l(r)}\}$ determine obviously the same ζ .

(F2) Assume that $\{\tilde{\eta}_r\}$, $\{\tilde{\eta}'_r\}$ have the property that to each r there exists an index $l(r)$ such that $P'_{l(r)} \subset P_r$ and $j_*\tilde{\eta}'_{l(r)} = \tilde{\eta}_r$ ($j: (Q, P'_{l(r)}) \subset (Q, P_r)$), then $\{\tilde{\eta}_r\}$ and $\{\tilde{\eta}'_r\}$ determine the same ζ .

(F3) Let on the other hand ζ be determined by $\{\tilde{\eta}_r\}$, (ζ' by $\{\tilde{\eta}'_r\}$) and $\zeta = \zeta'$, then we can find to each P_r an index $l(r)$ as in (F2) such that $k_*\zeta = \tilde{\eta}_r = j_*k'_*\zeta' = j_*\tilde{\eta}'_{l(r)}$, with inclusions $k: (Q, X) \subset (Q, P_r)$, $k': (Q, X) \subset (Q, P'_{l(r)})$.

As a result we have that $\zeta = \zeta'$ if and only if (F2) holds.

The assignment $\tilde{\alpha}$ can be defined by using the families $\{\tilde{\eta}_r\}$ instead of $\{\zeta_i\}$: We have the family $\{\alpha\tilde{\eta}_r\}$ which in turn determines a family $\{\zeta'_i \in H'_{n+1}(\bar{X}_i, \bar{A}_i)\}$ and finally the element $\tilde{\alpha}\bar{\zeta}$.

These remarks enable us to prove

(a) $\tilde{\alpha}$ is natural and compatible with suspensions.

Proof. Let $f: (X, x_0) \rightarrow (Y, y_0)$ be continuous. We can extend f to a $F: (Q, X) \rightarrow (Q, Y)$. Denoting the corresponding sequence of approximating ANRs by $P'_1 = Q \supset P_2 \supset \dots, \bigcap_{i=1}^{\infty} P'_i = Y$, there corresponds to

each index i a minimal $l(i)$ such that $F(P_i) \subset P'_{l(i)}$. We have $F': (Q, P_i) \rightarrow (Q, P'_{l(i)})$. So we can assign to each $\tilde{\eta}_r$ a $\tilde{\eta}'_{l(r)} = F'_* \tilde{\eta}_r \in H_{n+1}(Q, P'_{l(i)})$, in such a way that $\{\tilde{\eta}'_{l(r)}\}$ is a family which determines $f_* \zeta$. The naturality of $\tilde{\alpha}$ requires the commutativity of the diagram

$$\begin{array}{ccc} H_n(X) & \xrightarrow{\tilde{\alpha}} & H'_n(X) \\ f_* \downarrow & & \downarrow f_* \\ H_n(Y) & \xrightarrow{\tilde{\alpha}} & H'_n(Y). \end{array}$$

Since α itself is supposed to be natural, this can be deduced by looking upon the two chains of correspondences for a fixed $\zeta \in H_n(X)$:

$$\begin{array}{ccccccc} & & \{\tilde{\eta}_r\} & \mapsto & \{\tilde{\eta}'_{l(r)}\} & \mapsto & \{\alpha \eta'_{l(r)}\} & \mapsto & \tilde{\alpha} f_* \zeta \\ & \nearrow & & & & & & & \parallel \\ \zeta & & & & & & & & \\ & \searrow & & & & & & & \\ & & \{\tilde{\eta}_r\} & \mapsto & \{\alpha \tilde{\eta}_r\} & \mapsto & \{F'_* \alpha \tilde{\eta}_r\} & \mapsto & f_* \tilde{\alpha} \zeta \\ & & & & \downarrow & & & & \\ & & & & \tilde{\alpha} \zeta & & & & \end{array}$$

where the vertical “ $=$ ” indicates a relation of type (F1).

The second assertion in (a) follows immediately by construction.

(b) $\tilde{\alpha}$ is a homomorphism.

Proof. This follows because the whole construction process of $\tilde{\alpha}$ is compatible with the group structure:

Let $\bar{\zeta}_1, \bar{\zeta}_2 \in H_n(X)$ be given, then we have $\partial(\bar{\zeta}_1 + \bar{\zeta}_2) = \bar{\zeta}_1 + \bar{\zeta}_2$, moreover $\{\eta_r^{(1)} + \eta_r^{(2)}\}$ is a family for $\bar{\zeta}_1 + \bar{\zeta}_2$ (in the sense of the preceding construction) etc.

(c) $\tilde{\alpha}$ is by α uniquely determined.

Proof. Let $\beta: H_n(X) \rightarrow H'_n(X)$ be another extension of α , then for given $\bar{\zeta} \in H_n(X)$, the family $\{\beta \tilde{\eta}_r\}$ is easily seen to determine $\beta \bar{\zeta} \in H'_n(X)$. Since $\alpha = \beta$ for all $\tilde{\eta}_r$, we have $\{\alpha \tilde{\eta}_r\} = \{\beta \tilde{\eta}_r\}$, where, by definition, the first family determines $\tilde{\alpha} \bar{\zeta}$ and the second one $\beta \bar{\zeta}$. Hence (c) follows. \square

(d) $\tilde{\alpha}$ is an isomorphism.

Proof. Follows directly from (c) or can be seen by putting α^{-1} instead of α into the construction process of $\tilde{\alpha}$. \square

This completes the proof of 5.3. \square

We could of course easily adjust the preceding proof to establish the following slightly more general result:

5.6. COROLLARY. *Let H_* , H'_* be homology theories on $\underline{\text{Com}}$ (in the sense of Definition 5.1) $\alpha: H_* \rightarrow H'_*$ a natural transformation on \underline{P}_0 , then there exists a unique extension $\tilde{\alpha}: H_* \rightarrow H'_0$ of α over $\underline{\text{Com}}$.*

Concerning the relations to $\overline{\underline{\text{Com}}}$ we have:

5.7. COROLLARY. *Let (1) $H_*: \underline{\text{Com}} \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ be any homology theory on $\underline{\text{Com}}$ (2) \underline{E} a CW spectrum such that \underline{E}_* is a homology theory on $\underline{\text{Com}}$ (in the sense of Definition 5.1, which implies in particular that the cluster axiom holds) (3) $\alpha: H_* \approx \underline{E}_*$ a natural isomorphism on the category \underline{P}_0 . Then there exists an extension $\overline{H}_*: \overline{\underline{\text{Com}}} \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ of H_* over the shape category $\overline{\underline{\text{Com}}}$ and an isomorphism $\tilde{\alpha}: H_* \approx \underline{E}_*$.*

This is an immediate consequence of 5.2 and 5.3.

In the next section we need the category $\underline{\text{Com}}^f$ of finite dimensional compact metric spaces. The proof of 5.3 immediately carries over to this case, so that we can restate 5.3, 5.6, 5.7 for $\underline{\text{Com}}^f$.

5.8. THEOREM. *The conclusions of 5.3, 5.6, 5.7 are still valid after replacing $\underline{\text{Com}}$ by $\underline{\text{Com}}^f$.*

REMARK. As pointed out in [3] p. 209 (proof of Theorem 3.2), the arguments which lead to the construction of $\tilde{\alpha}$ (in the course of the proof of Theorem 5.3) are still valid whenever only the homology theory H'_* (but not necessarily H_*) satisfies a clusteraxiom: We need the clusteraxiom solely to paste together a sequence $\{\alpha\zeta_i\}$ (or alternatively $\{\alpha\tilde{\eta}_r\}$) in order to obtain a $\tilde{\alpha}\tilde{\zeta}$. The clusteraxiom is not needed to break a given element $\tilde{\zeta}$ into appropriate pieces $\{\zeta_i\}$.

6. Applications. We collect together some consequences of the preceding existence and uniqueness theorems. Some of these results are already known, but with different proofs.

We summarize some of the results of §§1–3 in:

6.1. THEOREM. *Let \underline{E} be an CW spectrum, then the homology theory $\underline{E}_*: \underline{P}_{1h} \rightarrow \mathbf{Ab}^{\mathbf{Z}}$ allows, up to an isomorphism, a unique extension $\tilde{\underline{E}}_*: \underline{K}_h \rightarrow \mathbf{Ab}^{\mathbf{Z}}$, $\tilde{\underline{E}}_*(\) = \underline{E}_*(|\bar{S}(\)|)$, such that the Whitehead axiom **W** (Definition 1.2) holds. Moreover $\tilde{\underline{E}}_*$ is of compact support (Definition 1.7). For $\underline{E} \in \mathbf{CSpec}$ and X being an s -continuum, one has a natural isomorphism $\bar{\underline{E}}_*(X) \approx \tilde{\underline{E}}_*(X)$.*

Proof. This follows immediately from 1.8, 2.2, 3.3.

6.2. THEOREM. *For any abelian group G , Steenrod-Sitnikov homology theory $H_*^S(\ ; G)$ turns out to be naturally isomorphic to $\overline{K(G)}_*(\)$ on the category $\underline{\mathbf{Com}}$. Hence $H_*^S(\ ; G)$ can be uniquely extended over $\overline{\mathbf{Com}}$.*

Proof. As well $H_*^S(\ ; G)$ as $\overline{K(G)}_*(\)$ are fulfilling the Milnor axioms §5 (A1)–(A4): As far as $H_*^S(\ ; G)$ is concerned this can be found in [10], while 4.3 and 5.2 together confirm that for $\overline{K(G)}_*(\)$. Since $H_*^S(\ ; G) = H_*(\ ; G)$ and $\overline{K(G)}_*(\)$ are isomorphic on \underline{P}_0 , Theorem 6.2 can be immediately deduced from 5.3 and 5.7. \square

6.3. COROLLARY. *Let G be a finitely generated abelian group, X an s -continuum, then there exists a natural isomorphism between $H_*(|\bar{S}(X)|; G)$ and $H_*^S(X; G)$.*

Proof. This is an immediate consequence of 6.1 and 6.2.

REMARK. The last corollary is a generalization of Theorem 7.7 in [2] (where this was proved by different methods for $G = \mathbf{Z}$). Observe that the s -connectedness of X is not necessary for 6.2 but enters into our considerations through 6.1.

6.4. THEOREM. *Let \hat{H}_* be an ordinary homology theory on $\underline{\mathbf{Com}}$ (i.e. one for which §5 (A1)–(A4) and in addition a dimension axiom is valid) with coefficient group $\hat{H}_0(\text{point}) \approx G$. Then $\hat{H}_*(\)$ is naturally isomorphic to Steenrod-Sitnikov homology $H_*^S(\ ; G)$.*

Proof. By the Eilenberg-Steenrod uniqueness theorem, we have $\hat{H}_*(\) \approx H_*(\ ; G)$ on \underline{P}_0 (resp. on polyhedral pairs). Now the assertion follows by application of 6.2.

REMARK. This is nothing else than J. Milnor's uniqueness theorem in [12].

For the sake of completeness we restate Corollary 4.2 in [4] which follows now from 6.1, the fact that $\underline{K}(G) \in \underline{\text{CSpec}}$ for finitely generated G and homological algebra:

6.5. COROLLARY. *Let G be finitely generated, abelian; $X \in \underline{\text{Com}}$ s -connected (i.e. a s -continuum), then we have a natural universal coefficient sequence:*

$$0 \rightarrow \underline{K}(\underline{Z})_n(X) \otimes G \rightarrow \underline{K}(G)_n(X) \rightarrow \text{Tor}(\underline{K}(\underline{Z})_{n-1}(X), G) \rightarrow 0.$$

REMARK. This result cannot be extended to an arbitrary coefficient groups G (which is not finitely generated) because of a result in [10] asserting that a non-trivial homology theory satisfying a clusteraxiom does not admit a universal coefficient sequence for *all* abelian G . As a consequence we are able to provide for any non-finitely generated G an s -continuum X such that

$$\underline{K}(G)_*(X) \neq \overline{K(G)}_*(X).$$

As can be easily realized, the s -continuum $X = \text{Cl}_{i=1}^\infty S_i^2$ ($S_i^2 = S^2$ for any $i = 1, 2, \dots$) has the desired property.

In [6], [8], [9] the authors deal with a functor

$$\text{Ext}: \underline{\text{Com}}_n \rightarrow \underline{\text{Ab}}$$

which is defined by functional analytic methods. This functor gives rise to a homology theory

$$\epsilon_n(X) = \begin{cases} \text{Ex}(X) & \text{if } n \equiv 1 \pmod{2} \\ \text{Ext}(\Sigma X) & \text{if } n \equiv 0 \pmod{2} \end{cases}$$

on the category $\underline{\text{Com}}$ which has very far reaching applications in functional analysis.

Concerning the definition of Ext and the verification of its basic properties the reader is referred to [6]. In [5] as well as in [7] the problem of the uniqueness of Steenrod extensions has been raised for the first time.

In [8] Remark 7.7 the authors deduce a non-canonical version of Theorem 5.7 for homology theories with coefficient groups of finite type.

Furthermore it is proved in [6] (Theorem 7.3) that for Ext and consequently for ϵ_* the clusteraxiom holds.

This makes sense either for the based version of ϵ_* or by talking about "spaces X_i , $i = 1, 2, \dots$, any two of them having only a fixed point b in common."

Details of all this can be found in [6], [8].

We have the following isomorphism theorem:

6.6. THEOREM. *On the category \underline{P}_0 there exists an isomorphism $\alpha: \varepsilon_* \approx \underline{\text{BU}}_*$ which allows a unique extension to an isomorphism*

$$\bar{\alpha}: \varepsilon_* \approx \overline{\text{BU}}_* \text{ on } \underline{\text{Com}}^f.$$

Proof. The existence of an isomorphism α on \underline{P}_0 is the content of Theorem 7.7 in [6]. Both functors ε_* and $\overline{\text{BU}}_*$ are homology theories on $\underline{\text{Com}}^f$ in the sense of Definition 5.1: In the case of ε_* this has been readily proved in [6]. As far as $\overline{\text{BU}}_*$ is concerned, this follows from 4.4 and 5.2. Hence we can apply Theorem 5.7, establishing the desired conclusion.

As a corollary we obtain as a consequence of 5.8:

6.7. COROLLARY. *The homology theory $\varepsilon_*: \underline{\text{Com}}_h^f \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$ allows an extension over $\overline{\text{Com}}_h^f$. In other words: There exists a homology theory*

$$\bar{\varepsilon}_*: \overline{\text{Com}}_h^f \rightarrow \underline{\text{Ab}}^{\mathbb{Z}}$$

such that

$$\bar{\varepsilon}_* h = \varepsilon_*.$$

REMARK. (1) Corollary 6.7 allows of course a direct proof, which uses only the explicit definition of Ext.

(2) Theorems 6.6 and 6.7 give a precise meaning to the statement that “ ε_* is complex K -homology in the strong shape category”. Moreover it gives an explanation to the fact stated in [9], Corollary 5.10, that $\varepsilon_*(X) \approx \varepsilon_*(Y)$ for two spaces X, Y in $\underline{\text{Com}}^f$ having the same (Borsuk-)shape: Because ε_* is an invariant of strong shape theory, we conclude that $\varepsilon_*(X) \approx \varepsilon_*(Y)$ for spaces X, Y being equivalent in $\underline{\text{Com}}_h$. However two compact metric spaces X, Y are equivalent in $\underline{\text{Com}}_h$ if and only if they are equivalent in the Borsuk shape category.

(3) The restriction to finite dimensional compact metric spaces in 6.6 stems from 4.4 which we were not able to establish without that additional assumption. The same restriction is imposed on the corresponding statement in [8]. Whether Theorem 6.6 is valid for $\underline{\text{Com}}$ (instead of merely $\underline{\text{Com}}^f$) is an open question.

7. Survey of the shape construction. This section is *not* devoted to a rigorous treatment of the strong shape category $\underline{\bar{K}}$ resp. the related homotopy category $\underline{\bar{K}}_h$. This can be found in [4]. We only intend to record

some facts without proofs in order to make this paper as independent as possible. A different approach to strong shape theory can be found in [7].

(1) A 2-category \underline{K} is an ordinary category (i.e. a 1-category) such that $\underline{K}(X, Y)$, $X, Y \in \text{ob } \underline{K}$, carries again the structure of a category (whose morphism are called 2-morphisms).

A 2-functor $\Phi: \underline{K} \rightarrow \underline{L}$ between two 2-categories is a functor *up to 2-morphisms* between the underlying 1-categories (this means e.g. that one has only a 2-morphism $\omega: \Phi(gf) \rightarrow \Phi(g)\Phi(f)$, whenever the compositions are defined, which is not necesesarly the identity). These definitions are completed by requiring various compatibility conditions.

By the way, the concept of a n -functor (resp. a n -category) can be established inductively.

In practice we work with categories of topological spaces, where the 3-category structure can be introduced by taking homotopies as 2-morphisms and homotopies between homotopies as 3-morphisms. Let $H_0, H_1: X \times I \rightarrow Y$ be two homotopies between mappings f_0, f_1 , then a homotopy between homotopies H_0, H_1 is a mapping

$$\xi: X \times I \times I \rightarrow Y$$

having the properties:

$$\begin{aligned} \xi(x, t, i) &= H_i(x, t), \\ \xi(x, i, s) &= f_i(s), \quad i = 0, 1. \end{aligned}$$

In order to equip $\underline{K}(X, Y)$ with the structure of a 2-category, one has to adjust a little the concept of a homotopy; a point, which among other things, we are not going to explain, referring to [4] for the details.

(2) Let \underline{K} be a category of topological spaces (e.g. $\underline{K} = \text{Top}$, $\underline{K} = \text{Com}$, \underline{K} = based or unbased metric spaces and continuous maps) and \underline{P} a subcategory of “good” spaces (e.g. all ANE spaces, all CW-spaces, all compact CW-spaces etc.) The strong shape category $\overline{\underline{K}}$ depends upon the particular choice of this \underline{P} ; it has the same objects as \underline{K} .

Let $X \in \underline{K}$ be any space, then we have a 2-category \underline{P}_X having (1) mappings $g: X \rightarrow P \in \underline{P}$ as objects, (2) pairs $(r, \omega): g_1 \rightarrow g_2$ as morphisms, where $r \in \underline{P}$ is continuous and $\omega: rg_1 \approx g_2$ a given homotopy.

A 2-morphism $(\nu, \xi): (r_1, \omega_1) \approx (r_2, \omega_2): g_1 \rightarrow g_2$ consists of a pair where $\nu: r_1 \approx r_2$ is an ordinary homotopy and $\xi: \omega_2 \circ \nu g_1 \approx \omega_1$ a 2-homotopy. The latter is not simply a homotopy between homotopies, but a *homotopy class* of such a homotopy. This in turn requires the definition of a 3-homotopy which is established analogously to that of a 2-homotopy.

A 2-functor $\bar{f}: \underline{P}_Y \rightarrow \underline{P}_X$, $X, Y \in \underline{K}$ is defined to be a shape mapping $\bar{f} \in \bar{K}(X, Y)$ whenever the following conditions are fulfilled:

$\bar{f}1$) $g: Y \rightarrow P \Rightarrow \bar{f}(g): X \rightarrow P$

$\bar{f}2$) $(r, \omega): g_1 \rightarrow g_2 \Rightarrow \bar{f}(r, \omega) = (r, \omega_1)$ for suitable homotopy ω_1 .

$\bar{f}3$) A corresponding condition involving the 2-morphisms in \underline{P}_Y .

Moreover we require that for a 2-morphism $(\nu, \xi): (r_1, \omega_1) \approx (r_2, \omega_2)$, $(r_i, \omega_i): g_1 \rightarrow g_2$ with $g_i = b_i a$, $a \in \underline{P}_Y$, $\xi = \xi' a$, $\bar{f}(b_i, 1) = (b_i, \gamma_i)$ one has $\bar{f}(\nu, \xi) = (\nu, \tilde{\xi})$ with

$$\tilde{\xi} = \delta_2^{-1} \circ (\gamma_2(\xi' \bar{f}(a)))(r_1 \gamma_1)^{-1} \delta_1$$

and with 2 homotopies

$$\delta_i: \tilde{\omega}_1 \approx \gamma_2(\omega'_i \bar{f}(a))(r_1 \gamma_1)^{-1},$$

$(\bar{f}(r_i, \omega_i) = (r_i, \tilde{\omega}_i), \omega_i = \omega'_i a)$ being explained before.

Finally we require

$\bar{f}4$) For any $(r_1, \omega_1) \in \underline{P}_Y(g_1, g_2)$, $(r_2, \omega_2) \in \underline{P}_Y(g_2, g_3)$. The connecting 2-morphism (whose existence is required in the definition of a 2-functor) is of the form

$$(1, \eta): \bar{f}((r_1, \omega_1)(r_2, \omega_2)) \rightarrow \bar{f}(r_1, \omega_1) \bar{f}(r_2, \omega_2).$$

Suppose now that $(r, \omega): g_1 \rightarrow g_2$ is a given 1-morphism such that $\omega = \omega' a$, $g_i = b_i a$, then we have a 2-morphism

$$(\omega, 1): (r, \omega)(b_1, 1) \rightarrow (b_2, 1)$$

and a connecting morphism

$$(1, \eta)^{-1}: \bar{f}(r, \omega) \bar{f}(b_1, 1) \rightarrow \bar{f}((r, \omega)(b_1, 1)).$$

The composition of these 2-morphisms yields a 2-morphism

$$(r, \tilde{\omega}) = \bar{f}(r, \omega) \xrightarrow{\approx} \bar{f}(b_2, 1) \bar{f}(b_1, 1)^{-1}$$

whose second component δ is recognized as a 2-homotopy between $\tilde{\omega}$ and $\gamma_2(\omega' \bar{f}(a))(r_1 \gamma_1)^{-1}$ (using the previous notation $\bar{f}(b_i, 1) = (b_i, \gamma_i)$).

REMARK. This is a *2-stage strong shape category* because it involves only 2-morphisms (and therefore 2-homotopies but no 3-, 4- etc. homotopies. For compact metric spaces this turns out to be sufficient, while for more general spaces one has to go over to ∞ -categories \underline{P}_Y and ∞ -functors. These objects are much more involved: In case of a ∞ -functor one does not only have connecting morphisms $\Phi(gf) \rightarrow \Phi(g)\Phi(f)$ but also morphisms regulating non existing associativities on *all* levels.

A model for a ∞ -stage strong shape category (in fact: The homotopy category of such a category) has recently been proposed by Ju. Lisica and S. Mardesic respectively (generalizing the considerations in [4]) independently by the present author at the Leningrad topological conference.

Like in \underline{K} the homotopy category \overline{K}_h is established by means of mappings $\bar{F} \in \overline{K}(X \times I, Y)$.

We have a functor $h: \underline{K} \rightarrow \overline{K}$ being defined by

$$h(X) = X, \quad X \in \underline{K}$$

on the objects and by

$$h(f)(g) = gf$$

for a morphism $f \in \underline{K}(X, Y)$. Moreover for $Y \in \underline{P}$ we have an assignment

$$h': \overline{K}(X, Y) \rightarrow \underline{K}(X, Y)$$

defined by

$$h'(\bar{f}) = \bar{f}(1_Y)$$

such that the following properties are satisfied:

- (1) $h'hf = f, \quad f \in \underline{K}(X, Y), \quad Y \in \underline{P},$
- (2) $hh'\bar{f} \approx \bar{f}, \quad \bar{f} \in \overline{K}(X, Y), \quad Y \in \underline{P},$

together with corresponding relations for the higher morphisms. We need the following conclusion:

7.1. PROPOSITION. *Let $\bar{f} \in \overline{K}(X, Y)$ be any shape morphism for $Y \in \underline{P}$, then there exists a continuous $f \in \underline{K}(X, Y)$ such that $h(f) \approx \bar{f}$.*

Proof. Set $f = h'(\bar{f})$.

(3) In order to define homology in \overline{K} we need a new smash-product $X \overline{\wedge} Y$ for $X, Y \in \underline{K}$, which actually leads us out of the category \overline{K} : The new smash-product $X \overline{\wedge} Y$ of two based spaces $(X, x_0), (Y, y_0)$ is a 2-category $\underline{P}_X \overline{\wedge} \underline{P}_Y$ whose objects are factorizations

$$\left(X \wedge Y \xrightarrow[g_1 \wedge g_2]{} P_1 \wedge P_2 \xrightarrow[r]{} P \right)$$

of objects $(g: X \wedge Y \rightarrow P) \in \underline{P}_{X \wedge Y}$, with $P_i \in \underline{P}$. The 1- and 2-morphisms are defined similarly.

One can extend the concept of a shape mapping $\bar{f}: X \rightarrow Y$ to that of a shape mapping

$$\bar{f}: X_1 \overline{\wedge} \cdots \overline{\wedge} X_n \rightarrow Y_1 \overline{\wedge} \cdots \overline{\wedge} Y_m$$

as being a 2-functor (more precisely: an equivalence class of such 2-functors)

$$\bar{f}: \underline{P}_{Y_1} \overline{\wedge} \cdots \overline{\wedge} \underline{P}_{Y_m} \rightarrow \underline{P}_{X_1} \overline{\wedge} \cdots \overline{\wedge} \underline{P}_{X_n}$$

which fulfills the conditions $\bar{f}(1) - \bar{f}(3)$, adapted to the new situation. Details can be found in [4].

Let \underline{E} be any CW-spectrum, then we can define homology groups with coefficients in \underline{E} by

$$\bar{E}_n(X) = \varinjlim \bar{K}_h(S^{n+l}, E_l \overline{\wedge} X), \quad n \in \mathbb{Z}.$$

In [4] §3 we explain, among other things, that for compact metric X this $\bar{E}_n(\)$ is in fact a homology functor which fulfills the Eilenberg-Steenrod axioms for a generalized, reduced homology theory ([4] Theorem 3.1). The crucial problem is embodied in the construction of induced morphisms $\bar{E}_*(\bar{f})$, $\bar{f} \in \bar{K}(X, Y)$, which, in turn, requires the existence of a $\bar{f} \overline{\wedge} 1_Z: X \overline{\wedge} Z \rightarrow Y \overline{\wedge} Z$. According to [4] Propositions 2.1 this can be accomplished for a CW-space Z and a metric space Y (or, alternatively, for all X, Y, Z and continuous $f \in \underline{K}$).

We have to refer to the techniques for the explicit construction of a shape mapping, laid down in [4]. In our case it suffices to consider a $\bar{f}: S^{n+l} \rightarrow E_l \overline{\wedge} X$, where $\underline{E} = \{E_l\}$ is a CW-spectrum (which implies, that all E_l are “good” spaces in \underline{P}) and where X is supposed to be a compact metric space. We embed X into a Hilbert cube Q and consider a decreasing sequence of compact ANEs: $Q = P_1 \supset P_2 \supset \cdots$, with $\bigcap P_i = X$. It turns out to be sufficient to evaluate \bar{f} only on mappings $1_{E_l} \wedge g_i: E_l \wedge X \rightarrow E_l \wedge P_i$, where $g_i: X \rightarrow P_i$ is the inclusion. This is a consequence of [4] Proposition A7. We need this for example in the course of the proof of Proposition 4.1.

At several occasions we use the shape singular complex $\bar{S}(X)$ (resp. $S(X \overline{\wedge} Y)$) of a topological space $X \in \underline{K}$ (resp. of a product $X \overline{\wedge} Y$). This is defined in complete analogy to the classical case as a simplicial set whose simplexes are shape mappings $\bar{\sigma}^n \in \bar{K}((\Delta^n)^+, (X, x_0))$ where Δ^n denotes the standard simplex. The boundary and degeneracy operators are defined like those for the ordinary singular complex $S(X)$. We have that

$$\bar{S}: \bar{K} \rightarrow \mathfrak{S}_E \quad (\text{category of Kan-complexes})$$

is a functor, which appears (as in ordinary topology) together with a natural transformation

$$\bar{\omega}_X: |\bar{S}(X)| \rightarrow X, \quad X \in \underline{K}$$

where $|\cdots|$ denotes, as usual, the geometric realization. We need the fact that:

7.2. PROPOSITION ([2] Theorem 5.1(c)). *The shape mapping $\bar{\omega}_X$ is a weak homotopy equivalence (hence it induces an isomorphism $\bar{\pi}_*(\bar{\omega}_X)$, $\bar{\pi}_*$ being the shape homotopy group functor, $\bar{\pi}_n(X, x_0) = \underline{K}_h(S^n, X)$).*

All spaces are supposed to be equipped with a basepoint, all mappings and homotopies are assumed to be base point preserving, although for the mere definition of \underline{K} this is not explicitly needed.

When talking about \underline{K} , \underline{K} etc. in §1–§3, we assume \underline{K} to be the category Top_0 . Later on we have to restrict ourselves to the full subcategories $\underline{\text{Com}}$ (= category of compact metric (which always means of course: *metrizable*) spaces) or even to $\underline{\text{Com}}^f$ (= category of finite dimensional spaces in $\underline{\text{Com}}$). In the latter case, every $X \in \underline{\text{Com}}^f$ can already be embedded in some sphere S^N and we can assume that all $P_i \in \underline{P}$ occurring in the preceding remark are already lying in this S^N .

As we mentioned already there are many approaches to strong shape theory for compacta (cf. [7]), all of them turn out to be equivalent as homotopy categories (cf. [11], also for further references). The preceding construction leads to individual mappings (rather than to homotopy classes right-away). However it turns out to be a matter of convenience and taste what particular construction somebody is using in order to accomplish a given aim.

Concerning the different kinds of homology theories which appeared in literature, Theorem 5.7 assures us that the homology theory in [7] and the homology theory sh_* of [8] are isomorphic: They are both satisfying the Milnor axioms and they agree on finite CW spaces. The existence of a non-canonical isomorphisms between these homology theories has already been mentioned in [7] §8.

Finally we must point out that for a CW-spectrum $\underline{E} = \{E_i\}$ we understand the cohomology $\underline{E}^n(X)$ as $\varinjlim_k [\Sigma^k X, E_{n+k}]$ and *not* in the sense of the Boardman category [1]. This definition is for our purposes more adequate because it corresponds to Čech cohomology (for $\underline{E} = \underline{K}(G)$) (see [3], [13]).

REFERENCES

- [1] F. Adams, *Stable homotopy and generalized homology*, Chicago Lecture Notes in Mathem. The Univ. of Chicago Press (1974).
- [2] F. W. Bauer, *A shape theory with singular homology*, Pacific J. Math., **62** (1976), 25–65.

- [3] ———, *Under what conditions are shape homology \overline{E}_* and Steenrod homology sE_* isomorphic?* Shape Theory and Geometric Topology Proceedings, Dubrovnik 1981 Lecture Notes in Math., Vol. 870.
- [4] ———, *Duality in manifolds*, Annali di Matematica Pura ed Applicata (IV) (XXXV), (1984), 241–302.
- [5] L. G. Brown, *Characterizing $\text{Ext}(X)$ “K-Theory and Operator Algebras”* Athens, Georgia, Lecture Notes in Math., **575** (1975), 10–18.
- [6] L. G. Brown, R. G. Douglas and P. A. Fillmore, *Extensions of C^* -algebras and K-homology*, Ann. of Math., (2) **105** (1977), 265–324.
- [7] D. A. Edwards and H. M. Hastings, *Čech and Steenrod Homotopy Theories with Applications to Geometric Topology*, Lecture Notes in Math., **542** (1976).
- [8] D. S. Kahn, J. Kaminker and C. Schochet, *Generalized homology theories on compact metric spaces*, Michigan Math. J., **24** (1977), 203–224.
- [9] J. Kaminker and C. Schochet, *K-theory and Steenrod homology; Applications to the Brown-Douglas-Fillmore theory of operator algebras*, Trans. Amer. Math. Soc., **227** (1977), 63–107.
- [10] M. Kernchen, *Bemerkungen zur Borel-Moore-Homologie*, Manuscripta Math., **39** (1982), 111–118.
- [11] Ju. Lisica, *Strong theory of shapes and Steenrod-Sitnikov Homology*, Sib. Math. J., **24** No. 4, (1983), 557–572.
- [12] J. Milnor, *On the Steenrod homology theory*, Berkeley, (unpublished manuscript) (1960).
- [13] K. Morita, *Čech cohomology and covering dimension for topological spaces*, Fund. Math., **LXXXVII** (1975), 31–51.

Received November 22, 1985.

JOHANN WOLFGANG GOETHE UNIVERSITÄT
 ROBERT MAYER STRASSE 6-10
 600 FRANKFURT AM MAIN 11
 WEST GERMANY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

H. SAMELSON
Stanford University
Stanford, CA 94305

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Anthony Peter Bahri and Peter Gilkey , The eta invariant, Pin^c bordism, and equivariant Spin^c bordism for cyclic 2-groups	1
Friedrich-Wilhelm Bauer , Extensions of generalized homology theories	25
Marilyn Breen , A characterization theorem for compact unions of two starshaped sets in R^3	63
Boniface Ihemotuonye Eke , Special generating sets of purely inseparable extension fields of unbounded exponent	73
Robert William Gilmer, Jr. and William James Heinzer, Jónsson ω_0 -generated algebraic field extensions	81
Guido Lupaciolu , Holomorphic continuation in several complex variables	117
Douglas C. McMahon, Jaap C. S. P. van der Woude and Ta-Sun Wu , Connectedness related to almost periodicity of compositions of flow homomorphisms	127
Sophocles Mercourakis , Some characterizations of analytic metric spaces	149
Peter Frederick Stiller , The Picard numbers of elliptic surfaces with many symmetries	157
David J. Winter , Reducible complements of Lie algebra radicals	191
Rade Živaljević , On a cohomology theory based on hyperfinite sums of microsimplexes	201