

Pacific Journal of Mathematics

**VECTOR-VALUED SINGULAR INTEGRAL OPERATORS ON
 L^p -SPACES WITH MIXED NORMS AND APPLICATIONS**

DICESAR LASS FERNANDEZ

VECTOR-VALUED SINGULAR INTEGRAL OPERATORS ON L^p -SPACES WITH MIXED NORMS AND APPLICATIONS

DICESAR LASS FERNANDEZ

We establish L^p and $L^p(l^q)$ estimates for singular integral operators with variable operator-valued product kernels. Application to the strong maximal function, double Hilbert transform, Littlewood-Paley inequalities and Fourier multipliers for L^p -spaces with mixed norm are given.

Introduction. A classical theorem due to Hardy and Littlewood and the improvement given by Fefferman-Stein [6] assert that the maximal function

$$Mf(x) = \sup_I |I|^{-1} \int_I |f(u)| du$$

is bounded on L^p , $1 < p < \infty$, and has a vectorial extension $\tilde{M}(f_j) = (Mf_j)$ bounded in $L^p(l^q)$, $1 < p, q < \infty$. On the other hand, another classical theorem, due to M. Riesz, which asserts that the Hilbert transform

$$Hf(x) = \text{p.v.} \int_{-\infty}^{\infty} \frac{f(y)}{x-y} dy$$

is bounded on L^p , $1 < p < \infty$, was improved by Burkholder [5] for L^p -function with values in Banach spaces with the so called UMD property. In particular, the Hilbert transform has a vectorial extension $\tilde{H}(f_j) = (Hf_j)$ bounded in $L^p(l^q)$, $1 < p, q < \infty$. Recently Rubio de Francia-Ruiz-Torrea [13] and [14] have shown that the maximal operator and the Hilbert transform are operators of same kind: vector-valued singular integral operators. Actually, they improved a theorem on vector-valued convolution operators due to Benedek-Calderón-Panzone [2].

Let us now consider the *rectangular* (strong) maximal function

$$Mf(x, y) = \sup_{I, J} |I \times J|^{-1} \int_J \int_I |f(u, v)| du dv$$

and the double Hilbert transform

$$Hf(x, y) = \text{p.v.} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{f(x - u, y - v)}{(x - u)(y - v)} du dv.$$

By iteration, we see at once that rectangular maximal function and the double Hilbert transform are bounded on $L^p(\mathbf{R}^2)$, $1 < p < \infty$. But, this is not the case if we replace the usual L^p spaces by the $L'(L^s)$ spaces with mixed norm of Benedek-Panzone [1]. The boundedness now does not follow from a simple iteration. The $L'(L^s)$ -norm estimate for the strong maximal function was stated by Stöckert [17], but a very nice proof was given by E. Hernández [9]; on the other hand, for the double Hilbert transform it goes back to M. Cotlar.

Our concern here is to establish a theory of vector singular integral operators with variable product kernels. This will be done in the mold of Rubio de Francia-Ruiz-Torrea [13] and [14], and in such a way to be possible to handle with it the scalar strong maximal function and double Hilbert transform as well as its sequential extensions. As applications we also obtain an inequality of Littlewood-Paley type for $L'(L^s)$ -spaces and derive a multiplier theorem of Marcinkiewicz-Lizorkin type in a simple and natural way.

1. Vector-valued singular integral operators with product kernels.

We begin by recalling the Rubio de Francia-Ruiz-Torrea theorem on vector-valued singular integral operators (see [13]).

1.1. THEOREM. *Suppose E and F denote Banach spaces. For $\Delta = \{(x, y) \in \mathbf{R}^n \times \mathbf{R}^n, x = y\}$, let $k \in L^1_{\text{loc}}(\mathbf{R}^n \times \mathbf{R}^n - \Delta, L(E, F))$ be an operator-valued kernel which satisfies*

$$(1) \quad \int_{|y' - x| \geq 2|y' - y|} \|k(x, y) - k(x, y')\|_{L(E, F)} dx \leq C,$$

and

$$(2) \quad \int_{|y - x'| \geq 2|x - x'|} \|k(x, y) - k(x', y)\|_{L(E, F)} dy \leq C.$$

Let T be a linear bounded operator from $L^r(\mathbf{R}^n, E)$ into $L^r(\mathbf{R}^n, F)$, for some r with $1 < r < \infty$, such that

$$(3) \quad Tf(x) = \int_{\mathbf{R}^n} k(x, y)f(y) dy,$$

for all $f \in L^\infty_c(\mathbf{R}^n, E)$ (the linear space of all E -valued measurable functions which are essentially bounded and have compact support) and $x \notin \text{supp } f$.

Then, for all p with $1 < p < \infty$, we have

$$(4) \quad \|Tf\|_{L^p(\mathbf{R}^n, F)} \leq C\|f\|_{L^p(\mathbf{R}^n, E)}, \quad f \in L_c^\infty(\mathbf{R}^n, E).$$

Moreover, for all q with $1 < q < \infty$, we also have

$$(5) \quad \|TF\|_{L^p(\mathbf{R}^n, l^q(F))} \leq C\|F\|_{L^p(\mathbf{R}^n, l^q(E))}, \quad F = (f_j) \in L_c^\infty(\mathbf{R}^n, l^q(E)).$$

We shall rely on the above theorem to give a version of it for product kernels $k(x, u, y, v) = k_2(y, v)k_1(x, u)$ and the $L^p = L^{p_2}(L^{p_1})$ spaces with mixed norms of Benedek-Panzone [1], from where we take notations. The proof we shall give follows an idea by Benedek-Calderón-Panzone [2].

1.2. THEOREM. Suppose E, F and G denote Banach spaces. Let us consider operator-valued kernels k_1 and k_2 in $L^1_{\text{loc}}(\mathbf{R}^m \times \mathbf{R}^n - \Delta, L(E, F))$ and $L^1_{\text{loc}}(\mathbf{R}^n \times \mathbf{R}^n - \Delta, L(F, G))$, respectively, which satisfy

$$(1) \quad \int_{|z-w'| \geq 2|w-w'|} \|k_j(z, w) - k_j(z, w')\|_{L_j} dz \leq C_j, \quad j = 1, 2,$$

and

$$(2) \quad \int_{|z'-w| \geq 2|z'-z|} \|k_j(z, w) - k_j(z', w)\|_{L_j} dw \leq C_j, \quad j = 1, 2,$$

where $L_1 = L(E, F)$ and $L_2 = L(F, G)$. Let T_1 and T be linear bounded operators from $L^p(\mathbf{R}^m, E)$ into $L^p(\mathbf{R}^m, F)$ and from $L^p(\mathbf{R}^n \times \mathbf{R}^n, E)$ into $L^p(\mathbf{R}^m \times \mathbf{R}^n, G)$, for all p with $1 < p < \infty$, respectively. Suppose also that T_1 and T satisfy

$$(3) \quad T_1 f(x) = \int_{\mathbf{R}^m} k_1(x, u) f(u) du$$

for all $f \in L_c^\infty(\mathbf{R}^m, E)$ and $x \notin \text{supp } f$, and

$$(4) \quad Tf(x, y) = \int_{\mathbf{R}^n} \int_{\mathbf{R}^n} k_2(y, v) k_1(x, u) f(u, v) du dv$$

for all $f \in L_c^\infty(\mathbf{R}^m \times \mathbf{R}^n, E)$ and $(x, y) \notin \text{supp } f$. Then, for all $P = (p_1, p_2)$ with $1 < p_1, p_2 < \infty$, the linear operator T can be extended to all $L^P(\mathbf{R}^m \times \mathbf{R}^n, E)$ into $L^P(\mathbf{R}^m \times \mathbf{R}^n, G)$ such that

$$(5) \quad \|Tf\|_{L^P(\mathbf{R}^m \times \mathbf{R}^n, G)} \leq C\|f\|_{L^P(\mathbf{R}^m \times \mathbf{R}^n, E)},$$

for all $f \in L^P(\mathbf{R}^m \times \mathbf{R}^n, E)$.

Proof. Step 1. Let f be in $M(\mathbf{R}^{m+n}, E)$, the set of all E -valued measurable functions on \mathbf{R}^{m+n} . For each $y \in \mathbf{R}^n$ and each $f \in M(\mathbf{R}^{m+n}, E)$ we associate the functions $f = f_y \in M(\mathbf{R}^m, E)$ defined

by $\pi f = f_y(\cdot) = f(\cdot, y)$. Thus, we shall have $L^p(\mathbf{R}^{m+n}, E) = L^p(\mathbf{R}^n, L^p(\mathbf{R}^m, E))$, $L^p(\mathbf{R}^{m+n}, E) = L^{p_2}(\mathbf{R}^n, L^{p_1}(\mathbf{R}^m, E))$, $P = (p_1, p_2)$ and

$$\|\pi f\|_{L^{p_2}(\mathbf{R}^n, L^{p_1}(\mathbf{R}^m, E))} = \|f\|_{L^p(\mathbf{R}^{m+n}, E)}.$$

Moreover, if $f \in L_c^\infty(\mathbf{R}^{m+n}, E)$ we shall have $f_y \in L_c^\infty(\mathbf{R}^m, E)$.

Step 2. For all $f \in M(\mathbf{R}^{m+n}, E)$ and $\lambda > 0$, we define $H_\lambda f$ by

$$(H_\lambda f)(x, y) = \chi_\lambda(x) f(x, y),$$

where χ_λ is the characteristic function of the set $\{|x| < \lambda\}$. Now, if $y \in \mathbf{R}^n$, we define $K_\lambda(y) \in L(L^p(\mathbf{R}^m, E), L^p(\mathbf{R}^m, G))$ by

$$[K_\lambda(y, v)h](u) = \chi_\lambda k_2(y, v)[T_1(\chi_\lambda h)](u).$$

Then, if $\|\cdot\|$ denotes the operator norm on $L(L^p(\mathbf{R}^m, E), L^p(\mathbf{R}^m, F))$, since the singular integral operator $T_1 f$ is bounded from $L^p(\mathbf{R}^m, E)$ into $L^p(\mathbf{R}^m, F)$, it follows that

$$\begin{aligned} \|K_\lambda(y, v)\| &= \sup\{\|K_\lambda(y, v)h\|_{L^p(G)}; \|h\|_{L^p(E)} \leq 1\} \\ &\leq \sup\{\|k_2(y, v)T_1(\chi_\lambda h)\|_{L^p(G)}; \|h\|_{L^p(E)} \leq 1\} \\ &\leq \|k_2(y, v)\|_{L(F, G)} \sup\{\|T_1(\chi_\lambda h)\|_{L^p(G)}; \|h\|_{L^p(E)} \leq 1\} \\ &\leq \|k_2(y, v)\|_{L(F, G)} \sup\{C\|\chi_\lambda h\|_{L^p(E)}; \|h\|_{L^p(E)} \leq 1\} \\ &\leq C\|k_2(y, v)\|_{L(F, G)}, \end{aligned}$$

which shows that $K_\lambda(y, v) \in L_{\text{loc}}^1(\mathbf{R}^n, L(L^p(E), L^p(G)))$. Moreover, we have

$$\begin{aligned} &\int_{|y-y'| > 2|v-v'|} \|K_\lambda(y, v) - K_\lambda(y, v')\| dy \\ &\leq C_1 \int_{|y-y'| > 2|v-v'|} \|k_2(y, v) - k_2(y, v')\| dy \leq C_1 C_2 \end{aligned}$$

and

$$\begin{aligned} &\int_{|v-y'| > 2|y-y'|} \|K_\lambda(y, v) - K_\lambda(y', v)\| dv \\ &\leq C_1 \int_{|v-y'| > 2|y-y'|} \|k_2(y, v) - k_2(y', v)\| dv \leq C_1 C_2. \end{aligned}$$

Now, for $F \in L_c^\infty(\mathbf{R}^n, L^p(\mathbf{R}^m, E))$, we get

$$T_\lambda F(y) = \int_{\mathbf{R}^n} K_\lambda(y, v) F(v) dv.$$

Since $\pi H_\lambda TH_\lambda f = T_\lambda \pi f$, due to hypothesis (4) we get

$$\begin{aligned} \|T_\lambda \pi f\|_{L^p(\mathbf{R}^n, L^p(\mathbf{R}^m, G))} &= \|\pi H_\lambda TH_\lambda f\|_{L^p(\mathbf{R}^n, L^p(\mathbf{R}^m, G))} \\ &= \|H_\lambda TH_\lambda f\|_{L^p(\mathbf{R}^m \times \mathbf{R}^n; G)} \\ &\leq \|TH_\lambda f\|_{L^p(\mathbf{R}^{m+n}, G)} \leq C \|H_\lambda f\|_{L^p(\mathbf{R}^{m+n}, E)} \\ &\leq C \|f\|_{L^p(\mathbf{R}^{m+n}, E)} = C \|\pi f\|_{L^p(\mathbf{R}^n, L^p(\mathbf{R}^m, E))}. \end{aligned}$$

Consequently, setting $p = p_1$, $A = L^{p_1}(\mathbf{R}^m, E)$ and $B = L^{p_1}(\mathbf{R}^m, G)$, we have

$$\|T_\lambda F\|_{L^{p_1}(\mathbf{R}^n, B)} \leq C \|F\|_{L^{p_1}(\mathbf{R}^n, A)},$$

for all F given by $F = \pi f$, $f \in L_c^\infty(\mathbf{R}^m \times \mathbf{R}^n, E)$. Since $L_c^\infty(\mathbf{R}^{m+n}, E) = L_c^\infty(\mathbf{R}^n, L_c^\infty(\mathbf{R}^m, E))$ is dense in $L_c^\infty(\mathbf{R}^n, L^p(\mathbf{R}^m, E)) = L_c^\infty(\mathbf{R}^n, A)$, in the norm of $L^p(\mathbf{R}^n, A)$, we have 1.1(4) for all $F \in L_c^\infty(\mathbf{R}^n, A)$. Now, the Rubio de Francia-Ruiz-Torrea theorem yields

$$\|T_\lambda F\|_{L^{p_2}(\mathbf{R}^n, B)} \leq C \|F\|_{L^{p_2}(\mathbf{R}^n, A)},$$

for all $F \in L^{p_2}(\mathbf{R}^n, A)$ and $1 < p_2 < \infty$.

Step 3. We shall have

$$\begin{aligned} \|H_\lambda TH_\lambda f\|_{L^p(\mathbf{R}^{m+n}, G)} &= \|\pi H_\lambda TH_\lambda f\|_{L^{p_2}(\mathbf{R}^n, L^{p_1}(\mathbf{R}^m, G))} \\ &= \|T_\lambda \pi f\|_{L^{p_2}(\mathbf{R}^n, L^{p_1}(\mathbf{R}^m, G))} \\ &\leq C \|\pi f\|_{L^{p_2}(\mathbf{R}^n, L^{p_1}(\mathbf{R}^m, E))} = C \|f\|_{L^p(\mathbf{R}^{m+n}, E)}. \end{aligned}$$

Finally, since $H_\lambda f = f$, for λ large enough, we get

$$\begin{aligned} \|Tf\|_{L^p(\mathbf{R}^{m+n}, G)} &\leq \liminf_{\lambda \rightarrow \infty} \|H_\lambda TH_\lambda f\|_{L^p(\mathbf{R}^{m+n}, G)} \\ &\leq C \|f\|_{L^p(\mathbf{R}^{m+n}, E)}, \end{aligned}$$

for all $f \in L_c^\infty(\mathbf{R}^{m+n}, E)$ and consequently for all $f \in L^p(\mathbf{R}^{m+n}, E)$.

The proof is complete.

1.3. Let us recall that $l^Q(X)$, $Q = (q_1, q_2)$, denotes the linear space of all X -valued double sequences (C_{ij}) such that

$$(1) \quad \|(C_{ij})\|_{l^Q(X)} = \|(C_{ij})\|_{l^{q_2}(l^{q_1}(X))} = \left\{ \sum_j \left\{ \sum_i |C_{ij}|_X^{q_1} \right\}^{q_2/q_1} \right\}^{1/q_2} < \infty.$$

1.4. COROLLARY. Suppose E, F and G denote Banach spaces and let us consider operator-valued kernels k_1 and k_2 as in Theorem 1.2. We define the kernels $\tilde{k}_1(x) \in L(l^Q(E), l^Q(F))$, and $\tilde{k}_2(y) \in L(l^Q(F), l^Q(G))$ by

$$(1) \quad \tilde{k}_1(x)(a_{ij})_{ij} = (k_1(x)a_{ij})_{ij}$$

and

$$(2) \quad \tilde{k}_2(y)(b_{ij})_{ij} = (k_2(y)b_{ij})_{ij}.$$

We define, for $f = (f_{ij}) \in L_c^\infty(\mathbf{R}^m, l^Q(E))$,

$$(3) \quad \tilde{T}_1 f(x) = \tilde{T}_1(f_{ij}) = \int_{\mathbf{R}^m} \tilde{k}_1(x, u)(f_{ij}(u)) du$$

and for $f \in L_c^\infty(\mathbf{R}^{m+n}, l^Q(E))$

$$(4) \quad \tilde{T}f(x, y) = \tilde{T}(f_{ij}) = \int_{\mathbf{R}^n} \int_{\mathbf{R}^m} \tilde{k}_2(y, v)\tilde{k}_1(x, u)(f_{ij}(u, v)) du dv.$$

We shall assume that, for all p with $1 < p < \infty$, the operators \tilde{T}_1 and \tilde{T} are bounded from $L^p(\mathbf{R}^m, l^Q(E))$ into $L^p(\mathbf{R}^m, l^Q(F))$ and from $L^p(\mathbf{R}^{m+n}, l^Q(E))$ into $L^p(\mathbf{R}^{m+n}, l^Q(G))$, respectively. Then, for all $P = (p_1, p_2)$ with $1 < p_1, p_2 < \infty$, the linear operators T can be extended to all $L^P(\mathbf{R}^{m+n}, l^Q(E))$.

Moreover, we shall have

$$(5) \quad \|Tf\|_{L^P(\mathbf{R}^{m+n}, l^Q(G))} \leq C\|f\|_{L^P(\mathbf{R}^{m+n}, l^Q(E))}.$$

Proof. Since, for $\nu = 1, 2$,

$$\|\tilde{k}_\nu(z, w) - \tilde{k}_\nu(z, w')\|_{L(l^Q(E), l^Q(F))} \leq \|k_\nu(z, w) - k_\nu(z, w')\|_{L(E, F)},$$

and

$$\|\tilde{k}_\nu(z, w) - \tilde{k}_\nu(z', w)\|_{L(l^Q(F), l^Q(G))} \leq \|k_\nu(z, w) - k_\nu(z', w)\|_{L(F, G)},$$

it follows that the hypotheses of Theorem 1.2 are fulfilled with the kernels k_1 and k_2 and the spaces E, F and G replaced by the kernels \tilde{k}_1 and \tilde{k}_2 and the spaces $l^Q(E), l^Q(F)$ and $l^Q(G)$, respectively. Hence, the corollary follows as desired.

1.5. REMARK. When $k_\nu(z, w) = k_\nu(z - w)$, $\nu = 1, 2$, we have the singular integral operators of convolution type. This particular type of singular integral operators was studied, also in the product case, by the author in [8].

2. The double Hilbert transform.

2.1. Our concern here is to obtain L^P and $L^P(l^Q)$ estimates for the *double Hilbert transform* which is defined by

$$(1) \quad Hf(x, y) = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{f(u, v)}{(x - u)(y - v)} du dv.$$

This is an integral singular operator of convolution type and the mixed L^P -estimate goes back to M. Cotlar. The mixed $L^P(l^Q)$ -estimate seems new.

2.2. THEOREM. *The double Hilbert transform given by 2.1(1) is an integral singular operator bounded in the spaces $L^P(\mathbf{R}^2)$, where $P = (p_1, p_2)$ with $1 < p_1, p_2 < \infty$, i.e.*

$$(1) \quad \|Hf\|_{L^P} \leq C_P \|f\|_{L^P},$$

for all $f \in L^P(\mathbf{R}^2) = L^{p_2}(L^{p_1})$.

Proof. Step 1. The kernels $k_j(z, w) = 1/(z - w)$ satisfy conditions 1.2(1)–(2).

Step 2. The integral singular operator T_1 associated with the kernel k_1 is bounded in $L^p(\mathbf{R})$, $1 < p < \infty$.

Step 3. The integral given by 2.1(1) is well defined for all $f \in L_c^\infty(\mathbf{R}^2)$ and $(x, y) \notin \text{supp } f$.

Step 4. By iteration we see that H is bounded in $L^p(\mathbf{R}^2)$, $1 < p < \infty$.

Hence, the conditions of Theorem 1.2 are satisfied and the assertion follows.

2.3. COROLLARY. *In the conditions of Theorem 2.2, for all $F = (f_{ij}) \in L^P(l^Q)$, $1 < Q = (q_1, q_2) < \infty$, we have*

$$(1) \quad \|(Hf_{ij})\|_{L^P(l^Q)} \leq C \|(f_{ij})\|_{L^P(l^Q)}.$$

Proof. It follows from Corollary 1.3.

3. The rectangular maximal function of F. Zó's type. The following version of a theorem due to F. Zó [19] will be needed to obtain the maximal inequality which we are looking for.

3.1. THEOREM. Suppose $\varphi \in L^1(\mathbf{R}^m)$ and $\psi \in L^1(\mathbf{R}^n)$. For $s > 0$ and $t > 0$, let us set $\varphi_s(x) = s^{-m}\varphi(s^{-1}x)$ and $\psi_t(y) = t^{-n}\psi(t^{-1}y)$. We shall also suppose that

$$(1) \quad \int_{|x|>4|u|} \sup_{s>0} |\varphi_s(u-x) - \varphi_s(x)| dx \leq C_1, \quad u \in \mathbf{R}^m$$

and

$$(2) \quad \int_{|y|>4|v|} \sup_{t>0} |\psi_t(v-y) - \psi_t(y)| dy \leq C_2, \quad v \in \mathbf{R}^n.$$

Now, for $f \in L_c^\infty(\mathbf{R}^{m+n})$, we get

$$(3) \quad M_{\varphi\psi}f = \sup\{|\psi_t\varphi_s * f|; s > 0, t > 0\}.$$

Then, for all $P = (p_1, p_2)$ with $1 < p_1, p_2 < \infty$, we have

$$(4) \quad \|M_{\varphi\psi}f\|_{L^P} \leq C\|f\|_{L^P},$$

for all $f \in L^P(\mathbf{R}^{m+n})$.

Proof. Step 1. If $f \in L^1 \cap L^\infty$, the mappings $(s, t) \rightarrow \psi_t\varphi_s * f(x, y)$ are uniformly continuous. Indeed since the mapping $(s, t) \rightarrow \psi_t\varphi_s$ is continuous from \mathbf{R}^2 , into $L^1(\mathbf{R}^2)$ we have

$$|\psi_{t'}\varphi_{s'} * f(x, y) - \psi_{t''}\varphi_{s''} * f(x, y)| \leq \|f\|_{L^\infty} \|\psi_{t'}\varphi_{s'} - \psi_{t''}\varphi_{s''}\|_{L^1}.$$

Therefore, it is enough to prove (4) taking the supremum on $Q_+ \times Q_+$.

Step 2. Fixing an enumeration of the positive rationals, let us denote by Q_j the set of rationals with indices in $\{1, \dots, j\}$. For c and d in \mathbf{N} , let us set

$$(5) \quad M_{cd}f = \sup\{|\psi_t\varphi_s * f|; (s, t) \in Q_c \times Q_d\}.$$

If l_j^∞ and l_{ij}^∞ stand for the complex euclidean spaces \mathbf{C}^j and \mathbf{C}^{i+j} , respectively, equipped with the sup-norm, the non-linear operator M_{cd} can be viewed as a vector-valued linear operator

$$(6) \quad f \in L_c^\infty(\mathbf{R}^{m+n}) \rightarrow N_{cd}f = \{\psi_t\varphi_s * f\}_{(s,t) \in Q_c \times Q_d} \in L^\infty(\mathbf{R}^{m+n}, l_{cd}^\infty).$$

The kernel k_{cd} of this operator is a product of the two kernels:

$$k_c^1(x) \in L(\mathbf{C}, l_c^\infty) \simeq l_c^\infty, \quad x \in \mathbf{R}^m,$$

and

$$k_d^2(y) \in L(l_c^\infty, l_{cd}^\infty), \quad y \in \mathbf{R}^n,$$

given by

$$k_c^1(x)a = \{\varphi_s(x)a; s \in Q_c\}$$

and

$$k_d^2(y)(b_j) = \{\psi_t(y)b_j; t \in Q_d, 1 \leq j \leq c\}.$$

Hence

$$k_{cd}(x, y) = k_d^2(y)k_c^1(x)z = \{\psi_t(y)\varphi_s(x)z; s \in Q_c, t \in Q_d\}.$$

On the other hand, we have

$$\begin{aligned} \|k_c^1(x)\| &= \sup_{|z|=1} \|k_c^1(x)z\|_{l^\infty} \\ &= \sup_{|z|=1} \sup_{s \in Q_c} |\varphi_s(x)z| = \sup_{s \in Q_c} |\varphi_s(x)| \end{aligned}$$

and

$$\begin{aligned} \|k_d^2(y)\| &= \sup_{\|(a_j)\|_\infty=1} \|k_d^2(y)(a_j)\|_{l^\infty} \\ &= \sup_{\|(a_j)\|_\infty=1} \sup_{t \in Q_d} |\psi_t(y)a_j| = \sup_{t \in Q_d} |\psi_t(y)|. \end{aligned}$$

Therefore k_c^1 and k_d^2 are locally integrable. Moreover

$$\begin{aligned} &\int_{|x|>4|u|} \|k_c^1(u-x) - k_c^1(x)\| dx \\ &= \int_{|x|>4|u|} \sup_z \|[k_c^1(u-x) - k_c^1(x)]z\|_{l^\infty} dx \\ &= \int_{|x|>4|u|} \sup_z \sup_s |\varphi_s(u-x) - \varphi_s(x)| |z| dx \leq C_1 \end{aligned}$$

and

$$\begin{aligned} &\int_{|y|>4|v|} \|k_d^2(v-y) - k_d^2(y)\| dy \\ &= \int_{|y|>4|v|} \sup_{(a_j)} \|[k_d^2(v-y) - k_d^2(y)](a_j)\|_{l^\infty} dy \\ &= \int_{|y|>4|v|} \sup_{(a_j)} \sup_{t,j} |\psi_t(v-t) - \psi_t(y)| |a_j| dy \\ &= \int_{|y|>4|v|} \sup_t |\psi_t(v-y) - \psi_t(y)| dy \leq C_2. \end{aligned}$$

Step 3. Due to Zó's result, for all p with $1 < p < \infty$, we have

$$\begin{aligned} \|M_{cd}f\|_{L_{xy}^p} &\leq \|M_{\varphi\psi}f\|_{L_{xy}^p} \leq \|M_{\varphi}M_{\psi}f\|_{L_{xy}^p} \leq C\| \|M_{\psi}f\|_{L_x^p}\|_{L_y^p} \\ &\leq C\| \|M_{\psi}f\|_{L_y^p}\|_{L_x^p} \leq C^1\| \|f\|_{L_y^p}\|_{L_x^p} = C^1\|f\|_{L_{xy}^p}, \end{aligned}$$

where L_{xy}^p , L_x^p and L_y^p have obvious meaning.

Step 4. Since the operator $M_{cd}f = k_{cd} * f$ satisfies the hypotheses of Theorem 1.2 it follows that

$$(7) \quad \|M_{cd}f\|_{L^p(\mathbf{R}^{m+n}, l_{cd}^{\infty})} \leq C\|f\|_{L^p(\mathbf{R}^{m+n})}$$

for all $f \in L_c^{\infty}(\mathbf{R}^{m+n})$. But $M_{cd}f$ has also a sense for all $f \in L^p(\mathbf{R}^{m+n})$ and it is not hard to see that the extension \tilde{M}_{cd} of M_{cd} to all $L^p(\mathbf{R}^{m+n})$ coincides with M_{cd} . Thus (7) holds for all $f \in L^p(\mathbf{R}^{m+n})$. Finally, letting $|(c, d)| \rightarrow \infty$, the monotone convergence theorem yields (4).

The proof is complete.

3.2. THEOREM. *Let φ_s and ψ_t be as in Theorem 3.1. For $f = (f_{ij})$ in $L_c^{\infty}(\mathbf{R}^{m+n}, l^Q)$, where $Q = (q_1, q_2)$ is given with $1 < q_1, q_2 < \infty$, let us consider the vectorial rectangular maximal function*

$$(1) \quad \tilde{M}_{\varphi\psi}(f_{ij})_{ij} = (M_{\varphi\psi}f_{ij})_{ij}$$

where $M_{\varphi\psi}f_{ij}$ is the maximal function given by 2.1(3). Then, if $P = (p_1, p_2)$ is given with $1 < p_1, p_2 < \infty$, we have

$$(2) \quad \|\tilde{M}_{\varphi\psi}(f_{ij})\|_{L^p(\mathbf{R}^{m+n}, l^Q)} \leq C\|(f_{ij})\|_{L^p(\mathbf{R}^{m+n}, l^Q)},$$

for all $f = (f_{ij}) \in L^p(\mathbf{R}^{m+n}, l^Q)$.

Proof. As before, let us replace the maximal function $M_{\varphi\psi}g$ by $M_{cd}g$ and let us consider the vectorial linear operator

$$\begin{aligned} \tilde{T}_{cd}: (f_{ij}) \in L_c^{\infty}(\mathbf{R}^{m+n}, l^Q) \\ \rightarrow \tilde{T}_{cd}(f_{ij}) = ((\psi_t\varphi_s * f_{ij})_{st})_{ij} \in L^{\infty}(\mathbf{R}^{m+n}, l^Q(l_{cd}^{\infty})). \end{aligned}$$

The kernel of this operator is a product kernel $k(x, y) = k_2(y)k_1(x)$, where

$$k_1(x): (a_{ij}) \in l^Q \rightarrow k_1^1(a_{ij}) = (\varphi_s(x)a_{ij}) \in l^Q(l_c^{\infty})$$

and

$$k_2(y): (b_{ijs}) \in l^Q(l_c^{\infty}) \rightarrow k_2(y)(b_{ijs}) = (\psi_t(y)b_{ijs}) \in l^Q(l_{st}^{\infty}).$$

If $\| \cdot \|$ denotes the norm on $L(I^Q, I^Q(I_c^\infty))$, we shall have

$$\begin{aligned} & \int_{|x|>4|u|} \|k_c^1(u-x) - k_c^1(x)\| dx \\ &= \int_{|x|>4|u|} \sup_{\| (a_{ij}) \|_{l^Q} = 1} \|([\varphi_s(u-x) - \varphi_s(x)] a_{ij})\|_{l^Q(I_c^\infty)} dx \\ &\leq \int_{|x|>4|u|} \sup_{s \in Q_s} |\varphi_s(u-x) - \varphi_s(x)| dx \leq C_1, \end{aligned}$$

and if $\| \cdot \|$ denotes the norm on $L(I^Q(I_c^\infty), I^Q(I_{cd}^\infty))$, we also have

$$\begin{aligned} & \int_{|y|>4|v|} \|k_d^2(v-y) - k_d^2(y)\| dy \\ &\leq \int_{|y|>4|v|} \sup_{\| (b_{ijs}) \|_{l^Q(I_{cd}^\infty)} = 1} \|([\psi_t(v-y) - \psi_t(y)] b_{ijs})\|_{l^Q(I_{cd}^\infty)} dy \\ &\leq \int_{|y|>4|v|} \sup_{t \in Q_d} \|\psi_t(v-y) - \psi_t(y)\| \sup_{\|b\|=1} (b_{ijs})_{l^Q(I_c^\infty)} dy \\ &\leq \int_{|y|>4|v|} \sup_{t \in Q_d} |\psi_t(v-y) - \psi_t(y)| dy \leq C_2. \end{aligned}$$

Now, it remains to prove that, for all p with $1 < p < \infty$, we have

$$(3) \quad \|\tilde{T}_{cd}(f_{ij})\|_{L^p(I^Q(I^\infty))} \leq C \| (f_{ij}) \|_{L^p(I^Q)},$$

where C is a constant independent of c, d and p . Let us consider the partial operators \tilde{M}_c and \tilde{T}_d given by

$$\tilde{M}_c(f_{ij}) = (M_c f_{ij}) = \left(\sup_{s \in Q_c} |\varphi_s * _y f_{ij}| \right)_{ij}$$

and

$$\tilde{T}_d(f_{ij}) = (T_d f_{ij}) = (f_{ij} * _y \psi_t).$$

We shall have

$$T_{cd} f_{ij} \leq T_d M_c f_{ij}.$$

Thus, due to Zó's result (and Fubini's theorem) we see that \tilde{M}_c and \tilde{T}_d are bounded operators from $L_{xy}^p(I_{ij}^Q)$ into $L_{xy}^p(I_{ij}^Q)$ and from $L_{xy}^p(I_{ij}^Q(I_c^\infty))$ into $L_{xy}^p(I_{ij}^Q)$, respectively. Consequently, if $(f_{ij}) \in L^p(I^Q)$ we shall have $(g_{ij}) \in L^p(I^Q)$, where

$$g_{ij} = M_c f_{ij} = \sup_{s \in Q_c} |\varphi_s * _x f_{ij}|.$$

Hence

$$\begin{aligned} \|\tilde{T}_{cd}(f_{ij})\|_{L^p(I^Q(I^\infty))} &\leq \|(T_d M_c f_{ij})\|_{L^p(I^Q(I^\infty))} \\ &\leq C\|(M_c f_{ij})\|_{L^p(I^Q)} \leq C\|(f_{ij})\|_{L^p(I^Q)}. \end{aligned}$$

The proof is complete.

4. The rectangular maximal function. We are now ready to state inequalities for the rectangular maximal functions of Hardy-Littlewood and of Fefferman-Stein type for L^P spaces with mixed norms.

4.1. THEOREM. *Suppose $f \in L^1_{\text{loc}}(\mathbf{R}^{m+n})$ and let us consider the rectangular maximal function Mf defined by*

$$(1) \quad Mf(x, y) = \sup_{I, J} |I \times J|^{-1} \int_J \int_I |f(x - u, y - v)| \, du \, dv,$$

where I and J are (hyper)-cubes centered at the origin of \mathbf{R}^m and \mathbf{R}^n , respectively. Then, if $f \in L^P(\mathbf{R}^{m+n})$, where $P = (p_1, p_2)$ with $1 < p_1, p_2 < \infty$, $Mf(x, y)$ is finite for a.e. $(x, y) \in \mathbf{R}^{m+n}$. Moreover, there is a constant $C > 0$ such that

$$(2) \quad \|Mf\|_{L^P(\mathbf{R}^{m+n})} \leq C\|f\|_{L^P(\mathbf{R}^{m+n})}$$

for all $f \in L^P(\mathbf{R}^{m+n})$.

Proof. Let I_1 and J_1 be the unit cubes on \mathbf{R}^m and \mathbf{R}^n , and let us consider the dilated cubes I_s and J_t with side length s and t , respectively. Now, let $\varphi \in C_c^\infty(\mathbf{R}^m)$ and $\psi \in C_c^\infty(\mathbf{R}^n)$ such that $\varphi(x) = \psi(y) = 1$, for $x \in I$, and $y \in J$, respectively. Then

$$|I_s \times J_t|^{-1} \chi_{I_s \times J_t} \leq \varphi_s \psi_t$$

and

$$\begin{aligned} Mf(x, y) &= \sup_{s, t} \int_{\mathbf{R}^m} \int_{\mathbf{R}^n} |f(x - u, y - v)| |I_s \times J_t|^{-1} \chi_{I_s \times J_t}(u, v) \, du \, dv \\ &\leq M_{\varphi\psi} f(x, y). \end{aligned}$$

Now, from Theorem 3.1, the maximal inequality (2) follows at once.

4.2. THEOREM. *Suppose $f = (f_{ij}) \in L^1_{\text{loc}}(\mathbf{R}^{m+n}, I^Q)$, where $Q = (q_1, q_2)$ is given with $1 < q_1, q_2 < \infty$. The vectorial rectangular maximal function is given by*

$$(1) \quad \tilde{M}(f_{ij})_{ij} = (Mf_{ij})_{ij}$$

where Mf_{ij} is the rectangular maximal function. Then, there is a constant $C > 0$ such that for all $P = (p_1, p_2)$ with $1 < p_1, p_2 < \infty$ and $f = (f_{ij}) \in L^P(\mathbf{R}^{m+n}, l^Q)$ we have

$$(2) \quad \|\tilde{M}(f_{ij})_{ij}\|_{L^P(\mathbf{R}^{m+n}, l^Q)} \leq C\|(f_{ij})\|_{L^P(\mathbf{R}^{m+n}, l^Q)}.$$

Proof. It follows at once from Theorem 2.2 as in the proof of Theorem 3.1.

4.3. REMARK. The inequality 4.1(2) in the case $m = n = 1$ was stated by Stöckert [17]. But the inequality 4.2(2) seems new and it was proved by the author in [8]. (However see Schmeisser [15] and the references quoted there.)

5. Application to the Littlewood-Paley theorem.

5.1. PROPOSITION. For $f \in S(\mathbf{R}^2)$ and I and J numerical intervals, the (iterated) partial sum operator is defined by

$$(1) \quad (S_{I \times J} f)^\wedge(s, t) = \chi_I(s)\chi_J(f)\hat{f}(s, t)$$

and we have

$$(2) \quad \|S_{I \times J} f\|_{L^P} \leq C\|f\|_{L^P}, \quad 1 < P = (p_1, p_2) < \infty,$$

for all $f \in S(\mathbf{R}^2)$, with C independent of f . Moreover, $S_{I \times J}$ can be extended to all $L^P(\mathbf{R}^2)$.

Proof. If $I = J = (0, \infty)$, then

$$(3) \quad S_{I \times J} f = (1/4)(f + iH_{10}f + iH_{01} - H_{11}f)$$

where $H_{10}f$, $H_{01}f$ and $H_{11}f$ are the partial and double Hilbert transform. In this case we have obviously (2). The general case follows by modifications of (3) as in the one-dimensional case.

As the partial and double Hilbert transform have an l^Q -extension, Theorem 5.1 has the following extension.

5.2. THEOREM. Let $(I_i \times J_j)_{i,j \in \mathbf{N}}$ be a double sequence of intervals in \mathbf{R}^2 and (f_{ij}) be a double sequence of functions in $S(\mathbf{R}^2, l^Q)$. Then

$$(1) \quad \|(S_{I_i \times J_j} f_{ij})\|_{L^P(l^Q)} \leq C\|(f_{ij})\|_{L^P(l^Q)}, \quad 1 < P, Q < \infty,$$

where C is independent of $(I_i \times J_j)$ and (f_{ij}) . Moreover, the operator $S(f_{ij}) = (S_{I_i \times J_j} f_{ij})$ can be extended continuously to all $L^P(\mathbf{R}^d)$.

We shall reverse inequality 5.2(1) for $Q = (2, 2)$ and the family of dyadic intervals, i.e. we shall obtain the Littlewood-Paley inequalities for mixed norms. We shall need some preliminaries.

5.3. LEMMA. *Let $\varphi \in S(\mathbf{R})$ be given with $\hat{\varphi}(0) = 0$ and $\hat{\varphi}(t) = 1$ if $t \in [1/2, 1]$. Setting $\varphi_j(x) = 2^j\varphi(2^jx)$, $j \in \mathbf{Z}$, we have*

$$(1) \quad \sum_{j \in \mathbf{Z}} |\hat{\varphi}_j(t)|^2 \leq C;$$

$$(2) \quad \sum_{j \in \mathbf{Z}} |\varphi_j(x)|^2 \leq C|x|^{-2};$$

$$(3) \quad \left(\sum_{j \in \mathbf{Z}} |\varphi_j(x - y) - \varphi_j(x)|^2 \right)^{1/2} \leq C|y|/|x|^2, \quad \text{if } |x| \geq 2|y|.$$

Proof. See [13] or [14].

5.4. THEOREM. *Let φ and ψ be given as in Lemma 5.3. Then*

$$(1) \quad \|(\varphi_i\psi_j * f)_{ij}\|_{L^{p_1}(l^{p_2})} \leq C\|f\|_{L^p}$$

for all $f \in L^p(\mathbf{R}^2)$, with $1 < P = (p_1, p_2) < \infty$.

Proof. We consider the operator

$$(2) \quad T: f \in S(\mathbf{R}^2) \rightarrow Tf = (\varphi_i\psi_j * f)_{ij} \in M(\mathbf{R}^2, l^2).$$

We have to show that T is bounded from L^P into $L^P(l^2)$.

Step 1. T is well defined. Indeed, by 5.3(1), we have

$$\begin{aligned} \int \int \sum_j \sum_i |\varphi_i\psi_j * f|^2 dx dy &= \sum_j \sum_i \int \int |\psi_j\varphi_i * f|^2 dx dy \\ &= \sum_j \sum_i \int \int |\hat{\psi}_j(t)\hat{\varphi}_i(s)\hat{f}(s, t)|^2 ds dt \leq C \int \int |\hat{f}(s, t)|^2 ds dt. \end{aligned}$$

i.e., we have $\sum_{ij} |\psi_j\varphi_i * f(x, y)|^2 < \infty$, a.e., and $Tf(x, y) \in l^2(\mathbf{Z}^2)$.

Step 3. Tf is measurable. Since $l^2(\mathbf{Z}^2)$ is separable it is enough to show that Tf is weakly measurable. But for all $\alpha = (\alpha_{ij}) \in l^2(\mathbf{Z}^2)$ we have

$$\langle Tf(x, y), \alpha \rangle = \sum_{ij} \alpha_{ij} \varphi_i\psi_j * f(x, y)$$

which is obviously measurable.

Step 4. T has a bounded extension from L^2 into $L^2(l^2)$ because (3) holds.

Step 5. The kernel k_φ defined by

$$k_\varphi(x): \lambda \in \mathbf{C} \rightarrow k_\varphi(x) = (\varphi_i(x)\lambda)_i \in l^2(\mathbf{Z})$$

is well defined, belongs to $L^1_{\text{loc}}(\mathbf{R} - \{0\}, L(\mathbf{C}, l^2))$ and verifies Hörmander's condition.

Step 6. The kernel k_ψ defined by

$$k_\psi(y) = (\alpha_i)_i \in l^2(\mathbf{Z}) \rightarrow k_\psi(y)(\alpha_i)_i = (\psi_j(y)\alpha_i)_{ij} \in l^2(\mathbf{Z}^2)$$

is well defined by 5.3(2). On the other hand, the mapping

$$(x, y) \rightarrow \sum_j \sum_i \alpha_{ij} \varphi_i(x) \psi_j(y)$$

is measurable for all $\alpha = (\alpha_{ij}) \in l^2(\mathbf{Z}^2)$. Thus, k_ψ is measurable, belongs to $L^1_{\text{loc}}(\mathbf{R} - \{0\}, L(l^2(\mathbf{Z}), l^2(\mathbf{Z}^2)))$, and satisfies Hörmander's condition.

Step 7. The above results clearly also hold for the cut operators T_{mn} and the respective kernels k_φ^m and k_ψ^n . Thus, since $T_{mn}f = (\psi_j \varphi_i * f; 1 \leq i \leq m, 1 \leq j \leq n)$ is a sequence in l^2 we have

$$T_{mn}f(x, y) = \int \int k_\psi^n(y - v) k_\varphi^m(x - u) f(u, v) du dv.$$

Step 8. The operators T_{mn} are bounded from $L^p(\mathbf{R}^2)$ into $L^p(\mathbf{R}^2, l^2(\mathbf{Z}^2))$, with operator norms bounded by a constant independent of m and n .

Step 9. From Theorem 1.2 and Corollary 1.4 we obtain

$$(3) \quad \|T_{mn}f\|_{L^p(l^2)} \leq C \|f\|_{L^p},$$

with the constant C independent of m and n .

Step 10. The monotone convergence theorem applied to (3) yields (1), as desired.

5.5. REMARK. For a related result, but with a different proof, of Theorem 5.4 see Bordin-Fernandez [3].

5.6. Let Δ_1 be the set of all dyadic intervals in \mathbf{R} , and let $\Delta = \Delta_1 \times \Delta_1$ be the set of dyadic bi-dimensional intervals.

5.7. THEOREM. *If $f \in L^P(\mathbf{R}^2)$, $1 < P = (p_1, p_2) < \infty$, then*

$$(1) \quad c_P \|f\|_{L^P} \leq \left\| \left[\sum_{I \in \Delta} |S_I f|^2 \right]^{1/2} \right\|_{L^P} \leq C_P \|f\|_{L^P}$$

where c_P and C_P are independent of f .

Proof. Let $\varphi \in S(\mathbf{R}^2)$ and let φ and ψ be as in Lemma 5.2. Since $\hat{\varphi}_i(s) = \hat{\varphi}(2^i s)$ and $\hat{\psi}_j(t) = \hat{\psi}(2^j t)$, we have $\hat{\varphi}_i(s) = 1$ if $s \in [2^{i-1}, 2^i]$ and $\hat{\psi}_j(t) = 1$ if $t \in [2^{j-1}, 2^j]$. Hence

$$(2) \quad S_I f = S_{I_1 \times I_2} f = S_{I_1 \times I_2} (\varphi_i \psi_j * f).$$

Now, Theorems 5.2 and 5.4 yield

$$(3) \quad \left\| \left[\sum_{I \in \Delta} |S_I f|^2 \right]^{1/2} \right\|_{L^P} \leq C \|f\|_{L^P}.$$

Finally, to reverse (3) we use polarization and duality as in the well known cases (see [16], [14] and [18]).

6. Multiplier theorems.

6.1. DEFINITION. A scalar valued measurable function φ , defined in $\mathbf{R} \times \mathbf{R}$, is said to be of *bounded V-variation* if there exists a positive constant M and consequences $(C_{km}; k \in \mathbf{Z}, m \in \mathbf{N})$, $(a_{km}; k \in \mathbf{Z}, m \in \mathbf{N})$ and $(b_{km}; k \in \mathbf{Z}, m \in \mathbf{N})$, which satisfy

$$(1) \quad \lim_{m \rightarrow \infty} \sum_{k \in \mathbf{Z}} C_{km} \chi_{(-\infty, a_{km})} \chi_{(-\infty, b_{km})} = \varphi, \quad \text{a.e.}$$

and

$$(2) \quad \sum_k |C_{km}| \leq M, \quad \text{for all } m.$$

We shall write $V(\varphi)$ for the infimum of such constants M .

6.2. THEOREM. *Let (φ_{mn}) be a (double-)sequence of uniformly bounded V-variations, i.e.*

$$(1) \quad V(\varphi_{mn}) \leq M, \quad \text{for all } m \text{ and } n.$$

For $g \in S(\mathbf{R}^2)$, let T_{mn} denote the operator defined by

$$(2) \quad (T_{mn} g)^\wedge = \varphi_{mn} \hat{g}.$$

Then, if (f_{mn}) is a sequence in $S(\mathbf{R}^2)$, we have

$$(3) \quad \|(T_{mn}f_{mn})\|_{L^p(I^Q)} \leq C\|(f_{mn})\|_{L^p(I^Q)}.$$

Proof. Let us suppose $f_{mn} \equiv 0$, for m and/or n large. Let us set $N = (m, n)$, and let

$$h_{mN} = \sum_k C_{kmN} \chi_{I_{kmN}} \quad (I_{kmN} = (-\infty, a_{km_i}) \times (-\infty, b_{km_j}))$$

be a function which satisfies 6.1(1)–(2) and $h_{mN} \rightarrow \varphi_N$, as $m \rightarrow \infty$. Next, define $(S_{mN}f_N)^\wedge = h_{mN}\hat{f}_N$. We claim that

$$(4) \quad \|(S_{mN}f_N)\|_{L^p(I^Q)} \leq C\|(f_N)\|_{L^p(I^Q)}.$$

In fact, by Hölder’s inequality

$$(5) \quad |S_{mN}f_N|^{q_1} = \left| \sum_k C_{kmN} (\chi_{I_{kmN}} \hat{f}_N)^\vee \right|^{q_1} \\ \leq M \sum_k |C_{kmN}| \left| (\chi_{I_{kmN}} \hat{f}_N)^\vee \right|^{q_1};$$

hence, recalling 5.2(1) and hypothesis (1), we have

$$\|(S_{mN}f_N)\|_{L^p(I^Q)} = \left\| \left[\sum_j \left[\sum_i |S_{mN}f_N|^{q_1} \right]^{q_2/q_1} \right]^{1/q_2} \right\|_{L^p} \quad (N = (i, j)) \\ \leq C \left\| \left[\sum_j \left[\sum_{isk} |C_{kmN}| \left| (\chi_{I_{kmN}} \hat{f}_N)^\vee \right|^{q_1} \right]^{q_2/q_1} \right]^{1/q_2} \right\|_{L^p} \\ \leq C \left\| \left[\sum_j \left[\sum_{ik} \left(|\chi_{I_{kmN}}| C_{kmN} \right)^{1/q_1} \hat{f}_N \right]^{q_1} \right]^{q_2/q_1} \right]^{1/q_2} \right\|_{L^p} \\ \leq C \left\| \left[\sum_j \left[\sum_{ik} |C_{kmN}| |f_N|^{q_1} \right]^{q_2/q_1} \right]^{1/q_2} \right\|_{L^p} \\ \leq C \left\| \left[\sum_j \left[\sum_i \left(\sum_k |C_{kmN}| |f_N|^{q_1} \right) \right]^{q_2/q_1} \right]^{1/q_2} \right\|_{L^p} \\ \leq C\|(f_N)\|_{L^p(I^Q)}.$$

Finally, by an application of the Lebesgue dominated theorem and Fatou’s lemma, from (4) we get (3) as desired.

The following lemma is well known (see [5, Th. 4.2–3 and 5]) and will play a major role in the multiplier theorem we shall state.

6.3. LEMMA. Let m be a bounded measurable function which has continuous derivatives of order (α, β) , $\alpha = 0, 1$ and $\beta = 0, 1$, away from the axis, and satisfies

$$(1) \quad |x^\alpha y^\beta D^{\alpha\beta} m(x, y)| \leq M, \quad x \neq 0, y \neq 0.$$

Then, the V -variation of the restriction of m to the dyadic intervals are uniformly bounded, i.e.

$$(2) \quad V(\chi_K m) \leq N,$$

for all dyadic intervals $K = I \times J$ in \mathbf{R}^2 .

Finally, as a consequence of the foregoing results we obtain a multiplier theorem of Lizorkin type.

6.4. THEOREM. Let m be a scalar-valued function in \mathbf{R}^2 given as in Lemma 6.3. Let T_m be the multiplier operator defined on $\varphi \in S(\mathbf{R}^2)$ by

$$(1) \quad (T_m \varphi)^\wedge = m \hat{\varphi}.$$

Then, T_m has an extension to all $L^p(\mathbf{R}^2)$ such that

$$(2) \quad \|T_m f\|_{L^p} \leq C \|f\|_{L^p},$$

for all $f \in L^p(\mathbf{R}^2)$, where the constant C depends on p only.

Proof. By the Littlewood-Paley inequalities, Theorem 6.2 and Lemma 6.3 we shall have

$$\begin{aligned} \|T_m f\|_{L^p} &\leq C \left\| \left[\sum_{K \in \Delta_2} |S_K(T_m f)|^2 \right]^{1/2} \right\|_{L^p} \\ &= C \left\| \left[\sum_{K \in \Delta_2} |(\chi_K m \chi_K \hat{f})^\vee|^2 \right]^{1/2} \right\|_{L^p} \\ &\leq C \left\| \left[\sum_{K \in \Delta_2} |(\chi_K \hat{f})^\vee|^2 \right]^{1/2} \right\|_{L^p} \leq C \|f\|_{L^p}. \end{aligned}$$

REFERENCES

- [1] A. Benedek and R. Panzone, *The space L^p , with mixed norm*, Duke Math. J., **28** (1961), 301–324.
- [2] A. Benedek, A. P. Calderón and R. Panzone, *Convolution operators on Banach space valued functions*, Proc. Nat. Acad. Sci., **48** (1962), 356–365.
- [3] B. Bordin and D. L. Fernandez, *On a Littlewood-Paley type theorem*, Ann. Real Acad. Ci. Madrid, (to appear).

- [4] J. Bourgain, *Some remarks on Banach spaces in which martingale difference sequences are unconditional*, Ark. Mat., **21** (1983), 163–168.
- [5] D. Burkholder, *A geometric characterization that implies the existence of certain singular integrals of Banach spaces-valued functions*, Proc. Conf. Harmonic analysis in Honor of A. Zygmund, Chicago, 1982, Wadsworth, Belmont, 1983.
- [6] C. Fefferman and E. M. Stein, *Some maximal inequalities*, Amer. J. Math., **83** (1971), 107–115.
- [7] ———, *H^p -spaces of several variables*, Acta Math., **129** (1972), 137–193.
- [8] D. L. Fernandez, *Vector valued convolution operators and maximal inequalities for L^p spaces with mixed norms*, Proc. 19^o. Sem. Bras. Analise, S. J. Campos (SP) (1984), 247–268.
- [9] E. Hernández, Personal communication (1985).
- [10] W. Littman, C. McCarthy and N. Riviere, *L^p -multiplier theorems*, Studia Math., **30** (1968), 193–217.
- [11] P. I. Lizorkin, *Multipliers of Fourier integrals and bounds of convolution in spaces with mixed norm*, Applications Math. Ivestia, **4** (1970), 225–255.
- [12] J. Peetre, *On spaces of Triebel-Lizorkin type*, Arkiv för Mat., **14** (1975), 123–130.
- [13] J. L. Rubio de Francia, F. J. Ruiz and J. L. Torrea, *Les operateurs de Calderón-Zygmund Vectoriels*, C.R.A. Sc. Paris, Sér. I, **297** (1983), 477–480.
- [14] ———, *Calderón-Zygmund theory for operator-valued kernels*, Advances in Math., **62** (1986), 7–48.
- [15] H. J. Schmeisser, *Maximal inequalities and Fourier multipliers for spaces with mixed quasi-norms*, Applications Z. für Anal. und ihre Anwendungen, **3** (1984), 153–166.
- [16] E. M. Stein, *Singular Integrals and Differentiability Properties of Functions*, Princeton Univ. Press, 1970. Princeton, N. J.
- [17] B. Stöckert, *Ungleichungen von Plancherel-Polya-Nikol'skij typ in gewichteten L_p^{Ω} -Räumen mit gemischten Norm*, Math. Nach., **86** (1978), 19–32.
- [18] J. L. Torrea, *Integrales Singulares Vectoriales*, Notas de Algebra y Analysis No. 12. Bahia Blanca, 1984.
- [19] F. Zó, *A note on the approximation of the identity*, Studia Math., **55** (1976), 111–122.

Received December 26, 1985. This work was partially supported by a grant of FAPESP (Proc. 84/1686-5) and part of it was done when the author was a visiting fellow of the Groupe d'Analyse Harmonique d'Orsay–Université de Paris Sud (Nov. 84–Mar. 85).

UNIVERSIDADE ESTADUAL DE CAMPINAS
CAIXA POSTAL 6065
13.100-CAMPINAS-SP-BRASIL

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720
C. C. MOORE
University of California
Berkeley, CA 94720
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906-1982)

SUPPORTING INSTITUTIONS

| | |
|------------------------------------|-----------------------------------|
| UNIVERSITY OF ARIZONA | UNIVERSITY OF OREGON |
| UNIVERSITY OF BRITISH COLUMBIA | UNIVERSITY OF SOUTHERN CALIFORNIA |
| CALIFORNIA INSTITUTE OF TECHNOLOGY | STANFORD UNIVERSITY |
| UNIVERSITY OF CALIFORNIA | UNIVERSITY OF HAWAII |
| MONTANA STATE UNIVERSITY | UNIVERSITY OF TOKYO |
| UNIVERSITY OF NEVADA, RENO | UNIVERSITY OF UTAH |
| NEW MEXICO STATE UNIVERSITY | WASHINGTON STATE UNIVERSITY |
| OREGON STATE UNIVERSITY | UNIVERSITY OF WASHINGTON |

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1987 by Pacific Journal of Mathematics

| | |
|--|-----|
| Pere Ara , Matrix rings over $*$ -regular rings and pseudo-rank functions | 209 |
| Lindsay Nathan Childs , Representing classes in the Brauer group of quadratic number rings as smash products | 243 |
| Dicesar Lass Fernandez , Vector-valued singular integral operators on L^p -spaces with mixed norms and applications | 257 |
| Louis M. Friedler, Harold W. Martin and Scott Warner Williams , Paracompact C -scattered spaces | 277 |
| Daciberg Lima Gonçalves , Fixed points of S^1 -fibrations | 297 |
| Adolf J. Hildebrand , The divisor function at consecutive integers | 307 |
| George Alan Jennings , Lines having contact four with a projective hypersurface | 321 |
| Tze-Beng Ng , 4-fields on $(4k + 2)$ -dimensional manifolds | 337 |
| Mei-Chi Shaw , Eigenfunctions of the nonlinear equation $\Delta u + \nu f(x, u) = 0$ in R^2 | 349 |
| Roman Svirsky , Maximally resonant potentials subject to p -norm constraints | 357 |
| Lowell G. Sweet and James A. MacDougall , Four-dimensional homogeneous algebras | 375 |
| William Douglas Withers , Analysis of invariant measures in dynamical systems by Hausdorff measure | 385 |