

Pacific Journal of Mathematics

**WEIGHTED NORM INEQUALITIES FOR THE FOURIER
TRANSFORM ON CONNECTED LOCALLY COMPACT GROUPS**

JEFFREY A. HOGAN

WEIGHTED NORM INEQUALITIES FOR THE FOURIER TRANSFORM ON CONNECTED LOCALLY COMPACT GROUPS

JEFFREY A. HOGAN

Let G be a locally compact connected group. If G is also either compact or abelian, sufficient conditions on a non-negative pair of measurable functions T and V are given to imply that there exists a constant c independent of f for which an inequality of the form

$$\left(\int_{\Gamma} |\hat{f}(\gamma)|^{p'} T(\gamma) dm_{\Gamma}(\gamma) \right)^{1/p'} \leq c \left(\int_G |f(x)|^p V(x) dm_G(x) \right)^{1/p}$$

holds for all integrable f on G ($1 < p \leq 2, p' = p/(p-1)$). Here, \hat{f} denotes the Fourier transform of f defined on Γ (the dual of G) and m_G, m_{Γ} are Haar measures on G, Γ respectively. Conditions on T, V are also given to ensure that the inequality holds with p' replaced by a more general exponent on a more restricted class of groups.

1. Introduction. In [6], sufficient conditions are placed on non-negative pairs of functions T, V to imply

$$\left(\int_{\mathbf{R}^n} |\hat{f}(x)|^q T(x) dx \right)^{1/q} \leq c \left(\int_{\mathbf{R}^n} |f(x)|^p V(x) dx \right)^{1/p}$$

for all integrable functions f on \mathbf{R}^n , with c a constant independent of f . Our aims here are to first make sense of such inequalities in a more abstract setting (in particular, for functions defined on a fairly broad class of locally compact groups G and their Fourier transforms defined on the suitable dual object Γ with m_G, m_{Γ} being the appropriate measures on G and Γ respectively), and then to produce conditions on T, V analogous to those found in [6] so that the inequality will be valid. In the case $1 < p \leq 2$ and $q = p' = p/(p-1)$, the main technical difficulty is the proof of the existence of a measurable function $W: G \rightarrow \mathbf{R}^+ (= [0, \infty))$ with the properties

$$(1.1) \quad \begin{aligned} V(x) &\leq 2W(x) < 2V(x) && \text{for all } x \in G \text{ and} \\ m_G\{x \in G; W(x) = \alpha\} &= 0 && \text{for all } \alpha > 0. \end{aligned}$$

In the case $q \neq p'$ however, it is necessary to have a measurable function $S: \Gamma \rightarrow \mathbf{R}^+$ satisfying

$$(1.2) \quad \begin{aligned} T(\gamma) &\leq 2S(\gamma) < 2T(\gamma) && \text{for all } \gamma \in \Gamma \text{ and} \\ m_{\Gamma}\{\gamma \in \Gamma; S(\gamma) = \alpha\} &= 0 && \text{for all } \alpha > 0. \end{aligned}$$

Let G be a locally compact abelian (LCA) group. Let Γ denote the dual group of G . Equip G and Γ with Haar measures m_G and m_Γ normalized so that the Plancherel identity is valid. If $\gamma \in \Gamma$ and $f \in L^1(G)$, the Fourier transform of f is defined by

$$\hat{f}(\gamma) = \int_G f(x)\overline{\gamma(x)} dm_G(x).$$

The Hausdorff-Young theorem then states that if $1 \leq p \leq 2$ and $f \in L^p(G)$, then $\hat{f} \in L^{p'}(\Gamma)$ and

$$\left(\int_\Gamma |\hat{f}(\gamma)|^{p'} dm_\Gamma(\gamma) \right)^{1/p'} \leq \left(\int_G |f(x)|^p dm_G(x) \right)^{1/p}.$$

Now let G be any group. A representation of G is a homomorphism $U: G \rightarrow \text{GL}(\mathcal{H}_U)$ where $\text{GL}(\mathcal{H}_U)$ is the group of invertible endomorphisms of some finite-dimensional linear space \mathcal{H}_U . \mathcal{H}_U is the representation space of U and the dimension of \mathcal{H}_U is defined to be the dimension of the representation U ($= d(U)$).

A unitary representation of G is a homomorphism $U: G \rightarrow \mathcal{U}(\mathcal{H}_U)$ where $\mathcal{U}(\mathcal{H}_U)$ is the group of unitary endomorphisms of some finite-dimensional Hilbert space \mathcal{H}_U .

If G is compact, let \hat{G} denote a maximal set of pairwise inequivalent continuous unitary representations of G . If $f \in L^1(G)$ and $U \in \hat{G}$, define $\hat{f}(U) \in \text{End}(\mathcal{H}_U)$ by

$$\hat{f}(U) = \int_G f(x)U(x) dm_G(x).$$

Let $E(\hat{G}) = \prod_{U \in \hat{G}} \text{End}(\mathcal{H}_U)$. If $\psi \in E(\hat{G})$, define

$$\|\psi\|_p = \left\{ \sum_{U \in \hat{G}} d(U)\|\psi(U)\|_{\phi_p}^p \right\}^{1/p} \quad \text{if } p \in [1, \infty),$$

$$\|\psi\|_\infty = \sup_{U \in \hat{G}} \|\psi(U)\|_{\phi_\infty},$$

where $\|\cdot\|_{\phi_p}$ denotes the p th von Neumann norm on $\text{End}(\mathcal{H}_U)$ ([5] Sec. 28.34). Now let $E^p(\hat{G})$ be the linear space of $\psi \in E(\hat{G})$ for which $\|\psi\|_p < \infty$. In this context, the Hausdorff-Young theorem states that if $1 \leq p \leq 2$ and $f \in L^p(G)$ then $\hat{f} \in E^{p'}(\hat{G})$ and

$$\left(\sum_{U \in \hat{G}} d(U)\|\hat{f}(U)\|_{\phi_{p'}}^{p'} \right)^{1/p'} \leq \left(\int_G |f(x)|^p dm_G(x) \right)^{1/p}.$$

The sufficiency conditions for the case $q = p'$ are as follows.

THEOREM 1.1. *Let G be an LCA, non-compact, connected group with dual group Γ . Let T be a non-negative, measurable function on Γ and V a non-negative, measurable function on G . Suppose $1 < p \leq 2$ and there exist positive constants A, B such that*

$$(1.3) \quad \left(\int_{T(\gamma) > Br} T(\gamma) \, dm_{\Gamma}(\gamma) \right) \left(\int_{V(x) < r^{p-1}} V(x)^{-1/(p-1)} \, dm_G(x) \right) \leq A$$

for all $r > 0$. Then, for every integrable function f on G ,

$$(1.4) \quad \left(\int_{\Gamma} |\hat{f}(\gamma)|^{p'} T(\gamma) \, dm_{\Gamma}(\gamma) \right)^{1/p'} \leq c \left(\int_G |f(x)|^p V(x) \, dm_G(x) \right)^{1/p}$$

with c dependent on A, B, p but not on f .

If G is compact and abelian, Γ is discrete and m_{Γ} is just the counting measure.

THEOREM 1.2. *Let G be a compact, connected, abelian group with dual group Γ . Let T be a non-negative function on Γ and V a non-negative, measurable function on G . Suppose $1 < p \leq 2$ and there exist positive constants A, B such that*

$$(1.5) \quad \left(\sum_{T(\gamma) > Br} T(\gamma) \right) \left(\int_{V(x) < r^{p-1}} V(x)^{-1/(p-1)} \, dm_G(x) \right) \leq A$$

for all $r > 0$. Then for every integrable function f on G ,

$$(1.6) \quad \left(\sum_{\gamma \in \Gamma} |\hat{f}(\gamma)|^{p'} T(\gamma) \right)^{1/p'} \leq c \left(\int_G |f(x)|^p V(x) \, dm_G(x) \right)^{1/p}$$

with c dependent on A, B, p but not on f .

If G is compact and non-abelian, the statement of the theorem is slightly more complicated.

THEOREM 1.3. *Let G be a compact, connected, non-abelian group with dual object \hat{G} . Let T be a non-negative function on \hat{G} and V a non-negative, measurable function on G . Suppose $1 < p \leq 2$ and there exist positive constants A, B such that*

$$(1.7) \quad \left(\sum_{T(U) > Br} T(U) d(U)^2 \right) \left(\int_{V(x) < r^{p-1}} V(x)^{-1/(p-1)} \, dm_G(x) \right) \leq A$$

for all $r > 0$. Then for every integrable function f on G ,

$$(1.8) \quad \left(\sum_{U \in \hat{G}} \|\hat{f}(U)\|_{\phi_r}^{p'} T(U) d(U) \right)^{1/p'} \leq c \left(\int_G |f(x)|^p V(x) dm_G(x) \right)^{1/p}$$

with c dependent on A, B, p but not on f .

Note that the statement of Theorem 1.3 collapses to that of Theorem 1.2 if G is abelian.

The author wishes to express his appreciation to John Price and Michael Cowling of the University of New South Wales for many helpful conversations.

2. Non-compact abelian case. To prove the theorems, several lemmas are required leading to the existence of a function W on G satisfying (1.1).

LEMMA 2.1 (Price, Hogan). *Let (X, m_X) be a σ -finite measure space. Let V be a non-negative, measurable function on X which is greater than 0 m_X -a.e. For each $r > 0$, let $B_r = \{x \in X; V(x) = r\}$ and $R = \{r > 0; m_X(B_r) > 0\}$. Then R is countable.*

Proof. $X = \bigcup_{n \in \mathbf{Z}} K_n, m_X(K_n) < \infty$ for each $n \in \mathbf{Z}$. Let $R_n = \{r \in R; m_X(B_r \cap K_n) > 0\}$ and $R_n^m = \{r \in R; m_X(B_r \cap K_n) > 1/m\}$ for $m \in \mathbf{Z}^+ (= \{n \in \mathbf{Z}; n > 0\})$. Then $m_X(B_r \cap K_n) \leq m_X(K_n) < \infty$ for $n \in \mathbf{Z}$. So $\#R_n^m (= \text{cardinality of } R_n^m) < \infty$. But $R_n = \bigcup_{m=1}^\infty R_n^m$, hence R_n is countable. Let $r \in R, 0 < m_X(B_r) \leq \sum_{n \in \mathbf{Z}} m_X(B_r \cap K_n)$. Therefore, $r \in R_n$ for some $n \in \mathbf{Z}$, i.e. $R \subseteq \bigcup_{n \in \mathbf{Z}} R_n$ and hence, R is countable. \square

The utility of this lemma lies in the observation that if G is a connected LCA group then by Theorem 2.4.1 of [8],

$$(2.1) \quad G \simeq H \times \mathbf{R}^n \quad (n \geq 0)$$

with H compact and abelian. (Here \simeq denotes a topological isomorphism.) Of course, if G is non-compact then $n > 0$ in (2.1). G is then σ -compact. The regularity of the measure m_G implies the σ -finiteness of the measure space (G, m_G) , so Lemma 2.1 applies to G .

LEMMA 2.2. *If G is a connected, non-compact, LCA group and V is a non-negative, measurable function on G which is greater than 0 m_G -a.e., then there exists a measurable function W on G satisfying (1.1).*

Proof. $G \simeq H \times \mathbf{R}^n$ with H compact and $n \geq 1$. Let m_H denote the unique Haar measure on H satisfying $m_H(H) = 1$. Let m_n denote

ordinary Lebesgue measure on \mathbf{R}^n . Then $m_G = m_H \times m_n$ is a Haar measure on G . (Any other Haar measure on G will be just a constant multiple of this one, so sets of measure zero are independent of the choice involved.) Denote the Euclidean norm on \mathbf{R}^n by $\| \cdot \|$. Let $B_r = \{(h, x) \in G; V(h, x) = r\}$ ($r > 0$), a Borel subset of G . Let $E = \bigcup_{m_G(B_r) > 0} B_r$. By Lemma 2.1, E is also a Borel set (being the countable union of Borel sets). Let $S_r(y)$ be the sphere in \mathbf{R}^n with centre y , radius r . Let $G(h, x) = \|x\|$ and $g(h, x) = \sin\|x\|$.

$$\begin{aligned} G(h_1, x_1) = G(h_2, x_2) &\Leftrightarrow \|x_1\| = \|x_2\| \quad (= r \text{ say}) \\ &\Leftrightarrow (h_1, x_1), (h_2, x_2) \in H \times S_r(0). \end{aligned}$$

Therefore,

$$\begin{aligned} m_G\{(h, x) \in G; G(h, x) = r\} &= m_G(H \times S_r(0)) \\ &= m_H(H)m_n(S_r(0)) = 0 \quad \text{for all } r > 0. \end{aligned}$$

Now, $\{(h, x) \in G; g(h, x) = \alpha\}$ is either empty (if $|\alpha| > 1$), or a countable union of sets of the form $H \times S_r(0)$, so $m_G\{(h, x) \in G; g(h, x) = \alpha\} = 0$ for all $\alpha > 0$. Let

$$W(h, x) = \begin{cases} 3V(h, x)/4 & \text{on } G \setminus E \\ 3r/4 + rg(h, x)/8 & \text{on } B_r. \end{cases}$$

W satisfies (1.1) and the measurability of V and continuity of g imply the measurability of W . □

In what follows, χ_E denotes the characteristic function of a measurable set E and we adopt the convention $0 \cdot \infty = 0$.

Proof of Theorem 1.1. If $V(x) = 0$ on a set of positive measure then (1.3) implies $T(\gamma) = 0$ m_{Γ} -a.e., so that (1.4) is true trivially. If $m_G\{x \in G; V(x) = \infty \text{ and } f(x) \neq 0\} > 0$, then the right side of (1.4) is infinite so that (1.2) is true trivially. From now on we assume $V(x) > 0$ m_G -a.e. and $m_G\{x \in G; V(x) = \infty \text{ and } f(x) \neq 0\} = 0$. Let

$$(2.2) \quad I = \int_{\Gamma} |\hat{f}(\gamma)|^{p'} T(\gamma) dm_{\Gamma}(\gamma) \leq 2^{p'}(I_1 + I_2)$$

where

$$I_1 = \sum_{j=-\infty}^{\infty} \int_{2^j B < T(\gamma) \leq 2^{j+1} B} T(\gamma) \left| \int_{V(x) \geq 2^{j-p'-1}} f(x) \overline{\gamma(x)} dm_G(x) \right|^{p'} dm_{\Gamma}(\gamma)$$

and

$$I_2 = \sum_{j=-\infty}^{\infty} \int_{2^j B < T(\gamma) \leq 2^{j+1} B} T(\gamma) \left| \int_{V(x) < 2^{jp-j-1}} f(x) \overline{\gamma(x)} dm_G(x) \right|^{p'} dm_{\Gamma}(\gamma).$$

$$I_1 \leq 2B \sum_{j=-\infty}^{\infty} 2^j \int_{\Gamma} |(f \chi_{V \geq 2^{jp-j-1}})^{\wedge}(\gamma)|^{p'} dm_{\Gamma}(\gamma)$$

$$\leq 2B \sum_{j=-\infty}^{\infty} 2^j \left(\int_G |f(x)|^p \chi_{V \geq 2^{jp-j-1}}(x) dm_G(x) \right)^{p'/p}$$

(by the Hausdorff-Young inequality)

$$\leq 2B \left(\int_G |f(x)|^p h(x) dm_G(x) \right)^{p'/p}$$

(since $p'/p \geq 1$) where $h(x) = \sum_{j=-\infty}^{\infty} 2^{jp/p'} \chi_{V \geq 2^{jp-j-1}}(x)$. For any x , $h(x) = \sum_{j=-\infty}^J 2^{jp/p'}$ where J is the largest integer satisfying $V(x) \geq 2^{Jp-J-1}$. Estimating the geometric series defining h , we see $h(x) \leq 2V(x)/(1 - 2^{1-p})$. Therefore, there exists a constant c such that

$$(2.3) \quad I_1 \leq c \left(\int_G |f(x)|^p V(x) dm_G(x) \right)^{p'/p}.$$

Now, $I_2 \leq \int_{\Gamma} T(\gamma) \left(\int_{2V(x) < (T(\gamma)/B)^{p-1}} |f(x)| dm_G(x) \right)^{p'} dm_{\Gamma}(\gamma)$. By Lemma 2.2, there exists a measurable function W satisfying (1.1). Let J be the least integer such that $2^J \geq \|f\|_1$. Let $r_j = \infty$ and for $j < J$, choose r_j such that

$$3 \cdot 2^{j-2} < \int_{2W(x) < r_j^{p-1}} |f(x)| dm_G(x) \leq 2^j.$$

$$I_2 \leq \sum_{j=-\infty}^J \int_{Br_{j-1} < T(\gamma) \leq Br_j} T(\gamma) \left(\int_{2W(x) < r_j^{p-1}} |f(x)| dm_G(x) \right)^{p'} dm_{\Gamma}(\gamma).$$

Let

$$D_j = \{x \in G; r_{j-2}^{p-1} \leq 2W(x) < r_{j-1}^{p-1}\}.$$

Then

$$\int_{2W(x) < r_{j-1}^{p-1}} |f(x)| dm_G(x) \leq 8 \int_{D_j} |f(x)| dm_G(x) \quad \text{and}$$

$$I_2 \leq 8^{p'} \sum_{j=-\infty}^J \int_{Br_{j-1} < T(\gamma) \leq Br_j} T(\gamma) \left(\int_{D_j} |f(x)| dm_G(x) \right)^{p'} dm_{\Gamma}(\gamma).$$

Since $V(x) > 0$ m_G -a.e. and $f = 0$ m_G -a.e. on the set where $V = \infty$, we

may write

$$|f(x)| = (|f(x)|V(x)^{1/p})V(x)^{-1/p}$$

and apply Hölder's inequality on D_j to get

$$I_2 \leq 8^{p'} \sum_{j=-\infty}^J \int_{B_{r_{j-1}} < T(\gamma)} T(\gamma) dm_{\Gamma}(\gamma) \left(\int_{D_j} V(x)^{-p'/p} dm_G(x) \right) \\ \times \left(\int_{D_j} |f(x)|^p V(x) dm_G(x) \right)^{p'/p}.$$

However, $D_j \subseteq \{x \in G; V(x) < r_{j-1}^{p-1}\}$, so

$$(2.4) \quad I_2 \leq 8^{p'} \sum_{j=-\infty}^J \int_{B_{r_{j-1}} < T(\gamma)} T(\gamma) dm_{\Gamma}(\gamma) \left(\int_{V(x) < r_{j-1}^{p-1}} V(x)^{-p'/p} dm_G(x) \right) \\ \times \left(\int_{D_j} |f(x)|^p V(x) dm_G(x) \right)^{p'/p} \\ \leq 8^{p'} A \left(\sum_{j=-\infty}^J \int_{D_j} |f(x)|^p V(x) dm_G(x) \right)^{p'/p} \\ \text{(by the condition (1.3) and the fact that } p'/p \geq 1) \\ = 8^{p'} A \left(\int_G |f(x)|^p V(x) dm_G(x) \right)^{p'/p}.$$

Combining (2.2), (2.3) and (2.4) gives (1.4). \square

3. Compact abelian case. The decomposition (2.1) is trivial if G is compact in the sense that no \mathbf{R}^n factor appears. A different argument is required.

LEMMA 3.1. *If G is a compact, connected, abelian group with Haar measure m_G and V is greater than 0 m_G -a.e., then there exists a measurable function W on G satisfying (1.1).*

Proof. Choose a non-trivial $\gamma \in \Gamma$ and let $\ker \gamma = H$. H is a proper closed subgroup of G , hence $m_G(H) = 0$ (since G is compact and connected). Now $\gamma(x) = \gamma(y)$ if and only if $xH = yH$ so that γ is constant on cosets of H and assumes different values on different cosets. Let $\phi(x) = \operatorname{Re}(\gamma(x))$, i.e. $\phi(x) = \frac{1}{2}(\gamma(x) + \overline{\gamma(x)})$. If $\phi(x) = \cos \theta$ for

some $\theta \in [0, 2\pi)$,

$$\begin{aligned} m_G\{x \in G; \phi(x) = \cos \theta\} \\ &= m_G\{x \in G; \gamma(x) = \cos \theta + i \sin \theta \text{ or } \cos \theta - i \sin \theta\} \\ &= m_G\{z_1H \cup z_2H\} \quad \text{with } z_1H, z_2H \text{ cosets of } H \text{ in } G \\ &\leq m_G(z_1H) + m_G(z_2H) = 0. \end{aligned}$$

Let $B_r = \{x \in G; V(x) = r\}$ ($r > 0$) and $E = \bigcup_{m_G(B_r) > 0} B_r$. E and each B_r are measurable. Let

$$W(x) = \begin{cases} 3V(x)/4 & \text{on } G \setminus E \\ 3r/4 + r\phi(x)/8 & \text{on } B_r. \end{cases}$$

W satisfies (1.1) and the measurability of V and continuity of ϕ imply the measurability of W . □

Proof of Theorem 1.2. This is exactly as for Theorem 1.1 except that the measure on Γ is now counting measure so that all integrals over Γ (and its subsets) are replaced by the corresponding sums.

4. Compact non-abelian case. If G is compact and non-abelian, there is a further complication due to the fact that the representations of G are not necessarily one-dimensional, so that an argument like that of Lemma 3.1 will not hold. We rely instead on a decomposition of G involving Lie groups.

LEMMA 4.1. *Let G be a compact, connected Lie group and d the invariant Riemannian metric on G . For each $y \in G$, let $S_r(y) = \{x \in G; d(x, y) = r\}$ ($r \geq 0$). Then $m_G(S_r(y)) = 0$.*

Proof. Omitted.

LEMMA 4.2. *If G is a compact, connected, non-abelian group with Haar measure m_G and V is a non-negative, measurable function on G which is greater than 0 m_G -a.e., then there exists a measurable function W on G satisfying (1.1).*

Proof. If G is compact and connected, then

$$G \simeq \left(T \times \prod_{i \in I} G_i \right) / H$$

where T is a compact, connected, abelian group, $\{G_i; i \in I\}$ is a family of compact, simply connected, simple Lie groups and H is a totally disconnected, closed subgroup of $Z(T \times \prod_{i \in I} G_i)$ (the centre of $T \times \prod_{i \in I} G_i$). For a discussion of this decomposition, see §6.5.6 of [7].

G is non-abelian, so I must be non-empty and at least one G_i must be non-trivial. Let G_1 be non-trivial, $T_1 = T \times \prod_{i \in I, i \neq 1} G_i$ and $G_2 = G_1 \times T_1$. G_1 , being a connected, compact Lie group, possesses an invariant Riemannian metric d yielding the topology.

Define an invariant pseudo-metric \bar{d} on G_2 by

$$\bar{d}(x, y) = d(x_1, y_1)$$

where x_1, y_1 are the G_1 -components of $x, y (\in G_2)$ respectively. Define $D: G \times G \rightarrow \mathbf{R}^+$ by

$$D(xH, yH) = \inf_{a, b \in H} \bar{d}(xa, yb).$$

In particular, $D(yH, H) = \inf_{a \in H} d(y_1, a_1)$ (by the invariance of d). Now $H \subseteq Z(G_1 \times T_1) \simeq Z(G_1) \times Z(T_1)$ and $Z(G_1)$ is finite since G_1 is simple. Therefore, the infimum in the definition of $D(yH, H)$ is taken over a finite set (i.e. $d(y_1, a_1)$ takes only finitely many values as a varies over H). So we may write

$$D(yH, H) = \min_{a \in H} d(y_1, a_1).$$

We shall need Theorem C §63 of [3] which implies that if $\pi: G_2 \rightarrow G_2/H = G$ is the canonical homomorphism then $m_{G_2} \circ \pi^{-1}$ is a Haar measure on G . (There exists a constant $\alpha > 0$ for which $m_G = \alpha m_{G_2} \circ \pi^{-1}$.) So

$$\begin{aligned} & m_G \{ yH \in G; D(yH, H) = \beta \} \\ &= \alpha m_{G_2} \left\{ y \in G_2; \min_{a \in H} d(y_1, a_1) = \beta \right\} \\ &\leq \alpha m_{G_2} \left\{ y \in G_2; d(y_1, a_1) = \beta \text{ for some } a \in H \right\} \\ &\leq \alpha m_{G_2} \left\{ \bigcup_{i=1}^n (S_\beta(a_i^1) \times T_1) \right\} \\ &\quad \left(\text{where } \{ a_1^1, \dots, a_1^n \} \text{ is the finite set consisting} \right. \\ &\quad \left. \text{of the projection of } H \text{ onto } Z(G_1) \right) \\ &\leq \alpha \sum_{i=1}^n m_{G_1}(S_\beta(a_i^1)) m_{T_1}(T_1) = 0 \end{aligned}$$

where the last step follows by Lemma 4.1 and the compactness of T_1 . Let $g(yH) = \sin D(yH, H)$. As before, let $B_r = \{ x \in G; V(x) = r \} (r > 0)$ and $E = \bigcup_{m_G(B_r) > 0} B_r$. E and B_r are measurable. Let

$$W(x) = \begin{cases} 3V(x)/4 & \text{on } G \setminus E \\ 3r/4 + rg(x)/8 & \text{on } B_r. \end{cases}$$

For $1 \leq i \leq n$ define $h_i: G_2 \rightarrow \mathbf{R}^+$ by $h_i(y) = d(y, a_i)$. Each h_i is continuous, hence measurable, and $D(yH, H) = \min_{1 \leq i \leq n} h_i(y)$, so D is measurable as a function on G and hence g is measurable also. The measurability of V and g now implies the measurability of W . Also, W satisfies (1.1). \square

Proof of Theorem 1.3. First note that if U is a representation of G , then $d(U) \geq 1$. Suppose $V(x) = 0$ on a set of positive measure. Then the second integral in (1.7) is infinite for all $r > 0$. Therefore the first integral (sum) in (1.7) must be zero for all $r > 0$ and this implies that $T(U) = 0$ for all $U \in \hat{G}$. If this were the case then (1.8) would be true trivially.

If $m_G\{x \in G; V(x) = \infty \text{ and } f(x) \neq 0\} > 0$, then the second integral in (1.8) would be infinite and once again (1.8) would be true trivially. From now on we assume that $V(x) > 0$ m_G -a.e. and $m_G\{x \in G, f(x) \neq 0 \text{ and } V(x) = \infty\} = 0$. Let

$$(4.1) \quad I = \sum_{U \in \hat{G}} \|\hat{f}(U)\|_{\phi_p}^{p'} T(U) d(U) \leq 2^{p'}(I_1 + I_2)$$

where

$$I_1 = \sum_{j=-\infty}^{\infty} \sum_{2^j B < T(U) \leq 2^{j+1} B} T(U) d(U) \left\| \int_{V(x) \geq 2^{jp-j-1}} f(x)U(x) dm_G(x) \right\|_{\phi_p}^{p'}$$

and

$$I_2 = \sum_{j=-\infty}^{\infty} \sum_{2^j B < T(U) \leq 2^{j+1} B} T(U) d(U) \left\| \int_{V(x) < 2^{jp-j-1}} f(x)U(x) dm_G(x) \right\|_{\phi_p}^{p'}$$

(using the fact that $\|A + B\|_{\phi_p}^{p'} \leq 2^{p'}(\|A\|_{\phi_p}^{p'} + \|B\|_{\phi_p}^{p'})$ for $A, B \in \text{End}(\mathcal{H}_U)$).

$$(4.2) \quad I_1 \leq 2B \sum_{j=-\infty}^{\infty} 2^j \sum_{U \in \hat{G}} \|(f\chi_{V \geq 2^{jp-j-1}})^{\wedge}(U)\|_{\phi_p}^{p'} d(U)$$

$$\leq 2B \sum_{j=-\infty}^{\infty} 2^j \left(\int_G |f(x)|^p \chi_{V \geq 2^{jp-j-1}}(x) dm_G(x) \right)^{p'/p}$$

(by the Hausdorff-Young inequality)

$$\leq c \left(\int_G |f(x)|^p V(x) dm_G(x) \right)^{p'/p}$$

(estimating the integral as in the proof of Theorem 1.1).

$$I_2 \leq \sum_{U \in \hat{G}} T(U) d(U) \left\| \int_{2V(x) < (T(U)/B)^{p-1}} f(x)U(x) dm_G(x) \right\|_{\phi_p}^{p'}$$

By Lemma 4.2, there exists a measurable function W satisfying (1.1). Let J be the least integer satisfying $2^J \geq \|f\|_1$, let $r_j = \infty$ and for $j < J$, choose r_j such that

$$3 \cdot 2^{j-2} < \int_{2W(x) < r_j^{p-1}} |f(x)| dm_G(x) \leq 2^j.$$

Let $D_j = \{x \in G; r_{j-2}^{p-1} \leq 2W(x) < r_{j-1}^{p-1}\}$.

$$\begin{aligned} I_2 &\leq \sum_{j=-\infty}^J \sum_{Br_{j-1} < T(U) \leq Br_j} T(U) d(U) \left\| \int_{2W(x) < r_j^{p-1}} f(x)U(x) dm_G(x) \right\|_{\phi_p}^{p'} \\ &\leq \sum_{j=-\infty}^J \sum_{Br_{j-1} < T(U) \leq Br_j} T(U) d(U) \left(\int_{2W(x) < r_j^{p-1}} |f(x)| \|U(x)\|_{\phi_p} dm_G(x) \right)^{p'} \\ &= \sum_{j=-\infty}^J \sum_{Br_{j-1} < T(U) \leq Br_j} T(U) d(U)^2 \left(\int_{2W(x) < r_j^{p-1}} |f(x)| dm_G(x) \right)^{p'} \\ &\quad \left(\|U(x)\|_{\phi_p} = d(U)^{1/p'} \text{ since } U(x) \in \mathcal{U}(\mathcal{H}_U) \right) \\ &\leq 8^{p'} \sum_{j=-\infty}^J \sum_{Br_{j-1} < T(U) \leq Br_j} T(U) d(U)^2 \left(\int_{D_j} |f(x)| dm_G(x) \right)^{p'}. \end{aligned}$$

We may write $|f(x)| = (|f(x)|V(x)^{1/p})V(x)^{-1/p}$ and apply Hölder's inequality on D_j to get

$$\begin{aligned} I_2 &\leq 8^{p'} \sum_{j=-\infty}^J \sum_{Br_{j-1} < T(U)} T(U) d(U)^2 \left(\int_{D_j} V(x)^{-p'/p} dm_G(x) \right) \\ &\quad \times \left(\int_{D_j} |f(x)|^p V(x) dm_G(x) \right)^{p'/p}. \end{aligned}$$

However $D_j \subseteq \{x \in G; V(x) < r_{j-1}^{p-1}\}$ and combining this with the assumption (1.7) gives

$$\begin{aligned} (4.3) \quad I_2 &\leq 8^{p'} A \sum_{j=-\infty}^J \left(\int_{D_j} |f(x)|^p V(x) dm_G(x) \right)^{p'/p} \\ &\leq 8^{p'} A \left(\int_G |f(x)|^p V(x) dm_G(x) \right)^{p'/p} \end{aligned}$$

(since $p'/p \geq 1$). Combining (4.1), (4.2) and (4.3) gives (1.8). □

The requirement in the above theorems that the group G be connected may be relaxed slightly as follows.

PROPOSITION 4.3. *Suppose G is a topological group with identity component G_0 such that G/G_0 is countable. If either of the conditions*

- (i) G_0 is non-compact LCA
- (ii) G_0 is compact abelian
- (iii) G_0 is compact non-abelian

hold, and V is a non-negative, measurable function on G which is greater than 0 m_G -a.e., then there exists a measurable function W on G satisfying (1.1).

Proof. Choose coset representatives x_j ($j \in \mathbf{Z}^+$) for G/G_0 . Let $x_j G_0$ be a coset of G_0 and define $V_j: G_0 \rightarrow \mathbf{R}^+$ by $V_j(x) = V(x_j x)$ ($x \in G_0$). By either Lemma 2.2, 3.1 or 4.2 (depending on whether (i), (ii) or (iii) is satisfied), construct a measurable function W_j on G_0 with the properties

$$V_j(x) \leq 2W_j(x) < 2V_j(x) \quad \text{for all } x \in G_0 \quad \text{and}$$

$$m_{G_0}\{x \in G_0; W_j(x) = \alpha\} = 0 \quad \text{for all } \alpha > 0.$$

Now let $W^j: x_j G_0 \rightarrow \mathbf{R}^+$ be given by $W^j(x_j x) = W_j(x)$, and $W: G \rightarrow \mathbf{R}^+$ by $W(y) = W^j(y)$ for $y \in x_j G_0$. W is measurable and satisfies (1.1). \square

So if, in Proposition 4.3, G satisfies (i), (ii) or (iii), then the conclusions of Lemmas 2.2, 3.1 or 4.2 (respectively) hold, and hence Theorems 1.1, 1.2 or 1.3 (respectively) will apply to G .

5. General exponents. In this section, a supplementary result is stated which loosens the restrictions on the exponents involved in the inequality at the expense of additional restrictions on the class of groups for which such an inequality may hold.

Given weight functions V on G and T on \hat{G} (the dual object of G), the proof requires the existence of measurable functions W on G and S on \hat{G} satisfying (1.1) and (1.2) respectively. If G is compact, however, (1.2) is clearly impossible since \hat{G} is discrete. The proof will work for

$$(5.1) \quad G \simeq H \times J \times \mathbf{R}^n \quad (n > 0)$$

where H is a countable abelian group and J is a compact abelian metric group. The decomposition (5.1) then implies

$$(5.2) \quad \hat{G} = \Gamma \simeq \hat{H} \times \hat{J} \times \mathbf{R}^n \quad (n > 0)$$

where \hat{H} (the dual group of H) is a compact, abelian, metric group and \hat{J} (the dual group of J) is a countable, abelian group. Hence, $H \times J$ and $\hat{H} \times \hat{J}$ are σ -compact and we may use Lemma 2.1 and the proof of Lemma 2.2 to construct functions W and S satisfying (1.1) and (1.2) respectively. (Note that Lemma 2.2 requires only that $G \simeq K \times \mathbf{R}^n$ with $n > 0$ and K σ -compact.)

If T is a measurable function on a measure space M , then T^* : $[0, \infty] \rightarrow [0, \infty]$ is the non-increasing rearrangement of T , a discussion of which may be found in Chapter 5 §3 of [9].

THEOREM 5.1. *Let G be an LCA group of the form (5.1) with dual group Γ . Suppose V is a non-negative, measurable function on G and T is a non-negative, measurable function on Γ . If $1 < p < 2$, $p \leq q < p'$ and there exist positive constants A, B for which*

$$(5.3) \quad \left(\int_{(tT^*(t))^{p'/q} > Bt} T^*(t) dt \right) \left(\int_{V(x) < r^{p-1}} V(x)^{-1/(p-1)} dm_G(x) \right)^{q/p'} \leq A$$

holds for all $r > 0$, then for every integrable f on G

$$(5.4) \quad \left(\int_{\Gamma} |\hat{f}(\gamma)|^q T(\gamma) dm_{\Gamma}(\gamma) \right)^{1/q} \leq c \left(\int_G |f(x)|^p V(x) dm_G(x) \right)^{1/p}$$

with c independent of f . If $2 < q < \infty$, $q' < p \leq q$ and there exist positive constants A, B for which

$$(5.5) \quad \left(\int_{(tW(t))^{q/p'} > Bt} W(t) dt \right) \left(\int_{rT(\gamma) > 1} T(\gamma) dm_{\Gamma}(\gamma) \right)^{p/q'} \leq A$$

holds for all $r > 0$ (with $W(t) = (V^{-1/(p-1)})^*(t)$), then for every integrable f on G , (5.4) holds with c independent of f .

By considering the case $G = \mathbf{R}^n$, the sufficiency conditions given here may be shown to be not necessary (as in [6]). However, if T and V are assumed to be radial and, as functions of $|x|$, T is non-increasing while V is non-decreasing, then the conditions are also necessary.

REFERENCES

[1] N. E. Aguilera and E. O. Harboure, *On the search for weighted norm inequalities for the Fourier transform*, Pacific J. Math., **104** (1983), 1–14.
 [2] R. E. Edwards, *Integration and Harmonic Analysis on Compact Groups*, Cambridge University Press, Cambridge (1972).
 [3] P. R. Halmos, *Measure Theory*, Van Nostrand, Princeton, New Jersey (1950).
 [4] S. Helgason, *Differential Geometry and Symmetric Spaces*, Academic Press, New York, London (1962, 70).

- [5] E. Hewitt and K. A. Ross, *Abstract Harmonic Analysis I and II*, Springer-Verlag, New York (1963, 70).
- [6] B. Muckenhoupt, *Weighted norm inequalities for the Fourier transform*, Trans. Amer. Math. Soc., **276** (1983), 729–742.
- [7] J. F. Price, *Lie Groups and Compact Groups*, Cambridge University Press, Cambridge (1977).
- [8] W. Rudin, *Fourier Analysis on Groups*, Interscience, New York, London (1967).
- [9] E. Stein and G. Weiss, *Introduction to Fourier Analysis on Euclidean Spaces*, Princeton University Press, Princeton, New Jersey (1971).

Received September 21, 1986. This research was undertaken with the assistance of an Australian Postgraduate Research Award.

UNIVERSITY OF NEW SOUTH WALES
KENSINGTON, N.S.W. 2033 AUSTRALIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1988 by Pacific Journal of Mathematics

Selman Akbulut and Henry Churchill King , Polynomial equations of immersed surfaces	209
Alberto Baidier and Richard C. Churchill , The Campbell-Hausdorff group and a polar decomposition of graded algebra automorphisms	219
Wayne C. Bell and John William Hagood , Separation properties and exact Radon-Nikodým derivatives for bounded finitely additive measures	237
Dennis J. Garity, James P. Henderson and David G. Wright , Menger spaces and inverse limits	249
B. Brent Gordon , Algebraically defined subspaces in the cohomology of a Kuga fiber variety	261
Jeffrey A. Hogan , Weighted norm inequalities for the Fourier transform on connected locally compact groups	277
Guojun Liao , A study of regularity problem of harmonic maps	291
Chin-pi Lu , Modules satisfying ACC on a certain type of colons	303
Kunio Murasugi , Jones polynomials of periodic links	319
Hans Schoutens , Approximation properties for some non-Noetherian local rings	331
Peter Sjögren , Convergence for the square root of the Poisson kernel	361
Alexandru Ion Suciu , The oriented homotopy type of spun 3-manifolds	393