

Pacific Journal of Mathematics

**ON HOMOMORPHISMS OF MATRIX ALGEBRAS OF
CONTINUOUS FUNCTIONS**

MARIUS DADARLAT

ON HOMOMORPHISMS OF MATRIX ALGEBRAS OF CONTINUOUS FUNCTIONS

MARIUS DĂDĂRLAT

If X is a topological space we denote by $C(X) \otimes M_n$ the algebra of continuous functions from X to the algebra M_n of $n \times n$ complex matrices. A complete characterization of those topological spaces Y is given (in terms of vector bundles on Y) such that each unital algebra-homomorphism $\Phi: C(X) \otimes M_n \rightarrow C(Y) \otimes M_{kn}$ is of the form $\alpha \circ (\Phi' \otimes \text{id}_n)$ for some homomorphism $\Phi': C(X) \rightarrow C(Y) \otimes M_k$ and some suitable inner (or $C(Y)$ -linear) automorphism α of the algebra $C(Y) \otimes M_{kn}$. In particular this decomposition is assured provided that Y is a finite CW-complex of dimension $\leq 2k$ and $K^0(Y)$ does not have n -torsion.

Our interest in such homomorphisms arose in connection with a question of E. G. Effros [1] concerning the structure of inductive limits of C^* -algebras of the form $C(X) \otimes M_n$. In this context certain classes of homomorphisms related to a covering $X \rightarrow Y$ have been considered by C. Pasnicu [5]. When restricted to the case of automorphisms our results give nothing new (see [4], [6] and [7]).

1. Preliminaries. Let $\text{GL}_n(C)$ be the general linear group (nonsingular $n \times n$ matrices over the complex field) and denote by 1_n its unit. Let $\text{Vect}_m(Y)$ denote the set of isomorphism classes of complex vector bundles of rank m on the topological space Y . In $\text{Vect}_m(Y)$ we have one naturally distinguished element—the class of the trivial bundle of rank m . Let $T_n \text{Vect}_m(Y)$ be the subset of $\text{Vect}_m(Y)$ given by all vector bundles E such that the direct sum $E \oplus E \oplus \cdots \oplus E$ (n -times) is isomorphic to the trivial bundle of rank nm .

If A, B are unital complex algebras we denote by $\text{Hom}(A, B)$ the set of all unital algebra-homomorphisms from A to B . Two homomorphisms $\Phi_1, \Phi_2 \in \text{Hom}(A, B)$ are said to be inner equivalent if $\Phi_2 = u\Phi_1u^{-1}$ for some invertible element $u \in B$. Let $\text{Hom}(A, B)/\sim$ be the set of classes of inner equivalent homomorphisms from A to B .

We need some elementary sheaf cohomology. Let G be a Lie group and let H be a closed subgroup of G . For each topological space Y the fibration $H \rightarrow G \rightarrow G/H$ induces the following exact sequence of

pointed cohomology sets:

$$H^0(Y, H) \rightarrow H^0(Y, G) \rightarrow H^0(Y, G/H) \xrightarrow{\delta} H^1(Y, H) \rightarrow H^1(Y, G).$$

We have $H^0(Y, H) = C(Y, H)$ (continuous maps from Y to H) and $H^0(Y, G) = C(Y, G)$. These sets are pointed by the constant map $f = 1_G$ given by the unity of G . Similarly $H^0(Y, G/H) = C(Y, G/H)$ is pointed by the constant map $f = \{H\}$. The cohomology sets $H^1(Y, H)$ and $H^1(Y, G)$ are pointed by the trivial cocycles $(Y, 1_H)$ and $(Y, 1_G)$ respectively [2]. Given $f \in C(Y, G/H)$ the cocycle $\delta(f)$ represents the obstruction for lifting f to a function in $C(Y, G)$. By the exactness of the above sequence f has a continuous lifting if and only if $\delta(f) = (Y, 1_H)$. The action of G on G/H induces an action of $C(Y, G)$ on $C(Y, G/H)$. If $f_1, f_2 \in C(Y, G/H)$ then $\delta(f_1) = \delta(f_2)$ if and only if $f_2 = gf_1$ for some $g \in C(Y, G)$.

2. Results.

PROPOSITION 1. *Let Y be a topological space. Then there is a bijection $\text{Hom}(M_n, C(Y) \otimes M_{kn}) / \sim \rightarrow T_n \text{Vect}_k(Y)$.*

Proof. We describe the exact sequence induced by the following fibration:

$$\text{GL}_k(C) \xrightarrow{\gamma} \text{GL}_{kn}(C) \xrightarrow{j} \text{GL}_{kn}(C) / \text{GL}_k(C)$$

where the imbedding γ is given by

$$\gamma(u) = u \otimes 1_n, \quad M_{kn} = M_k \otimes M_n.$$

There is a commutative diagram of pointed sets:

$$\begin{array}{ccccccc} C(Y, \text{GL}_{kn}(C)) & \rightarrow & C(Y, \text{GL}_{kn}(C) / \text{GL}_k(C)) & \xrightarrow{\delta} & H^1(Y, \text{GL}_k(C)) & \xrightarrow{\gamma_*} & H^1(Y, \text{GL}_{kn}(C)) \\ \parallel & & \downarrow \alpha & & \downarrow \beta & & \downarrow \beta_1 \\ C(Y, \text{GL}_{kn}(C)) & \rightarrow & \text{Hom}(M_n, C(Y) \otimes M_{kn}) & \xrightarrow{\delta'} & \text{Vect}_k(Y) & \xrightarrow{\gamma'} & \text{Vect}_{kn}(Y) \end{array}$$

The vertical arrows are bijections. To describe α recall that

$$\text{Hom}(M_n, M_{kn}) \simeq \text{GL}_{kn}(C) / \text{GL}_k(C)$$

as topological spaces, the homeomorphism being induced by the map $\eta: \text{GL}_{kn}(C) \rightarrow \text{Hom}(M_n, M_{kn})$ given by $\eta(v)(a) = v(1_k \otimes a)v^{-1}$, $a \in M_n$. Let η_1 be the map

$$C(Y, \text{GL}_{kn}(C) / \text{GL}_k(C)) \rightarrow C(Y, \text{Hom}(M_n, M_{kn}))$$

induced by η . By definition we set $\alpha = \alpha_1 \eta_1$ where

$$\alpha_1: C(Y, \text{Hom}(M_n, M_{kn})) \rightarrow \text{Hom}(M_n, C(Y) \otimes M_{kn})$$

is given by $\alpha_1(\Psi)(a)(y) = \Psi(y)(a)$, $a \in M_n, y \in Y$. If in

$$\text{Hom}(M_n, C(Y) \otimes M_{kn})$$

we distinguish the homomorphism $a \mapsto a \otimes 1_k$, α will be an isomorphism of pointed sets. The maps β and β_1 are the natural ones. Namely if (U_i, g_{ij}) is a GL_k -cocycle, then $\beta(U_i, g_{ij})$ is the isomorphism class of the vector bundle obtained by clutching the trivial bundles $U_i \times C^k$ with the transition functions (g_{ij}) . The map β_1 is defined in a similar way. The other maps are defined to make the diagram commutative. If $v \in C(Y, \text{GL}_{kn}(C))$ then $j'(v): M_n \rightarrow C(Y) \otimes M_{kn}$ is defined by

$$j'(v)(a)(y) = v(y)(1_k \otimes a)v(y)^{-1}, \quad a \in M_n, y \in Y.$$

The map γ' takes the vector bundle E to the direct sum $E \oplus E \oplus \dots \oplus E$ (n -times). After the above identifications, it follows that two homomorphisms $\Phi_1, \Phi_2 \in \text{Hom}(M_n, C(Y) \otimes M_{kn})$ are inner equivalent if and only if $\delta'(\Phi_1) = \delta'(\Phi_2)$. The isomorphism class of the vector bundle $\delta'(\Phi_1)$ represents the obstruction for lifting Φ_1 to an invertible element in $C(Y) \otimes M_{kn}$. Also, by the exactness of the second row in the above diagram, the image of δ' is equal to $T_n \text{Vect}_k(Y)$.

THEOREM 2. *Let X, Y be topological spaces. Then the following assertions are equivalent:*

- (i) *The set $T_n \text{Vect}_k(Y)$ reduces to the trivial bundle of rank k .*
- (ii) *Each homomorphism $\Phi \in \text{Hom}(C(X) \otimes M_n, C(Y) \otimes M_{kn})$ is inner equivalent to a homomorphism of the form $\Phi' \otimes \text{id}_n$ for some $\Phi' \in \text{Hom}(C(X), C(Y) \otimes M_k)$.*

Proof. The implication (i) \Rightarrow (ii) follows easily from Proposition 1. Indeed, if we choose a point x in X and a homomorphism Φ_1 in $\text{Hom}(M_n, C(Y) \otimes M_{kn})$ which is not inner equivalent to the homomorphism $a \mapsto 1_k \otimes a$ then the homomorphism $C(X) \otimes M_n \ni F \mapsto \Phi_1(F(x)) \in C(Y) \otimes M_{kn}$ failed to satisfy (ii).

To prove the other implication we assume, as a preliminary step, that Φ acts on matrices as an amplification:

$$\Phi(1_{C(X)} \otimes a) = 1_{C(Y)} \otimes 1_k \otimes a, \quad a \in M_n.$$

Under this assumption we get

$$\begin{aligned} \Phi(f \otimes a) &= \Phi(f \otimes 1_n)\Phi(1 \otimes a) = \Phi(1 \otimes a)\Phi(f \otimes 1_n) \\ &= 1 \otimes 1_k \otimes a \cdot \Phi(f \otimes 1_n), \quad a \in M_n, f \in C(X). \end{aligned}$$

The previous computation shows us that the algebra $\Phi(C(X) \otimes 1_n)$ lies in the relative commutant of $1_{C(Y)} \otimes 1_k \otimes M_n$ in $C(Y) \otimes M_k \otimes M_n$ which is equal to $C(Y) \otimes M_k \otimes 1_n$. It follows that there is a unique homomorphism $\Phi' \in \text{Hom}(C(X), C(Y) \otimes M_k)$ such that $\Phi(f \otimes 1_n) = \Phi'(f) \otimes 1_n$. Using again our assumption on Φ we get $\Phi = \Phi' \otimes \text{id}_n$.

Consider now an arbitrary homomorphism Φ and let Φ_1 denote its restriction to M_n . Using (i) it follows by Proposition 1 that there is some invertible element $u \in C(Y) \otimes M_{kn}$ such that

$$\Phi_1(a) = \Phi(1 \otimes a) = u(1 \otimes 1_k \otimes a)u^{-1}, \quad a \in M_n.$$

Hence the homomorphism $u^{-1}\Phi u$ acts on matrices as an amplification.

REMARK 3. The assertion (i) in the above theorem holds provided that Y is homotopy equivalent to a finite CW-complex of dimension $\leq 2k$ and the K -theory group $K^0(Y)$ does not have n -torsion. This follows from the stability properties of vector bundles (see [3, Ch. 8, Th. 1.5]).

Note that $T_n \text{Vect}_1(Y)$ is a subgroup of the group $(\text{Vect}_1(Y), \otimes)$. We have a natural action of $T_n \text{Vect}_1(Y)$ on $T_n \text{Vect}_k(Y)$ given by $(L, E) \mapsto L \otimes E$. By similar methods one can prove the following

THEOREM 4. *Let X, Y be topological spaces. Then the following assertions are equivalent:*

- (i) $T_n \text{Vect}_1(Y)$ acts transitively on $T_n \text{Vect}_k(Y)$.
- (ii) For any homomorphism $\Phi \in \text{Hom}(C(X) \otimes M_n, C(Y) \otimes M_{kn})$ there is an automorphism α of $C(Y) \otimes M_{kn}$ which is $C(Y)$ -linear such that $\alpha \circ \Phi = \Phi' \otimes \text{id}_n$ for some homomorphism $\Phi' \in \text{Hom}(C(X), C(Y) \otimes M_k)$.

REFERENCES

- [1] E. G. Effros, *On the structure of C^* -algebras: Some old and some new problems*, in *Operator Algebras and Applications*. Proc. Symp. Pure Math., Amer. Math. Soc., Providence, RI, 1982.
- [2] F. Hirzebruch, *Topological Methods in Algebraic Geometry*, Springer-Verlag, New York, 1965.
- [3] D. Husemoller, *Fibre Bundles*, McGraw-Hill Book Company, 1966.
- [4] R. V. Kadison and J. R. Ringrose, *Derivations and automorphisms of operator algebras*, *Commun. Math. Phys.*, **4** (1967), 32–63.
- [5] C. Pasnicu, *On certain inductive limit C^* -algebras*, *Indiana Univ. Math. J.*, **35** (1986), 269–288.
- [6] J. Phillips and I. Raeburn, *Automorphisms of C^* -algebras and second Cech cohomology*, *Indiana Univ. Math. J.*, **29** (1980), 799–822.

- [7] K. Thomsen, *Automorphisms of homogeneous C^* -algebras*, Bull. Austral. Math. Soc., **33** (1986), 145–154.

Received December 15, 1986.

THE NATIONAL INSTITUTE FOR SCIENTIFIC AND TECHNICAL CREATION
BDUL PACII 220
79622 BUCHAREST, ROMANIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906-1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1988 by Pacific Journal of Mathematics

Jeffery Marc Bergen and Luisa Carini, A note on derivations with power central values on a Lie ideal	209
Alfonso Castro and Sumalee Unsurangsi, A semilinear wave equation with nonmonotone nonlinearity	215
Marius Dadarlat, On homomorphisms of matrix algebras of continuous functions	227
A. Didierjean, Quelques classes de cobordisme non orienté refusant de se fibrer sur des sphères	233
Edward George Effros and Zhong-Jin Ruan, On matricially normed spaces	243
Sherif El-Helaly and Taqdir Husain, Orthogonal bases are Schauder bases and a characterization of Φ -algebras	265
Edward Richard Fadell and Peter N-S Wong, On deforming G -maps to be fixed point free	277
Jean-Jacques Gervais, Stability of unfoldings in the context of equivariant contact-equivalence	283
Douglas Martin Grenier, Fundamental domains for the general linear group	293
Ronald Scott Irving and Brad Shelton, Loewy series and simple projective modules in the category $\mathcal{O}_{\mathcal{S}}$	319
Russell Allan Johnson, On the Sato-Segal-Wilson solutions of the K-dV equation	343
Thomas Alan Keagy and William F. Ford, Acceleration by subsequence transformations	357
Min Ho Lee, Mixed cusp forms and holomorphic forms on elliptic varieties	363
Charles Livingston, Indecomposable surfaces in 4-space	371
Geoffrey Lynn Price, Shifts of integer index on the hyperfinite II_1 factor	379
Andrzej Śladek, Witt rings of complete skew fields	391