

Pacific Journal of Mathematics

ON DEFORMING G -MAPS TO BE FIXED POINT FREE

EDWARD RICHARD FADELL AND PETER N-S WONG

ON DEFORMING G -MAPS TO BE FIXED POINT FREE

EDWARD FADELL AND PETER WONG

When $f: M \rightarrow M$ is a self-map of a compact manifold and $\dim M \geq 3$, a classical theorem of Wecken states that f is homotopic to a fixed point free map if, and only if, the Nielsen number $n(f)$ of f is zero. When M is simply connected, and $\dim M \geq 3$ the NASC becomes $L(f) = 0$, where $L(f)$ is the Lefschetz number of f . An equivariant version of the latter result for G -maps $f: M \rightarrow M$, where M is a compact G -manifold, is due to D. Wilczyński, under the assumption that M^H is simply connected of dimension ≥ 3 for any isotropy subgroup H with finite Weyl group WH . Under these assumptions, f is G -homotopic to a fixed point free map if, and only if, $L(f^H) = 0$ for any isotropy subgroup H (WH finite), where $f^H = f|_{M^H}$ and M^H represents those elements of M fixed by H . A special case of this result was also obtained independently by A. Vidal via equivariant obstruction theory. In this note we prove the analogous equivariant result without assuming that the M^H are simply connected, assuming that $n(f^H) = 0$, for all H with WH finite. There is also a codimension condition. Here is the main result.

THEOREM. *Let G denote a compact Lie group and M a compact, smooth G -manifold. Let $(H_1), \dots, (H_k)$ denote an admissible ordering of the isotropy types of M , $M_i = \{x \in M: (G_x) = (H_j), j \leq i\}$ the associated filtration. Also, let \mathcal{F} denote the set of integers i , $1 \leq i \leq k$, such that the Weyl group $WH_i = NH_i/H_i$ is finite. Suppose that for each $i \in \mathcal{F}$, $\dim M^{H_i} \geq 3$ and the codimension of $M_{i-1} \cap M^{H_i}$ in M^{H_i} is at least 2. Then, a G -map $f: M \rightarrow M$ is G -homotopic to a fixed point free G -map $f': M \rightarrow M$ if, and only if, the Nielsen number $n(f^{H_i}) = 0$ for each $i \in \mathcal{F}$.*

1. Preliminaries. Throughout this note G will denote a compact Lie group and M will denote a compact, smooth G -manifold. For any closed subgroup H in G , we denote by NH the normalizer of H in G and by $WH = NH/H$, the Weyl group of H in G . The conjugacy class of H , denoted by (H) , is called the orbit type of H . If $x \in M$ then G_x denotes the isotropy subgroup of x , i.e. $G_x = \{g \in G | gx = x\}$. For each subgroup H of G , $M^H = \{x \in M | hx = x \text{ for all } h \in H\}$ and $M_H = \{x \in M | G_x = H\}$. Let $\{(H_j)\}$ denote the (finite) set of isotropy

types of M . If (H_j) is subconjugate to (H_i) , we write $(H_j) \leq (H_i)$. We can choose an *admissible* ordering on $\{(H_j)\}$ so that $(H_j) \leq (H_i)$ implies $i \leq j$ (see [1]). Then we have a filtration of G -subspaces $M_1 \subset M_2 \subset \cdots \subset M_k = M$, where $M_i = \{x \in M : (G_x) = (H_j), j \leq i\}$.

We now recall the definition of the *local* Nielsen number. If U is an open subset of M and $f: U \rightarrow M$ is a compactly fixed map (not necessarily a G -map), then the Nielsen number $n(f, U)$ is defined [3] using the equivalence relation on the fixed point set $\text{Fix } f$ as follows. Two fixed points x and y in U are equivalent if there is a path α in U such that $f(\alpha)$ and α are endpoint homotopic in M . The remainder of the local theory proceeds along the lines of the global theory. Note that if $\dim M \geq 3$, the path α above may be taken as a simple path in U and assuming (as we may) that $\text{Fix } f$ is finite, α may be chosen to avoid all fixed points different from x and y . Then in a small closed tubular neighborhood $T \subset U$ of α f may be altered in the interior T^0 of T (via the Wecken method [1] or the “Whitney trick” see [4]) to obtain a map $f': U \rightarrow M$ so that f' is an extension of $f|U - T^0$, $f' \sim f$ and f' has only one fixed point in U . This is the technique of coalescing fixed points. Note that T is a closed n -ball. If this remaining fixed point has *index* 0, f' may be altered within T^0 to remove it, thus obtaining $f'' \sim f$ such that $f''|U - T^0 = f|U - T^0$ and f'' has no fixed points in T . A key point here is that f is altered in the interior of a closed *contractible* neighborhood of α .

2. The Proof of Main Theorem. We first note that the G -map $f: M \rightarrow M$ preserves the filtration $M_1 \subset \cdots \subset M_k$, i.e., $f(M_i) \subset M_i$. Also, $W_i = WH_i$ acts on M^{H_i} and freely on $M^{H_i} - M_{i-1}$. Furthermore $f^{H_i} = f|M^{H_i}: M^{H_i} \rightarrow M^{H_i}$ is a W_i -map. We will set $f_i = f|M_i$. We then let \mathcal{F} denote the indices i , $1 \leq i \leq k$, such that W_i is finite. Whenever A is a G -set, \tilde{A} will denote the corresponding set of orbits, i.e., $\tilde{A} = A/G$. Similarly $\tilde{f}: \tilde{A} \rightarrow \tilde{B}$, will denote the map induced by a G -map $f: A \rightarrow B$. Finally, $\text{Fix } f$ is a G -set, and each orbit in $\text{Fix } f$ will be referred to as a *fixed orbit*.

2.1. LEMMA. (Controlled Homotopy Extension). *Let (X, A) denote a G -pair such that all orbits in $X - A$ have the same orbit type G/H and $f: (X, A) \rightarrow (X, A)$ a G -map (of pairs). Let V denote a closed G -neighborhood of A and $f^H: X^H \rightarrow X^H$ the restriction of f to X^H . Then, any (NH/H) -homotopy f_t^H , relative to V^H , with $f_0^H = f^H$ extends to a G -homotopy f_t , relative to V , with $f_0 = f$. Furthermore if f_1^H is fixed point free on $(X - A)^H$, then so is f_1 on $X - A$.*

Proof. The proof is an easy consequence of II.5.12 in Bredon [1]. Let $W = NH/H$ and $Y = X - A$. The homotopy f_t is defined on Y by setting

$$f_t(y) = gf_t^H(g^{-1}y), \quad G_y = gHg^{-1}.$$

Then, f_t is defined on Y and on $V \cap Y$, $f_t = f_0$, $0 \leq t \leq 1$. Thus, setting $f_t = f$ on V extends f_t to all of X . Note that

$$gf_1^H(g^{-1}y) = y \Leftrightarrow f_1^H(g^{-1}y) = g^{-1}y, \quad y \in Y,$$

which verifies the last assertion of the lemma.

2.2. PROPOSITION (Inductive Step when W_i is finite). *Let $f: M \rightarrow M$ denote a G -map such that*

- (1) $f_{i-1}: M_{i-1} \rightarrow M_{i-1}$ is fixed point free,
- (2) $W_i = NH_i/H_i$ is finite,
- (3) f_i has a finite number of fixed orbits on $U_i = M_i - M_{i-1}$,
- (4) $n(f_i^H, U_i^H) = 0$.

Then, f is G -homotopic to $f': M \rightarrow M$, relative to M_{i-1} so that f'_i is fixed point free.

Proof. Consider the map $f_i: M_i \rightarrow M_i$ and let $H = H_i$ and $W = W_i$ for notational convenience. Now focus attention on $f_i^H: M_i^H \rightarrow M_i^H$ and let \mathcal{O}_1 and \mathcal{O}_2 denote two fixed W -orbits in U_i^H . Call the fixed orbits \mathcal{O}_1 and \mathcal{O}_2 Nielsen equivalent if for some $x \in \mathcal{O}_1$ and $y \in \mathcal{O}_2$, x and y are Nielsen equivalent in U_i^H (see [2]). We will coalesce two Nielsen equivalent orbits into one fixed orbit as follows. Suppose $\mathcal{O}_1 = Wx$ and $\mathcal{O}_2 = Wy$ with x and y Nielsen equivalent in U_i^H , i.e., there is a path α from x to y in U_i^H so that $f\alpha \sim \alpha$ (in M_i^H and with ends fixed). Because of assumption (3) we may assume that α avoids all other points of $\text{Fix } f_i$ other than x and y . Project α to $\tilde{\alpha}$ in U_i^H/W by the orbit map $\eta: U_i^H \rightarrow U_i^H/W$ and let $\tilde{\beta}$ denote a simple path homotopic (relative to end points) to $\tilde{\alpha}$. Then $\tilde{\beta}$ lifts to a simple path β from x to y . If $N(\tilde{\beta})$ is a closed ball neighborhood of $\tilde{\beta}$ in U_i^H/W , then, since $N(\tilde{\beta})$ is contractible, $\eta^{-1}(N(\tilde{\beta})) = WN(\beta)$, where $N(\beta)$ is the corresponding ball neighborhood of β . Thus, $\eta^{-1}(N(\tilde{\beta}))$ consists of disjoint translates of $N(\beta)$ by W . The local Nielsen number $n(f_i^H, N(\beta))$ is at most one (see [3]). Applying the local Wecken theorem or the ‘‘Whitney trick’’ in $N(\beta)$ (see [3] or [4]), we can obtain a homotopy $H: N(\beta) \times I \rightarrow M_i^H$ such that $H_t|_{\partial N(\beta)} = f_i$ for all t , $0 \leq t \leq 1$, $H_0 = f_i|_{N(\beta)}$ and H_1 has at most one fixed point in the interior of $N(\beta)$. H has the extension $H(wx, t) = wH(x, t)$ to

$(WN(\beta)) \times I$ and to all of $M_i^H \times I$ by using f_i outside of $WN(\beta) \times I$. Then $H: M_i^H \times I \rightarrow M_i$ is a W -homotopy, with $H_1 = \varphi_i: M_i^H \rightarrow M_i^H$ having one less or two less fixed orbits. Continuing in this manner we obtain a W -map $\varphi'_i: M_i^H \rightarrow M_i^H$, W -homotopic to f_i with finitely many fixed orbits no two of which are Nielsen equivalent. If x belongs to one of the remaining orbits Wx and D is a sufficiently small neighborhood of x , then the local indices $i(\varphi'_i, D)$ and $i(\varphi'_i, wD)$, $w \in W$, are the same and $i(\varphi'_i, WD) = |W|i(\varphi'_i, D)$. Since $n(f_i^H, U_i^H) = 0$, we must have $i(\varphi'_i, D) = 0$, since Wx is the union of Nielsen classes. We can now remove x as a fixed point via a homotopy relative to ∂D and extend (as above) to a W -map $\varphi''_i: M_i^H \rightarrow M_i^H$, W -homotopic to φ'_i , with Wx eliminated as a fixed orbit. Continuing in this manner we arrive at a W -map $\psi_i: M_i^H \rightarrow M_i^H$ which is fixed point free and W -homotopic to f_i relative to some closed neighborhood of M_{i-1}^H . By Lemma 2.1 this map ψ_i extends to a fixed point free G -map $f'_i: M_i \rightarrow M_i$, G -homotopic to f_i (relative to M_{i-1}) and the G -homotopy extension theorem provides the required extension f' of f'_i .

2.3. PROPOSITION (*Inductive Step when $\dim W_i > 0$*). Let $f_i: M_i \rightarrow M_i$ denote a G -map such that

- (1) f_i^H is fixed point free on M_{i-1}^H ,
- (2) $\dim W_i > 0$, $W_i = NH_i/H_i$.

Then, f_i is G -homotopic relative to A to a G -map $f'_i: M_i \rightarrow M_i$ such that f'_i is fixed point free.

Proof. This proposition follows from Lemma 3.3 in [6].

Proof of Theorem. We assume (inductively) that $f: M \rightarrow M$ is a G -map such that $f_{i-1}: M_{i-1} \rightarrow M_{i-1}$ is fixed point free. As a first step, choose a closed G -neighborhood V of M_{i-1} in M_i so that M_{i-1} is a G -deformation retract of V . Then, f is G -homotopic, relative to M_{i-1} , to a map f' such that $f'_i: M_i \rightarrow M_i$ has no fixed points in V_i . Thus, we may assume that f itself has this property so that f_i is compactly fixed on $U_i = M_i - M_{i-1}$. In particular f_i^H is compactly fixed on U_i^H and the local Nielsen number $n(f_i^H, U_i^H)$ is defined. We consider two cases.

Case 1. $W_i = NH_i/N_i$ is finite, i.e., $i \in \mathcal{F}$.

In this case, the codimension condition applies to yield $n(f_i^H, U_i^H) = n(f_i^H) = 0$. This is because any path in M_i^H from x to y , $x \cup y \in U_i^H$ may be deformed (ends fixed) to a path in U_i^H , i.e., to one avoiding the submanifold M_{i-1}^H . Let V' denote a closed G -neighborhood of M_{i-1}^H

in M_i^H so that the fixed points $\text{Fix } f_i^H$ of f_i^H are in $M_i^H - V'$. Choose a closed G -neighborhood $Q \subset U_i^H$ of $\text{Fix } f_i$ so that $f_i^H(Q) \subset U_i^H$. Working in the orbit space U_i^H/W_i , we deform $\tilde{f}_i|_{\tilde{Q}}$, relative $\partial\tilde{Q}$, to a map \tilde{f}'_i with finitely many fixed points. Since W_i acts freely on U_i^H we may apply the covering homotopy theorem to conclude that f_i^H is homotopic relative to V' to $\varphi: M_i^H \rightarrow M_i^H$ where φ has finitely many fixed W_i -orbits and the homotopy is compactly fixed. Thus, $n(\varphi, U_i^H) = 0$ and we may apply Proposition 2.2 to conclude that f is G -homotopic, relative to M_{i-1} , to a map $f': M \rightarrow M$ with $f'_i: M_i \rightarrow M_i$ fixed point free.

Case 2. $\dim W_i > 0$. We apply Proposition 2.3 and then the G -homotopy extension theorem to conclude that f is G -homotopic, relative to M_{i-1} , to a map $f': M \rightarrow M$ with $f'_i: M_i \rightarrow M_i$ fixed point free.

Applying induction completes the proof of the sufficiency. The necessity is clear.

REFERENCES

- [1] G. Bredon, *Introduction to Compact Transformation Groups*, Academic Press, New York 1972.
- [2] R. F. Brown, *The Lefschetz Fixed Point Theorem*, Scott-Foresman 1971.
- [3] E. Fadell and S. Husseini, *Local fixed point index theory for non-simply connected manifolds*, Illinois J. Math., **25** (1981), 673-699.
- [4] Boju Jiang, *Fixed Point Classes From a Differential Viewpoint*, Lecture Notes #886, Springer-Verlag 1981, 163-170.
- [5] A. Vidal, *Equivariant Obstruction Theory for G -Deformations* (in German), Dissertation 1985, Universität Heidelberg.
- [6] D. Wilczyński, *Fixed point free equivariant homotopy classes*, Fund. Math., **123** (1984), 47-60.

Received April 6, 1987. The first author was supported in part by the National Science Foundation under Grant No. DMS-8320099.

UNIVERSITY OF WISCONSIN
MADISON, WI 53706

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906-1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA, RENO

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF HAWAII

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to *Pacific Journal of Mathematics*, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1988 by Pacific Journal of Mathematics

Jeffery Marc Bergen and Luisa Carini, A note on derivations with power central values on a Lie ideal	209
Alfonso Castro and Sumalee Unsurangsi, A semilinear wave equation with nonmonotone nonlinearity	215
Marius Dadarlat, On homomorphisms of matrix algebras of continuous functions	227
A. Didierjean, Quelques classes de cobordisme non orienté refusant de se fibrer sur des sphères	233
Edward George Effros and Zhong-Jin Ruan, On matricially normed spaces	243
Sherif El-Helaly and Taqdir Husain, Orthogonal bases are Schauder bases and a characterization of Φ -algebras	265
Edward Richard Fadell and Peter N-S Wong, On deforming G -maps to be fixed point free	277
Jean-Jacques Gervais, Stability of unfoldings in the context of equivariant contact-equivalence	283
Douglas Martin Grenier, Fundamental domains for the general linear group	293
Ronald Scott Irving and Brad Shelton, Loewy series and simple projective modules in the category $\mathcal{O}_{\mathcal{S}}$	319
Russell Allan Johnson, On the Sato-Segal-Wilson solutions of the K-dV equation	343
Thomas Alan Keagy and William F. Ford, Acceleration by subsequence transformations	357
Min Ho Lee, Mixed cusp forms and holomorphic forms on elliptic varieties	363
Charles Livingston, Indecomposable surfaces in 4-space	371
Geoffrey Lynn Price, Shifts of integer index on the hyperfinite II_1 factor	379
Andrzej Śladek, Witt rings of complete skew fields	391