

Pacific Journal of Mathematics

**ISOMORPHIC BANACH-STONE THEOREMS AND
ISOMORPHISMS WHICH ARE CLOSE TO ISOMETRIES**

EHRHARD BEHREND

ISOMORPHIC BANACH-STONE THEOREMS AND ISOMORPHISMS WHICH ARE CLOSE TO ISOMETRIES

EHRHARD BEHREND

A Banach space X is said to have the *isomorphic Banach-Stone property* if for locally compact Hausdorff spaces K and L one always can conclude that K and L are homeomorphic provided that the Banach spaces $C_0(K, X)$ and $C_0(L, X)$ (=the continuous X -valued functions on K resp. L which vanish at infinity) are isomorphic with sufficiently small Banach-Mazur distance.

Our main results are a characterization of the finite-dimensional spaces with this property, and we also get an abundance of new finite- and infinite-dimensional examples.

These results appear as corollaries to general theorems about isomorphisms between certain spaces of continuous vector-valued functions. They enable us also to conclude that, for certain spaces X , and all compact K all isomorphisms T on $C_0(K, X)$ with $(1/(1+\tau))\|f\| \leq \|Tf\| \leq (1+\tau)\|f\|$ for small τ can be approximated by isometries.

1. Introduction. The *classical Banach-Stone theorem* asserts that two locally compact Hausdorff spaces K and L are homeomorphic if and only if the Banach spaces C_0K and C_0L are isometrically isomorphic (C_0K =the space of continuous scalar-valued functions which vanish at infinity, provided with the supremum norm). Several *generalizations* of this theorem have been studied in the literature. For example, there have been investigated Banach spaces X such that “scalar-valued” can be replaced by “ X -valued” in the assertion of the Banach-Stone theorem. X is then said to have the *Banach-Stone property*. This generalization is well understood; the interested reader is referred to [4] and [5] where it is shown that in a sense which can be made precise one knows all possibilities to determine spaces with the Banach-Stone property.

Another generalization is due to Amir [2] and Cambern [9] who proved independently that in the assertion of the theorem one may replace “ C_0K and C_0L are isometrically isomorphic” by “the Banach spaces C_0K and C_0L are isomorphic, and the Banach-Mazur distance is smaller than two”. Later it was shown that also in this version

one may replace the scalars by certain Banach spaces if the Banach-Mazur distance of the function spaces under consideration is sufficiently small. The most far-reaching results into this direction are those of Cambern [11] and Jarosz [15].

1.1. DEFINITION. Let X be a Banach space. We will say that X has the *isomorphic Banach-Stone property* if there is a $\delta > 0$ such that the following holds: whenever K and L are locally compact Hausdorff spaces such that there exists an isomorphism $T: C_0(K, X) \rightarrow C_0(L, X)$ (these are the continuous X -valued functions on K resp. L which vanish at infinity, and this function space is provided with the supremum norm), then K and L are homeomorphic provided that $\|T\| \|T^{-1}\| \leq 1 + \delta$.

Cambern proved that uniformly convex spaces have this property, and by Jarosz's result a certain rather technical condition concerning the dual of X is sufficient.

Given K and L and a homeomorphism $\varphi: K \rightarrow L$ one can try to define $T: C_0(K, X) \rightarrow C_0(L, X)$ by $(Tf)(\varphi(k)) = T_k[f(k)]$, where T_k is an isomorphism from X to X for every k . A moment's reflection shows that T will be an isomorphism with $\|T\| \|T^{-1}\|$ close to one if the T_k have uniformly this property, and it even suffices that the $(Tf)(\varphi(k))$ are close to the $T_k[f(k)]$. One might ask whether every isomorphism T can be approximated in this way. To be more precise:

1.2. DEFINITION. X is said to have the *strong isomorphic Banach-Stone property* if for every $\varepsilon > 0$ there is a $\delta > 0$ such that the following holds: whenever K , L , and T are given as in 1.1, then there exist a homeomorphism $\varphi: K \rightarrow L$ and a family $(T_k)_{k \in K}$ of isomorphisms on X such that

$$\begin{aligned} \|T_k\| \|T_k^{-1}\| &\leq 1 + \varepsilon \quad \text{and} \\ \|(Tf)(\varphi(k)) - T_k[f(k)]\| &\leq \varepsilon \end{aligned}$$

for all $k \in K$ and all $f \in C_0(K, X)$ with $\|f\| \leq 1$.

There exist two fairly general approaches to decide whether a given X satisfies 1.1 or 1.2. They are completely independent, but both rely on an isomorphic adaption of certain concepts of M -structure theory (which gave the most far-reaching results in the isometrical case). The *first approach* depends on the behaviour of certain M -ideals under isomorphisms. Using this it can be shown that every X such that l_2^1 (the two-dimensional L^1 -space) is bounded away uniformly from every two-dimensional subspace of X' (=the dual of X) has the

isomorphic Banach-Stone property, a result which will be published in a joint paper of M. Cambern and the author [7].

Here we will study *another approach* which depends only in part on M -structure (namely on the behaviour of certain operators which are just the isomorphic variants of multipliers). In cases where both [7] and the present results apply the admissible δ 's which are determined by [7] are usually larger. In the present paper, however, we are provided also with spaces satisfying 1.2, and in the finite-dimensional case we even obtain necessary and sufficient conditions.

In contrast to the isometric vector-valued theory several interesting questions are open for isomorphic vector-valued Banach-Stone theorems. In particular there is nothing known about the range of the possible constants δ for a Banach space X with the isomorphic Banach-Stone property. The only exception is the case of one-dimensional X : here precisely the $\delta < 1$ are admissible ([10], [12]).

Our results will be derived as *special cases of a much more general theory* concerning isomorphisms between certain spaces of continuous vector-valued functions. To be more precise, we start with

1.4. DEFINITION. Let $K_1, \dots, K_r, L_1, \dots, L_s$ be locally compact Hausdorff spaces and $X_1, \dots, X_r, Y_1, \dots, Y_s$ Banach spaces.

(i) $C_0(K_1, \dots, K_r; X_1, \dots, X_r)$ denotes the space $\prod_{\rho=1}^r C_0(K_\rho, X_\rho)$, the L^∞ -direct product of the $C_0(K_\rho, X_\rho)$ (this is just the product of these spaces, provided with the supremum norm). The elements will be regarded as "functions" f defined on $K := K_1 \dot{\cup} \dots \dot{\cup} K_r$, where for $k \in K_\rho$ the element $f(k)$ lies in X_ρ .

(ii) Let $\tau \geq 0$ and

$$T: C_0(K_1, \dots, K_r; X_1, \dots, X_r) \rightarrow C_0(L_1, \dots, L_s; Y_1, \dots, Y_s)$$

be an isomorphism. T will be called a τ -isomorphism if

$$\frac{1}{1+\tau} \|f\| \leq \|Tf\| \leq (1+\tau) \|f\|$$

for every f (clearly every isomorphism T is—up to a constant—a τ -isomorphism for a suitable τ , and we may assume that $\|T\| \|T^{-1}\| = (1+\tau)^2$; the advantage of considering τ -isomorphisms lies in the fact T and T^{-1} satisfy the same norm conditions so that in the sequel one only will have to prove the assertions concerning T ; those concerning T^{-1} will then be true by symmetry).

We now are going to explain the results in some detail. In §2 we assign to each finite family X_1, \dots, X_r of Banach spaces a number

$\alpha(X_1, \dots, X_r)$. The definition is rather technical, and there is certainly no hope to calculate precise values in the general situation. However, we are able to prove the following:

THEOREM (TH. 2.4). *Suppose that $\alpha(X_1, \dots, X_r) + \alpha(Y_1, \dots, Y_s) < 1$. Then there exists a $\tau > 0$ such that the existence of a τ -isomorphism*

$$T: C_0(K_1, \dots, K_r; X_1, \dots, X_r) \rightarrow C_0(L_1, \dots, L_s; Y_1, \dots, Y_s)$$

implies that there is a homeomorphism

$$\varphi: K_1 \dot{\cup} \dots \dot{\cup} K_r \rightarrow L_1 \dot{\cup} \dots \dot{\cup} L_s.$$

We even can show more. If we denote, for Banach spaces X and Y , the Banach-Mazur distance

$$\inf\{\|T\| \|T^{-1}\| \mid T: X \rightarrow Y \text{ an isomorphism}\}$$

by $d(X, Y)$, then we have the

THEOREM (TH. 2.5). *Suppose that $d(X_\rho, X_{\rho'})$, $d(Y_\sigma, Y_{\sigma'}) > 1$ for all $\rho \neq \rho'$, $\sigma \neq \sigma'$. We further assume that $\alpha(X_1, \dots, X_r)$ and $\alpha(Y_1, \dots, Y_s)$ are sufficiently small. Then there exists a $\tau > 0$ such that the existence of a τ -isomorphism implies that $r = s$, that—up to rearrangement— K_ρ and L_ρ are homeomorphic, and $d(X_\rho, Y_\rho)$ is close to one for every ρ .*

Of course these theorems are of no interest as long as we have no possibilities to give estimates for the constants $\alpha(X_1, \dots, X_r)$, $\alpha(Y_1, \dots, Y_s)$. We develop two possibilities to get such estimates. For the first one we introduce a new constant which measures, in a sense, how large the segments in the unit sphere are:

1.5. DEFINITION. Let X be a Banach space. By $\alpha_1(X)$ we denote the infimum of those numbers $\alpha > 0$ for which there exists an $\varepsilon > 0$ such that the following holds: if x_0, x are vectors in X with

$$1 - \varepsilon \leq \|x_0\| \leq 1 + \varepsilon$$

$$\|x_0 + \Theta x\| \leq 1 + \varepsilon \text{ for all scalars } \Theta \text{ with } |\Theta| = 1,$$

then $\|x\| \leq \alpha$.

In the case of finite-dimensional spaces $\alpha_1(X)$ is just the maximal radius of one-dimensional balls contained in the unit sphere.

$\alpha_1(X) = 0$ just means that X is uniformly convex (which has to be understood in the sense of [14] for complex spaces).

The *second possibility* uses isomorphic variants of multipliers:

1.6. DEFINITION. Let X be a Banach space and $R: X \rightarrow X$ an operator

(i) Suppose first that $\|R\| \leq 1$. Then R is called a *multiplier* (or *M -bounded*) if

$$\|Rx - x_0\| \leq 1$$

whenever x_0, x are vectors such that $\|\Theta x - x_0\| \leq 1$ for all scalars Θ with $|\Theta| = 1$. Operators with $\|R\| > 1$ will be called multipliers if $R/\|R\|$ has the above property.

The collection of all multipliers is denoted by $\text{Mult}(X)$ (the *multiplier of X*), and this set is said to be *trivial* if it contains only the operators $\mathbf{K} \cdot \text{Id}$; as usual \mathbf{K} stands for the scalar field, $\mathbf{K} = \mathbf{R}$ or $\mathbf{K} = \mathbf{C}$.

(ii) For $\varepsilon \geq 0$ we say that R belongs to $\text{Mult}_\varepsilon(X)$ if R satisfies the following condition:

$$\|Rx - x_0\| \leq 1 + \varepsilon$$

for all vectors with $\|\Theta x - x_0\| \leq 1$ (all Θ with $|\Theta| = 1$).

Note. For a survey of properties of $\text{Mult}(X)$ the reader is referred to [1], [4], and [6]. For our purposes it is important to note that there are many classes of spaces known where $\text{Mult}(X)$ is trivial, e.g.

X strictly convex [16], [19]

X smooth [6]

$X = L^1(\mu)$, and $\dim X > 2$ [6].

X is finite-dimensional and contains no nontrivial M -summand (i.e. no subspace J different from $\{0\}$ and X with an L^∞ -complement; [6]).

Trivially $\text{Mult}_0(X)$ consists of the R in $\text{Mult}(X)$ with $\|R\| \leq 1$. We need to know how close the $R \in \text{Mult}_\varepsilon(X)$, which behave nearly as multipliers, are to the trivial elements of $\text{Mult}(X)$, the operators $\mathbf{K} \text{Id}$. This gives rise to the next definition; there we will use the symbol $(M)_\alpha$ to denote, for a subset M of a normed space, the set $\{x \mid \text{there exists an } m \in M \text{ with } \|x - m\| \leq \alpha\}$.

1.7. DEFINITION. $\alpha_2(X)$ denotes the infimum of those numbers α for which there is an $\varepsilon > 0$ such that

$$\text{Mult}_\varepsilon(X) \subset (\mathbf{K} \text{Id})_\alpha.$$

A systematic study of $\text{Mult}_\varepsilon(X)$ and $\alpha_2(X)$ is not intended in this paper. We only collect together in §3 those properties which are of

importance for our purposes. In particular we prove (Th. 3.2 and Th. 3.3):

THEOREM.

- (i) *If X is finite-dimensional and $\text{Mult}(X)$ is trivial then $\alpha_2(X) = 0$.*
- (ii) *$\alpha_2(X) = 0$ for every uniformly smooth space.*

Our *main result*, by which we are in a position to apply the investigations of §2, is then the following theorem. Its proof and the first consequences are discussed in §4:

THEOREM. *Let X_1, \dots, X_r be Banach spaces. Then*

$$\alpha(X_1, \dots, X_r) \leq \max_{\rho} \min\{\alpha_1(X_{\rho}), 2\alpha_2(X_{\rho})\}.$$

It is then fairly easy to derive various isomorphic Banach-Stone theorems. In §5 we prove:

- uniformly convex and uniformly smooth spaces have the strong isomorphic Banach-Stone property;
- let X be finite-dimensional; we write X as $\prod_{\rho=1}^{r\infty} X_{\rho}^{n_{\rho}}$, where the X_{ρ} are subspaces which are pairwise not isometrically isomorphic and have trivial $\text{Mult}(X_{\rho})$; such a representation is always possible and is—up to rearrangement—unique (cf. [6]); then X has the isomorphic Banach-Stone property

$$\text{iff } \min n_{\rho} = 1, \quad \text{and}$$

X has the strong isomorphic Banach-Stone property

$$\text{iff } r = 1 \quad \text{and} \quad n_1 = 1$$

(which, by [3], is the same characterization as in the isometrical theory).

Another application of the results of §4 is discussed in §6: how close can τ -isomorphisms on a space be approximated by isometries?

1.8. DEFINITION. We say that a Banach-space X has the *AI-property* (which stands for “approximation by isometries”) if for every $\varepsilon > 0$ there is a $\tau > 0$ such that every τ -isomorphism lies in $(\text{Iso}_0(X))_{\varepsilon}$ (where $\text{Iso}_0(X)$ is the group of all isometries).

We prove that, for certain X , all $C(K, X)$ for arbitrary compact Hausdorff spaces K have the AI-property. These results seem to be new even in the case $X = \mathbf{K}$ (where, however, there is some overlap with the work of Benyamini [8]; cf. also theorem 6.1 in Jarosz [17]).

Some of our investigations, particularly in §2 and §4, are rather technical. It is suggested to the reader to consider first the special case $r = s = 1$ and the case of compact instead of locally compact spaces.

NOTATION. Banach spaces are always real or complex and not reduced to zero, operators are linear and continuous. Topological spaces are assumed to be Hausdorff spaces. The symbol " \cong " means "isometrically isomorphic" when applied to Banach spaces and "homeomorphic" in the case of topological spaces. Isomorphisms are always onto isomorphisms. The spaces $C_0(K, \mathbf{K})$ are abbreviated by C_0K for locally compact K and CK for compact spaces. Finally, $B(x, r)$ stands for the closed ball with center x and radius r in a Banach space.

Acknowledgment. The author wishes to express his gratitude to Michael Cambern and his colleagues from the University of California at Santa Barbara for their hospitality during the author's stay in the summer of 1986.

2. The constant $\alpha(X_1, \dots, X_n)$. Let $T: C_0(K_1, \dots, K_r; X_1, \dots, X_r) \rightarrow C_0(L_1, \dots, L_s; Y_1, \dots, Y_s)$ be a τ -isomorphism. We will abbreviate

$$V := C_0(K_1, \dots, K_r; X_1, \dots, X_r),$$

$$W := C_0(L_1, \dots, L_s; Y_1, \dots, Y_s),$$

$$K := K_1 \dot{\cup} \dots \dot{\cup} K_r,$$

$$L := L_1 \dot{\cup} \dots \dot{\cup} L_s,$$

and we will associate certain points of K with points of L .

2.1. DEFINITION. Let $\alpha \geq 0$.

(i) For $k \in K$ and $l \in L$ we write $k \xrightarrow{\alpha}_T l$ if $\|(Tf)(l)\| \leq \alpha\|f\|$ for every $f \in V$ such that $f(k) = 0$.

(ii) $K \xrightarrow{\alpha}_T L$ means that for every k there is an l with $k \xrightarrow{\alpha}_T l$.

2.2. PROPOSITION.

(i) Suppose that $k \xrightarrow{\alpha}_T l$. Then

$$\|(Tf)(l)\| \leq \alpha\|f\| + (1 + \tau)\|f(k)\|$$

for every $f \in V$.

(ii) Suppose that $k \xrightarrow{\alpha}_T l$ and $l \xrightarrow{\beta}_{T^{-1}} \tilde{k}$. Then $k = \tilde{k}$ provided that $(\alpha + \beta)(1 + \tau) < 1$. Similarly $l \xrightarrow{\beta}_{T^{-1}} k$, $k \xrightarrow{\alpha}_T \tilde{l}$ yield $l = \tilde{l}$ under the same condition.

(iii) Suppose that $K \xrightarrow[T]{\alpha} L$ and $L \xrightarrow[T^{-1}]{\beta} K$ and that $(\alpha + \beta)(1 + \tau) < 1$. Then for every k (resp. l) there is precisely one l (resp. k) such that $k \xrightarrow[T]{\alpha} l$ (resp. $l \xrightarrow[T^{-1}]{\beta} k$). This induces mappings $\varphi: K \rightarrow L$ and $\psi: L \rightarrow K$.

(iv) Under the conditions of (iii) we have $\varphi = \psi^{-1}$, and φ and ψ are continuous (i.e. $K \cong L$).

Proof.

(i) Let $f \in V$ be given and $\eta > 0$. We choose $\tilde{f} \in V$ with $\|f\| = \|\tilde{f}\|$, $\|f - \tilde{f}\| \leq \|f(k)\| + \eta$, $\tilde{f}(k) = 0$. It follows that $\|(T\tilde{f})(l)\| \leq \alpha\|f\|$ and $\|Tf - T\tilde{f}\| \leq (1 + \tau)(\|f(k)\| + \eta)$ so that

$$\|(Tf)(l)\| \leq \alpha\|f\| + (1 + \tau)(\|f(k)\| + \eta).$$

This proves the assertion since η was arbitrary.

(ii) Suppose that $k \neq \tilde{k}$. We choose $f \in V$ with $\|f\| = 1$, $f(k) = 0$, $\|f(\tilde{k})\| = 1$. By $k \xrightarrow[T]{\alpha} l$ it follows that $\|(Tf)(l)\| \leq \alpha$, and (i) yields

$$\begin{aligned} 1 &= \|f(\tilde{k})\| = \|T^{-1}(Tf)(\tilde{k})\| \\ &\leq \beta\|Tf\| + (1 + \tau)\|Tf(l)\| \\ &\leq \beta(1 + \tau) + (1 + \tau)\alpha, \end{aligned}$$

a contradiction.

(iii) Let k, l_1, l_2 be given with $k \xrightarrow[T]{\alpha} l_1$, $k \xrightarrow[T]{\alpha} l_2$, and $l_1 \neq l_2$. Choose $g \in W$ with $\|g\| = 1$, $g(l_1) = 0$, $\|g(l_2)\| = 1$. (ii) yields $\|(T^{-1}g)(k)\| \leq \beta$ so that, also by (ii),

$$\begin{aligned} 1 &= \|g(l_2)\| = \|T(T^{-1}g)(l_2)\| \\ &\leq \alpha\|T^{-1}g\| + (1 + \tau)\|T^{-1}g(k)\| \\ &\leq (\alpha + \beta)(1 + \tau) \end{aligned}$$

which contradicts our assumption.

(iv) The assertion $\varphi = \psi^{-1}$ is a reformulation of (ii). Now let $k_0 \in K$ be given and U a neighbourhood of $\varphi(k_0)$. Choose a $g \in W$ with $\|g\| = \|g(\varphi(k_0))\| = 1$, $g(l) = 0$ for $l \notin U$, and consider

$$\tilde{U} := \{k \mid \|(T^{-1}g)(k)\| > \beta\}.$$

$l \notin U$ yields $\|(T^{-1}g)(\varphi^{-1}(k))\| \leq \beta$ by definition so that $\varphi(\tilde{U}) \subset U$.

We also have $k_0 \in \tilde{U}$ since otherwise

$$1 = \|g(l_0)\| = \|T(T^{-1}g)(l_0)\| \leq \alpha\|T^{-1}g\| + (1 + \tau)\beta \\ \leq (\alpha + \beta)(1 + \tau).$$

This proposition motivates the following

2.3. DEFINITION. Let Y_1, \dots, Y_s be Banach spaces. By $\alpha(Y_1, \dots, Y_s)$ we mean the infimum of the numbers $\alpha > 0$ with the following property: There exists a $\tau > 0$ such that $K \xrightarrow[T]{\alpha} L$ for all Banach spaces X_1, \dots, X_r , all locally compact spaces $K_1, \dots, K_r, L_1, \dots, L_s$ and all τ -isomorphism $T: V \rightarrow W$ (notation as above).

As an immediate reformulation of 2.2 we then have

2.4. THEOREM. Suppose that $\alpha(X_1, \dots, X_r) + \alpha(Y_1, \dots, Y_s) < 1$. Then there exists a $\tau > 0$ such that, whenever there is a τ -isomorphism

$$T: C_0(K_1, \dots, K_r; X_1, \dots, X_r) \rightarrow C_0(L_1, \dots, L_s; Y_1, \dots, Y_s),$$

then necessarily $K_1 \dot{\cup} \dots \dot{\cup} K_r \cong L_1 \dot{\cup} \dots \dot{\cup} L_s$.

Proof. Choose $\alpha > \alpha(X_1, \dots, X_r)$, $\beta > \alpha(Y_1, \dots, Y_s)$ such that $\alpha + \beta < 1$. By definition there is a $\tau > 0$ such that the existence of $T: V \rightarrow W$ yields $K \xrightarrow[T]{\alpha} L$ and $L \xrightarrow[T^{-1}]{\beta} K$. Without loss of generality we may assume that $(\alpha + \beta)(1 + \tau) < 1$ so that Proposition 2.2 applies.

We now are going to investigate further properties of the mapping φ of 2.2.

Let T be given as in 2.2, suppose that $(\alpha + \beta)(1 + \tau) < 1$, and apply 2.2 (iii) to obtain φ .

Let $k \in K$ be arbitrarily given, say $k \in K_\rho$ and $\varphi(k) =: l \in L_\sigma$. We choose a fixed $h \in C_0 K$ with support in K_ρ and $\|h\| = h(k) = 1$ and define

$$S_k: X_\rho \rightarrow Y_\sigma$$

by $S_k(x) := (Tx^\sim)(l)$, where x^\sim denotes the function

$$x^\sim(k) := \begin{cases} h(k)x & \text{for } k \in K_\rho, \\ 0 & \text{otherwise.} \end{cases}$$

Clearly S_k is an operator with $\|S_k\| \leq 1 + \tau$.

Similarly we define $R_l: Y_\sigma \rightarrow X_\rho$, this time using a fixed $\hat{h} \in C_0 L$ and T^{-1} instead of T . Fix an $x \in X_\rho$. We have

$$[T(x^\sim) - S_k(x)^\sim](l) = 0$$

so that

$$\|T^{-1}[T(x^\sim) - S_k(x^\sim)](k)\| \leq \beta \|Tx^\sim - S_k(x^\sim)\| \leq 2(1 + \tau)\beta \|x\|.$$

But this means just

$$\|x - R_l S_k x\| \leq 2(1 + \tau)\beta \|x\|,$$

and similarly $\|y - S_k R_l y\| \leq 2(1 + \tau)\alpha \|y\|$ is valid. It is then a simple exercise to show that this implies that S_k is an isomorphism with

$$\|S_k\| \|S_k^{-1}\| \leq \hat{\alpha} := (1 + \tau)^2 \min\{(1 - 2(1 + \tau)\alpha)^{-1}, (1 - 2(1 + \tau)\beta)^{-1}\},$$

provided that α and β are smaller than $1/[2(1 + \tau)]$. Thus $d(X_\rho, Y_\sigma) \leq \hat{\alpha}$, and consequently $d(Y_\sigma, Y_{\sigma'}) \leq \hat{\alpha}^2$ if $k, k' \in K_\rho$ are given such that $\varphi(k) \in L_\sigma$, $\varphi(k') \in L_{\sigma'}$. In particular, if the $d(Y_\sigma, Y_{\sigma'})$ are all greater than $\hat{\alpha}^2$, then all $k \in K_\rho$ are mapped to the same L_σ , i.e. $\varphi(K_\rho) \subset L_\sigma$. Considering φ^{-1} we get $\varphi^{-1}(L_\sigma) \subset K_\rho$, so that φ induces a homeomorphism between K_ρ and L_σ . Summing up, we have proved the

2.5. THEOREM. *Suppose that*

$$\alpha_X := \alpha(X_1, \dots, X_r), \quad \alpha_Y := \alpha(Y_1, \dots, Y_l) < \frac{1}{2}$$

and that

$$d(X_\rho, X_{\rho'}), d(Y_\sigma, Y_{\sigma'}) > \min\{(1 - 2\alpha_X)^{-2}, (1 - 2\alpha_Y)^{-2}\} =: \hat{\alpha}_0^2.$$

Then for every $\varepsilon > 0$ there is a $\tau > 0$ such that the existence of a τ -isomorphism $T: V \rightarrow W$ implies that $r = s$ and that there exists a bijection $\omega: \{1, \dots, r\} \rightarrow \{1, \dots, s\}$ such that $K_\rho \cong L_{\omega(\rho)}$ and $d(X_\rho, Y_{\omega(\rho)}) \leq \hat{\alpha}_0 + \varepsilon$ for every ρ .

Proof. Choose $\alpha > \alpha_X$, $\beta > \beta_Y$, $\tau > 0$ such that the above considerations apply (with $\hat{\alpha} = \hat{\alpha}_0 + \varepsilon$).

3. Operators which are nearly multipliers and the constant $\alpha_2(X)$. In this section we discuss the class $\text{Mult}_\varepsilon(X)$ (see Def. 1.6(ii)). We determine $\alpha_2(X)$ for some classes of spaces and prepare the proof of our main results in §4. The following assertions are immediate:

3.1. LEMMA.

- (i) $\text{Mult}_\varepsilon(X)$ is a closed subset of the ball with radius $1 + \varepsilon$.
- (ii) $\bigcap_{\varepsilon > \varepsilon_0} \text{Mult}_\varepsilon(X) = \text{Mult}_{\varepsilon_0}(X)$ for every $\varepsilon_0 \geq 0$, so that in particular $\bigcap_{\varepsilon > 0} \text{Mult}_\varepsilon(X) \subset \text{Mult}(X)$.

We start by investigating two classes of examples.

3.2. THEOREM. *Let X be finite-dimensional.*

(i) *For every $\alpha > 0$ there is an $\varepsilon > 0$ such that*

$$\text{Mult}_\varepsilon(X) \subset (\text{Mult}(X))_\alpha.$$

(ii) *Suppose that in addition $\text{Mult}(X)$ is trivial. Then $\alpha_2(X) = 0$.*

Proof. (i) follows from a simple compactness argument by using the compactness of $\{R \mid R \in \text{Mult}(X), \|R\| \leq 1\}$ and 3.1(ii). (ii) is a consequence of (i).

3.3. THEOREM. *Let X be uniformly smooth; this means that for every $\eta > 0$ there is an $\tilde{\varepsilon} > 0$ such that $\|x - y\| \leq \tilde{\varepsilon}$ implies that $\|x + y\| \geq \|x\| + \|y\| - \eta\|x - y\|$ (see [13], p. 147). Then $\alpha_2(X) = 0$.*

Proof. The proof is sketched in several steps. Details are left to the reader

1. For every two-dimensional Banach space X_0 there exist $x_0, y_0 \in X_0$ with norm one such that

$$\{\tilde{x} \mid \|\tilde{x}\| \leq 1\} \subset \{\alpha x_0 + \beta y_0 \mid |\alpha|, |\beta| \leq 1\}$$

(choose x_0, y_0 to be an Auerbach-system; see 1.c.3 in [16]).

2. Let $R: X \rightarrow X$ be an operator on any Banach space X such that, for a fixed $\varepsilon \geq 0$, we have $\|Rx - \lambda_x x\| \leq \varepsilon$ for each x in the unit ball, where $\lambda_x \in \mathbf{K}$ may depend on x (i.e. every x is nearly an eigenvector). Then there is a $\lambda \in \mathbf{K}$ such that $\|R - \lambda \text{Id}_X\| \leq 18\varepsilon$. (First let x, y be vectors as in “1.”. With $\lambda_x, \lambda_y, \lambda \in \mathbf{K}$ such that

$$\|Rx - \lambda_x x\|, \quad \|Ry - \lambda_y y\|, \quad \left\| R \left(\frac{x+y}{\|x+y\|} \right) - \lambda \left(\frac{x+y}{\|x+y\|} \right) \right\| \leq \varepsilon$$

it follows that $\|(\lambda_x - \lambda)x + (\lambda_y - \lambda)y\| \leq 4\varepsilon$ so that $|\lambda_x - \lambda_y| \leq 8\varepsilon$.)

Now let x, y be arbitrary with $\|x\| = \|y\| = 1$. We choose λ_x, λ_y with $\|Rx - \lambda_x x\|, \|Ry - \lambda_y y\| \leq \varepsilon$, and we claim that $|\lambda_x - \lambda_y| \leq 18\varepsilon$.

To this end, let $x_0, y_0 \in \text{lin}\{x, y\}$ be as in “1.” (we note that without loss of generality x and y are linearly independent). By the first part of the proof we know that $|\lambda_{x_0} - \lambda_{y_0}| \leq 8\varepsilon$ is true for the associated λ 's. Therefore the numbers $\|(R - \lambda_{x_0} \text{Id}_X)(z)\|$ are bounded by 8ε for z in $\{\alpha x_0 + \beta y_0 \mid |\alpha|, |\beta| \leq 1\}$, and in particular we have

$$\|Rx - \lambda_{x_0} x\|, \quad \|Ry - \lambda_{x_0} y\| \leq 8\varepsilon$$

so that $|\lambda_x - \lambda_y| \leq 18\varepsilon$.

Now choose any x_0 with $\|x_0\| = 1$ and define $\lambda := \lambda_{x_0}$. Then obviously $\|R - \lambda \text{Id}_X\| \leq 18\varepsilon$.

3. Let $\varepsilon \geq 0$. For any x we define

$$I(x, \varepsilon; X) := \bigcap B(x_0, r(1 + \varepsilon)),$$

where the intersection runs over all $x_0 \in X$, $r \geq 0$ such that $\Theta x \in B(x_0, r)$ for every Θ with $|\Theta| = 1$. Then clearly $R \in \text{Mult}_\varepsilon(X)$ iff $Rx \in I(x, \varepsilon; X)$ for every x with $\|x\| \leq 1$.

4. Let X be a Banach space. Suppose that for every $\alpha > 0$ there is an $\varepsilon > 0$ such that

$$I(x, \varepsilon; X) \subset (\mathbf{K}x)_\alpha$$

for any x with $\|x\| \leq 1$. Then $\alpha_2(X) = 0$ (as an immediate consequence of "2." and "3.").

5. Let Y be a two-dimensional smooth Banach space. Then for every $\alpha > 0$ there is an $\varepsilon > 0$ with

$$I(x, \varepsilon; Y) \subset (\mathbf{K}x)_\alpha \quad (\text{all } x \text{ with } \|x\| \leq 1).$$

This follows from a compactness argument and the fact that $I(x, 0; Y) \subset \mathbf{K}x$ for smooth spaces.

6. Let $\tilde{\varepsilon}:]0, 1] \rightarrow]0, 1]$ be a fixed function. In \mathcal{B} , the set of all two-dimensional Banach spaces provided with the distance $\log d(\cdot, \cdot)$, we consider the subset $\mathcal{B}_{\tilde{\varepsilon}}: Y \in \mathcal{B}_{\tilde{\varepsilon}}$ iff

$$\|x - y\| \leq \tilde{\varepsilon}(\eta) \Rightarrow \|x + y\| \geq \|x\| + \|y\| - \eta\|x - y\|$$

(all $\eta \in]0, 1]$, $x, y \in Y$).

Then $\mathcal{B}_{\tilde{\varepsilon}}$ is closed in \mathcal{B} and thus compact.

7. Let $\tilde{\varepsilon}$ be as in "6.". Then for every $\alpha > 0$ there is an $\varepsilon > 0$ such that

$$I(x, \varepsilon; Y) \subset (\mathbf{K}x)_\alpha$$

uniformly for all Y in $\mathcal{B}_{\tilde{\varepsilon}}$ and $x \in Y$ with $\|x\| \leq 1$ (this follows from "5." and "6." and a continuity argument).

8. Now let X be a uniformly smooth Banach space. Then there is, by definition, a function $\tilde{\varepsilon}:]0, 1] \rightarrow]0, 1]$ such that all two-dimensional subspaces of X belong to $\mathcal{B}_{\tilde{\varepsilon}}$. Let $\alpha > 0$ be given and ε as in "7.". Then, for $x \in X$ with $\|x\| \leq 1$, we have

$$I(x, \varepsilon; Y) \subset (\mathbf{K}x)_\alpha$$

for every two-dimensional subspace which contains x . But this implies $I(x, \varepsilon; X) \subset (\mathbf{K}x)_\alpha$.

Now a combination of "8." and "4." proves the theorem.

We return to the spaces

$$C(K_1, \dots, K_r; X_1, \dots, X_r), \quad C(L_1, \dots, L_s; Y_1, \dots, Y_s),$$

and we adopt the notation of §2. Since for every bounded and continuous $h: K \rightarrow \mathbb{K}$ with $\|h\| \leq 1$ the operator $M_h: V \rightarrow V$, defined by $f \mapsto hf$, lies in $\text{Mult}_0(V)$, the operator TM_hT^{-1} lies in $\text{Mult}_{2\tau+\tau^2}(W)$ for every τ -isomorphism T . The proof of these facts is straightforward.

This observation which will be of crucial importance later necessitates the study of the $\text{Mult}_\varepsilon(V)$, $\text{Mult}_\varepsilon(W)$.

3.4. PROPOSITION. *Let $\varepsilon \geq 0$ and $R \in \text{Mult}_\varepsilon(V)$.*

(i) *Fix $\rho \in \{1, \dots, r\}$ and $k \in K_\rho$. With x^\sim as in §2 (cf. the discussion leading to Th. 2.5) we define $R_k: X_\rho \rightarrow X_\rho$ by $x \mapsto (Rx^\sim)(k)$. Then $R_k \in \text{Mult}_\varepsilon(X_\rho)$.*

(ii) *For $f \in V$ with $f(k) = 0$ we have $\|(Rf)(k)\| \leq \varepsilon\|f\|$.*

(iii) *For every f and every k we have*

$$\|R_k f(k)\| - 2\varepsilon\|f\| \leq \|(Rf)(k)\| \leq \|R_k f(k)\| + 2\varepsilon\|f\|.$$

Proof.

(i) $\|\Theta x - x_0\| \leq 1$ (all $|\Theta| = 1$) yields $\|\Theta x^\sim - x_0^\sim\| \leq 1$ and thus $\|Rx^\sim - x_0^\sim\| \leq 1 + \varepsilon$. Evaluating this at k gives $\|R_k x - x_0\| \leq 1 + \varepsilon$.

(ii) Let $x \in X_\rho$ with $\|x\| = 1$ be given. For $\eta > 0$ we choose an $h \in C_0 K$ with support in $K_\rho \cap \{\|f\| < \eta\}$ such that $h(k) = \|h\| = 1 - \eta$ (where $f \in V$, $f(k) = 0$, $\|f\| \leq 1$). Define hx by

$$(hx)(k') = \begin{cases} h(k')x & \text{if } k' \in K_\rho, \\ 0 & \text{otherwise.} \end{cases}$$

Then $\|\Theta f - hx\| \leq 1$ for every Θ with $|\Theta| = 1$ so that $\|Rf - hx\| \leq 1 + \varepsilon$. The special choice $x := (Rf)(k)/\|Rf(k)\|$ gives, by considering $(Rf - hx)(k)$,

$$1 - \eta + \|Rf(k)\| \leq 1 + \varepsilon.$$

This proves that $\|Rf(k)\| \leq \varepsilon$ since $\eta > 0$ was arbitrary.

(iii) Let $f \in V$ and $k \in K$ be given. $f - f(k)^\sim$ vanishes at k so that by (ii) $\|(Rf)(k) - R_k f(k)\| \leq 2\varepsilon\|f\|$.

Using this preparation we can prove that property of the $R \in \text{Mult}_\varepsilon(X)$ which will be very important in §4.

3.5. COROLLARY. *Let $\alpha, \eta, \varepsilon > 0$ be given and $k \in K$. We assume that*

(i) $R \in \text{Mult}_\varepsilon(V)$,

(ii) *there is a $g_0 \in V$ such that $\|g_0\| = 1$, $\|Rg_0(k)\| \geq 1 - \eta$,*

(iii) $\text{Mult}_\varepsilon(X_\rho) \subset (\mathbf{K}\text{Id}_{X_\rho})_\alpha$.

Then $\|(Rf)(k)\| \geq \|f(k)\| - (2\alpha + 4\varepsilon + \eta)\|f\|$ for every $f \in V$ (roughly: with f also Rf is large at k , and this can be asserted at every k where Rg_0 is large for at least one g_0).

Proof. 3.4(iii), applied for g_0 , gives $\|R_k\| \geq 1 - \eta - 2\varepsilon$. Since $R_k \in \text{Mult}_\varepsilon(X_\rho) \subset (\mathbf{K}\text{Id}_{X_\rho})_\alpha$ we find a $\lambda \in \mathbf{K}$ with $\|R_k - \lambda \text{Id}_{X_\rho}\| \leq \alpha$. Here necessarily $|\lambda| \geq 1 - \eta - 2\varepsilon - \alpha$. Now let $f \in V$ be given. Again by 3.4(iii) we have

$$\begin{aligned} \|(Rf)(k)\| &\geq \|R_k f(k)\| - 2\varepsilon\|f\| \\ &\geq \|\lambda f(k)\| - 2\varepsilon\|f\| - \alpha\|f(k)\| \\ &\geq \|f(k)\| - (2\alpha + 4\varepsilon + \eta)\|f\|. \end{aligned}$$

4. An estimate of the constant $\alpha(X_1, \dots, X_r)$ and some important consequences. The following theorem is doubtlessly the most important one in this paper. Only by using this we are able to apply the results of §2 (by having in mind, of course, our knowledge about the $\alpha_1(X)$, $\alpha_2(X)$).

4.1. THEOREM. $\alpha(X_1, \dots, X_r) \leq \max_\rho \min\{\alpha_1(X_\rho), 2\alpha_2(X_\rho)\}$.

Proof. For the sake of convenience we prove the theorem for a family Y_1, \dots, Y_s (which avoids the change of notation).

Let $\alpha > \max_\sigma \min\{\alpha_1(Y_\sigma), 2\alpha_2(Y_\sigma)\}$ arbitrary. We assume that the Y_σ are given such that

$$\begin{aligned} \alpha &> \alpha_1(Y_\sigma) \quad \text{for } \sigma = 1, \dots, \hat{s}, \quad \text{and} \\ \alpha &> 2\alpha_2(Y_\sigma) \quad \text{for } \sigma = \hat{s} + 1, \dots, s. \end{aligned}$$

It has to be shown that there is a $\tau > 0$ such that $K \xrightarrow[\tau]{\alpha} L$ whenever $X_1, \dots, X_r, K_1, \dots, K_r, L_1, \dots, L_s$ are given and there exists a τ -isomorphism $T: V \rightarrow W$ (notation as in §2).

At first we choose any α', α'' with $\alpha_2(Y_\sigma) < \alpha' < \alpha'' < \alpha/2$ for $\sigma = \hat{s} + 1, \dots, s$ and $\alpha_1(Y_\sigma) < 2\alpha''$ for $\sigma = 1, \dots, \hat{s}$, and then an $\varepsilon > 0$ with

$$— 2\alpha' + 4\varepsilon < 2\alpha'',$$

- $1 - \varepsilon \leq \|x_0\| \leq 1 + \varepsilon$, $\|x_0 + \Theta x\| \leq 1 + \varepsilon$ for all $|\Theta| = 1$ implies $\|x\| \leq 2\alpha''$; all $x_0, x \in Y_\sigma$, where $\sigma = 1, \dots, \hat{s}$,
- $\text{Mult}_\varepsilon(Y_\sigma) \subset (\mathbf{K} \text{Id}_{Y_\sigma})_{\alpha'}$ for $\sigma = \hat{s} + 1, \dots, s$.

Finally we suppose that $\tau > 0$ is so small that

- $(1 + \tau)^2 \leq 1 + \varepsilon$,
- $1 - 2\tau - \tau^2/(1 + \tau) \geq 1 - \varepsilon$,
- $(2\alpha' + 4\varepsilon + \eta_\tau + \tau)(1 + \tau) \leq 2\alpha''$,
where $\eta_\tau := (1 - 2\tau - \tau^2)/(1 + \tau)^2 - 1$.

Now let X_1, \dots, X_r , K_1, \dots, K_r , L_1, \dots, L_s and a τ -isomorphism $T: V \rightarrow W$ be given. We claim that $K \xrightarrow[T]{\alpha} L$, i.e.

$$\alpha_k := \inf_l \sup \{ \|(Tf)(l)\| \mid f \in V, f(k) = 0, \|f\| \leq 1 \} < \alpha$$

for every $k \in K$. We will show that $t < \alpha_k$ implies $t \leq 2\alpha''$ so that $\alpha_k \leq 2\alpha'' < \alpha$. Let k and such a t be given and fix any function $f_0 \in V$ with $\|f_0\| = \|f_0(k)\| = 1$. The set $\tilde{L}_\tau := \{l \mid \|Tf_0(l)\| \geq \tau\}$ is compact so that we find $f_1, \dots, f_n \in V$ with $\|f_i\| \leq 1$, $f_i(k) = 0$, $\max_i \|Tf_i(l)\| > t$ for every $l \in \tilde{L}_\tau$.

Since all these f_i vanish at k_0 it is possible to choose an $h \in C_0 K$ with $\|hf_i\| \leq \tau$, $\|h\| = -h(k) = 1$, $\|\Theta f_i - hf_0\| \leq 1 + \tau$ for $i = 1, \dots, n$ and $|\Theta| = 1$.

We have $\|f_0 - hf_0\| = 2$ so that there is an $l_0 \in L$ with $\|[Tf_0 - T(hf_0)](l_0)\| \geq 2/(1 + \tau)$ which yields

$$\|[T(hf_0)](l_0)\|, \quad \|(Tf_0)(l_0)\| \geq (1 - 2\tau - \tau^2)/(1 + \tau)$$

(note that $\|T(hf_0)\|, \|Tf_0\| \leq 1 + \tau$). Therefore in particular $l_0 \in \tilde{L}_\tau$, and we find an $i_0 \in \{1, \dots, n\}$ with $\|(Tf_{i_0})(l_0)\| > t$. l_0 lies in one of the L_σ , and we consider two cases.

Case 1. $\sigma \in \{1, \dots, \hat{s}\}$.

We know that $\|\Theta f_{i_0} - hf_0\| \leq 1 + \tau$ for all $|\Theta| = 1$, i.e. with $x_0 := [T(hf_0)](l_0)$, $x := (Tf_{i_0})(l_0)$ we have

$$\begin{aligned} (1 - 2\tau - \tau^2)/(1 + \tau) &\leq \|x_0\| \leq (1 + \tau)^2 \\ &t < \|x\| \\ \|\Theta x - x_0\| &\leq (1 + \tau)^2 \quad (\text{all } |\Theta| = 1). \end{aligned}$$

By the choice of τ this implies $\|x\| \leq 2\alpha''$, and thus $t \leq 2\alpha''$ as claimed.

Case 2. $\sigma \in \{\hat{s} + 1, \dots, s\}$.

The operator $M_h: V \rightarrow V$, $f \mapsto hf$, belongs to $\text{Mult}_0(V)$ so that $R := TM_h T^{-1}$ lies in $\text{Mult}_{2\tau+\tau^2}(W) \subset \text{Mult}_\varepsilon(W)$.

With $g_0 := Tf_0/\|Tf_0\|$ we have $Rg_0 = T(hf_0)/\|Tf_0\|$ and consequently $\|(Rg_0)(l_0)\| \geq (1 - 2\tau - \tau^2)/(1 + \tau)^2 = 1 - \eta_\tau$. On the other hand we know that $\|hf_{i_0}\| \leq \tau$ so that $\|R(Tf_{i_0})\| = \|T(hf_{i_0})\| \leq \tau(1 + \tau)$.

Now Corollary 3.5 comes into play. Since $R \in \text{Mult}_\varepsilon(W)$ and $\text{Mult}_\varepsilon(Y_\sigma) \subset (\mathbf{K}\text{Id}_{Y_\sigma})_{\alpha'}$ it follows that

$$\begin{aligned} t < \|Tf_{i_0}(l_0)\| &\leq (2\alpha' + 4\varepsilon + \eta_\tau)\|Tf_{i_0}\| + \|R(Tf_{i_0})\| \\ &\leq (2\alpha' + 4\varepsilon + \eta_\tau + \tau)(1 + \tau) \leq 2\alpha'', \end{aligned}$$

and this completes the proof.

We now turn to some interesting *consequences* which can be derived very simply by combining Theorem 2.4 or Theorem 2.5 with Theorem 3.2, Theorem 3.3, the remarks following Definition 1.5 and, of course, Theorem 4.1.

4.2. THEOREM. *Suppose that $\min\{\alpha_1(X), 2\alpha_2(X)\} < \frac{1}{2}$ for $X = X_\rho$ ($\rho = 1, \dots, r$) and $X = Y_\sigma$ ($\sigma = 1, \dots, s$).*

Then there is a $\tau > 0$ such that the existence of a τ -isomorphism

$$T: C_0(K_1, \dots, K_r; X_1, \dots, X_r) \rightarrow C_0(L_1, \dots, L_s; Y_1, \dots, Y_s)$$

implies that $K_1 \dot{\cup} \dots \dot{\cup} K_r \cong L_1 \dot{\cup} \dots \dot{\cup} L_s$.

4.3. THEOREM. *Suppose that $\min\{\alpha_1(X), 2\alpha_2(X)\} < \frac{1}{2}$. Then there is a $\delta > 0$ such that*

$$d(C_0(K, X), C_0(L, X)) \geq 1 + \delta$$

for arbitrary K, L with $K \not\cong L$.

4.4. THEOREM. *Suppose that $\alpha_1(X) = \alpha_2(X) = 0$ for $X = X_\rho$ ($\rho = 1, \dots, r$) and $X = Y_\sigma$ ($\sigma = 1, \dots, s$). Suppose further that $d(X_\rho, X_{\rho'}), d(Y_\sigma, Y_{\sigma'}) > 1$ for $\rho \neq \rho', \sigma \neq \sigma'$. Then for every $\delta > 0$ there exists a $\tau > 0$ such that the following holds: whenever*

$$T: C_0(K_1, \dots, K_r; X_1, \dots, X_r) \rightarrow C_0(L_1, \dots, L_s; Y_1, \dots, Y_s)$$

is a τ -isomorphism, then $r = s$, and there is a bijection $\omega: \{1, \dots, r\} \rightarrow \{1, \dots, s\}$ with $K_\rho \cong L_{\omega(\rho)}$ and $d(X_\rho, Y_{\omega(\rho)}) \leq 1 + \delta$ for every ρ .

Note. In particular this can be applied to such situations where for every $\tau > 0$ τ -isomorphisms exist. Then $d(X_\rho, Y_{\omega(\rho)}) = 1$ for all ρ .

5. Isomorphic Banach-Stone theorems. We now state our main results on isomorphic Banach-Stone theorems. They appear as special cases of our more general approach.

5.1. THEOREM. *Each of the following conditions implies that X has the isomorphic Banach-Stone property:*

- (i) $\min\{\alpha_1(X), 2\alpha_2(X)\} < \frac{1}{2}$;
- (ii) X is of the form $\prod_{\rho=1}^{\infty} X_{\rho}^{n_{\rho}}$, where $d(X_{\rho}, X_{\rho'}) \geq 1 + \delta_0$ for a suitable $\delta_0 > 0$ and all $\rho \neq \rho'$. Further $\min n_{\rho} = 1$ and

$$\max_{\rho} \min\{\alpha_1(X_{\rho}), 2\alpha_2(X_{\rho})\} < 1 - (1 + \delta_0)^{-1/2} \approx \delta_0/2.$$

Proof. One only has to note that $C_0(K, \prod X_{\rho}^{n_{\rho}})$ can be identified with $C_0(n_1 K, \dots, n_r K; X_1, \dots, X_r)$; here nK stands for the disjoint union of n copies of K .

5.2. THEOREM. *Each of the following conditions implies that X has the strong isomorphic Banach-Stone property:*

- (i) X is uniformly smooth;
- (ii) X is uniformly convex;
- (iii) X is finite-dimensional with $\text{Mult}(X) = \mathbf{K} \text{Id}_X$;
- (iv) $\min\{\alpha_1(X), 2\alpha_2(X)\} = 0$;
- (v) $\alpha(X) = 0$.

Proof. In view of Theorem 3.3, the remark following Definition 1.5, Theorem 3.2(ii), and Theorem 4.1 it suffices to prove (v). Let $\varepsilon > 0$ be given and $\alpha > 0$ so small that $(1 + \alpha)^2/(1 - 2\alpha(1 + \alpha)) \leq 1 + \varepsilon$. Choose $\tau > 0$ for this α according to 2.3, where $\tau \leq \alpha$. Now let any τ -isomorphism $T: C_0(K, X) \rightarrow C_0(L, X)$ be given. With φ and $(S_k)_{k \in K}$ as in the discussion leading to Theorem 2.5 we define $T_k := S_k$.

The T_k are isomorphisms, and

$$\|T_k\| \|T_k^{-1}\| \leq (1 + \tau)^2/(1 - 2(1 + \tau)\alpha) \leq 1 + \varepsilon.$$

It remains to prove that $\|Tf(\varphi(k)) - T_k f(k)\| \leq \varepsilon$ for all k and f with $\|f\| \leq 1$. Let such f and k be given. With $\varphi(k) =: l$ we have $k \xrightarrow{\alpha} l$ and $[f(k)^{\sim} - f](k) = 0$. Thus

$$\|T_k(f(k)) - Tf(\varphi(k))\| = \|T[f(k)^{\sim} - f](l)\| \leq 2\alpha \leq \varepsilon$$

which completes the proof.

5.3. THEOREM. *Let X be a Banach space which can be written as $X = \prod_{\rho=1}^r X_{\rho}^{n_{\rho}}$, where $d(X_{\rho}, X_{\rho'}) > 1$ for $\rho \neq \rho'$ and $\alpha(X_{\rho}) = 0$ for $\rho = 1, \dots, r$. We emphasize that this is possible for every finite-dimensional space (combine Theorem 3.2 (ii) with Proposition 5.1 in [6]) but also surely for many other spaces.*

- (i) X has the isomorphic Banach-Stone property iff $\min n_p = 1$.
- (ii) X has the strong isomorphic Banach-Stone property iff $r = n_1 = 1$.

Proof. (i) If $\min n_p = 1$ then X has the Banach-Stone property by Theorem 5.1. If $\min n_p > 1$ then X does not even have the usual Banach-Stone property since there are non-homeomorphic K, L with $nK \cong nL$ for every $n \geq 2$ (so that for these K, L we have $K \not\cong L$ but $C_0(K, X) \cong C_0(L, X)$); see [4], Corollary 11.15.

(ii) One implication is proved in Theorem 5.2.

Conversely, if $r > 1$ or $r = 1$ and $n_1 > 1$ then X can be written as $J \oplus_\infty J^\perp$ for suitable nontrivial subspaces J, J^\perp (\oplus_∞ denotes the L^∞ -direct sum). With $K = L = \{1, 2\}$ we identify $C_0(K, X)$ and $C_0(L, X)$ with $J \oplus_\infty J \oplus_\infty J^\perp \oplus_\infty J^\perp$. An isometry T on this space is defined by $(j_1, j_2, j_1^\perp, j_2^\perp) \mapsto (j_1, j_2, j_2^\perp, j_1^\perp)$. It is plain that for no choice of $\varphi: \{1, 2\} \rightarrow \{1, 2\}$ and isomorphism T_k can the map T be approximated such that $Tf(\varphi(k)) - T_k(f(k))$ is small for all $k \in \{1, 2\}$ and f with $\|f\| = 1$ (since $Tf(\varphi(k))$ can be large with $f(k)$ small). Therefore X (and more generally every X with a nontrivial M -summand) fails to have the strong isomorphic Banach-Stone property.

6. Isomorphisms which are close to isometries. Let X be a Banach space with $\alpha(X) = 0$ and K a compact space.

We choose an $\alpha > 0$ and an admissible $\tau > 0$ for this α (as in 2.3) and suppose that $T: C(K, X) \rightarrow C(K, X)$ is a τ -isomorphism. We are going to investigate whether T is close to an isometry. Let φ be defined as in Theorem 2.4, and for simplicity we will assume for the moment that $\varphi = \text{Id}_K$. Thus T is a τ -isomorphism with

$$f(k) = 0 \Rightarrow \|(Tf)(k)\| \leq \alpha \|f\|.$$

We consider maps $T_k: X \rightarrow X$, $x \mapsto (T\mathbf{x})(k)$, where \mathbf{x} denotes the constant function x (the T_k are just the S_k of §2, where $h = 1$). Then, as it is shown in §2, the T_k are isomorphisms on X with

$$\|T_k\| \leq 1 + \tau, \quad \|T_k^{-1}\| \leq \frac{1 + \tau}{1 - 2(1 + \tau)\alpha}.$$

Now consider the operator $\prod T_k$.

Since $k \mapsto T_k$ is continuous with respect to the strong operator topology (this can not be guaranteed in the general situation of §2) this operator is well-defined on $C(K, X)$.

Suppose it would be possible to find isometries I_k on X with $T_k \approx I_k$ and $k \mapsto I_k$ strongly continuous (so that $\prod I_k$ would be a well-defined isometry on $C(K, X)$). Then $T \approx \prod T_k$ and $\prod T_k \approx \prod I_k$, so that T would be close to an isometry.

Thus the main difficulty in this approximation problem is to approximate a continuous family of near-isometries continuously by isometries.

To treat this more precisely it is useful to introduce the following

6.1. DEFINITION. Let X be a Banach space and $\eta > 0$. By $\text{Iso}_\eta(X)$ we mean the collection of all isomorphisms $T: X \rightarrow X$ which are η -isomorphisms, i.e. $(1/(1+\eta))\|x\| \leq \|Tx\| \leq (1+\eta)\|x\|$.

We suppose that for every $\varepsilon > 0$ there is an $\eta > 0$ with

$$\text{Iso}_\eta(X) \subset (\text{Iso}_0(X))_\varepsilon,$$

which is certainly true for finite-dimensional spaces (note that $\bigcap_{\eta>0} \text{Iso}_\eta(X) = \text{Iso}_0(X)$ and apply a compactness argument).

Then returning to the preceding discussion, we could choose $I_k \in \text{Iso}_0(X)$ for every k such that $\|T_k - I_k\| \leq \varepsilon$ (provided that τ was so small that $T_k \in \text{Iso}_\eta(X)$). There seems, however, to be no general strategy to guarantee that the choice of these I_k leads to a *continuous* family $k \mapsto I_k$.

We restrict ourselves by giving a sufficient condition:

6.2. THEOREM. Let X be a Banach space with $\alpha(X) = 0$. Further suppose that for every $\tilde{\varepsilon} > 0$ there is an $\eta > 0$ such that

- (i) $\text{Iso}_\eta(X) \subset (\text{Iso}_0(X))_{\tilde{\varepsilon}}$,
- (ii) there is a mapping $\mu: \text{Iso}_\eta(X) \rightarrow \text{Iso}_0(X)$ which is continuous with respect to the strong operator topology and for which $\|\mu(R) - R\| \leq 2\tilde{\varepsilon}$ for every R .

Then $C(K, X)$ has the AI-property for every compact space K .

Proof. Let $\varepsilon > 0$ be given. We choose $\mu > 0$ and η for $\tilde{\varepsilon} := \varepsilon/3$.

Now we select $\alpha > 0$ with

$$1 + \alpha \leq 1 + \eta, \quad \frac{1 + \alpha}{1 - 2\alpha(1 + \alpha)} \geq \frac{1}{1 + \eta}, \quad 2\alpha \leq \frac{\varepsilon}{3}$$

and finally $\tau > 0$ with $\tau \leq \alpha$ for this α as in Definition 2.3. Let T_0 be a τ -isomorphism on $C(K, X)$. We define $T := T_0 \circ I_\varphi$, where φ is the homeomorphism defined by T_0 and I_φ denotes the isometry $f \mapsto f \circ \varphi^{-1}$ on $C(K, X)$. Then T is a τ -isomorphism and we have $k \xrightarrow[T]{\alpha} k$ for every k .

Consider the T_k as above. Then $T_k \in \text{Iso}_\eta(X)$ by construction, and $k \mapsto I_k := \mu(T_k)$ is a continuous family (continuous w.r.t. the strong operator topology) of isometries such that $\|\prod T_k - \prod I_k\| \leq \frac{2}{3}\varepsilon$. But $\|Tf(k) - T_k(f(k))\| \leq 2\alpha\|f\|$ (since $(f - \mathbf{f}(\mathbf{k}))(k) = 0$) so that $\|T - \prod T_k\| \leq 2\alpha \leq \varepsilon/3$. This proves that $\|T - \prod I_k\| = \|T_0 - (\prod I_k)I_\varphi^{-1}\| \leq \varepsilon$, and $(\prod I_k)(I_\varphi)^{-1}$ is an isometry on $C(K, X)$.

Therefore $C(K, X)$ has the AI-property.

It seems to be difficult to find examples. We only mention the following

6.3. THEOREM. *Let X be a finite-dimensional Banach space such that there are isometries I_1, \dots, I_n with $\text{Iso}_0(X) = \cup_i \{\Theta I_i \mid |\Theta| = 1\}$. (This is for example the case if the unit ball of X has only finitely many extreme points.) Further assume that $\text{Mult}(X)$ is trivial so that $\alpha(X) = 0$.*

Then $C(K, X)$ has the AI-property for every compact space K .

Proof. We first note: For every I with $\|I\| = 1$ and $\eta > 0$ there is a continuous map

$$\mu_I: (\{\Theta I \mid |\Theta| = 1\})_\eta \rightarrow \{\Theta I \mid |\Theta| = 1\}$$

with $\|\mu_I(R) - R\| \leq 2\eta$ for every R (this is trivial if $\mathbf{K} = \mathbf{R}$ and easy if $\mathbf{K} = \mathbf{C}$).

It is now immediate how Theorem 6.2 can be applied: choose a minimal number of I_1, \dots, I_n and consider only such $\tilde{\varepsilon}$ where the $(\{\Theta I_i \mid |\Theta| = 1\})_{\tilde{\varepsilon}}$ are pairwise disjoint so that μ can be defined by gluing together the μ_{I_i} .

As an important special case we mention the case $X = \mathbf{K}$. Here, however, it is easy to be more explicit, and one obtains

6.4. COROLLARY. *For every τ -isomorphism T on a CK-space there is an isometry I with $\|T - I\| \leq 3(\tau + \tau^2)$.*

Finally it should be noted that there seems to be no possibility to treat similarly the case of $C_0(K, X)$ with locally compact K . The difficulties arise from the fact that in the definition of the T_k we have to choose an element h of $C_0(K)$ which is one at k and this is not the same h for all k in the non-compact case. Of course something can be said if one has partitions of unity, but nothing up to now for

arbitrary K . It is even not obvious how to treat the case $X = \mathbf{K}$ in general. If $\mathbf{K} = \mathbf{R}$ one has an analogue of Corollary 6.4 for arbitrary C_0K -spaces; the case $\mathbf{K} = \mathbf{C}$ however leads to the following problem: Let K be locally compact and $T: C_0K \rightarrow C_0K$ a τ -isomorphism. Suppose that for every compact $D \subset K$ there is a continuous function $h_D: D \rightarrow \{\Theta \mid |\Theta| = 1\}$ with $\|Tf|_D - h_D f|_D\| \leq \varepsilon \|f\|$ for every f . Is it possible to find a continuous $h: K \rightarrow \{\Theta \mid |\Theta| = 1\}$ with $\|Tf - hf\| \leq 2\varepsilon \|f\|$ for all f ?

Acknowledgments. I am grateful to the referee for drawing my attention to Jarosz' paper [17]. Globevnik's paper [14] was made available to me by D. Werner. The results proved there—together with our theorems—have the surprising consequence that all complex L^1 -spaces have the strong isomorphic Banach-Stone property.

REFERENCES

- [1] E. M. Alfsen and E. G. Effros, *Structure in real Banach spaces* I, II, Ann. of Math., **96** (1972), 98–173.
- [2] D. Amir, *On isomorphisms of continuous function spaces*, Israel J. Math., **3** (1965), 205–210.
- [3] E. Behrends, *On the Banach-Stone theorem*, Math. Annalen, **233** (1978), 261–272.
- [4] ———, *M-Structure and The Banach-Stone Theorem*, Lecture Notes in Mathematics 736, Springer-Verlag 1979.
- [5] ———, *How to obtain vector-valued Banach-Stone theorems by using M-structure methods*, Math. Annalen, **261** (1982), 387–398.
- [6] ———, *Multiplier representations and an application to the problem whether $A \otimes_\varepsilon X$ determines A and/or X* , Math. Scand., **52** (1983), 117–144.
- [7] E. Behrends and M. Cambern, *On isomorphic Banach-Stone theorems*, (to appear: Studia Math.).
- [8] Y. Benyamini, *Small into-isomorphisms between spaces of continuous functions*, Proc. Amer. Math. Soc., **83** (1981), 479–485.
- [9] M. Cambern, *On isomorphisms with small bound*, Proc. Amer. Math. Soc., **18** (1967), 1062–1066.
- [10] ———, *Isomorphisms of $C_0(Y)$ onto $C(X)$* , Pacific J. Math., **35** (1970), 307–312.
- [11] ———, *Isomorphisms of spaces of norm-continuous functions*, Pacific J. Math., **116** (1985), 243–254.
- [12] H. B. Cohen, *A bound-two isomorphism between $C(X)$ Banach spaces*, Proc. Amer. Math. Soc., **50** (1975), 215–217.
- [13] M. M. Day, *Normed Linear Spaces*, 3rd ed., Springer-Verlag, 1973.
- [14] J. Globevnik, *On complex strict and uniform convexity*, Proc. Amer. Math. Soc., **47** (1975), 175–178.

- [15] K. Jarosz, *A generalization of the Banach-Stone theorem*, *Studia Math.*, **73** (1982), 33–39.
- [16] ———, *Multipliers in complex Banach spaces and structure of the unit ball*, preprint.
- [17] ———, *Perturbations of Banach Algebras*, *Lecture Notes in Mathematics* 1120, Springer-Verlag 1985.
- [18] J. Lindenstrauss and L. Tzafriri, *Classical Banach Spaces I*, Springer-Verlag, 1977.
- [19] G. Wodinski, *Multiplikatoren in komplexen Banachräumen*, Dissertation, Freie Universität Berlin, 1986.

Received April 29, 1987 and in revised form August 10, 1987.

I. MATHEMATISCHES INSTITUT
DER FREIEN UNIVERSITÄT BERLIN
ARNIMALLEE 3
D 1000 BERLIN 33
FEDERAL REPUBLIC OF GERMANY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1988 by Pacific Journal of Mathematics

William Charles Bauldry, Attila Mate and Paul Nevai, Asymptotics for solutions of systems of smooth recurrence equations	209
Ehrhard Behrends, Isomorphic Banach-Stone theorems and isomorphisms which are close to isometries	229
Fernanda Maria Botelho, Rotation sets of maps of the annulus	251
Edward Graham Evans, Jr. and Phillip Alan Griffith, Binomial behavior of Betti numbers for modules of finite length	267
Andrei Iordan, Pseudoconvex domains with peak functions at each point of the boundary	277
Zyun'iti Iwase, Dehn-surgery along a torus T^2 -knot	289
Marko Kranjc, Embedding 2-complexes in \mathbf{R}^4	301
Aloys Krieg, Eisenstein-series on real, complex, and quaternionic half-spaces	315
Masato Kuwata, Intersection homology of weighted projective spaces and pseudo-lens spaces	355
Carl Pomerance, András Sárközy and Cameron Leigh Stewart, On divisors of sums of integers. III	363
Martin Schechter, Potential estimates in Orlicz spaces	381