

Pacific Journal of Mathematics

DEHN-SURGERY ALONG A TORUS T^2 -KNOT

ZYUN'ITI IWASE

DEHN-SURGERY ALONG A TORUS T^2 -KNOT

ZYUN'ITI IWASE

A T^2 -knot means a 2-torus embedded in a 4-manifold. We define *torus T^2 -knots* in the 4-sphere S^4 as a generalization of torus knots in S^3 . We classify them up to equivalence and study the manifolds obtained by Dehn-surgery along them.

1. Introduction. Dehn-surgery, which was introduced by Dehn [1], plays an important role in knot theory and 3-dimensional manifold theory. The classical Dehn-surgery is the operation of cutting off the tubular neighborhood $N = S^1 \times D^2$ of a knot in S^3 and of pasting it back via an element of $\pi_0 \text{Diff } \partial N$, which is isomorphic to $\text{GL}_2\mathbf{Z}$. Gluck-surgery [2] along a 2-knot in S^4 is a 4-dimensional version of Dehn-surgery. In this version, $N = S^2 \times D^2$ and $\pi_0 \text{Diff } \partial N = (\mathbf{Z}/2)^3$. $(\mathbf{Z}/2)^2$ corresponds to the orientation reversing diffeomorphisms of S^2 and ∂D^2 . Therefore Gluck-surgery yields at most one new manifold from one 2-knot and it is a homotopy 4-sphere (see [2]). Another 4-dimensional version is Dehn-surgery along a 2-torus embedded in S^4 [7], which we call a T^2 -knot in this paper. In this version, $N = T^2 \times D^2$ and $\pi_0 \text{Diff } \partial N = \text{GL}_3\mathbf{Z}$. Countably many manifolds are obtained from one T^2 -knot. A manifold obtained by Gluck-surgery is also obtained by Dehn-surgery along a T^2 -knot (see Proposition 3.5). Dehn-surgery along an unknot is studied in [7], [9]. See also [3].

In this paper, we define a *torus T^2 -knot* which is analogous to the torus knots in the classical knot theory, and classify them up to equivalence. Then we study the manifolds obtained by Dehn-surgeries along them.

Dehn-surgery along a torus knot is studied by Moser [8].

THEOREM 1.1. (*Moser [8], Propositions 3.1, 3.2, 4.*) *Assume that a Dehn-surgery of type (α, β) is performed along $k(p, q)$, the torus knot of type (p, q) . Put $|\sigma| = |pq\beta - \alpha|$. The manifold obtained is denoted by M .*

(i) *If $|\sigma| \neq 0$, then M is a Seifert manifold with fibers of multiplicities $p, q, |\sigma|$.*

(ii) *If $|\sigma| = 1$, then M is a lens space $L(|\alpha|, \beta q^2)$.*

(iii) If $|\sigma| = 0$, then M is the connected sum of two lens spaces $L(p, q) \# L(q, p)$.

Our results are the following. Theorem 1.3 is a generalization of Theorem 1.1.

Let $k(p, q)$ be the torus knot of type (p, q) in S^3 , and B be a 3-ball contained in $S^3 - k(p, q)$. Define $S(k(p, q))$ and $\tilde{S}(k(p, q))$ by

$$\begin{aligned} (S^4, S(k(p, q))) &= ((S^3, k(p, q)) - \text{Int } B) \times S^1 \cup_{\text{id}} S^2 \times D^2, \\ (S^4, \tilde{S}(k(p, q))) &= ((S^3, k(p, q)) - \text{Int } B) \times S^1 \cup_{\tau} S^2 \times D^2 \end{aligned}$$

where id is the natural identification of $\partial B \times S^1$ with $S^2 \times \partial D^2$ and τ is the map $(u, v) \mapsto (uv, v)$ where we identify S^2 with the Riemann sphere and D^2 with the unit disk in \mathbb{C} .

PROPOSITION 1.2. (Lemma 2.6 and Proposition 2.9.) Any torus T^2 -knot is equivalent to one and only one of the following:

- (i) $S(k(p, q))$, $1 < p < q$, $\text{gcd}(p, q) = 1$;
- (ii) $\tilde{S}(k(p, q))$, $1 < p < q$, $\text{gcd}(p, q) = 1$;
- (iii) unknotted T^2 -knot.

THEOREM 1.3. (See Proposition 3.6, Remark 3.7, Proposition 3.9, Corollary 3.10, Proposition 3.11.) Assume that a Dehn-surgery of type (α, β, γ) (see Definition 3.2) is performed along $S(k(p, q))$ or $\tilde{S}(k(p, q))$. Put $\sigma = |pq\beta - \alpha|$. The manifold obtained is denoted by M .

(i) If $\sigma \neq 0$, then M is the total space of a good torus fibration over S^2 with one twin singular fiber of multiplicity p and two multiple tori of multiplicity q and σ .

(ii) If $\sigma = 1$, then M is $L_{|\alpha|}$ or $L'_{|\alpha|}$.

(iii) If $\sigma = 0$, then M is an irrational connected sum along circles [5] of either L_m or L'_m and $L(n, r) \times S^1$ for some m, n, r .

(iv) If $\gamma = 0$, then $M = (M_0 - \text{Int } B^3) \times S^1 \cup_h S^2 \times D^2$ where M_0 is the manifold obtained by a Dehn-surgery of type (α, β) along the torus knot of type (p, q) and $h = \text{id}$ (if $K = S(k(p, q))$), $h = \tau$ ($K = \tilde{S}(k(p, q))$). Especially if $(\alpha, \beta, \gamma) = (pq, 1, 0)$, $M = L_p \# L_q$.

(L_m and L'_m are the manifold defined in [9]. See also [3].)

We use standard notations. $N(X)$ means the tubular neighborhood of X . All the homology groups are with coefficients in \mathbb{Z} unless otherwise indicated.

2. Torus T^2 -knots.

DEFINITION 2.1. Let M^n (resp. N^{n-2}) be an n - (resp. $(n - 2)$ -) dimensional manifold. A submanifold K in M^n is called an N^{n-2} -knot in M^n if K is diffeomorphic to N^{n-2} . Let K, K' be two N^{n-2} -knots in M^n . K and K' are *equivalent* if there exists a diffeomorphism $h: (M^n, K) \rightarrow (M^n, K')$.

We will be mainly concerned with T^2 -knots in S^4 .

Recall that a T^1 -knot (i.e. a classical knot) K in S^3 is called unknotted if K bounds a disk D^2 in S^3 .

DEFINITION 2.2. A T^2 -knot K in S^4 is called *unknotted* if K bounds a solid torus $S^1 \times D^2$ in S^4 .

Any two unknotted T^2 -knots are equivalent.

REMARK 2.3. There exist three isotopy classes of embeddings $T^2 \rightarrow S^4$ such that their images are unknotted (see Theorem 5.3 in [7]). But we are considering a T^2 -knot itself, not its embedding map.

Recall that a T^1 -knot K in S^3 is called a torus knot if K is essentially embedded in $\partial N(U)$, where U is an unknotted T^1 -knot in S^3 .

DEFINITION 2.4. A T^2 -knot K in S^4 is called a *torus knot* if K is incompressibly embedded in $\partial N(U)$, where U is an unknotted T^2 -knot in S^4 .

LEMMA 2.5. *Let K, K' be incompressible 2-tori in T^3 such that $[K] = [K']$ in $H_2(T^3)$. Then, there is an ambient isotopy which carries K to K' .*

Proof. We may assume that $(T^3, K) = (S^1 \times S^1 \times S^1, S^1 \times S^1 \times \{*\})$ without loss of generality.

By Theorem VI.34 and VI.17 in [4], there exists a diffeomorphism $f: (T^3, K') \rightarrow (T^3, K)$. Since $f_*([K']) = [K]$, there exists a diffeomorphism $g: (T^3, K) \rightarrow (T^3, K)$ with $f_* = g_*$. Since $f^{-1} \cdot g: (T^3, K) \rightarrow (T^3, K')$ satisfies $(f^{-1} \cdot g)_* = \text{id}$, it is isotopic to the identity map. \square

Let U be an unknotted T^2 -knot in S^4 . $\overline{S^4 - N(U)}$ is a *twin* (see [7]). We denote the twin by the symbol Tw . A twin consists of two $S^2 \times D^2$'s plumbed at two points with opposite signs. Let R, S be the cores of two $S^2 \times D^2$'s. They generate $H_2(Tw)$. Let $D(r), D(s)$ be 2-disks properly embedded in Tw such that $R \cdot D(r) = S \cdot D(s) = 1$ and $R \cdot D(s) = S \cdot D(r) = 0$. $\partial D(r)$ and $\partial D(s)$ are circles in $\partial(Tw)$. We

call them r and s respectively. Their homology classes in $H_1(\partial(Tw))$ are well-defined. Choose a circle l in $\partial(Tw)$ such that $\langle l, r, s \rangle$ is an oriented basis of $H_1(\partial(Tw))$. Two l 's are mapped to each other by some diffeomorphism between Tw 's which fixes r and s (see Remark 2.5 in [3]). Next, we consider the manifold $D^2 \times T^2 \cong D^2 \times S^1 \times S^1$. Let $\bar{l}, \bar{r}, \bar{s}$ be the circles $\partial D^2 \times \{*\} \times \{*\}, \{*\} \times S^1 \times \{*\}, \{*\} \times \{*\} \times S^1$ in $\partial(D^2 \times T^2)$ respectively. $S^4 = Tw \cup_f T^2 \times D^2$ where $f_*[\bar{l} \bar{r} \bar{s}] = [l r s]$. Put $T_0 = T^2 \times D^2 \cap Tw \subset S^4$. Assume that K is a torus T^2 -knot contained in T_0 . Denote K by $K(p, q, q')$ if $[K] = p(r \times s) + q(s \times l) + q'(l \times r)$ in $H_2(\partial(Tw))$ where $p, q, q' \in \mathbf{Z}$. Note that by Lemma 2.5, p, q, q' determines the knot type of K .

Let $h: T_0 \rightarrow T_0$ be a diffeomorphism with $h_*[l r s] = [l r s]A^h$, where $A^h \in GL_3\mathbf{Z}$. There is a diffeomorphism $\bar{h}: S^4 \rightarrow S^4$ such that $\bar{h}|_{T_0} = h$ if and only if $A^h \in H$, where

$$H = \left\{ \begin{bmatrix} \pm 1 & 0 & 0 \\ 0 & a & b \\ 0 & c & d \end{bmatrix} \in GL_3\mathbf{Z} \mid a + b + c + d \equiv 0 \pmod{2} \right\}$$

(see Theorem 5.3 in [7] and Lemma 2.6 in [3]). If

$$h_*[l r s] = [l r s] \begin{bmatrix} \varepsilon & 0 & 0 \\ 0 & a & b \\ 0 & c & d \end{bmatrix},$$

then

$$h_*[r \times s \ s \times l \ l \times r] = [r \times s \ s \times l \ l \times r] \begin{bmatrix} ad - bc & 0 & 0 \\ 0 & \varepsilon d & -\varepsilon c \\ 0 & -\varepsilon b & \varepsilon a \end{bmatrix}.$$

Therefore if

$$\begin{bmatrix} p \\ q \\ q' \end{bmatrix} = \begin{bmatrix} ad - bc & 0 & 0 \\ 0 & \varepsilon d & -\varepsilon c \\ 0 & -\varepsilon b & \varepsilon a \end{bmatrix} \begin{bmatrix} p_1 \\ q_1 \\ q'_1 \end{bmatrix},$$

$K(p, q, q')$ and $K(p_1, q_1, q'_1)$ have the same knot type. Since

$$\begin{aligned} & GL_2\mathbf{Z} / \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a + b + c + d \equiv 0 \pmod{2} \right\} \\ &= \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\}, \end{aligned}$$

every torus knot is equivalent to $K(p, q, 0)$ or $K(p, q, q)$ for some non-negative integers p, q with $\gcd(p, q) = 1$.

It is easy to check the following:

LEMMA 2.6. $K(p, q, 0) = S(k(p, q))$ and $K(p, q, q) = \tilde{S}(k(p, q))$.

Therefore, every torus T^2 -knot is equivalent to $S(k(p, q))$ or $\tilde{S}(k(p, q))$ for some p, q . Since $k(p, q) = k(q, p)$, we have $K(p, q, 0) = K(q, p, 0)$ and $K(p, q, q) = K(q, p, p)$. It is clear that $\pi_1(S^3 - k(p, q)) \cong \pi_1(S^4 - K(p, q, 0)) \cong \pi_1(S^4 - K(p, q, q))$. The following theorem is known. For example, see [10], p. 54.

THEOREM 2.7. (*O. Schreier.*) *If $1 < p < q$ and $1 < p' < q'$, then $\pi_1(S^3 - k(p, q)) \cong \pi_1(S^3 - k(p', q'))$ if and only if $p = p', q = q'$. \square*

The exteriors of $K(p, q, 0)$ and $K(p, q, q)$ have the same homotopy type. But,

LEMMA 2.8. *If $p > 1$ and $q > 1$, then the exteriors of $K(p, q, 0)$ and $K(p, q, q)$ have different diffeomorphism types.*

Proof. Put $K = K(p, q, 0)$ or $K(p, q, q)$, $k = k(p, q)$. Recall that $S^4 - \text{Int } N(K) = (B^3 - \text{Int } N(k)) \times S^1 \cup S^2 \times D^2$. Let i be the inclusion map $\partial N(K) \rightarrow S^4 - \text{Int } N(K)$ and $*$ be a point in $\partial N(k)$. Then, $[\{*\} \times S^1] \in \pi_1(\partial N(K))$ generates $\text{Ker}(i_*: \pi_1(\partial N(K)) \rightarrow \pi_1(S^4 - \text{Int } N(K)))$.

Fact. Let X be a spin 4-manifold with $\partial X = T^3$. Let C_1, C_2 be two loops in ∂X with a diffeomorphism $\partial X \rightarrow S^1 \times S^1 \times S^1$ which maps C_1, C_2 to $\{*\} \times \{*\} \times S^1$ and $\{*\}' \times \{*\} \times S^1$ ($* \neq *'$). Assume that $[C_1] = [C_2] = 0$ in $H_1(X; \mathbf{Z}/2)$. Let D_i be a 2-chain in X such that (a) $\partial D_i \pmod{2} = C_i$ ($i = 1, 2$), (b) $[D_1] = [D_2]$ in $H_2(X, \partial X; \mathbf{Z}/2)$, (c) D_1 and D_2 meet transversely. Then, $D_1 \cdot D_2 \pmod{2}$ is determined by $[C_1]$ in $H_1(\partial X; \mathbf{Z}/2)$ (see the proof of Lemma 2.10 in [3]).

Put $Y([C_1]) = D_1 \cdot D_2 \in \mathbf{Z}/2$.

For $K(p, q, 0)$ (resp. $K(p, q, q)$), $Y([\{*\} \times S^1]) = 0$ (resp. 1). This completes the proof. \square

We have proved :

PROPOSITION 2.9. *Any torus T^2 -knot is equivalent to one and only one of the following:*

- (i) $K(p, q, 0)$, $1 < p < q, \text{gcd}(p, q) = 1$;
- (ii) $K(p, q, q)$, $1 < p < q, \text{gcd}(p, q) = 1$;
- (iii) *unknotted T^2 -knot.* \square

For the definition of *good torus fibrations (GTF)*, see [6] or [3], §3.

LEMMA 2.10. *The exteriors of $K(p, q, 0)$ and $K(p, q, q)$ have structures of good torus fibrations with one twin singular fiber of multiplicity p and one multiple torus of multiplicity q .*

Proof. Recall that $S^4 = Tw \cup_f T^2 \times D^2$ where $f_*[\bar{l} \ \bar{r} \ \bar{s}] = [l \ r \ s]$ and that $T_0 = Tw \cap T^2 \times D^2 \subset S^4$. Let $\text{pr}: T^3 = T^1 \times T^2 \rightarrow T^1$ be the projection map to the first coordinate and let $h: T_0 \rightarrow T^3$ be a diffeomorphism such that $(\text{pr} \cdot h)_*$ maps l, r, s to $q\gamma, p\gamma, 0$ (resp. $q\gamma, p\gamma, p\gamma$) where γ is a generator of $H_1(T^1)$. Since $-pl + qr \mapsto 0$ and $s \mapsto 0$ (resp. $-pl + qr \mapsto 0$ and $s - r \mapsto 0$), $\text{pr} \cdot h$ has fiber $(-pl + qr) \times s = q(r \times s) + p(s \times l)$ (resp. $(-pl + qr) \times (s - r) = q(r \times s) + p(s \times l) + p(l \times r)$). By Proposition 3.12 and Definition 3.16 in [3], $\text{pr} \cdot h$ extends to a GTF $f: S^4 \rightarrow S^2$ with one twin singular fiber of multiplicity p and one multiple torus of multiplicity q . Since a general fiber in T_0 is $K(p, q, 0)$ (resp. $K(p, q, q)$), the lemma is proved. \square

3. Dehn-surgery along a torus T^2 -knot. In this section, K denotes $K(p, q, 0)$ or $K(p, q, q)$ and k denotes $k(p, q)$ (the classical torus knot of type (p, q) in S^3). We assume that K is embedded in S^4 as is in the proof of Lemma 2.10.

We choose a basis (m, l_1, l_2) of $H_1(\partial N(K))$ as follows. Let m denote the meridian curve of k , which can be regarded as the meridian curve of K . Let l_0 be the preferred longitude of k . Put $l_1 = l_0 \times \{*\} \subset (B^3 - \text{Int } N(k)) \times S^1 \cup S^2 \times D^2 = S^4 - \text{Int } N(K)$. We orient m and l_1 so that $pqm + l_1 = -pl + qr$ in T_0 . Put $l_2 = s$ for $K(p, q, 0)$ and $l_2 = s - r$ for $K(p, q, q)$. Note that l_2 generates the kernel of $\pi_1(\partial N(K)) \rightarrow \pi_1(S^4 - \text{Int } N(K))$.

REMARK 3.1. Note that for $K = K(p, q, 0)$ or $K(p, q, q)$ a general fiber contained in $\partial N(K)$ represents $(pqm + l_1) \times l_2$ in $H_2(\partial N(K))$.

DEFINITION 3.2. Let K be a torus T^2 -knot in S^4 . $N(K)$ is diffeomorphic to $T^2 \times D^2$. Let $i: \partial N(K) \rightarrow \partial(S^4 - \text{Int } N(K))$ be the natural identification and $h: \partial N(K) \rightarrow \partial N(K)$ be a diffeomorphism such that $i \cdot h(m) = \alpha m + \beta l_1 + \gamma l_2$. $M = S^4 - N(K) \cup_{i,h} N(K)$ is called the manifold obtained by *Dehn-surgery of type (α, β, γ) along K* .

Montesinos showed that any homotopy 4-sphere obtained by Dehn-surgery along an unknotted T^2 -knot in S^4 is diffeomorphic to S^4 (see [7], p. 187). Pao studied the 4-manifolds with effective T^2 -action in [9]. All the 4-manifolds obtained by Dehn-surgeries along an unknotted T^2 -knot are contained in his list. See also [3].

Note that the diffeomorphism type of M is determined by K and $i \cdot h(m)$ (see Remark 2.7. in [3]).

Dehn-surgery along any T^2 -knot can be defined in the same way, but the description of the type of the surgery might be a little complicated.

PROPOSITION 3.3. *Assume that a Dehn-surgery of type (α, β, γ) is performed along $K(p, q, 0)$ (resp. $K(p, q, q)$). Denote the manifold by M .*

(i) *If $\gcd(\alpha, \beta) = 1$, then $\pi_1(M)$ is isomorphic to the fundamental group of the manifold obtained by a Dehn-surgery of type (α, β) along a torus T^1 -knot of type (p, q) .*

(ii) $H_1(M) \cong \mathbf{Z}/\alpha$.

(iii) *M is spin if and only if $\beta\gamma \equiv 0$ (resp. $(1 - \beta)\gamma \equiv 0$) (mod 2) or $\alpha \equiv 1$ (mod 2).*

Proof. The proofs of (i) and (ii) are almost clear. For (iii), we need a lemma.

LEMMA 3.4. *Put $X = S^4 - \text{Int } N$ where N is a tubular neighborhood of a T^2 -knot (not necessarily a torus T^2 -knot) K . Let Y be the function defined in the proof of Lemma 2.8. Assume that $\text{Ker}(i_*: H_1(\partial X; \mathbf{Z}/2) \rightarrow H_1(X; \mathbf{Z}/2)) = \{0, e_1, e_2, e_3\}$ where $i: \partial X \rightarrow X$ is the inclusion map. Then, one of $Y(e_i)$'s is 1, the others are 0.*

Proof. Note that if $\{i, j, k\} = \{1, 2, 3\}$, then $e_k = e_i + e_j$ holds.

Let V be a 2-sided 3-dimensional submanifold of S^4 such that $\partial V = K$ and that $(N, N \cap V)$ is diffeomorphic to $(D^2 \times T^2, r \times T^2)$ where r is a radius of D^2 . Put $V_0 = \partial N \cap V$. Note that e_1, e_2, e_3 are represented by curves in V_0 . Assume that $i_*: H_1(V_0; \mathbf{Z}/2) \rightarrow H_1(V; \mathbf{Z}/2)$ is injective (i is the inclusion map). Then, the Mayer-Vietoris exact sequence shows that $H_2(V; \mathbf{Z}/2) \rightarrow H_2(V, V_0; \mathbf{Z}/2)$ is surjective. Since $H_3(V, V_0; \mathbf{Z}/2) \rightarrow H_2(V_0; \mathbf{Z}/2)$ is bijective, therefore $H_2(V; \mathbf{Z}/2) \rightarrow H_2(V, V_0; \mathbf{Z}/2)$ is injective. By Poincaré duality, $H_2(V, V_0; \mathbf{Z}/2)$ is isomorphic to $H^1(V; \mathbf{Z}/2)$, which is isomorphic to $H_1(V; \mathbf{Z}/2)$. Therefore $\chi(V) = 1$. Put $\bar{V} = V \cup_{\partial} S^1 \times D^2$. Then,

$$\chi(\bar{V}) = \chi(V) + \chi(S^1 \times D^2) - \chi(\partial V) = 1.$$

Since \bar{V} is a closed 3-manifold, this is a contradiction. Therefore $\text{Ker}(i_*: H_1(V_0; \mathbf{Z}/2) \rightarrow H_1(V; \mathbf{Z}/2))$ contains e_i , one element of $\{e_1, e_2, e_3\}$. Since V has trivial normal bundle, we can move V slightly into V' so that $V \cap V' = \emptyset$. Therefore $Y(e_i) = 0$.

Let m be the meridian curve of K and we consider the manifold $Q = X \cup_a Tw$ with the attaching map $a: \partial(Tw) \rightarrow \partial X$ satisfying $a_*[l \ r \ s] = [\tilde{m} \ \tilde{e}_j \ \tilde{e}_k]$ where $\langle \tilde{m}, \tilde{e}_j, \tilde{e}_k \rangle$ is a basis of $H_1(\partial X; \mathbf{Z})$ whose mod 2 reduction is $\langle m, e_j, e_k \rangle$. The Mayer-Vietoris sequence with coefficients in $\mathbf{Z}/2$

$$H_2(X) \oplus H_2(Tw) \xrightarrow{j_*} H_2(Q) \rightarrow H_1(\partial X) \xrightarrow{i_*} H_1(X) \oplus H_1(Tw)$$

shows that

$$H_2(Q; \mathbf{Z}/2) = \text{Im}(j_*) \oplus \langle [D(e_j) + D(r)], [D(e_k) + D(s)] \rangle,$$

where $D(c)$ is a mod 2 2-chain satisfying $\partial D(c) = c$ and $D(e_j), D(e_k) \subset X, D(r), D(s) \subset Tw$. The self-intersection number on $\text{Im}(j_*)$ is zero since X and Tw are subsets of S^4 . Since $[D(e_j) + D(r)]^2 = Y(e_j) + Y(r) = Y(e_j)$ and $[D(e_k) + D(s)]^2 = Y(e_k) + Y(s) = Y(e_k)$, we have

$$\begin{aligned} [D(e_i) + D(r + s)]^2 &= [D(e_j) + D(e_k) + D(r) + D(s)]^2 \\ &= [D(e_j) + D(r)]^2 + [D(e_k) + D(s)]^2 \\ &= Y(e_j) + Y(e_k). \end{aligned}$$

On the other hand, $[D(e_i) + D(r + s)]^2 = Y(e_i) + Y(r + s) = 0 + 1 = 1$. Therefore $Y(e_j) = 0, Y(e_k) = 1$ or $Y(e_j) = 1, Y(e_k) = 0$. □

We now continue with the proof of Proposition 3.3.

If α is odd, $H_2(M; \mathbf{Z}/2)$ is zero. Therefore M is spin. Assume that α is even. The Mayer-Vietoris sequence with coefficients in $\mathbf{Z}/2$

$$\begin{aligned} H_2(D^2 \times T^2) \oplus H_2(X) &\xrightarrow{j_*} H_2(M) \rightarrow H_1(\partial(D^2 \times T^2)) \\ &\xrightarrow{i_*} H_1(D^2 \times T^2) \oplus H_1(X) \end{aligned}$$

shows that $H_2(M; \mathbf{Z}/2) = \text{Im}(j_*) \oplus \langle [D_M + D(\alpha m + \beta l_1 + \gamma l_2)] \rangle$ where X is the knot exterior and D_M is the meridian disk of $D^2 \times T^2$ and $D(\alpha m + \beta l_1 + \gamma l_2)$ is a mod 2 2-chain in X with $\partial D(\alpha m + \beta l_1 + \gamma l_2) = \alpha m + \beta l_1 + \gamma l_2$. For $K = K(p, q, 0)$ (resp. $K(p, q, q)$), $Y(l_2) = 0$ (resp. 1) and $Y(l_1) = 0$. By Lemma 3.4, $Y(l_1 + l_2) = 1$ (resp. 0). Therefore $[D_M + D(\alpha m + \beta l_1 + \gamma l_2)]^2 = \beta\gamma \pmod 2$ (resp. $(1 - \beta)\gamma \pmod 2$). Since the self-intersection number on $\text{Im}(j_*)$ is zero, this completes the proof. □

PROPOSITION 3.5. *If a closed 4-manifold M is obtained by Gluck-surgery along an S^2 -knot in S^4 , then M is also obtained by Dehn-surgery along a T^2 -knot in S^4 .*

Proof. Here K denotes the S^2 -knot. Identify $N(K) \cong S^2 \times D^2$ with $\hat{C} \times D$ where \hat{C} is the Riemann sphere and D is the unit disk in \mathbf{C} . Recall that $M = (S^4 - \text{Int } N(K)) \cup_{\tau} N(K)$ is the manifold obtained by Gluck-surgery along K where $\tau(u, v) = (uv, v)$. Observe that $\tau(\{|u| = c\} \times \partial D) = \{|u| = c\} \times \partial D$ for any $c \in \mathbf{R} \cup \{\infty\}$. Embed $(D^3 \times D^1, D^3 \times \partial D^1)$ in $(S^4 - \text{Int } N(K), \partial N(K))$ with $D^3 \times \{-1\} \subset D_0 \times \partial D^2$ and $D^3 \times \{1\} \subset D_{\infty} \times \partial D^2$ where $D_0 = \{|z| \leq 1/9\} \subset \hat{C}$ and $D_{\infty} = \{|z| \geq 9\} \subset \hat{C}$. Let H denote its image. One can consider H as a 1-handle attached to $N(K)$. Verify that there exists an annulus A_0 (resp. A_{∞}) properly embedded in $D_0 \times D^2$ (resp. $D_{\infty} \times D^2$) such that $K' = (\hat{C} - \text{Int } D_0 - \text{Int } D_{\infty}) \times \{0\} \cup A_0 \cup U \times D^1 \cup A_{\infty}$ is an embedded torus where U is an unknot in D^3 . Especially, if K is unknotted, then so is K' .

K' has a tubular neighborhood $N(K')$ such that

$$N(K')|_{(\hat{C} - \text{Int}(D_0 \cup D_{\infty}))} = N(K)|_{(\hat{C} - \text{Int}(D_0 \cup D_{\infty}))}$$

and $N(K') \subset N(K) \cup H$ and $N(K')|(U \times \{*\}) = N_0(U) \times \{*\}$ where $N_0(U)$ is a tubular neighborhood of U in D^3 .

Let $f: \partial N(K') \rightarrow \partial N(K')$ be a diffeomorphism such that

$$f|_{(\hat{C} - \text{Int}(D_0 \cup D_{\infty})) \times \partial D} = \tau|_{(\hat{C} - \text{Int}(D_0 \cup D_{\infty})) \times \partial D}$$

and

$$f(\partial N(K') \cap D^3 \times \{*\}) = \partial N(K') \cap D^3 \times \{*\}.$$

Put $M' = (S^4 - \text{Int } N(K')) \cup_f N(K')$.

Construct a diffeomorphism $F: M' \rightarrow M$ as follows. Put

$$F|(S^4 - (D_0 \cup D_{\infty}) \times D - H) = \text{id}.$$

$M'_* = (D^3 - \text{Int } N_0(U)) \times \{*\} \cup_f N_0(U) \times \{*\}$ is the manifold obtained by Dehn-surgery of type $(1, 1)$ or $(1, -1)$ along U in D^3 . Therefore there exists a diffeomorphism $F_*: M'_* \rightarrow D^3 \times \{*\}$ with $F_*|_{\partial} = \text{id}$. Put $F|M'_* = F_*$. Finally, extend $F|_{\partial}(D_0 \times D)$ and $F|_{\partial}(D_{\infty} \times D)$ to $D_0 \times D$ and $D_{\infty} \times D$.

This completes the proof. □

PROPOSITION 3.6. *If a Dehn-surgery of type (α, β, γ) is performed along $K(p, q, 0)$ or $K(p, q, q)$ and $\sigma = |pq\beta - \alpha| \neq 0$, then the manifold obtained is the total space of a good torus fibration over S^2 with one twin singular fiber of multiplicity p and two multiple tori of multiplicity q and σ .*

Proof. Put $K = K(p, q, 0)$ or $K(p, q, q)$. By Lemma 2.10, the exterior of K has the structure of GTF. The intersection number of

$i \cdot h(m) = \alpha m + \beta l_1 + \gamma l_2$ and the fiber $(pqm + l_1) \times l_2$ in $\partial N(K)$ is $\pm(\alpha - pq\beta)$. Therefore, after surgery, a fiber is homologous to $\pm(\alpha - pq\beta)C$ in $N(K)$, where C is the core of $N(K)$. Now the proposition is proved. (See Definition 3.16 in [3].) □

REMARK 3.7. If $\sigma = 1$, then the manifold has only one twin singular fiber and one multiple torus. Therefore it is diffeomorphic to $L_{|\alpha|}$ or $L'_{|\alpha|}$ by the Main Theorem of [3] and Proposition 3.3.(ii). If $\alpha \equiv 0 \pmod{2}$, then it is $L_{|\alpha|}$ if and only if either $K = K(p, q, 0)$ and $\gamma \equiv 0 \pmod{2}$ or $K = K(p, q, q)$. See Proposition 3.3.(iii).

COROLLARY 3.8. *The manifold obtained by the Gluck-surgery along an untwisted spun (S^2 -) knot of any torus (S^1 -) knot is the 4-sphere.*

Proof. By the proof of Proposition 3.5, the manifold is diffeomorphic to the one obtained by a Dehn-surgery of type $(1, 0, \pm 1)$ along a torus T^2 -knot. Since $L_1 = S^4$, Corollary is proved. □

The following Proposition is almost clear.

PROPOSITION 3.9. *If a Dehn-surgery of type $(\alpha, \beta, 0)$ is performed along $K(p, q, 0)$ (resp. $K(p, q, q)$), then the manifold obtained is $(M_0 - \text{Int } B^3) \times S^1 \cup_h S^2 \times D^2$ where M_0 is the manifold obtained by a Dehn-surgery of type (α, β) along $k(p, q)$ and $h = \text{id}$ (resp. τ).*

COROLLARY 3.10. *If a Dehn-surgery of type $(pq, 1, 0)$ is performed along $K(p, q, 0)$ or $K(p, q, q)$, then the manifold obtained is $L_p \# L_q$.*

Proof. The manifold obtained is $(L - \text{Int } B^3) \times S^1 \cup S^2 \times D^2$ where $L = L(p, q) \# L(q, p)$. Corollary 4.10 in [3] completes the proof. □

PROPOSITION 3.11. *If a Dehn-surgery of type (α, β, γ) is performed along $K(p, q, 0)$ or $K(p, q, q)$ and $\sigma = |pq\beta - \alpha| = 0$, then the manifold obtained is an irrational connected sum along circles of either L_m or L'_m and $L(n, r) \times S^1$ for some m, n, r .*

Proof. In this case, the meridian of 4-dimensional solid torus sT^4 is attached in a fiber of a GTF of the knot exterior. Recall that the knot exterior X is made of Tw and $D^2 \times T^2$ pasted together along A , where A is diffeomorphic to $D^1 \times T^2$.

Put $\partial(D^2 \times T^2) - \text{Int } A = B$ and $\partial(Tw) - \text{Int } A = C$. B and C are diffeomorphic to $T^2 \times D^1$. Let $h: \partial(sT^4) \rightarrow B \cup C$ be the attaching

map. Since $B \cap C$ is a disjoint union of two fibers, we may assume that $h^{-1}(B) = \partial D^2 \times S^1 \times [*, **] \subset \partial D^2 \times S^1 \times S^1 = \partial(D^2 \times T^2)$, $h^{-1}(C) = \partial D^2 \times S^1 \times [**, *] \subset \partial D^2 \times S^1 \times S^1 = \partial(D^2 \times T^2)$. Put $V = D^2 \times S^1 \times [*, **]$ and $V' = D^2 \times S^1 \times [**, *]$. $V \cup V' = sT^4$. The manifold obtained by the Dehn-surgery is

$$\begin{aligned} M &= D^2 \times T^2 \cup Tw \cup sT^4 \\ &= D^2 \times T^2 \cup Tw \cup (V \cup V') = (D^2 \times T^2 \cup V) \cup (Tw \cup V'). \end{aligned}$$

Note that $\partial(D^2 \times T^2 \cup V) = \partial(Tw \cup V') = S^2 \times S^1$. If we attach $D^3 \times S^1$ to $D^2 \times T^2 \cup V$ (resp. $Tw \cup V'$) in the natural way, $V \cup D^3 \times S^1$ (resp. $V' \cup D^3 \times S^1$) is diffeomorphic to $D^2 \times T^2$.

It is easy to show that $D^2 \times T^2 \cup D^2 \times T^2$ is $L(n, r) \times S^1$ for some n, r . The proof of Theorem 4.1 in [3] says that $Tw \cup D^2 \times T^2$ is L_m or L'_m . This completes the proof. \square

REFERENCES

- [1] M. Dehn, *Über die Topologie des dreidimensionalen Raumes*, Math. Ann., **69** (1910), 137–168.
- [2] H. Gluck, *The embedding of two-spheres in the four-sphere*, Trans. Amer. Math. Soc., **104** (1962), 308–333.
- [3] Z. Iwase, *Good torus fibrations with twin singular fibers*, Japan. J. Math., **10** (1984), 321–352.
- [4] W. Jaco, *Lectures on three-manifold topology*, Regional conference series in mathematics; no. 43, the American Mathematical Society, 1980.
- [5] R. Mandelbaum, *Irrational connected sums and the topology of algebraic surfaces*, Trans. Amer. Math. Soc., **247** (1979), 137–156.
- [6] Y. Matsumoto, *Good torus fibrations*, Contemporary Math., **35** (1984), 375–397.
- [7] J. M. Montesinos, *On twins in the four-sphere I*, Quart. J. Math. Oxford, **34-2** (1983), 171–199.
- [8] L. Moser, *Elementary surgery along a torus knot*, Pacific J. Math., **38** (1971), 737–745.
- [9] P. S. Pao, *The topological structure of 4-manifolds with effective torus actions. I*, Trans. Amer. Math. Soc., **227** (1977), 279–317.
- [10] D. Rolfsen, *Knots and links*, Mathematics Lecture Series 7, Publish or Perish, Inc. Berkeley, 1976.

Received August 27, 1986 and in revised form July 15, 1987. The author is partially supported by Fellowships of the Japan Society for the Promotion of Science for Japanese Junior Scientists and Grant-in-Aid for Encouragement of Young Scientist, No. 61790113.

UNIVERSITY OF TOKYO
HONGO, TOKYO (113)
JAPAN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906-1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1988 by Pacific Journal of Mathematics

William Charles Bauldry, Attila Mate and Paul Nevai, Asymptotics for solutions of systems of smooth recurrence equations	209
Ehrhard Behrends, Isomorphic Banach-Stone theorems and isomorphisms which are close to isometries	229
Fernanda Maria Botelho, Rotation sets of maps of the annulus	251
Edward Graham Evans, Jr. and Phillip Alan Griffith, Binomial behavior of Betti numbers for modules of finite length	267
Andrei Iordan, Pseudoconvex domains with peak functions at each point of the boundary	277
Zyun'iti Iwase, Dehn-surgery along a torus T^2-knot	289
Marko Kranjc, Embedding 2-complexes in \mathbf{R}^4	301
Aloys Krieg, Eisenstein-series on real, complex, and quaternionic half-spaces	315
Masato Kuwata, Intersection homology of weighted projective spaces and pseudo-lens spaces	355
Carl Pomerance, András Sárközy and Cameron Leigh Stewart, On divisors of sums of integers. III	363
Martin Schechter, Potential estimates in Orlicz spaces	381