

Pacific Journal of Mathematics

A BONNESEN-STYLE INRADIUS INEQUALITY IN 3-SPACE

JANE SANGWINE-YAGER

A BONNESEN-STYLE INRADIUS INEQUALITY IN 3-SPACE

J. R. SANGWINE-YAGER

A Bonnesen-style inradius inequality for convex bodies in E^3 is obtained using the method of inner parallel bodies. The inequality involves the volume, surface area and mean-width of the body.

I. Introduction. By a convex body we mean a compact convex set with non-empty interior. Let K be a planar convex body with area A , perimeter L , inradius r , and circumradius R . An inequality of Bonnesen states:

$$L^2 - 4\pi A \geq \pi^2(R - r)^2.$$

This inequality follows from

$$(1) \quad 0 \geq A - xL + x^2\pi, \quad r \leq x \leq R.$$

Equality holds in (1), at $x = r$, for the “sausage” bodies, that is, those bodies which are the Minkowski sum of a line segment and a ball (with radius r). At $x = R$ equality only holds for balls. For proofs of these inequalities see Eggleston [5, pp. 108–110].

An extension of Bonnesen’s inradius inequality in the plane to higher dimensions began with the conjecture by Wills [11] that

$$0 \geq V - rS + (n - 1)r^n\omega_n.$$

In this paragraph, V will represent the n -dimensional volume of a convex body in E^n , S its n -dimensional surface area, and ω_n the volume of the unit n -ball. The conjecture was proved simultaneously by Bokowski [1] and Diskant [4]. Equality holds only for the n -balls. Osserman [8] showed that

$$(2) \quad 0 \geq V - rS + (n - 1)r^2 \sqrt[n-1]{\omega_n(S/n)^{n-2}}$$

where equality also holds only for the n -balls. This inequality is the sharper because a translate of rB is contained in K .

The results of this paper will be limited to the case $n = 3$. The volume and surface area of the convex body K will be represented by $V(K)$ and $S(K)$. The unit 3-ball centered at the origin is denoted B and $V(B) = \omega$. The functional $M(K)$ will be proportional to the

mean-width of K , where $M(B) = 3\omega$. Often we will let $V = V(K)$, $S = S(K)$ and $M = M(K)$.

The following inequality for 3-dimensional bodies will be established:

$$(3) \quad 0 \geq V - rS + \frac{2}{3}r^2M,$$

where equality holds if and only if K is a cap body of a ball. A cap body of a ball is the convex hull of the ball and countably many points exterior to it such that the line segment joining any two of these points intersects the ball. The cap bodies are also the extremal bodies for the inequality of Minkowski, $M^2 \geq 3S\omega$. From this inequality it follows that (3) is sharper than (2).

The inequality (3) is a consequence of a more general inequality for inner parallel bodies to be proved below. In proving our result we will actually establish a lower bound for the volume of the inner parallel bodies of a given body. This bound will resolve a conjecture of Matheron [7].

II. Main results. The inner parallel body of a convex body K at a distance $\lambda \geq 0$ is defined to be the intersection of the closed supporting half-spaces of K moved in a distance λ . The inradius r is the radius of the largest ball contained in K . The inner parallel body at a distance $r - \lambda$, $0 \leq \lambda \leq r$, shall be denoted K_λ ; the inradius of K_λ is λ . The kernel, K_0 , is the locus of centers of balls of radius r contained in K , and $K = K_r$.

The extremal bodies in the inequalities which follow are cap bodies. For the definition of p -extreme directions, $p = 0, 1$ or 2 , see Schneider [9, p. 135]. A supporting hyperplane is p -extreme if its outer normal is a p -extreme direction. A 0-extreme direction is simply called extreme [5, p. 27]; the dual of the Krein-Milman Theorem states that every convex body is the intersection of its extreme supporting half-spaces. A convex body K is a $(2-p)$ -tangential body of the body C if every p -extreme supporting hyperplane of K also supports C . A 1-tangential body is a cap body; an alternate definition of a cap body of a ball is given above. A theorem of Favard [6, pp. 273–274] states that K is a cap body of C if and only if the mixed volumes $V(K, K, K)$, $V(K, K, C)$ and $V(K, C, C)$ are equal. The definition and properties of mixed volumes in general may be found in [5, p. 84]. In particular, $V(K) = V(K, K, K)$, $S(K) = 3V(K, K, B)$, $M(K) = 3V(K, B, B)$, and the mixed volumes are linear in each argument.

We shall now prove the following

THEOREM. *Let K be a convex body in E^3 with non-empty interior and inradius r . For all λ , $0 \leq \lambda \leq r$,*

$$(4) \quad V(K_\lambda) \geq V - (r - \lambda)S + (r - \lambda)^2[2M + M(K_\lambda)]/3$$

where equality holds for all λ , $0 \leq \lambda \leq r$, if and only if K is a cap body of $K_0 + rB$.

Proof of the Theorem. Fix λ , $0 < \lambda < r$. For all t , $\lambda \leq t \leq r$, let

$$f(t) = V(K_\lambda) - V(K_t) + (t - \lambda)S(K_t) - (t - \lambda)^2[2M(K_t) + M(K_\lambda)]/3.$$

Bol [3] proved many results about functions of inner parallel bodies and their right- and left-hand derivatives. For example, $V(K_t)$ and $S(K_t)$ are absolutely continuous. If $F'(t)$ denotes the left-hand derivative of any function $F(t)$ with respect to t , then

$$(5) \quad V'(K_t) = S(K_t),$$

$$(6) \quad S'(K_t) \geq 2M(K_t).$$

The function $M(K_t)$ is a concave function of t which implies

$$(7) \quad M(K_t) - M(K_\lambda) \geq (t - \lambda)M'(K_t).$$

Using (5) we find

$$f'(t) = (t - \lambda)[[S'(K_t) - 2M(K_t)] + 2[M(K_t) - M(K_\lambda) - (t - \lambda)M'(K_t)]]/3.$$

Therefore by (6) and (7) $f'(t) \geq 0$; $f(\lambda) = 0$, implies $f(r) \geq 0$. This establishes the inequality.

It remains to establish the equality condition. Favard's Theorem states that K is a cap body of $K_0 + rB$ if and only if

$$(8) \quad V(K, K_0 + rB, K_0 + rB) = V(K, K, K_0 + rB).$$

If equality holds in (4) for all λ , $0 \leq \lambda \leq r$, then for λ fixed, $0 < \lambda < r$, $f(t)$ must be identically zero. Therefore equality must hold in (6) and (7) for all λ and t , $0 < \lambda < t \leq r$. It should be noted that the conditions for equality in (6) remain unknown since they were first investigated by Bol [3]. Equality in (7) implies that $M(K_t)$ is a linear function of t . Since families of parallel bodies are concave, we have

$$(9) \quad \frac{t}{r}K + \left(1 - \frac{t}{r}\right)K_0 = K_t,$$

for all t , $0 \leq t \leq r$. The representation for K_t in (9) may be used to

expand and differentiate $S(K_t)$:

$$S(K_t) = \left(\frac{t}{r}\right)^2 S(K) + 6\frac{t}{r} \left(1 - \frac{t}{r}\right) V(K, K_0, B) + \left(1 - \frac{t}{r}\right)^2 S(K_0),$$

$$(10) \quad \frac{r}{2} S'(K_t) = \frac{t}{r} S(K) + 3 \left(1 - \frac{2t}{r}\right) V(K, K_0, B) - \left(1 - \frac{t}{r}\right) S(K_0).$$

At $t = r$,

$$\frac{r}{2} S'(K) = S(K) - 3V(K, K_0, B).$$

Substitution into (6), with equality, gives

$$S(K) - 3V(K, K_0, B) = rM(K)$$

or

$$(11) \quad V(K, K, B) = V(K, K_0 + rB, B).$$

The expression in (11) implies that the supporting functions of K and $K_0 + rB$ are equal on the support of the mixed area measure $S(K, B; \cdot)$. Results of Schneider [9, pp. 134–135] imply (8).

On the other hand, suppose K is a cap body of $K_0 + rB$. Since the inner parallel bodies of K are the intersection of the extreme supporting half-spaces of K moved inward, the extreme directions of an inner parallel body are extreme directions of K . Choose λ and t , $0 < \lambda < t < r$, let $\theta = (t - \lambda)/(r - \lambda)$, and let u be a unit vector. If $h(C, u)$ is the supporting function of a convex body C in the direction u , then

$$(12) \quad K_0 + tB \subseteq \theta K + (1 - \theta)K_\lambda \subseteq K_t \subseteq \{x: \langle x, u \rangle \leq h(K, u) - (r - t)\}.$$

An extreme direction of $\theta K + (1 - \theta)K_\lambda$ is either an extreme direction of K , an extreme direction of K_λ (and hence K), or a 1-extreme direction of both. If u is a 0- or 1-extreme direction of K , then $h(K, u) = h(K_0, u) + r$, and the supporting half-space in (12) also supports $K_0 + tB$. Therefore equality holds throughout (12), and the representation in (9) is valid. This implies equality in (7). The result of Favard (8) has many implications. It implies both (11) and $V(K, K, K_0) = V(K, K_0 + rB, K_0)$. Furthermore, since the support of the mixed area measure $S(B, K_\lambda; \cdot)$ is the closure of the set of extreme directions of K_λ , $V(K, B, K_\lambda) = V(K_0 + rB, B, K_\lambda)$. Hence $V(K, B, K_0) = V(K_0 + rB, B, K_0)$. Substitution into the right-hand side of (10) yields

$$\frac{r}{2} S'(K_t) = 3(tV(K, B, B) + (r - t)V(K_0, B, B)).$$

Equality holds in (6) also. This completes the proof of the theorem. \square

For $\lambda = 0$, (4) becomes the following Bonnesen-style inequality

$$0 \geq V - rS + r^2[2M + M(K_0)]/3$$

which implies (3). Equality holds if and only if K is a cap body of $K_0 + rB$, and therefore in (3) if and only if K is a cap body of a ball. The cap bodies appear in the equality condition for (3) because they are precisely those bodies for which $3V = rS = r^2M$.

Matheron [7] showed that in the plane

$$A(K_\lambda) \geq A - (r - \lambda)L + (r - \lambda)^2\pi$$

where equality holds for all λ , $0 \leq \lambda \leq r$, if and only if K is a sausage body. Notice that at $\lambda = 0$ this inequality is (1), for $x = r$, since the area of the kernel is zero.

The right-hand side of Matheron's inequality is similar to Steiner's formula for the outer parallel bodies of K , except that the signs alternate. Matheron conjectured that similar lower bounds existed for the volume of the inner parallel bodies in higher dimensional spaces. Weil [10] has established necessary and sufficient conditions for one body to be the summand of another. This result implies that if $K_\lambda + (r - \lambda)B = K$, for K in E^3 , then

$$V(K_\lambda) - [V - (r - \lambda)S + (r - \lambda)^2M - (r - \lambda)^3\omega] = 0.$$

The conjectured inequality does not exist because the difference above is positive, at $\lambda = r$, for the cubes and negative for cap bodies. Notice that the right-hand side of (4) is equal to

$$V - (r - \lambda)S + (r - \lambda)^2M - (r - \lambda)^3\omega + (r - \lambda)^2[M(K_\lambda + (r - \lambda)B) - M]/3$$

which is smaller than the alternating sum proposed by Matheron because $K_\lambda + (r - \lambda)B \subseteq K$.

III. Further remarks. Remarks will be made on two significant differences between (3) and (1): inequality (3) does not strengthen the isoperimetric inequality, and it is not valid if the inradius r is replaced with the circumradius R .

For $n > 2$, sharpened versions of the isoperimetric inequality have been established, and Osserman [8, pp. 24–25] shows that these inequalities imply (2). Inequality (3) does not yield either isoperimetric-style inequality

$$S^2 \geq 3VM \quad \text{or} \quad S^3 \geq 27V\omega.$$

For example, (3) implies

$$S^2 - 3VM \geq \left(\frac{V}{r}\right)^2 - \frac{5}{3}VM + \frac{4}{9}r^2M^2,$$

but the right-hand side is negative for pencil-like bodies. This raises the question of whether (3) is a strong inequality. It is if we view (1), for $x = r$, and (3) in the following sense: They provide the best possible lower bounds by weighted arithmetic means for rL and rS respectively. In (3) rS is bounded below by $\theta 3V + (1 - \theta)r^2M$. This bound is best when θ is minimized ($\theta = 1$ for the cube). The value of $\theta = 1/3$ is approached in the limit for a family of cylinders which converge to a disk in the plane.

Recently, Bonnesen-style circumradius inequalities involving three quermassintegrals in n -dimensional space have been found by Bokowski and Heil [2]. One inequality, for $n = 3$, is

$$0 \leq V - 4RS/3 + R^2M.$$

Equality holds if and only if K is homothetic to RB . Notice that the direction of the inequality is reversed from (1) and (3). The inequality is also a bound for RS by a weighted arithmetic mean, and it has been shown to be best.

REFERENCES

- [1] J. Bokowski, *Eine verschärfte Ungleichung zwischen Volumen, Oberfläche und Inkugelradius im R^n* , Elem. Math., **28** (1973), 43–44.
- [2] J. Bokowski and E. Heil, *Integral representations of quermassintegrals and Bonnesen-Style inequalities*, Arch. Math., to appear.
- [3] G. Bol, *Beweis einer Vermutung von H. Minkowski*, Abh. Math. Sem. Univ. Hamburg, **15** (1943), 37–56.
- [4] V. I. Diskant, *A generalization of Bonnesen's inequalities*, Soviet Math. Dokl., **14** no. 6 (1973), 1728–1731 (Transl. of Dokl. Akad. Nauk SSSR, **213** (1973), no. 3).
- [5] H. G. Eggleston, *Convexity*, Cambridge U. Press, Cambridge, 1969.
- [6] J. Favard, *Sur les corps convexes*, J. Math. Pur. Appl., IX. Sér., **12** (1933), 219–282.
- [7] G. Matheron, *La formule de Steiner pour les érosions*, J. Appl. Prob., **15** (1978), 126–135.
- [8] R. Osserman, *Bonnesen-Style isoperimetric inequalities*, Amer. Math. Monthly, **86**, no. 1, (1979), 1–29.
- [9] R. Schneider, *On the Aleksandrov-Fenchel inequality*, Discrete Geometry and Convexity, New York Academy of Sciences, (1985), 132–141.
- [10] W. Weil, *Decomposition of convex bodies*, Mathematika, **21** (1974), 19–25.
- [11] J. M. Wills, *Zum Verhältnis von Volumen zur Oberfläche bei konvexen Körpern*, Arch. Math., **21** (1970), 557–560.

Received July 6, 1987. This work was supported in part by the Saint Mary's College Faculty Development Fund.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

RAMESH A. GANGOLLI
University of Washington
Seattle, WA 98195

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Marco Abate, Annular bundles	1
Ralph Cohen, Wen Hsiung Lin and Mark Mahowald, The Adams spectral sequence of the real projective spaces	27
Harry Joseph D'Souza, Threefolds whose hyperplane sections are elliptic surfaces	57
Theodore Gerard Faticoni, Localization in finite-dimensional FPF rings	79
Daniel Hitt, Invariant subspaces of \mathcal{H}^2 of an annulus	101
Ellen Kirkman and James J. Kuzmanovich, On the global dimension of fibre products	121
Angel Rafael Larotonda and Ignacio Zalduendo, Homogeneous spectral sets and local-global methods in Banach algebras	133
Halsey Lawrence Royden, Jr., Invariant subspaces of \mathcal{H}^p for multiply connected regions	151
Jane Sangwine-Yager, A Bonnesen-style inradius inequality in 3-space	173
Stefano Trapani, Holomorphically convex compact sets and cohomology	179
Thomas Vogel, Uniqueness for certain surfaces of prescribed mean curvature	197