

Pacific Journal of Mathematics

ON THE ILIEFF-SENDOV CONJECTURE

JOHNNY E. BROWN

ON THE ILIEFF-SENDOV CONJECTURE

JOHNNY E. BROWN

The well-known Ilieff-Sendov conjecture asserts that for any polynomial $p(z) = X \prod_{k=1}^n (z - z_k)$ with $|z_k| < 1$, each of the disks $|z - z_k| \leq 1$ ($1 \leq k \leq n$) must contain a critical point of p . This conjecture is proved for polynomials of arbitrary degree n with at most four distinct zeros. This extends a result of Saff and Twomey.

1. Introduction. The Gauss-Lucas Theorem states that all the critical points of a polynomial $p(z)$ lie in the convex hull of its zeros. This is a result concerning the position of all the zeros of $p'(z)$ relative to all the zeros of $p(z)$. Suppose we focus attention on any arbitrarily fixed zero of $p(z)$ and ask for the location of a zero of $p'(z)$ relative to it. This leads to the well-known conjecture of Ilieff and Sendov [3; Problem 4.5] which asserts that if $p(z)$ has the form

$$(1) \quad p(z) = f \prod_{k=1}^n (z - z_k), \quad |z_k| < 1 \quad (1 \leq k \leq n)$$

then each of the disks $|z - z_k| \leq 1$ ($1 \leq k \leq n$) contains a zero of $p'(z)$. The polynomial $p(z) = z^n - 1$ shows that "1" is sharp. This conjecture is nearly a quarter of a century old and has been verified in some special cases, most notably if $p(z)$ has the form (1) and if

(A) $2 \leq n \leq 5$ [1, 6, 8],

(B) $p(0) = 0$ [10],

(C) $p(z) = z^n + a_{n-1}z^{n-1} + \dots + a_1z + a_0, \quad a_k < 0 \quad (0 < k < n - 1)$

[11],

(D) $p(z)$ has only real zeros [7], $(n_0 \leq n) \leq \frac{n_2}{n_m} \leq \binom{n}{3}$ or

(E) $p(z) = z^n + a_{n-1}z^{n-1} + \dots + a_1z + a_0, \quad (a_0 < n_1 < \dots < n_m) \leq \binom{n}{3}$, or

$P(z) = E^{\wedge=0} a_j^z$ —

(F) the vertices of the convex hull of the zeros of $p(z)$ all lie on $|z| = 1$ [10],

(G) the convex hull of the zeros of $p(z)$ is a triangular region [12],

(H) $p(z) = (z - z_1)(z - z_2)(z - z_3)^3$ [9].

The last case (H), due to Saff and Twomey, states that the conjecture is true for any polynomial of the form (1) with at most three distinct

zeros. The purpose of this paper is to establish the conjecture for any polynomial with at most four distinct zeros.

Observe that (H) follows immediately from (G). A problem posed by Schmeisser [12] related to (G) is to determine whether the conjecture is true if the convex hull of the zeros of $p(z)$ is a quadrangular region. Our result verifies a special case of his problem.

2. Main results. We can now state our main result.

THEOREM. *If $p(z) = (z - z_1)^{n_1}(z - z_2)^{n_2}(z - z_3)^{n_3}(z - z_4)^{n_4}$ with $|z_k| \leq 1$ ($1 \leq k \leq 4$), then each of the disks $|z - z_k| < 1$ ($1 \leq k \leq 4$) contains a zero of $p'(z)$.*

Before embarking on the proof we briefly illustrate the idea by giving a simple proof of the Saff-Twomey result. Suppose $p(z) = (z - z_1)^{n_1}(z - z_2)^{n_2}(z - z_3)^{n_3}(z - z_4)^{n_4}$ with $|z_k| < 1$. Let us distinguish one of the zeros, say z_3 , and for convenience let $z_3 = a$. Thus $p(z) = (z - a)^{n_3}(z - z_1)^{n_1}(z - z_2)^{n_2}(z - z_4)^{n_4}$. We wish to show that $|z - a| < 1$ contains a zero of $p'(z)$. If $n_3 > 1$ we are done, so suppose $n_3 = 1$. It is then clear that $p'(z) = n_1(z - z_1)^{n_1-1}(z - z_2)^{n_2-1}(z - z_4)^{n_4-1}(z - a)(n_1 + n_2 + 1 = n)$ and if we let $q(z) = p(z)/(z - a)$ then also $p'(z) = (z - a)g'(z) + q(z)$. Hence we see that $p'(a) = g(a)$ and so

$$n(a - z_1)^{n_1-1}(a - z_2)^{n_2-1}(a - z_4)^{n_4-1} = (a - z_1)^{n_1}(a - z_2)^{n_2}(a - z_4)^{n_4}$$

We may suppose that $|a - z_1| \leq |a - z_2|$ and get

$$n|a - z_1|^2 \leq |a - z_1| |a - z_2| \leq 4.$$

Hence $|a - z_1| \leq 2/\sqrt{n}$. If $n \geq 4$, we are done; while if $n = 2$ or 3 we already know the conjecture is true. The proof of the theorem is based on this simple idea, however we need some preliminary results first.

Fix integers n and m with $n \geq m \geq 2$ and let $\mathcal{M}(n, m)$ denote the class of all monic polynomials of degree n with at most m distinct zeros in $|z| \leq 1$:

$$(2) \quad p(z) = \prod_{k=1}^m H(z - z_k), \quad |z_k| \leq 1 \quad (1 \leq k \leq m),$$

where $n_k \geq 1$ and $\sum_{k=1}^m n_k = n$. From (2) we get

$$(3) \quad p'(z) = n \left[\prod_{k=1}^m (z - z_k)^{n_k-1} \right] \left[\prod_{j=1}^m (z - \zeta_j) \right].$$

Let

$$I(z_k) = \begin{cases} \min_{1 \leq j \leq m-1} |z_k - \zeta_j|, & \text{if } n_k = 1, \\ \mathbf{0}, & \text{if } n_k > 1. \end{cases}$$

$$I(p) = \max_{l < k < m} I(z_k) \quad \text{and} \quad I(\mathcal{P}_n(m)) = \sup_{p \in \mathcal{P}_n(m)} I(p).$$

The Ilieff-Sendov conjecture says that $I(\mathcal{P}_n(m)) < 1$.

LEMMA 1. (i) *There exists an extremal polynomial $p^* \in \mathcal{P}_n(m)$ such that $I(p^*) = I(\mathcal{P}_n(m))$.*

(ii) *p^* has a zero on each subarc of $\mathbb{U} \setminus \mathbb{R}$ of length n .*

This lemma is essentially proved in [5, 7] but for completeness sake and a slightly easier proof we present it here.

Proof. Since $\forall p \in \mathcal{P}_n(m)$, where $\mathbb{U} \setminus \mathbb{R} \subseteq r$, for any $p \in \mathcal{P}_n(m)$, it is clear that $3\mathcal{P}_n(m)$ is a normal family in \mathbb{C} . Each polynomial in $3\mathcal{P}_n(m)$ is monic and has at most m distinct zeros and hence $\mathcal{P}_n(m)$ is compact. By definition there exists a sequence $\{p^k\} \subset 3\mathcal{P}_n(m)$ such that $I(p^k) \rightarrow I(\mathcal{P}_n(m))$ as $k \rightarrow \infty$. Choose a convergent subsequence (call it $\{p^k\}$ again) so that $p^k \rightarrow p^*$ uniformly on compact subsets of \mathbb{C} . Let

$$(4) \quad p^*(z) = \prod_{k=1}^m (z - z_k^*)^{n_k^*}, \quad |z_k^*| \leq 1, \quad \sum_{k=1}^m n_k^* = n$$

and so

$$(5) \quad p^{*'}(z) = n \left[\prod_{k=1}^m (z - z_k^*)^{n_k^* - 1} \right] \left[\prod_{j=1}^{m-1} (z - \zeta_j^*) \right].$$

Assume $I(p^*) < I(\mathcal{P}_n(m))$. Then $I(p^*) = I(3\mathcal{P}_n(m)) - 3e$, for some $e > 0$. Choose $0 < S < e$ so that $p^* \pm 0$ in $0 < |z| \leq 1$ ($7 \leq m$). Thus all the zeros of p^* lie in $Q = \{z \in \mathbb{U} \setminus \mathbb{R} : |z - \zeta_j^*| \leq I(p^*)\}$ and (by definition of $I(p^*)$) at least one zero of $p^{*'}$ lies in each of the disks $|z - \zeta_j^*| \leq I(p^*)$ ($1 \leq j \leq m$). By Hurwitz' Theorem, for all $k \geq KQ$ sufficiently large, all the zeros of p^k lie in Q and each disk $|z - \zeta_j^*| \leq I(p^*) + e$ will contain a zero of p^k . Hence we see that $I(p^k) \leq I(p^*) + e + S < I(p^*) + 2e = I(3\mathcal{P}_n(m)) - e$. Letting $k \rightarrow \infty$ we get $I(\mathcal{P}_n(m)) < I(\mathcal{P}_n(m)) - e$ and a contradiction is reached. This proves (i).

To prove (ii) we first assert that the extremal polynomial p^* of the form (4) must have at least one zero on $|z| = 1$. Suppose not. Then $r = \max_{1 \leq k \leq m} |z_k| < 1$. Define \tilde{p} by

$$\tilde{p}(z) = \frac{1}{r^n p^*(r/z)} r z = \prod_{k=1}^m \left(z - \frac{z_k^*}{r} \right)^{n_k}.$$

Clearly $\tilde{p} \in TPrSjri$ and since $\tilde{p}'(z) = -phrP^{*'}(rz)$, we see that

$$I(\tilde{p}) = \max_{\substack{1 \leq k \leq m \\ n_k = 1}} \left[\min_{1 \leq j < m-1} \left| \frac{z_k^*}{r} - \frac{z_j^*}{r} \right| \right] = \frac{1}{r} I(p^*) > i(p^*).$$

Contradiction.

Next, we assert that p^* must have at least two distinct roots on $|z| = 1$. Suppose not. Then p^* has the form (4), with say $|z^*| = 1$ and $|z_k^*| < 1$ ($1 \leq k \leq m - 1$). By a rotation, we can assume $z_m^* = 1$. Let $\delta = (1 - r)/2$, where $r = \max_{1 \leq k \leq m} |z_k^*|$. If $\tilde{p}(z) = p^*(z + s)$, then again $\tilde{p} \in TPrSjri$ and $I(\tilde{p}) = I(p^*)$. Hence \tilde{p} is also extremal. However, it is easy to see that its zeros $z_k^{\sim} = z_k^* - s$ satisfy $|z_k^{\sim}| < 1$ ($1 \leq k \leq m$). Contradiction.

We have shown that the extremal polynomial p^* has the form (4) with $|z^*| = |z_{m-1}^*| = 1$ and $|z_k^*| < 1$. Assume now that there is some arc on $|z| = 1$ of length $L > \frac{2\pi}{n}$, say $L = 2\pi/n - \theta$ for some $0 < \theta < \pi/2$, on which $p^* \neq 0$. By a rotation, we may suppose that $z_m^* = e^{ia}$ and $z_{m-1}^* = e^{-ia}$ for some $0 < a < \theta < \pi/2$. (Thus $p^*(e^{ie}) \neq 0$ for $a < |\theta| < \pi$.) By relabeling the zeros suppose that $|z_1^*| \leq |z_2^*| \leq \dots \leq |z_J^*| < 1$ and $|z_{J+1}^*| = \dots = |z_{m-2}^*| = 1$ (put $n_Q - 0$ if all the zeros of p^* lie on $|z| = 1$). Define r and δ as follows:

$$r = \begin{cases} 1, & \text{if } n_0 = 0, \\ \max_{1 \leq k \leq n_0} |z_k^*|, & \text{if } n_0 > 0, \end{cases}$$

and

$$s = \min \left\{ \cos a, \frac{1 - r}{2} \right\}$$

Again consider $\tilde{p}(z) = p^*(z + s)$ and note that $\tilde{p} \in TPrSjri$ with $I(\tilde{p}) = I(p^*)$. An easy check shows that the zeros $z_k^{\sim} = z_k^* - s$ of \tilde{p} all lie in $|z| < 1$. Contradiction. This completes the proof of the lemma. •

LEMMA 2. The Ilieff-Sendov conjecture is true if

- (i) $p(z) = \prod_{k=1}^n (z - e^{i\theta_k})^2$ and $2 < n < 4$,
- (ii) $p(z) = (z - e^{i\theta_0})^2 \prod_{k=1}^n (z - z_k)$ and $2 < n < 4$ ($d_0 \in \mathbb{R}$).

The first part of this lemma is known and there are various proofs, but the proof of (ii) gives (i) along the way so we include it here. This now makes the paper completely self-contained.

Proof. Let $f(z) = \prod_{k=1}^{n-1} (z - z_k)$ with $|z_k| < 1$. Then $p'(z) = n \prod_{k=1}^{n-1} (z - z_k)^{-1}$. Distinguish one of the zeros, say z_n (call it a) and by a rotation let

$$p(z) = \prod_{k=1}^{n-1} (z - z_k), \quad 0 < a < 1, \quad |z_k| < 1$$

We may also suppose that $0 < a < 1$. If $a = 0$, the conjecture is trivially true for this zero. If $a = 1$, we are also done. Indeed, if we let $q(z) = \prod_{k=1}^{n-1} (z - z_k)$. Then we see that $p''(z)/p'(z) = 2q'(z)/q(z)$ and so

$$(6) \quad \sum_{j=1}^{n-1} \frac{1}{1 - \zeta_j} = \sum_{k=1}^{n-1} \frac{1}{1 - z_k}.$$

Supposing $\operatorname{Re}\{1/(1 - \zeta_j)\} > \operatorname{Re}\{1/(1 - \zeta_{j+1})\}$ ($1 \leq j < n-1$) and we may, then from (6) we get

$$(n-1) \operatorname{Re} \left\{ \frac{1}{1 - \zeta_1} \right\} \geq 2 \sum_{k=1}^{n-1} \operatorname{Re} \left\{ \frac{1}{1 - z_k} \right\} \geq (n-1).$$

Hence $\operatorname{Re}\{1/(1 - \zeta_1)\} \geq 1$ or $|\zeta_1| < 1$ so certainly $|\zeta_1 - 1| < 1$ (cf. [2]). Hence suppose $0 < a < 1$.

If we put $z = T(w) = (w - a)/(aw - 1)$, we have

$$(7) \quad p(T(w)) = \tilde{p}(w)(aw - 1)^{-n},$$

where

$$(8) \quad \tilde{p}(w) = Aw [w^{n-1} + bn - iiv^{n-2} + \dots + *].$$

From (7), the zeros of $\tilde{p}(w)$ are $0, w_1, w_2, \dots, w_{n-1}$ where $w_k = T^{-1}(z_k)$ ($1 \leq k \leq n-1$). Thus we get

$$(9) \quad |b_1| = \prod_{k=1}^{n-1} |w_k| \leq 1$$

and

$$(10) \quad |v_1| = \left| \sum_{k=1}^{n-1} w_k \right| < n-1.$$

Differentiating (7) gives

$$\frac{dp\{T(w)\}}{dw} = \frac{dp\{T(w)\}}{dz} \frac{dz}{dw} = -a(aw-l)^{n-1} D_{1/a}p(w),$$

where

$$(11) \quad D_{1/a}\tilde{p}(w) = np(w) + (1/a - w)p'(w)$$

is the polar derivative of p with respect to $1/a$ (see Marden [4]). Hence we arrive at

$$(12) \quad p'(T(w)) = -a(aw - l)^{-1} \frac{dw}{dz} D_{1/a}\tilde{p}(w)$$

(where ' denotes differentiation with respect to z). A brief calculation using (8) and (11) gives

$$(13) \quad D_{1/a}\tilde{p}(w) = B \left[w^{n-1} + \dots + \left(\frac{b_1}{n + ab_{n-1}} \right) \right] = B \prod_{k=1}^{n-1} (w - \gamma_k),$$

where $|\gamma_1| < |\gamma_2| < \dots < |\gamma_{n-1}|$. It follows from (9), (10) and (13) that

$$(14) \quad \prod_{k=1}^{n-1} |\gamma_k| = \left| \frac{b_1}{n + ab_{n-1}} \right| \leq \frac{1}{n - a(n-1)}$$

Let us now suppose that $|\gamma_k| \leq \mu$. Then from (12) we have

$$D_{1/a}P(V) = \frac{a}{a} (\alpha\gamma_1 - 1)^{n+1} \wedge P \wedge (T(m)) = o.$$

Hence $|(Co) = 0$, where $Co = T(\gamma_x)$ and so $p'(z)$ has a zero Co such that

$$|\gamma_1| = |T^{-1}(\zeta_0)| = \left| \frac{Co - a}{aCo - 1} \right| \in \mathbb{R}, 0 < r < JX. \text{ This}$$

Thus we get $(Co - a)/i^a Co - 1) = r^{1/e_3}$ and we conclude that $-$ gives $Co = (\ll - re^{16})l(\wedge - are^{16})$

$$|\zeta_0 - a| = \frac{r(1 - a^2)}{|1 - are^{16}|} \leq \frac{\mu(1 - a^2)}{1 - a/Z}.$$

Hence if $\mu(1 - a^2)/(1 - a\mu) \leq 1$, or equivalently,

$$(15) \quad \mu < \frac{1}{1 + a - a^2},$$

then $Co = o_{\wedge} < 1$. It remains to show that (15) holds for the cases stated in the lemma.

Suppose first that $2 \leq n \leq 4$. Then from (14) we have

$$|y_1| \leq \left[\frac{1}{n-a(n-\lambda)_m} \right]^{1/(n-1)} \equiv \mu.$$

It is easy to check that (15) holds for this f_1 and $2 \leq n \leq 4$ (and any $0 < a < 1$). This proves (i).

Suppose next that $p(z)$ has the form stated in (ii). Then (for some $U, Q \in \mathbb{R}$) we conclude that $p(w)$ has a double root at $T^{-1}(e^{i\theta}) = T^{-1}(e^{i\theta})$. From (11) we see that $y_{n-\lambda} = e^m$ is a zero of $D_j a p(w)$. Using (14) we have the following

$$\prod_{k=1}^{n-1} \prod_{k=1}^{n-\lambda} \frac{1}{n-a(n-1)}$$

and so

$$|y_1| \leq \left[\frac{1}{n-a(n-1)} \right]^{1/(n-2)} \equiv \mu.$$

For this n and $2 \leq n \leq 7$, it is simple to check that (15) holds for any $0 < a < 1$.

We should point out that this proof can be slightly modified to give a proof of Laguerre's Theorem (see [4] for another proof).

Proof of theorem. For fixed $n \geq m \geq 3$, let $p(z)$ be of the form (2). From Lemma 1, an extremal polynomial for $\mathcal{P}_n^{\circ}(m)$ exists and without loss of generality we may assume p is extremal i.e., $I(p) = I(nim)$. Distinguish one of the zeros, say z_m and let $z_m = a$. We may suppose too that $0 \leq a \leq 1$. We want to show $|z - a| \leq 1$ contains a critical point of p . If $n_m > 1$, we are done so suppose $n_m = 1$. Thus we have

$$(16) \quad p(z) = (z - a) \prod_{k=\lambda}^{m-1} (z - z_k)^{n_k}, \quad |z_k| < 1, \quad \sum_{k=1}^{m-1} n_k = n - 1$$

and

$$(17) \quad p'(z) = n \left[\prod_{l=1}^{w-1} (z - z_k)^{n_k-1} \right] \left[\prod_{j=1}^{m-1} (z - \zeta_j) \right].$$

As in the proof of Lemma 2 we may suppose that $0 < a < 1$. Using (16) and (17), we obtain after cancellations

If we let $q(z) = YI^{1/2} \sim \dots$

$$n \prod_{j=1}^{m-\lambda} (a - \zeta_j) = \prod_{k=1}^{m-\lambda} (a - z_k).$$

Assuming that $|a - \zeta_j| \leq |a - \zeta_j|$ ($1 \leq j \leq m - 1$), as we may, we get

$$n|a - \zeta_i| \leq C \prod_{k=i}^{m-1} \frac{|a - z_k|}{|a - z_{j+1}|} 2^{m-k}$$

Hence we have the estimate

$$(18) \quad |a - \zeta_i| \leq \left[\frac{1}{n} \right]^{1/(m-1)}$$

where z_j and z_2 are any two distinct zeros of $p(z)$ other than $z = 1$ and

Suppose first that $n \geq 2^{m-1}$. From (18) we see that $|a - \zeta_i| < 1$ and we are done. Hence suppose $2 < n < 2^{m-1}$.

Now we restrict ourselves to $m = 4$. Thus we need only consider polynomials of the form (16) of degree $2 \leq n < 8$. By Lemma 2(i) we need only consider $n = 5, 6$ and 7 . Thus, our polynomial has the form

$$p(z) = (z - a)(z - z_1)^{n_1} (z - z_2)^{n_2} (z - z_3)^{n_3}$$

where $0 \leq n_1 \leq n_2 \leq n_3$ and $n_1 + n_2 + n_3 = n - 1$. Since $n = 5, 6$ or 7 , we must have $n_3 \geq 2$ in each case. Recall that $p(z)$ is an extremal polynomial for $\mathcal{P}_n^{\circ}(m)$ and hence by Lemma 1 it will have a zero on each subarc of $|z| = 1$ of length n . We are thus led to two cases:

Case 1. $p(z)$ has exactly two distinct zeros on $|z| = 1$.

Case 2. $p(z)$ has more than two distinct zeros on $|z| = 1$.

Case 2 is easily disposed of as follows. In this case (recall $0 < a < 1$), we must have $|z_1| = |z_2| = |z_3| = 1$ and since $n_3 \geq 2$, we see that p has a zero of order two on $|z| = 1$. Since $n \leq 7$, we may apply Lemma 2(ii) and we are done.

In order for Case 1 to hold, the two zeros must be negatives of each other, say ZQ and $-ZQ$. Making use of the estimate (18) we find that

$$|a - \zeta_i| \leq \left[\frac{2|a - z_0||a + z_0|}{n} \right]^{1/3} \leq \left[\frac{4}{n} \right]^{1/3}$$

and so $|a - \zeta_i| \leq 1$ for $n \geq 4$. By Lemma 2(i), the conjecture is true for $n = 2$ and 3 . The proof of the theorem is complete. D

3. Remarks. The Ilieff-Sendov conjecture can be interpreted physically in terms of force fields. Indeed, if C_0 is a critical point of $P(z) = \prod_{i=1}^n (z - z_i)$,

equilibrium point in the plane field in which the force exerted on a particle by a point charge at z_k (with charge n_k) is inversely proportional to its distance from z_k . The Ilieff-Sendov conjecture asserts that either the

disk of radius one about each zero z_k contains at least one of these equilibrium points or else $n_k > 1$. In general these cases are mutually exclusive. For example if $p(z) = (z - 1)^3(z - i)$, then the disk $|z - 1| \leq 1$ contains no such equilibrium point. Since the conjecture is trivially true for the zero z_k if $n_k > 1$, the conjecture is only interesting if z_k is a simple zero. In this case we are led to the problem of determining just how close a critical point can then be to a simple zero. We thus pose the following problem:

Let Q_n denote the set of all polynomials for the form $P(z) = z^n \prod_{k=1}^{n-1} (z - z_k)$, with $|z_k| > 1$ ($1 \leq k \leq n - 1$). Determine the largest constant $c_n > 0$ such that $p'(z) \neq 0$ in $|z| < c_n$ for all $p \in Q_n$.

Clearly $c_n < 1$ and we expect that the critical point nearest the zero $z = 0$ is an equilibrium point in the force field as described above. By concentrating all the charge at one point, together with a single charge at $z = 0$, it is reasonable to conjecture that $c_n = 1/n$ (consider $p(z) = z(z - e^{ie})^n$).

Finally, we must point out that recently a proof of the Ilieff-Sendov conjecture was announced (by title) in the *Abstracts of the American Mathematical Society* (June 1986) by V. I. Istrătescu. He claims his method uses a notion called "bare points" and "a Krein-Milman theorem". The proof of our main result is completely self-contained, relies on classical methods and indicates that a proof of the full conjecture must be delicate in nature.

REFERENCES

- [1] D. A. Brannan, *On a conjecture of Ilieff*, Proc. Cambridge Philos. Soc, 64 (1968), 83-85.
- [2] A. W. Goodman, Q. Rahman and J. Ratti, *On the zeros of a polynomial and its derivative*, Proc. Amer. Math. Soc, 21 (1969), 273-274.
- [3] W. K. Hayman, *Research Problems in Function Theory*, London (1967).
- [4] M. Marden, *Geometry of Polynomials*, Amer. Math. Soc, (1966).
- [5] _____, *Conjectures on the critical points of a polynomial*, Amer. Math. Monthly, 90(1980), 267-276.
- [6] A. Meir and A. Sharma, *On Ilieff's conjecture*, Pacific J. Math., 31 (1969), 459-467.
- [7] D. Phelps and R. Rodriguez, *Some properties of extremal polynomials for the Ilieff Conjecture*, Kodai Math. Sem. Report, 24 (1972), 172-175.
- [8] Z. Rubinstein, *On a problem of Ilieff*, Pacific J. Math., 26 (1968), 159-161.
- [9] E. B. Saff and J. Twomey, *A note on the location of critical points of polynomials*, Proc. Amer. Math. Soc, 27 (1971), 303-308.

- [10] G. Schmeisser, *Bermerkungen zueiner Vermutung von Ilieff*, Math. Z., 111 (1969), 121-125.
- [11]_____, *Zur Lage der kritischen Punkte eines Polynoms*, Rend. Sem. Mat. Univ. Padova.
- [12]_____, *On Ilieff's conjecture*, Math. Z., 156 (1977), 165-173.

Received June 9, 1987. Some of this work was done while the author was on sabbatical leave at the University of Delaware.

PURDUE UNIVERSITY
WEST LAFAYETTE, IN 47907

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024
HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721
VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720
C. C. MOORE
University of California
Berkeley, CA 94720
HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS E. F. BECKENBACH B. H. NEUMANN F. WOLF K. YOSHIDA
(1906-1982)

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the scheme of Math. Reviews, Index to Vol. 39. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (5 Vols., 10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) publishes 5 volumes per year. Application to mail at Second-class postage rates is pending at Carmel Valley, California, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Copyright © 1988 by Pacific Journal of Mathematics

Waleed A. Al-Salam and Mourad Ismail, q -beta integrals and the q -Hermite polynomials	209
Johnny E. Brown, On the Ilieff-Sendov conjecture	223
Lawrence Jay Corwin and Frederick Paul Greenleaf, Spectrum and multiplicities for restrictions of unitary representations in nilpotent Lie groups	233
Robert Jay Daverman, 1-dimensional phenomena in cell-like mappings on 3-manifolds	269
P. D. T. A. Elliott, A localized Erdős-Wintner theorem	287
Richard John Gardner, Relative width measures and the plank problem	299
F. Garibay, Peter Abraham Greenberg, L. Reséndis and Juan José Rivaud, The geometry of sum-preserving permutations	313
Shanyu Ji, Uniqueness problem without multiplicities in value distribution theory	323
Igal Megory-Cohen, Finite-dimensional representation of classical crossed-product algebras	349
Mirko Navara, Pavel Pták and Vladimír Rogalewicz, Enlargements of quantum logics	361
Claudio Nebbia, Amenability and Kunze-Stein property for groups acting on a tree	371
Chull Park and David Lee Skoug, A simple formula for conditional Wiener integrals with applications	381
Ronald Scott Irving and Brad Shelton, Correction to: "Loewy series and simple projective modules in the category \mathcal{O}_S "	395
Robert Tijdeman and Lian Xiang Wang, Correction to: "Sums of products of powers of given prime numbers"	396