

Pacific Journal of Mathematics

FANO BUNDLES OVER P^3 AND Q_3

MICHAŁ SZUREK AND JAROSŁAW WISNIEWSKI

FANO BUNDLES OVER P^3 AND Q_3

MICHAŁ SZUREK AND JAROSŁAW A. WIŚNIEWSKI

A vector bundle \mathcal{E} is called Fano if its projectivization $P(\mathcal{E})$ is a Fano manifold. In this article we prove that Fano bundles exist only on Fano manifolds and discuss rank-2 Fano bundles over the projective space P^3 and a 3-dimensional smooth quadric Q_3 .

Fano bundles appear naturally as we strive to construct examples of Fano manifolds of dimension ≥ 3 ; they form interesting yet accessible class of Fano n -folds. For example: among 87 types of Fano 3-folds with $b_2 \geq 2$ listed in [13] 22 types are ruled (i.e. obtained by projectivization of Fano bundles). Moreover some of the non-ruled manifolds listed there can be easily expressed as either finite covers of ruled 3-folds or divisors (or, more generally, complete intersections) in ruled Fano manifolds of higher dimension.

Let us mention another aspect of dealing with Fano bundles: it is how to determine whether or not a vector bundle is ample. This very fine property of a vector bundle cannot be determined by its numerical invariants, see [7]. Assuming the bundle to be stable helps to establish a sufficient condition for ampleness: [10], [17], which however is far from being necessary. In the present paper we take advantage of some already known facts about stable bundles with small Chern classes and determine that a bundle \mathcal{E} is not ample by finding its jumping lines or sections of $\mathcal{E}(-k)$.

Let us note that some results of this paper have already been published, see remarks after the proofs of Theorems (1.6) and (2.1).

1. Fano bundles; preliminaries. Let \mathcal{E} be a vector bundle of rank $r \geq 2$ on a smooth complex projective variety M . Let us recall that the tautological line bundle $\xi = \xi_{\mathcal{E}}$ on $V = P(\mathcal{E})$ is uniquely determined by the conditions $\xi|_F \approx \mathcal{O}_F(1)$ and $p_*\xi_{\mathcal{E}} = \mathcal{E}$. By p we have denoted the projection morphism of $V = P(\mathcal{E})$ onto M and by F —the fibre of p . Obviously, $F \cong P^{r-1}$ and $p: V \rightarrow M$ is a P^{r-1} -bundle. The Picard group of V can be expressed as a direct sum: $\text{Pic} V \cong \mathbb{Z} \cdot \xi_{\mathcal{E}} \oplus p^*(\text{Pic} M)$. Replacing \mathcal{E} by its twist with a line bundle \mathcal{L} on M does not affect

the projectivization and

$$\xi_{\mathcal{E} \otimes \mathcal{L}} = \xi_{\mathcal{E}} \otimes p^*(\mathcal{L}).$$

Moreover, \mathcal{E} is generated by global sections iff $\xi_{\mathcal{E}}$ is. We have the following relative Euler sequence on $V = P(\mathcal{E})$:

$$(1.1) \quad 0 \rightarrow \mathcal{O}_V \rightarrow p^*(\mathcal{E})^\vee \otimes \xi_{\mathcal{E}} \rightarrow T_{V|M} \rightarrow 0$$

where the latter bundle is the relative tangent bundle of p and fits in the exact sequence

$$(1.2) \quad 0 \rightarrow T_{V|M} \rightarrow TV \rightarrow p^*TM \rightarrow 0.$$

We then obtain

$$(1.3) \quad c_1 V = p^*(c_1 M - c_1 \mathcal{E}) + r \xi_{\mathcal{E}}$$

The theorem of Leray and Hirsch yields that in the cohomology ring of V the following holds

$$(1.4) \quad \xi_{\mathcal{E}}^r - p^*(c_1 \mathcal{E}) \xi_{\mathcal{E}}^{r-1} + p^*(c_2 \mathcal{E}) \xi_{\mathcal{E}}^{r-2} - \cdots \pm p^*(c_r \mathcal{E}) = 0.$$

From now on we assume in this section that \mathcal{E} is a rank- r Fano bundle on an n -fold M , i.e., that $\mathbb{P}(\mathcal{E})$ is a Fano manifold. We prove that such M must be Fano, as well.

(1.5) LEMMA. *Let $C \subset M$ be a rational curve with a normalization $\nu: P^1 \rightarrow C$. Assume that $\nu^*(\mathcal{E}) \cong \mathcal{O}(a_1) \oplus \mathcal{O}(a_2) \oplus \cdots \oplus \mathcal{O}(a_r)$, where $a_1 \leq a_2 \leq \cdots \leq a_r$. Then*

$$(c_1 M) \cdot C > \sum_{i=2}^r (a_i - a_1) \geq 0.$$

Proof. The right hand side inequality is obvious. To prove the left hand side inequality let us assume that $W = P(\nu^*\mathcal{E})$. The manifold W is then a P^{r-1} -bundle over P^1 , with a projection $\pi: W \rightarrow P^1$. We have a section C_0 of π associated to the epimorphism $\nu^*\mathcal{E} \rightarrow \mathcal{O}(a_1) \rightarrow 0$, such that

$$\xi_{\nu^*\mathcal{E}}|_{C_0} \cong \mathcal{O}_{P^1}(a_1).$$

The normalization map $\nu: P^1 \rightarrow M$ lifts to a map $\bar{\nu}: W \rightarrow V$, making the following diagram commute

$$\begin{array}{ccc} W & \xrightarrow{\bar{\nu}} & V \\ \pi \downarrow & & \downarrow p \\ P^1 & \xrightarrow{\nu} & M \end{array}$$

By the choice of C_0 we have

$$\bar{\nu}^*(\xi_{\mathcal{E}}) \cdot C_0 = a_1$$

and, since $c_1 V$ is ample, we obtain by (1.3)

$$\begin{aligned} 0 &< c_1 V \cdot \bar{\nu}(C_0) = \bar{\nu}^*(c_1 V) \cdot C_0 \\ &= r \cdot \bar{\nu}^*(\xi_{\mathcal{E}}) \cdot C_0 + (\pi \circ \nu)^*(c_1 M) \cdot C_0 - (\pi \circ \nu)^*(c_1 \mathcal{E}) \cdot C_0 \\ &= r \cdot a_1 + c_1 M \cdot C - \sum_{i=1}^r a_i \end{aligned}$$

which yields the desired inequality.

(1.6) **THEOREM.** *If \mathcal{E} is a Fano bundle on a manifold M then M is a Fano n -fold.*

Proof. As $c_1 V = -K_V$ is ample, the cone of curves on V is spanned by the classes of extremal curves (see [12] for definitions and Theorem 1.2 on the cone of curves of a Fano manifold). Let us denote these curves by l_0, l_1, \dots, l_v with l_0 contained in F , a fibre of the projection $p: P(\mathcal{E}) \rightarrow M$. We see that $p^*(c_1 M) \cdot l_0 = 0$ and for $i > 0$, $p(l_i)$ is a rational curve on M . Therefore from (1.5) it follows that

$$(1.7) \quad 0 < c_1 M \cdot p(l_i) = p^*(c_1 M) \cdot l_i$$

which means that $p^*(c_1 M)$ is numerically effective. Recall now (a conclusion from) the Kawamata-Shokurov contraction theorem, see (2.6) in [11]:

If D is nef and $aD - K$ is ample for some $a > 0$, then D is semiample, i.e., some power of D is generated by global sections.

It follows that $D := p^*(c_1 M)$ is semiample. Since $p: V \rightarrow M$ is a P^{r-1} -bundle, we have, for any integer k , $p_* p^*(\mathcal{O}(kc_1 M)) = \mathcal{O}(kc_1 M)$ and the images (under p_*) of global sections of $p^*(\mathcal{O}(kc_1 M))$ are global sections of $\mathcal{O}(kc_1 M)$. Therefore $c_1 M$ is semiample, hence to prove that it is ample it is enough to show that $c_1 M \cdot C > 0$ for any curve C in M .

Let C be an irreducible curve in M . Taking an appropriate component from an intersection of the inverse image $p^{-1}(C)$ with general $r - 1$ divisors from a very ample linear system, we can produce an irreducible curve $C_1 \subset V$, such that $p(C_1) = C$. Then C_1 is numerically equivalent to a linear combination $\sum a_i l_i$ with at least one a_i different from zero for $i > 0$. Let d be the degree of the map $p|_{C_1}: C_1 \rightarrow C$.

Now the inequality (1.7) gives

$$c_1 M \cdot C = \frac{1}{d} \cdot p^*(c_1 M) \cdot C_1 = \frac{1}{d} \cdot \left(\sum a_i \cdot p^*(c_1 M) \cdot l_i \right) > 0,$$

which concludes the proof of the theorem.

REMARK. Theorem (1.6) has already been known for bundles of rank 2 on surfaces [4] and 3-folds [1].

2. Rank-2 Fano bundles on P^3 . The results stated below (Theorem (2.1)) can be understood as one more example of an exceptional character of the null-correlation bundle (see e.g. [3] or [15] for the definition of the null-correlation bundle).

(2.1) THEOREM. *The only rank-2 Fano bundles with $c_1 = 0, -1$, on \mathbb{P}^3 are*

- (1) $\mathcal{E} = \mathcal{O} \oplus \mathcal{O}$,
- (2) $\mathcal{E} = \mathcal{O} \oplus \mathcal{O}(-1)$,
- (3) $\mathcal{E} = \mathcal{O}(-1) \oplus \mathcal{O}(+1)$,
- (4) $\mathcal{E} = \mathcal{O}(-2) \oplus \mathcal{O}(+1)$,
- (5) *the null-correlation bundle \mathcal{N} .*

Proof. Let $V = P(\mathcal{E})$. We then have

$$(2.2) \quad -K_V = 2\xi + (4 - c_1 \mathcal{E})H \\ = \begin{cases} 2\xi + 4H = 2\xi_{\mathcal{E}(1)} + 2H = 2\xi_{\mathcal{E}(2)} & \text{if } c_1 = 0, \\ 2\xi + 5H = 2\xi_{\mathcal{E}(2)} + H = 2\xi_{\mathcal{E}(3)} - H & \text{if } c_1 = -1, \end{cases}$$

and we see that any of the bundles listed above is Fano. Indeed, if \mathcal{E} is one of those listed as (1), (3) or (5) (respectively: (2) or (4)) then $c_1 \mathcal{E} = 0$ (resp. $c_1 \mathcal{E} = -1$) and $\mathcal{E}(1)$ (resp. $\mathcal{E}(2)$) is generated by its global sections. Now, since $\rho(V) = 2$, it follows from (2.2) that $c_1 V$ is ample as the sum of two non-proportional nef divisors.

An easy corollary follows.

(2.3) For a normalized Fano bundle \mathcal{E} of rank 2 on P^3 :

if $c_1 \mathcal{E} = 0$, then $\mathcal{E}(2)$ is ample,

if $c_1 \mathcal{E} = -1$, then $\mathcal{E}(3)$ is ample.

We shall discuss the two cases separately.

Case $c_1 = 0$. A straightforward consequence of the theorem of Leray and Hirsch (1.4) yields that in the cohomology ring of V the following holds

$$\xi^2 + c_2 H^2 = 0.$$

Since $H^4 = 0$ and $H^3\xi = 1$, the above formula then gives

$$(2.4) \quad H^2\xi^2 = 0, \quad H\xi^3 = -c_2, \quad \xi^4 = 0,$$

so that $(-K_V)^4 = (2\xi + 4H)^4 = 128(4 - c_2)$ and we see that $c_2 < 4$.

Assume first \mathcal{E} is not semistable, i.e., $H^0(\mathcal{E}(-1)) \neq 0$. Let s be a non-zero section of $\mathcal{E}(-1)$. We claim that s does not vanish anywhere. Indeed, if $Z = \{s = 0\}$ were not empty, then for a line L meeting Z in a finite number of points we would have

$$\mathcal{E}(-1)|L = \mathcal{O}(d) \oplus \mathcal{O}(e) \quad \text{with } d \geq 1, d + e = -2,$$

contradicting (2.3). Therefore s does not vanish and thus $\mathcal{E}(-1) = \mathcal{O} \oplus \mathcal{O}(-2)$, hence \mathcal{E} is as in (3) of the theorem.

Let now \mathcal{E} be semistable but not stable: $H^0(\mathcal{E}(-1)) = 0$, $H^0(\mathcal{E}) \neq 0$. If a non-zero section of \mathcal{E} does not vanish anywhere, \mathcal{E} must then be $\mathcal{O} \oplus \mathcal{O}$. Otherwise a section vanishes on a curve. If the curve is not a single line then cutting it by a line leads to a contradiction, as above. But if a single line L was a zero set of a section of \mathcal{E} then, by the adjunction formula, the degree of the canonical divisor of L would be

$$\deg(K_L) = (K_{P^3} + c_1\mathcal{E}) \cdot L = -4,$$

which is impossible. Because of Bogomolov's inequality $c_1^2 < 4c_2$ for stable bundles, [15], it remains then to study stable bundles with $c_1 = 0$ and $c_2 = 1, 2, 3$. In the first case \mathcal{E} is the null-correlation bundle \mathcal{N} , for which $\mathcal{N}(2)$ is ample; \mathcal{N} is then Fano.

In the remaining cases we know that \mathcal{E} has multiple jumping lines, i.e. such lines L for which $\mathcal{E}|L = \mathcal{O}_L(-2) \oplus \mathcal{O}_L(2)$, see [8], Proposition 9.11, and [18], respectively. In virtue of (2.3), such bundles cannot be Fano.

Case $c_1 = -1$. The multiplication table is now:

$$(2.5) \quad H^4 = 0, \quad H^3\xi = 1, \quad H^2\xi^2 = -1,$$

$$H\xi^3 = -c_2 + 1, \quad \xi^4 = 2c_2 - 1$$

and from

$$(-K_V)^4 = (2\xi + 5H)^4 = 32(-4c_2 + 17) > 0$$

we obtain that the only possible non-negative values for c_2 are 0, 2 or 4 (recall that Schwarzenberger's condition says $c_1c_2 \equiv 0 \pmod{2}$). Assume $H^0(\mathcal{E}(-1)) \neq 0$. As above, we show that no section $s \neq 0$

vanishes: if $Z = \{\text{zero}(s)\}$ were not empty, for a line L meeting Z at finitely many points we would have

$$\mathcal{E}(-1)|L = \mathcal{O}_L(d) \oplus \mathcal{O}_L(e) \quad \text{with } d \geq 1, d + e = -3,$$

contradicting (2.3). Therefore the sections $\mathcal{E}(-1)$ do not vanish anywhere, so that \mathcal{E} is as in (4) of Theorem (2.1).

Let then $H^0(\mathcal{E}(-1)) = 0$, $H^0(\mathcal{E}) \neq 0$. The zero set Z of a non-zero section is then a curve (if not empty). Again, if Z were anything different from a single line, for a line L that cuts Z at a finite number ≥ 2 of points we would have

$$\mathcal{E}|L = \mathcal{O}_L(d) \oplus \mathcal{O}_L(e), \quad d \geq 2, d + e = -1,$$

contradicting the ampleness of $\mathcal{E}(3)$. But $c_1 c_2$ is even so that the case $c_1 = -1$, $c_2 = 1$ does not hold, hence Z is not a line. The non-zero sections of \mathcal{E} do not vanish, hence $\mathcal{E} = \mathcal{O} \oplus \mathcal{O}(1)$.

It remains to exclude the cases of stable vector bundles with $c_1 = -1$ and $c_2 = 2$ or 4 . In the former case \mathcal{E} has multiple jumping lines, [9], Proposition 4.1, i.e., those for which $\mathcal{E}|L = \mathcal{O}_L(-3) \oplus \mathcal{O}_L(2)$, hence \mathcal{E} cannot be Fano in view of (2.3). In the latter one $\mathcal{E}(2)$ has a section, see [2], Lemma 1, and $2H + \xi$ is effective with

$$(2H + \xi)(c_1 P(\xi))^3 = (2H + \xi)(2\xi + 5H)^3 = -17.$$

These bundles are then not Fano.

REMARK. Theorem (2.1) (in a somewhat weaker form) was first announced by Artiushkin, [1]. His proof was, however, incorrect: in line 36 on page 14 if E is a normalized bundle on P^3 , then the tautological divisor $\xi_E = L$ in op. cit. need not to be effective, therefore $(-K)^3 \cdot L$ need not to be positive. Our actual proof is more complicated.

Let us conclude this section by proving that $P(\mathcal{N})$ has a P^1 -bundle structure over a 3-dimensional quadric Q_3 . To see this, first let us recall that $\mathcal{N}(1)$ can be defined as the bundle fitting in the following exact sequence on P^3

$$0 \rightarrow \mathcal{O} \rightarrow \Omega P^3(2) \rightarrow \mathcal{N}(1) \rightarrow 0.$$

Note that $P(\Omega P^3(2))$ is the incidence variety

$$I = \{(x, l) \in P^3 \times \text{Grass}(1, 3) : x \in l\}$$

and $\text{Grass}(1, 3)$ is isomorphic to a 4-dimensional quadratic. Now, from the above exact sequence it follows that $P(\mathcal{N}(1))$ is a divisor in I which is an inverse image of a hyperplane section of $\text{Grass}(1, 3)$.

Therefore:

(2.6) PROPOSITION. *The Fano 4-fold $P(\mathcal{N}(1))$ is a projectivization of a rank-2 vector bundle on smooth quadratic $Q_3 \subset \text{Grass}(1, 3)$, obtained by restricting to Q_3 the universal quotient bundle from $\text{Grass}(1, 3)$.*

3. Bundles over Q_3 . Let us recall that the cohomology ring of Q_3 is generated by the classes of $[H] \in H^2(Q_3, \mathbb{Z})$, $[L] \in H^4(Q_3, \mathbb{Z})$, and $[P] \in H^6(Q_3, \mathbb{Z})$ where H , L and P are a quadratic surface, a line and a point, respectively. There are the following relationships: $[H]^2 = 2L$, $[H][L] = [P]$ and hence $[H]^3 = 2[P]$. If \mathcal{F} is a coherent sheaf on Q_3 with the Chern polynomial

$$1 + c_1(\mathcal{F})[H]t + c_2(\mathcal{F})[L]t^2 + c_3(\mathcal{F})[P]t^3,$$

then the numbers c_i are called the Chern classes of \mathcal{F} .

Recall the Riemann-Roch formula for \mathcal{F} , [5]

$$\chi(\mathcal{F}) = \frac{1}{6}(2c_1^3 - 3c_1c_2 + 3c_3) + \frac{3}{2}(c_1^2 - c_2) + \frac{13}{6}c_1 + \text{rank } \mathcal{F}.$$

Let now \mathcal{E} be a rank-2 vector bundle on Q_3 . The theorem of Leray and Hirsch (1.4) gives the following relations between the generators of $\text{Pic}(P(\mathcal{E})) \cong \mathbb{Z} \oplus \mathbb{Z}$

$$\begin{cases} \text{if } c_1 = 0, \text{ then } \xi^2 + \frac{1}{2}c_2(\mathcal{E})H^2 = 0; \\ \text{if } c_1 = -1, \text{ then } \xi^2 + \xi H + \frac{1}{2}c_2(\mathcal{E})H^2 = 0. \end{cases}$$

Because $H^4 = 0$ and $H^3\xi = 2$, we obtain:

$$\text{if } c_1 = 0, \text{ then } H^2\xi^2 = 0, \quad H\xi^3 = -c_2, \quad \xi^4 = 0;$$

$$\text{if } c_1 = -1, \text{ then } H^2\xi^2 = -2, \quad H\xi^3 = 2 - c_2, \quad \xi^4 = 2c_2 - 2.$$

Let \mathcal{E} be a normalized rank-2 vector bundle on Q_3 and $V = P(\mathcal{E})$ its projectivization. We then have

$$(3.1) \quad c_1 V = -K_V = \begin{cases} 2\xi + 3H & \text{when } c_1 = 0, \\ 2\xi + 4H & \text{for } c_1 = -1. \end{cases}$$

Case of non-stable bundles. Assume \mathcal{E} is non-stable with $c_1(\mathcal{E}) = -1$. If a non-zero section from $H^0(\mathcal{E}(-1))$ vanishes at some point, let us consider a line L passing through this point and not contained in the zero set entirely. Then $\mathcal{E}(-1)|L = \mathcal{O}(d) \oplus \mathcal{O}(e)$ with $d \geq 1$, $d + e = -3$ that contradicts the ampleness of $\mathcal{E}(2)$, (3.1).

Assume $H^0(\mathcal{E}(-1)) = 0$, $H^0(\mathcal{E}) \neq 0$. Then a non-zero section of \mathcal{E} either does not vanish anywhere or it vanishes on a set of pure dimension 1. The divisor $\xi_{\mathcal{E}}$ is effective on $P(\mathcal{E})$ and

$$\xi \cdot (-K_V)^3 = 8\xi(\xi + 2H)^3 = 16(-2c_2 + 1),$$

and we see that $c_2 \leq 0$. But then sections of \mathcal{E} do not have zeros, hence $\mathcal{E} = \mathcal{O}(-1) \oplus \mathcal{O}$. Finally, we easily check that $\mathcal{O}(-1) \oplus \mathcal{O}$ is a Fano bundle (because $\mathcal{O}(1) \oplus \mathcal{O}(2)$ is ample).

In case $c_1 = 0$ we exclude non-semistable bundles in a very similar way. Finally, if \mathcal{E} is semistable but not stable, that is $H^0(\mathcal{E}) \neq 0 = H^0(\mathcal{E}(-1))$, the divisor $\xi_{\mathcal{E}}$ is effective and

$$0 < \xi(2\xi + 3H)^3 = 18(-2c_2 + 3)$$

so that $c_2 \leq 0$ (recall that $c_2 \equiv 0 \pmod{2}$, see [5], §1). If so, a non-zero section of \mathcal{E} does not vanish anywhere and \mathcal{E} must then be $\mathcal{O} \oplus \mathcal{O}$.

Case of stable bundles with $c_1 = 0$. From the condition $K^4 > 0$ we easily obtain that if $V = P(\mathcal{E})$ is Fano, then $c_2 \leq 4$, and since $c_2 \equiv 0 \pmod{2}$ it follows that either $c_2 = 2$ or 4 . We believe that there is no Fano bundle on Q_3 with $c_1 = 0$, $c_2 = 4$, however we do not have enough information on these bundles to prove it.

In case of $c_2 = 2$, one can easily check that the pull-back $\pi^*(\mathcal{N})$ of the null-correlation bundle, under a double covering $\pi: Q_3 \rightarrow P^3$, is Fano. Indeed, $\pi^*(\mathcal{N})(1)$ is then spanned on Q_3 , therefore $-K_{P(\pi^*(\mathcal{N}))} = 2\xi_{\pi^*(\mathcal{N})(1)} + H$ is ample. On the other hand we have

(3.2) PROPOSITION. *If \mathcal{E} is a stable bundle on Q_3 with $c_1 = 0$, $c_2 = 2$ such that $\mathcal{E}(1)$ is spanned by global sections then \mathcal{E} is a pull-back $\pi^*(\mathcal{N})$ of a null-correlation bundle \mathcal{N} , under a double covering $\pi: Q_3 \rightarrow P^3$.*

Proof. The argument is based on the following fact: for any two disjoint lines on P^3 there exists a section of a twisted null-correlation bundle $\mathcal{N}(1)$ vanishing exactly on these lines. Therefore, if we prove that a section of $\mathcal{E}(1)$ vanishes on a set being a pullback, via a double covering $\pi: Q_3 \rightarrow P^3$, of two disjoint lines on P^3 , then in view of Theorem 1.1 and Remark 1.1.1 from [8], $\mathcal{E}(1)$ is a pullback of $\mathcal{N}(1)$; if Z is the union of two disjoint lines and Y its pullback then it is easy to check that every isomorphism between $\omega_Q(-2)|_Y$ and ω_Y comes from $\omega_P(-2)|_Z \simeq \omega_Z$.

Assume \mathcal{E} is stable with $c_1(\mathcal{E}) = 0$, $c_2(\mathcal{E}) = 2$ on Q_3 . We easily compute the following cohomology table of $h^i(\mathcal{E}(-m))$

0	0	0	0	$\uparrow h^3$
1	1	0	0	h^2
0	0	1	1	h^1
0	0	0	0	h^0
$m = 3$	$m = 2$	$m = 1$	$m = 0$	\rightarrow

Indeed, vanishing of the lower and upper row is a consequence of the stability (plus Serre's duality) and the "spectrum" technique, namely Corollary 2.4 in [5], gives

$$h^1(\mathcal{E}(-2)) = h^1(\mathcal{E}(-3)) = h^2(\mathcal{E}) = h^2(\mathcal{E}(-1)) = 0$$

and the remaining part of the table follows from computing the Euler-Poincaré characteristic.

Since $\chi(\mathcal{E}(1)) = 5$ and $h^2(\mathcal{E}(1)) = h^1(\mathcal{E}(-4)) = 0$ by Corollary 2.4 in [5], we see $h^0(\mathcal{E}(1)) \geq 5$. Let Y be the zero of a generic section.

Since $H^0(\mathcal{E}) = 0$ and $\mathcal{E}(1)$ is assumed to be globally generated, Y is a smooth (not necessarily connected) curve. From the diagram

$$\begin{array}{ccccccc}
 & & & & 0 & & \\
 & & & & \downarrow & & \\
 0 & \rightarrow & \mathcal{O}(-2) & \rightarrow & \mathcal{E}(-1) & \rightarrow & J_Y \rightarrow 0 \\
 & & & & \downarrow & & \\
 & & & & \mathcal{O}_{Q_3} & & \\
 & & & & \downarrow & & \\
 & & & & \mathcal{O}_Y & & \\
 & & & & \downarrow & & \\
 & & & & 0 & &
 \end{array}$$

we calculate, with the aid of the cohomology table above, that $h^0(\mathcal{O}_Y) = 2$, i.e., Y consists of two connected components, say Y_1 and Y_2 .

Claim. Y_1 and Y_2 are conics.

Proof of claim. Since $c_2\mathcal{E}(1) = 4$ and both Y_i are smooth (therefore reduced) we have only to exclude the possibility that one of them is a line L . But then by the adjunction formula we would obtain

$$\deg(K_L) = (K_{Q_3} + c_1\mathcal{E}(1)) \cdot L = -1$$

which is impossible.

Let now H_i be the plane containing Y_i , $i = 1, 2$; clearly $Q_3 \cap H_i = Y_i$ and H_1, H_2 meet at one point in P^4 off Q_3 . Projecting $Q_3 \subset P^4$ from this point onto a hyperplane H in P^4 is a double covering of H and the images of Y_1 and Y_2 are two skew lines, say L_1 and L_2 . It then follows that $\mathcal{E}(1)$ is the pull-back of the null-correlation bundle $\mathcal{N}(1)$ corresponding to L_1 and L_2 .

REMARK. It is not entirely clear whether or not any stable bundle on Q_3 with $c_1 = 0$ and $c_2 = 2$ enjoys the property stated in (3.2).

Case of stable bundles with $c_1 = -1$, $c_2 = 1$. Here a more detailed description of Fano bundles can be given. Let \mathcal{E} be a stable bundle on Q_3 with $c_1 = -1$, $c_2 = 1$.

(3.3) The cohomology of such a bundle are the following:

- (1) $h^0(\mathcal{E}(m)) = 0$ for $m \leq 0$,
- (2) $h^0(\mathcal{E}(1)) = 4$,
- (3) $h^1(\mathcal{E}(m)) = h^2(\mathcal{E}(m)) = 0$ for all m ,
- (4) $h^3(\mathcal{E}(m)) = 0$ for $m \geq -2$.

Proof. (1) is a criterion of stability, (4) is dual to (1), (2) will follow from (3), (4) and the Riemann-Roch formula. Corollary 2.4 in [5] gives $h^1(\mathcal{E}(m)) = 0$ for $m \leq -1$. By duality, $h^2(\mathcal{E}(m)) = 0$ for $m \geq -1$ so that $h^1(\mathcal{E}) = \chi(\mathcal{E}) = 0$. The Castelnuovo criterion (see e.g. Lecture 14 in [14]) now yields that $\mathcal{E}(m)$ are generated by global sections if $m \geq 1$ and that all cohomology $H^i(\mathcal{E}(m))$ vanish for $i \geq 1$, $i + m \geq 1$. Now by duality (3) follows for any integer m .

Note that from the Castelnuovo criterion it follows that $\mathcal{E}(1)$ is spanned; therefore $\mathcal{E}(2)$ is ample and \mathcal{E} is Fano.

Now we prove that such \mathcal{E} is the one from (2.6). Since the bundle $\mathcal{E}(1)$ is spanned and $h^0(\mathcal{E}(1)) = 4$ it follows that the linear system $|H + \xi|$ is base point free and of dimension 3. Let $\varphi: P(\mathcal{E}) \rightarrow P^3$ be the map associated with this system.

(3.4). PROPOSITION. $\varphi: P(\mathcal{E}) \rightarrow P^3$ is a P^1 -bundle which is the projectivization of a null-correlation bundle.

Proof. First note that a general divisor D in the linear system $|2H + \xi|$ is a Fano 3-fold listed as $n^\circ 17$ in Table 2 [13]. The map $\varphi|_D$ is a blow-down morphism from D onto P^3 .

We claim that φ has no fibre of dimension ≥ 2 . Assume that S is such a fibre. Then $f := D \cap S$ is isomorphic to P^1 and $\mathcal{O}_f(H) \cong \mathcal{O}_{P^1}(1)$. In view of Theorem 2.1b', [6] we see that $S \cong P^2$ and $\mathcal{O}_S(H) \cong \mathcal{O}_{P^2}(1)$. But in this case $p: S \rightarrow Q_3$ is a plane embedding of P^2 in Q_3 , which is impossible.

Now any fibre of φ is numerically equivalent to $(H + \xi)^3$ and, since $H \cdot (H + \xi)^3 = 1$, it follows that it must be isomorphic to P^1 . The push-forward $\varphi_*(\mathcal{O}(H))$ is a rank-2 Fano bundle on P^3 . From the results of §2 we see that it is a null-correlation bundle.

COROLLARY. Any stable rank-2 bundle on Q_3 with Chern classes $c_1 = -1$, $c_2 = 1$ is a pull-back of the universal quotient bundle on

Grass(1, 3) via some hyperplane embedding

$$Q_3 \rightarrow \text{Grass}(1, 3) = Q_4 \subset P^5.$$

REMARK 1. The above example shows that the Horrocks splitting principle, as it stands on P^n (see e.g. [15]), cannot be applied literally to bundles on Q_3 (see [16] for an analogue of the Horrocks splitting principle on Q_n). Let us also notice that the bundle discussed above is uniform: its decomposition type is the same on all lines and smooth conics in Q_3 .

REMARK 2. It is proved in [19] that $V = P(\mathcal{N}) = P(\mathcal{E})$ (where \mathcal{E} is the bundle discussed above and \mathcal{N} is the null-correlation bundle on P^3) is the only ruled Fano 4-fold of index 2 obtained from a non-decomposable bundle.

Added in the proof. Together with Ignacio Sols we have concluded the case of rank-2 Fano bundles on Q_3 . Firstly, we have proved that the first twist of a stable bundle with $c_1 = 0$, $c_2 = 2$ is spanned by global sections (see Proposition (3.2) and the subsequent remark). Secondly, we have decided that bundles with $c_1 = 0$, $c_2 = 4$ are not Fano (see the discussion preceding (3.2)).

REFERENCES

- [1] I. Yu. Artiushkin, *Four-dimensional Fano manifolds representable as P^1 -bundles*, Vest. Mosk. Un-ta, ser 1, No. 1, (1987), 12–16, in Russian.
- [2] C. Banica and N. Manolache, *Rank 2 stable vector bundles on $P^3(C)$ with Chern classes $c_1 = -1$, $c_2 = 4$* . Math. Z., **190** (1985), 315–339.
- [3] W. Barth, *Some properties of stable rank-2 vector bundles on P_n* , Math. Ann., **226** (1977), 125–159.
- [4] I. V. Demin, *Three-dimensional Fano manifolds representable as line fiberings*, Izv. Akad. Nauk SSSR, **44**, n°4, (1980), 464–921, English translation in Math USSR Izv. 17 Addendum to this paper in Izv. Akad. Nauk SSSR 46 n°3, English translation in Math. USSR Izv. 20.
- [5] L. Ein and I. Sols, *Stable vector bundles on quadrics*, Nagoya Math. J., **96** (1984), 11–22.
- [6] T. Fujita, *On the structure of polarized varieties with Δ -genera zero*, J. Fac. Sci. Univ. Tokyo, **22** (1975), 103–115.
- [7] W. Fulton, *Ample vector bundles, Chern classes, numerical criteria*. Inv. Math., **32** (1978), 171–178.
- [8] R. Hartshorne, *Stable Vector Bundles of Rank 2 on P^3* , Math. Ann., **238** (1978), 229–280.
- [9] R. Hartshorne and I. Sols, *Stable rank 2 vector bundles on P^3 with $c_1 = -1$, $c_2 = 2$* . J. Reine und Angew. Math., **325** (1981), 145–152.
- [10] T. Hosoh, *Ample vector bundles on a rational surface*, Nagoya Math. J., **59** (1975), 135–148.

- [11] Y. Kawamata, *The cone of curves of algebraic varieties*, Ann. Math., **119** (1984), 603f–633.
- [12] Sh. Mori, *Threefolds whose canonical bundles are not numerically effective*, Ann. Math., **116** (1982), 133–176.
- [13] Sh. Mori and Sh. Mukai, *Classification of Fano 3-folds with $B_2 \geq 2$* , Manuscripta Math., **36** (1981), 147–162.
- [14] D. Mumford, *Lectures on curves on an algebraic surface*, Ann. of Math. Studies 59, Princeton Univ. Press 1966.
- [15] Ch. Okonek, M. Schneider and H. Spindler, *Vector Bundles on Complex Projective Spaces*, Birkhäuser 1981.
- [16] G. Ottaviani, *Critères de scindage pour les fibrès vectoriels sur les grassmanniennes et les quadriques*, Comptes Rendus Acad. Sci. Paris, **305** (1987), 257–260.
- [17] M. Schneider, *Stabile Vektorraumbündel vom rang 2 auf der projektiven Ebene*, Nachr. Akad. Wiss. Gottingen 1976, 83–86.
- [18] S. S. Tikhomirov, *On stable two-dimensional vector bundles on P^3 with Chern classes $c_1 = 0$, $c_2 = 3$ and 4*, Sbornik Nauch. Tr. Yarosl. Pedag. Inst. im. K. D. Ushinskogo, **180** (1979), 106–114, in Russian.
- [19] J. A. Wisniewski, *Ruled Fano 4-folds of index 2*, Proc. Amer. Math. Soc., **105** (1983), 55–61

Received January 26, 1988.

WARSAW UNIVERSITY

PKIN 9P. 00-901 WARSZAWA, POLAND

Jarosław A. Wiśniewski visiting at: The Johns Hopkins University
Baltimore, MD 21218, USA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Yusuf Abu-Muhanna and Abdallah Khalil Lyzzaik, The boundary behaviour of harmonic univalent maps	1
Lawrence Jay Corwin, Allen Moy and Paul J. Sally, Jr., Degrees and formal degrees for division algebras and GL_n over a p-adic field	21
Ulrich Dierkes and Gerhard Huisken, The n-dimensional analogue of the catenary: existence and nonexistence	47
Peter Gerard Dodds, C. B. Huijsmans and Bernardus de Pagter, Characterizations of conditional expectation-type operators	55
Douglas Dokken and Robert Ellis, The Poisson flow associated with a measure	79
Larry J. Santoni, Horrocks' question for monomially graded modules	105
Zbigniew Slodkowski, Pseudoconvex classes of functions. II. Affine pseudoconvex classes on R^N	125
Dean Ellis Smith, On the Cohen-Macaulay property in commutative algebra and simplicial topology	165
Michał Szurek and Jarosław Wisniewski, Fano bundles over P^3 and Q_3 ...	197