

Pacific Journal of Mathematics

UNITARY BORDISM OF CLASSIFYING SPACES OF QUATERNION GROUPS

ABDESLAM MESNAOUI

UNITARY BORDISM OF CLASSIFYING SPACES OF QUATERNION GROUPS

ABDESLAM MESNAOUI

Let Γ_k be the generalized quaternion group of order 2^k . In this article we determine a set of generators for the $U_*(pt)$ -module $\tilde{U}_*(B\Gamma_k)$ and give all linear relations between them. Moreover their orders are calculated.

0. Introduction. In this article we first study the case $\Gamma_k = \Gamma$ the quaternion group of order 8. We recall that

$$\Gamma = \{\pm 1, \pm i, \pm j, \pm k\}, \quad i^2 = j^2 = k^2 = -1, \quad ij = k, \quad jk = i, \quad ki = ij.$$

Γ acts on S^{4n-3} by using $(n+1)\eta$ where η denotes the following unitary irreducible representation of Γ : $i \rightarrow \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$, $j \rightarrow \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and we get the element $w_{4n+3} = [S^{4n+3}/\Gamma, q] \in \tilde{U}_{4n+3}(B\Gamma)$, q being the natural embedding: $S^{4n+3}/\Gamma \subset B\Gamma$. In [6] we have defined three elements of $\tilde{U}^2(B\Gamma)$ denoted by A, B, C as Euler classes for MU of irreducible representations of Γ of dimension 1 over \mathbb{C} . Let $u_{4n+1} \in \tilde{U}_{4n+1}(B\Gamma)$, $v_{4n+1} \in \tilde{U}_{4n+1}(B\Gamma)$ be respectively $A \cap w_{4n+3}$ and $B \cap w_{4n+3}$. Our first result is:

THEOREM 2.2. *The set $\{u_{4n+1}, v_{4n+1}, w_{4n+3}\}_{n \geq 0}$ is a system of generators for the $U_*(pt)$ -module $\tilde{U}_*(B\Gamma)$.*

Their orders are given by:

THEOREM 2.6. *We have: $\text{ord } w_{4n+3} = 2^{2n+3}$.*

THEOREM 2.8. *We have: $\text{ord } u_{4n+1} = \text{ord } v_{4n+1} = 2^{n+1}$.*

Now let Ω_* be $U^*(pt)[[Z]]$ graded by taking $\dim Z = 4$. If $P(Z) = \sum_{i \geq r} \alpha_i Z^i \in \Omega_n$ and $\alpha_r \neq 0$ then we denote $\nu(P) = 4r$. Let W, V_1, V_2 be the submodules of $\tilde{U}_*(B\Gamma)$ generated respectively by $\{w_{4n+3}\}_{n \geq 0}$, $\{u_{4n+1}\}_{n \geq 0}$, $\{v_{4n+1}\}_{n \geq 0}$. The following result gives the $U_*(pt)$ -module structure of $\tilde{U}_*(B\Gamma)$ and uses the elements $T(Z) \in \Omega_4$, $J(Z) \in \Omega_0$ as defined in [6], Section II.

THEOREM 2.4. (a) $\tilde{U}_*(B\Gamma) = W \oplus V_1 \oplus V_2$.

(b) In $\tilde{U}_{2p+1}(B\Gamma)$ we have $0 = a_0w_3 + a_1w_7 + \cdots + a_nw_{4n+3} = b_0u_1 + \cdots + b_mu_{4m+1}$ iff there are homogeneous polynomials $M(Z), M_2(Z)$ and homogeneous formal power series $N(Z), N_1(Z)$ of Ω_* satisfying: $b_mZ + b_{m-1}Z^2 + \cdots + b_0Z^{m+1} = M(Z)(2 + J(Z)) + N(Z)$, $a_nZ + a_{n-1}Z^2 + \cdots + a_0Z^{n+1} = M_1(Z)T(Z) + N_1(Z)$, $\nu(N) > 4(n+1)$, $\nu(N_1) > 4(n+1)$. Moreover $b_0u_1 + \cdots + b_mu_{4m+1} = 0$ iff $b_0v_1 + \cdots + b_mv_{4m+1} = 0$.

In Section III we consider $\tilde{U}_*(B\Gamma_k)$, $k \geq 4$. The generalized quaternion group Γ_k is generated by u, v with $u^t = v^2$, $t = 2^{k-2}$, $uvu = v$. Γ_k acts on S^{4n+3} by means of the irreducible unitary representation η_1 of Γ_k :

$$u \rightarrow \begin{pmatrix} \omega & 0 \\ 0 & \omega^{-1} \end{pmatrix}, \quad v \rightarrow \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

ω being a primitive 2^{k-1} th root of unity. We get:

$$w'_{4n+3} = [S^{4n+3}/\Gamma_k, q'] \in \tilde{U}_{4n+3}(B\Gamma_k), \quad q': S^{4n+3}/\Gamma_k \subset B\Gamma_k.$$

Now we use the elements $B'_k = B_k + G_k(D_k) \in \tilde{U}^2(B\Gamma_k)$, $C'_k = C_k + G_k(D_k) \in \tilde{U}^2(B\Gamma)$ (see [6], Theorem 3.14) to define $u'_{4n+1} = B'_k \cap w'_{4n+3} \in \tilde{U}_{4n+1}(B\Gamma_k)$, $v'_{4n+1} = C'_k \cap w'_{4n+3} \in \tilde{U}_{4n+1}(B\Gamma_k)$. Then we have Theorems 3.1, 3.2 identical respectively to the above Theorems 2.2, 2.4 where w_{4n+3} , u_{4n+1} , v_{4n+1} are replaced by w'_{4n+3} , u'_{4n+1} , v'_{4n+1} . However:

THEOREM 3.4. We have: $\text{ord } w'_{4n+3} = 2^{2n+k}$, $n \geq 0$.

THEOREM 3.5. We have: $\text{ord } u'_{4n+1} = \text{ord } v'_{4n+1} = 2^{n+1}$, $n \geq 0$, which are therefore independent of k .

The layout is as follows:

- I Preliminaries and notations.
- II Calculations in $\tilde{U}_*(B\Gamma)$: generators, orders and relations.
- III $\tilde{U}_*(B\Gamma_k)$, $k \geq 4$: generators, orders and relations.

We assume that the reader is acquainted with the notations and results of [6].

I. Preliminaries and notations. The notation U_* -AHSS will be used for the Atiyah-Hirzebruch spectral sequence corresponding to the homology theory determined by MU ; μ and μ' denote the edge homomorphisms $U^*(X) \rightarrow H^*(X)$ and $U_*(X) \rightarrow H_*(X)$ obtained from the

U_* -AHSS for a CW complex X . We have the following well-known result:

THEOREM 1.1. *Suppose X a CW-complex such that:*

- (a) *The U_* -AHSS for X collapses.*
- (b) *For each $n \geq 0$ there is a system (a_{in}) generating the group $H_n(X)$.*

Then for each $n \geq 0$ there is a system (A_{in}) such that:

- (a) *$A_{in} \in U_n(X)$, $\mu'(A_{in}) = a_{in}$ for every (i, n) .*
- (b) *The system (A_{in}) generates $U_*(X)$ as a $U_*(pt)$ -module.*

Moreover, (b) is valid for every system (A_{in}) such that $\mu'(A_{in}) = a_{in}$. \square

Consider the map of ring spectra $f: MU \rightarrow H$ (see [1]); by naturality of spectral sequences it follows that if X is a CW-complex then $f^*(X) = \mu$ and $f_*(X) = \mu'$ where $f^*(X): U^*(X) \rightarrow H^*(X)$, $f_*(X): U_*(X) \rightarrow H_*(X)$ denote the maps induced by f .

PROPOSITION 1.2. *If X is a CW-complex then the following diagram commutes:*

$$\begin{array}{ccc} U^m(X) \otimes U_n(X) & \xrightarrow{\cap} & U_{n-m}(X) \\ \mu \otimes \mu' \downarrow & & \downarrow \mu' \\ H^m(X) \otimes H_n(X) & \xrightarrow{\cap} & H_{n-m}(X) \end{array} \text{ commutes.}$$

Proof. Take $E = MU$. The cap product is the composite:

$$\tilde{E}_m(X^+) \otimes \tilde{E}_n(X^+) \xrightarrow{1 \otimes \Delta_*} \tilde{E}^m(X^+) \otimes \tilde{E}_n(X^+ \wedge X^+) \xrightarrow{\searrow} \tilde{E}_{n-m}(X^+),$$

\searrow being the slant product and $\Delta(x) = [x, x]$. Since Δ_* commutes with $f_*(-)$ we have to prove that the diagram:

$$\begin{array}{ccc} \tilde{E}^m(X^+) \otimes \tilde{E}_n(X^+ \wedge X^+) & \xrightarrow{\searrow} & \tilde{E}_{n-m}(X^+) \\ \downarrow f^*(-) \otimes f_*(-) & & \downarrow f_*(-) \\ \tilde{H}^m(X^+) \otimes \tilde{H}_n(X^+ \wedge X^+) & \xrightarrow{\searrow} & \tilde{H}_{n-m}(X^+) \end{array} \text{ commutes.}$$

More generally the diagram

$$\begin{array}{ccc} \tilde{E}^m(Y) \otimes \tilde{E}_n(Y \wedge Z) & \xrightarrow{\searrow} & \tilde{E}_{n-m}(Z) \\ \downarrow f^*(-) \otimes f_*(-) & & \downarrow f_*(-) \\ \tilde{H}^m(Y) \otimes \tilde{H}_n(Y \wedge Z) & \xrightarrow{\searrow} & \tilde{H}_{n-m}(Z) \end{array} \text{ commutes if } Y, Z$$

are pointed CW-complexes: indeed let x and y be any elements of $\tilde{E}^m(Y)$ and $\tilde{E}_n(Y \wedge Z)$ respectively represented by $g: Y \rightarrow \sum^m E$, $h: S^n \rightarrow E \wedge Y \wedge Z$. Then $f^\#(-)(x)$ is represented by the composite

$$g_1: Y \xrightarrow{g} \sum^m E \xrightarrow{\sum^m f} \sum^m H \quad \text{and} \quad f_\#(-)(y)$$

by the composite:

$$h_1: S^n \xrightarrow{h} E \wedge Y \wedge Z \xrightarrow{f \wedge 1 \wedge 1} H \wedge Y \wedge Z.$$

If we denote by T the transposition and k, k' the ring-spectra products then the diagram pictured on the next page commutes. Since the top line represents $x \setminus y$ and the bottom line

$$f^\#(-)(x) \setminus f_\#(-)(y)$$

we have $f_\#(-)(x \setminus y) = f^\#(-)(x) \setminus f_\#(-)(y)$. □

Let X be any CW-complex and ξ a complex vector bundle of \mathbb{C} -dimension n over X . If h denotes a map: $X \rightarrow BU(n)$ classifying ξ and $M(\xi)$ the Thom space of ξ , then $M(h): M(\xi) \rightarrow MU(n)$ determines an element $t_0(\xi) \in U^{2n}(M(\xi))$ which is a particular Thom class for ξ called the canonical Thom class for ξ . Moreover if $j: X \rightarrow M(\xi)$ is the zero section we have $j^*(t_0(\xi)) = cf_n(\xi)$, the highest Conner-Floyd characteristic class of ξ ; $j^*(t_0(\xi))$ is also called the Euler class $e(\xi)$ of ξ .

Fundamental classes for a U -manifold M^n for $E = MU$ or H may be obtained in the following manner: M^n can be embedded in S^{n+2k} for some large k and the normal bundle τ can be given a $U(k)$ -structure; let N be a tubular neighbourhood of M^n , which we identify with the total space of the normal disk bundle $D(\tau)$; we have the map $\pi: S^{n+2k} \rightarrow M(\tau)$ defined as follows: if $x \in N$ then $\pi(x)$ is the image of x by the projection $D(\tau) \rightarrow M(\tau)$ and if $x \in S^{n+2k} - \mathring{N}$, then $\pi(x) = *$ the base point of $M(\tau)$; let t be a Thom class of ξ for E ; we have the Thom-isomorphism $\phi_t: E_{2k+r}(M(\tau)) \rightarrow E_r(M^n)$ such that $\phi_t(x) = p_*(t \cap x)$, p being the projection $D(\tau) \rightarrow M^n$; let $u: S^0 \rightarrow E$ be the unit of E ; the map u is a map of spectra and is therefore a collection of maps $u_m: S^m \rightarrow E_m$ satisfying well-known axioms; then by [8], page 333, if $[u_{n+2k}]$ is the element of $\tilde{E}_{n+2k}(S^{n+2k})$ corresponding to u_{n+2k} , then the element $c(M) = \phi_t(\pi_*([u_{n+2k}])) \in E_n(M^n)$ is a fundamental class for M^n . Evidently the same method produces fundamental classes for the homology theory defined by the spectrum H .

$$\begin{array}{ccccccc}
S^{n-m} & \xrightarrow{\sum^{-m} h} & (\sum^{-m} E) \wedge Y \wedge Z & \xrightarrow{T \wedge 1} & Y \wedge \sum^{-m} E \wedge Z & \xrightarrow{g \wedge 1 \wedge 1} & \sum^m E \wedge \sum^{-m} E \wedge Z \xrightarrow{k \wedge 1} E \wedge Z \\
\parallel & & \sum^{-m} f \wedge 1 \wedge 1 \downarrow & & \downarrow 1 \wedge \sum^{-m} f \wedge 1 & & \sum^m f \wedge \sum^{-m} f \wedge 1 \downarrow f \wedge f \wedge 1 \downarrow \\
S^{n-m} & \xrightarrow{\sum^{-m} h_1} & (\sum^{-m} H) \wedge Y \wedge Z & \xrightarrow{T \wedge 1} & Y \wedge \sum^{-m} H \wedge Z & \xrightarrow{g_1 \wedge 1 \wedge 1} & \sum^m H \wedge \sum^{-m} H \wedge Z \xrightarrow{k' \wedge 1} H \wedge Z
\end{array}$$

From [8], page 335, §14-45, we have:

PROPOSITION 1.3. *If M^n is a closed U -manifold then $[M^n, 1] \in U_n(M^n) = E_n(M^n)$ is a fundamental class for M^n deduced from the canonical Thom class $t_0(\tau)$, τ being the normal bundle of an embedding $M^n \subset S^{n+2k}$, k large. \square*

PROPOSITION 1.4. *Let M^n be a closed U -manifold; then*

$$f_{\#}(-)([M^n, 1]) \in H_n(M^n)$$

is a fundamental class for M^n .

Proof. From 1.3 we have

$$[M^n, 1] = \phi_{t_0}(\pi_*[u_{n+2k}]) = c(M);$$

then

$$\begin{aligned} f_{\#}(-)(c(M)) &= f_{\#}(-)[\phi_{t_0}(\pi_*([u_{n+2k}]))] = f_{\#}(-)[p_*(t_0 \cap \pi_*([u_{n+2k}]))] \\ &= p_*[f_{\#}(-)(t_0 \cap \pi_*([u_{n+2k}]))] \\ &= p_*[f^{\#}(-)(t_0) \cap f_{\#}(-)(\pi_*([u_{n+2k}]))] \\ &= p_*[f^{\#}(-)(t_0) \cap \pi_*(f(-)([u_{n+2k}]))]. \end{aligned}$$

Since f is a map of spectra the unit of H is the composite $v: S^0 \xrightarrow{u} MU \xrightarrow{f} H$ and hence $f_{\#}(-)([u_{n+2k}]) = [v_{n+2k}]$. Now $f^{\#}(-)(t_0)$ is a Thom class t_1 for H and therefore

$$\begin{aligned} f_{\#}(-)(c(M)) &= p_*[t_1 \cap \pi_*([v_{n+2k}])] \\ &= \phi_{t_1}(\pi_*([v_{n+2k}])) = c_1(M^n) \in H_n(M^n) \end{aligned}$$

is a fundamental class for M^n . \square

The notation $c(M^n)$ will be for the fundamental class $[M^n, 1] \in U_n(M^n)$ and $c_1(M^n) \in H_n(M^n)$ will be the fundamental class $\mu'(c(M^n))$.

If PD or PD₁ denotes the Poincaré duality then we have:

PROPOSITION 1.5. *The following diagram commutes*

$$\begin{array}{ccc} U^m(M^n) & \xrightarrow{\text{PD}} & U_{n-m}(M^n) \\ \downarrow \mu & & \downarrow \mu' \\ H^m(M^n) & \xrightarrow{\text{PD}_1} & H_{n-m}(M^n) \end{array}$$

Proof. We have

$$\begin{aligned}\mu'(\text{PD}(x)) &= \mu'(x \cap c(M^n)) = \mu(x) \cap \mu'(c(M^n)) \\ &= \mu(x) \cap c_1(M^n) = (\text{PD})_1(\mu(x))\end{aligned}$$

by 1.2. □

Let N^m be a closed U -submanifold of a closed U -manifold M^n , and i the inclusion $N^m \subset M^n$; then the normal bundle τ of N^m in M^n is a complex-vector-bundle if $(n - m)$ is even and we have:

PROPOSITION 1.6. *If $(n - m)$ is even then $(\text{PD})^{-1}([N^m, i])$ is represented by:*

$$M^n \rightarrow M^n / (M^n - \overset{\circ}{N}) \simeq D(\tau) / S(\tau) = M(\tau) \xrightarrow{M(h)} MU(\tfrac{1}{2}(n - m)),$$

where h is a map classifying τ and N a tubular neighborhood of N^m homeomorphic to $D(\tau)$ (see [3], [7]). □

The generalized quaternion group Γ_k , $k \geq 4$, is generated by u, v subject to the relations $u^t = v^2$, $t = 2^{k-2}$, $uvu = v$. Consider the irreducible unitary representation η_1 of Γ_k : $u \rightarrow \begin{pmatrix} \omega & 0 \\ 0 & \omega^{-1} \end{pmatrix}$, $v \rightarrow \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$, ω being a primitive 2^{k-1} -th-root of unity. The group Γ_k acts on S^{4n+3} by means of $(n+1)\eta_1$ as a group of U -diffeomorphisms and we get a canonical U -structure on S^{4n+3}/Γ_k and a natural injection $S^{4n+3}/\Gamma_k \subset B\Gamma_k = \bigcup_{n \geq 0} S^{4n+3}/\Gamma_k$ (see [3], [10], page 508).

Let α be the complex vector bundle: $S^{4n+3} \times_{\Gamma_k} \mathbb{C}^2 \rightarrow S^{4n+3}/\Gamma_k$ where Γ_k acts on S^{4n+3} and \mathbb{C}^2 respectively by means of $(n+1)\eta_1$ and η_1 : if $a \in \Gamma_k$ and $(x, v) \in S^{4n+3} \times \mathbb{C}^2$ we have $a(s, w) = (as, av) = (sa^{-1}, av)$ and $S^{4n+3} \times_{\Gamma_k} \mathbb{C}^2 = (S^{4n+3} \times \mathbb{C}^2) / \Gamma_k$. Then by a result of R. H. Szczarba ([9]) we have $T(S^{4n+3}/\Gamma_k) + 1 = (n+1)\alpha$ where $T(S^{4n+3}/\Gamma_k)$ denotes the tangent bundle of S^{4n+3}/Γ_k . As an easy consequence we have:

PROPOSITION 1.7. *If i denotes the embedding $S^{4n+3}/\Gamma_k \subset S^{4n+7}/\Gamma_k$ such that*

$$i([z_1, z_2, \dots, z_{2n+2}]) = [z_1, z_2, \dots, z_{2n+2}, 0, 0],$$

then the normal bundle of S^{4n+3}/Γ_k in S^{4n+7}/Γ_k is isomorphic to the complex vector bundle α . □

We shall give a proof of the next result which can be found in [7]:

PROPOSITION 1.8. *If i denotes the embedding $S^{4n+3}/\Gamma_k \subset S^{4n+7}/\Gamma_k$ then $i^* \circ (\text{PD})^{-1}([S^{4n+3}/\Gamma_k, i]) = e(\alpha)$.*

Proof. Denote by τ the normal bundle of S^{4n+3}/Γ_k in S^{4n+7}/Γ_k and by h a classifying map: $S^{4n+3}/\Gamma_k \rightarrow BU(2)$ for τ . Then by 1.6, $(\text{PD})^{-1}([S^{4n+3}/\Gamma_k, i])$ is represented by the composite:

$$\begin{aligned} S^{4n+7}/\Gamma_k &\rightarrow (S^{4n+7}/\Gamma_k) / (S^{4n+7}/\Gamma_k - \overset{\circ}{N}) \\ &\simeq \frac{D(\tau)}{S(\tau)} = M(\tau) \xrightarrow{M(h)} MU(2), \end{aligned}$$

N being a tubular neighbourhood of S^{4n+3}/Γ_k homeomorphic to $D(\tau)$. Since the composite:

$$\begin{aligned} S^{4n+3}/\Gamma_k &\xrightarrow{i} S^{4n+7}/\Gamma_k \rightarrow S^{4n+7}/\Gamma_k / (S^{4n+7}/\Gamma_k - \overset{\circ}{N}) \\ &\simeq \frac{D(\tau)}{S(\tau)} = M(\tau) \end{aligned}$$

is the zero section: $S^{4n+3}/\Gamma_k \rightarrow M(\tau)$, it follows that

$$i^* \circ (P(D)^{-1})([S^{4n+3}/\Gamma_k, i]) = e(\tau).$$

Since τ and α are isomorphic as complex vector bundles by 1.7 the proposition is proved. \square

In Section III we shall use the following Euler classes for MU (see [6]):

$$\begin{aligned} A_k &= e(\xi_1) \in \tilde{U}^2(B\Gamma_k), & B_k &= e(\xi_2) \in \tilde{U}^2(B\Gamma_k), \\ C_k &= e(\xi_3) \in \tilde{U}^2(B\Gamma_k), & D_k &= e(\eta_1) \in \tilde{U}^4(B\Gamma_k) \end{aligned}$$

where $\xi_1, \xi_2, \xi_3, \eta_1$ are the complex vector bundles corresponding to the irreducible unitary representations $\xi_1: u \rightarrow 1, v \rightarrow -1$, $\xi_2: u \rightarrow -1, v \rightarrow 1$, $\xi_3: k \rightarrow -1, v \rightarrow -1$ and η_1 as defined above.

In order to calculate $U_*(B\Gamma_k)$ we first consider the case $k = 3$: $\Gamma_3 = \Gamma$, the quaternion group of order 8. We recall that $\Gamma = \{\pm 1, \pm i, \pm j, \pm k\}$ subject to the relations $i^2 = j^2 = k^2 = -1$, $ij = k$, $jk = i$, $ki = j$. The irreducible unitary representations of Γ are $1: i \rightarrow 1, j \rightarrow 1$, $\xi_i: i \rightarrow 1, j \rightarrow -1$, $\xi_j: i \rightarrow -1, j \rightarrow 1$, $\xi_k: i \rightarrow -1, j \rightarrow -1$ and $\eta: i \rightarrow \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, j \rightarrow \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. The character table of Γ is drawn on the next page.

The group Γ acts on S^{4n+3} by means of $(n+1)\eta$ as a group of U -diffeomorphisms; as with Γ_k we get a U -manifold $S^{4n+3}/\Gamma \subset B\Gamma = \bigcup_{n \geq 0} S^{4n+3}/\Gamma$. There will be no ambiguity if we use the same notation

conjugacy classes

	1	-1	$\pm i$	$\pm j$	$\pm k$	
1	1	1	1	1	1	
ξ_i	1	1	1	1	-1	
ξ_j	1	1	-1	1	-1	
ξ_k	1	1	-1	-1	-1	
η	2	2	0	0	0	

α as for Γ_k for the complex vector bundle $S^{4n+3} \times_{\Gamma} \mathbb{C}^2 \rightarrow S^{4n+3}/\Gamma$. Evidently the Propositions 1.6 and 1.7 are valid if Γ_k is replaced by Γ .

In Section II the following Euler class for MU will be of fundamental importance (see [6]):

$$\begin{aligned} A &= e(\xi_i) \in \tilde{U}^2(B\Gamma), & B &= e(\xi_j) \in \tilde{U}^2(B\Gamma), \\ C &= e(\xi_k) \in \tilde{U}^2(B\Gamma) \quad \text{and} \quad D &= e(\eta) \in \tilde{U}^4(B\Gamma). \end{aligned}$$

II. Calculation of $\tilde{U}_*(B\Gamma)$: generators, orders and relations. The reduced homology groups $\tilde{H}_*(B\Gamma)$ are such that:

$$\tilde{H}_{2n}(B\Gamma) = 0, \quad \tilde{H}_{4n+1}(B\Gamma) = \mathbb{Z}_2 \oplus \mathbb{Z}_2, \quad \tilde{H}_{4n+3}(B\Gamma) = \mathbb{Z}_8, \quad n \geq 0.$$

The \tilde{U}_* -AHSS of $B\Gamma$ collapses and we have a filtration of $\tilde{U}_n(B\Gamma)$:

$$J_{-1,n+1} = 0 \subset J_{0,n} \subset \cdots \subset J_{p,n-p} \subset \cdots \subset J_{n,0} = \tilde{U}_n(B\Gamma)$$

with $J_{p,q} = \text{Im}(\tilde{U}_{p+q}(X^p) \rightarrow \tilde{U}_{p+q}(B\Gamma))$, X^p being the p -skeleton of $B\Gamma$. Moreover $J_{p,q}/J_{p-1,q+1} = \tilde{H}_p(B\Gamma, U_q(pt))$.

PROPOSITION 2.1. (a) $\tilde{U}_{2n}(B\Gamma) = 0$, $\tilde{U}_{2n+1}(B\Gamma) = U_{2n+1}(B\Gamma)$, $U_{2n}(B\Gamma) = U_{2n}(pt)$.

(b) $\text{Ord}(\tilde{U}_{4n+3}(B\Gamma)) = 2^r$,

$$\begin{aligned} r &= 3 \left(\sum_{i=0}^n \text{Rank } U_{4i}(pt) \right) \\ &+ 2 \left(\sum_{i=0}^n \text{Rank } U_{4i+2}(pt) \right); \quad \text{Ord}(\tilde{U}_{4n+1}(B\Gamma)) = 2^s, \end{aligned}$$

$$s = 3 \left(\sum_{i=0}^{n-1} \text{Rank } U_{4i+2}(pt) \right) + 2 \left(\sum_{i=0}^n \text{Rank } U_{4i}(pt) \right).$$

Proof. (a) From the filtration $J_{-1,2n+1} = 0 \subset J_{0,2n} \subset \cdots \subset J_{p,2n-p} \subset \cdots \subset J_{2n,0}$, and $J_{p,2n-p}/J_{p-1,2n-p+1} = H_p(B\Gamma, U_{2n-p}(pt)) = 0$ it follows that $\tilde{U}_{2n}(B\Gamma) = 0$. Hence $U_{2n}(B\Gamma) = U_{2n}(pt)$ and $\tilde{U}_{2n+1}(B\Gamma) = U_{2n+1}(B\Gamma)$ because $U_{2n+1}(pt) = 0$.

(b) The orders are easy consequences of:

$$\begin{aligned} J_{4p+3,2q}/J_{4p+2,2q+1} &= H_{4p+3}(B\Gamma, U_{2q}(pt)) \\ &= \mathbb{Z}_8 \otimes U_{2q}(pt) = U_{2q}(pt)/8 \cdot U_{2q}(pt), \\ J_{4p+2,2q+1}/J_{4p+1,2q+2} &= 0, \\ J_{4p+1,2q+2}/J_{4p,2q+3} \\ &= U_{2q+2}(pt)/2U_{2q+2}(pt) \oplus U_{2q+2}(pt)/2U_{2q+2}(pt), \\ J_{4p,2q+3}/J_{4p-1,2q+4} &= 0. \end{aligned} \quad \square$$

Let $w_{4n+3} \in \tilde{U}_{4n+3}(B\Gamma)$ be $[S^{4n+3}/\Gamma, q]$, q being the inclusion $S^{4n+3}/\Gamma \subset B\Gamma$, $u_{4n+1} = A \cap w_{4n+3} \in \tilde{U}_{4n+1}(B\Gamma)$, $v_{4n+1} = B \cap w_{4n+3} \in \tilde{U}_{4n+1}(B\Gamma)$.

THEOREM 2.2. *The set $\{u_{4n+1}, v_{4n+1}, w_{4n+3}\}_{n \geq 0}$ is a system of generators for the $U_*(pt)$ -module $\tilde{U}_*(B\Gamma)$.*

Proof. Since the U_* -AHSS for $B\Gamma$ collapses we can use 1.1. If μ' denotes the edge homomorphism it is enough to prove that $\mu'(w_{4n+3})$, $\{\mu'(u_{4n+1}), \mu'(v_{4n+1})\}$ are systems of generators respectively for $\tilde{H}_{4n+3}(B\Gamma)$ and $\tilde{H}_{4n+1}(B\Gamma)$.

(a) Consider the following commutative diagram:

$$\begin{array}{ccc} \tilde{U}_{4n+3}(S^{4n+3}/\Gamma) & \xrightarrow{q_*} & \tilde{U}_{4n+3}(B\Gamma) \\ \mu' \downarrow & & \downarrow \mu' \\ \tilde{H}_{4n+3}(S^{4n+3}/\Gamma) & \xrightarrow{q_*} & \tilde{H}_{4n+3}(B\Gamma). \end{array}$$

We have $\mu'([S^{4n+3}/\Gamma, 1]) = c_1(S^{4n+3}/\Gamma)$, where $c_1(S^{4n+3}/\Gamma)$ denotes the fundamental class of S^{4n+3}/Γ (for H). Since $c_1(S^{4n+3}/\Gamma)$ is a generator of $\tilde{H}_{4n+3}(B\Gamma)$ it follows that $q_*(c_1(S^{4n+3}/\Gamma))$ is a generator of $\tilde{H}_{4n+3}(B\Gamma)$ because S^{4n+3}/Γ is the $(4n+3)$ -skeleton of $B\Gamma$. Now $q_*([S^{4n+3}/\Gamma, 1]) = [S^{4n+3}/\Gamma, q]$ and then $\mu'([S^{4n+3}/\Gamma, q])$ is a generator of $\tilde{H}_{4n+3}(B\Gamma)$.

(b) By [6], Section II, $\mu(A)$ and $\mu(B)$ generate the group $H^2(B\Gamma)$ and then if $A_1 = q^*(A) \in U^2(S^{4n+3}/\Gamma)$, $B_1 = q^*(B) \in U^2(S^{4n+3}/\Gamma)$, then the elements $\mu(A_1), \mu(B_1)$ generate $H^2(S^{4n+3}/\Gamma)$ because the following diagram commutes:

$$\begin{array}{ccc} U^2(B\Gamma) & \xrightarrow{q^*} & U^2(S^{4n+3}/\Gamma) \\ \mu \downarrow & & \downarrow \mu \\ H^2(B\Gamma) & \xrightarrow{q^*} & H^2(S^{4n+3}/\Gamma) \end{array}$$

and the bottom line is an isomorphism. Consider $t_{4n+3} = [S^{4n+3}/\Gamma, 1] \in U_{4n+3}(S^{4n+3}/\Gamma)$; then $\mu'(t_{4n+3}) = c_1(S^{4n+3}/\Gamma)$. Since the diagram:

$$\begin{array}{ccc} U^2(S^{4n+3}/\Gamma) & \xrightarrow{-\cap t_{4n+3}} & U_{4n+1}(S^{4n+3}/\Gamma) \\ \mu \downarrow & & \downarrow \mu' \\ H^2(S^{4n+3}/\Gamma) & \xrightarrow{-\cap c_1(S^{4n+3}/\Gamma)} & H_{4n+1}(S^{4n+3}/\Gamma) \end{array}$$

commutes by 1.5 and since the bottom line is an isomorphism it follows that $\mu'(A_1 \cap t_{4n+3})$ and $\mu'(B_1 \cap t_{4n+3})$ generate the group $H_{4n+1}(S^{4n+3}/\Gamma)$. Now by using the commutative diagram:

$$\begin{array}{ccc} U_{4n+1}(S^{4n+3}/\Gamma) & \xrightarrow{q_*} & U_{4n+1}(B\Gamma) \\ \downarrow \mu' & & \downarrow \mu' \\ H_{4n+1}(S^{4n+3}/\Gamma) & \xrightarrow{q_*} & H_{4n+1}(B\Gamma) \end{array}$$

we see that $q_*(A_1 \cap t_{4n+3})$ and $q_*(B_1 \cap t_{4n+3})$ generate the group $H_{4n+1}(B\Gamma)$. Since $q_*(A_1 \cap t_{4n+3}) = q_*(q^*(A) \cap t_{4n+3}) = A \cap q_*(t_{4n+3}) = A \cap w_{4n+3}$ and $q_*(B_1 \cap t_{4n+3}) = B \cap w_{4n+3}$ the assertion (b) has been proved. \square

(1) *Relations between the generators.* We first recall the definition of the pull back transfer. Let M^n be a closed U -manifold, N^m a closed U -submanifold of M^n with $(n - m)$ even and i the inclusion $N^m \subset M^n$. If $[V^r, f] \in U_r(M^n)$, then there is a weakly complex representative map $g: V^r \rightarrow M^n$ transversal to N^m . Hence $g^{-1}(N^m)$ is a smooth closed submanifold of V^r and $\dim g^{-1}(N^m) = r + m - n$. Since N^m is a U -submanifold of M^n the normal vector bundle τ of N^m is in fact a complex vector bundle and by transversality we have $T(W^{r+m-n}) + g_1^*(\tau) = j^*(T(V^r))$ (1) where $W^{r+m-n} = g^{-1}(N^m)$, $g_1 = g|g^{-1}(N^m)$, $j: W^{r+m-n} \subset V^r$ and $T(-)$ being the tangent vector

bundle. Since V^r is a U -manifold the stable tangent bundle of V^r has a complex structure and the above relation (1) determines a unique complex structure on the stable tangent bundle of W^{r+m-n} (see [5], page 16). Then we define $i! : U_r(M^n) \rightarrow U_{r+m-n}(N^m)$ by $i!([V^r, f]) = [W^{r+m-n}, g_1]$. Moreover, the following diagram is commutative:

$$\begin{array}{ccc} U^k(M^n) & \xrightarrow{i^*} & U^k(N^m) \\ \text{PD} \downarrow & & \downarrow \text{PD} \\ U_{n-k}(M^n) & \xrightarrow{i!} & U_{m-k}(N^m) \end{array}$$

PD being the Poincaré duality (see [2], [7]).

Now, there is a map $\Delta : \tilde{U}_*(B\Gamma) \rightarrow \tilde{U}_*(B\Gamma)$ defined by $\Delta(x) = D \cap x$, with $D = e(\eta)$, the Euler class of η . The map Δ is a homomorphism of graded $U_*(pt)$ -modules of degree -4 .

PROPOSITION 2.3. *We have*

$$\begin{aligned} \Delta(w_{4n+3}) &= w_{4(n-1)+3}, & \Delta(u_{4n+1}) &= u_{4(n-1)+1}, \\ \Delta(v_{4n+1}) &= v_{4(n-1)+1}, & n &\geq 0. \end{aligned}$$

Proof. Let p, r, s be respectively the inclusions $S^{4(n-1)+3}/\Gamma \subset S^{4n+3}/\Gamma$, $S^{4n+3}/\Gamma \subset S^{4n+7}/\Gamma$, $S^{4n+7}/\Gamma \subset B\Gamma$. Then

$$[S^{4n+3}/\Gamma, r] \in U_{4n+3}(S^{4n+7}/\Gamma).$$

We have the pull back transfer

$$r! : U_{4n+3}(S^{4n+7}/\Gamma) \rightarrow U_{4(n-1)+3}(S^{4n+3}/\Gamma)$$

and the commutative diagram:

$$\begin{array}{ccc} U^4(S^{4n+7}/\Gamma) & \xrightarrow{r^*} & U^4(S^{4n+3}/\Gamma) \\ \text{PD} \downarrow & & \downarrow \text{PD} \\ U_{4n+3}(S^{4n+7}/\Gamma) & \xrightarrow{r!} & U_{4(n-1)+3}(S^{4n+3}/\Gamma). \end{array}$$

The element $r!([S^{4n+3}/\Gamma, i])$ is $[g^{-1}(S^{4n+3}/\Gamma), g|g^{-1}(S^{4n+3}/\Gamma)]$ where g is the map: $S^{4n+3}/\Gamma \rightarrow S^{4n+7}/\Gamma$ defined by $g([z_1, z_2, \dots, z_{2n+2}]) = [z_1, z_2, \dots, z_{2n}, 0, 0, z_{2n+2}]$ because g is homotopic to r and transversal to S^{4n+3}/Γ . But $g^{-1}(S^{4n+3}/\Gamma) = S^{4(n-1)+3}/\Gamma$ and $g|g^{-1}(S^{4n+3}/\Gamma) = p$. It is easily seen that

$$r!([S^{4n+3}/\Gamma, r]) = [S^{4(n-1)+3}/\Gamma, p] \in U_{4(n-1)+3}(S^{4n+3}/\Gamma),$$

the U -structure on $S^{4(n-1)+3}/\Gamma$ being the canonical one (this result can be found in [7], Lemma 2.5, page 145). Now by 1.8 we have $r^* \circ (PD)^{-1}([S^{4n+3}/\Gamma, r]) = e(\alpha)$, α being \mathbb{C} -vector bundle $S^{4n+3} \times_{\Gamma} \mathbb{C}^2 \rightarrow S^{4n+3}/\Gamma$, Γ acting on S^{4n+3} and \mathbb{C}^2 respectively by using $(n+1)\eta$ and η (see Section I). Since $\alpha = (s \circ r)^*(\eta)(\eta: E \times_{\Gamma} \mathbb{C}^2 \rightarrow B\Gamma)$, we have $e(\alpha) = (s \circ r)^*(D)$ and then $r^* \circ (PD)^{-1}([S^{4n+3}/\Gamma, r]) = (s \circ r)^*(D)$. From the above diagram it follows that $(s \circ r)^*(D) = (PD)^{-1}([S^{4(n-1)+3}/\Gamma, p])$. The fundamental class of S^{4n+3}/Γ for MU involved in the Poincaré duality being $[S^{4n+3}/\Gamma, 1] \in U_{4n+3}(S^{4n+3}/\Gamma)$ (see 1.3) we have:

$$(s \circ r)^*(D) \cap [S^{4n+3}/\Gamma, 1] = [S^{4(n-1)+3}/\Gamma, p]$$

and consequently

$$\begin{aligned} w_{4(n-1)+3} &= (s \circ r)_*([S^{4(n-1)+3}/\Gamma, p]) \\ &= (s \circ r)_*[(s \circ r)^*(D) \cap [S^{4n+3}/\Gamma, 1]] \\ &= D \cap (s \circ r)_*([S^{4n+3}/\Gamma, 1]) \\ &= D \cap [S^{4n+3}/\Gamma, s \circ r] = D \cap w_{4n+3} = \Delta(w_{3n+3}). \end{aligned}$$

We have

$$\begin{aligned} \Delta(u_{4n+1}) &= \Delta(A \cap w_{4n+3}) = (D \cdot A) \cap (w_{4n+3}) \\ &= A \cap [D \cap w_{4n+3}] = A \cap w_{(n-1)+3} = u_{4(n-1)+1}. \end{aligned}$$

Similarly $\Delta(v_{4n-1}) = v_{4(n-1)+1}$. □

REMARK. The homomorphism Δ is sometimes called the Smith-homomorphism.

We recall from [6], Lemma 2.11 and Theorem 2.12, that if Λ_* denotes the $U^*(pt)$ -graded algebra $U^*(pt)[[X, Y, Z]]$, $\dim X = \dim Y = 2$, $\dim Z = 4$ and Ω_* the sub- $U^*(pt)$ -algebra $U^*(pt)[[Z]]$ then there is $T(Z) = 8Z + 2\lambda_2 Z^2 + \sum_{i \geq 3} \lambda_i Z^i \in \Omega_4$, $\lambda_2 \notin 2U^*(pt)$, such that: $M(D) = 0$ ($M(Z) \in \Omega_*$) iff $M(Z) \in T(Z)\Omega_*$. Moreover by [6], Lemmas 2.13, 2.15, there is

$$J(Z) = \mu_1 Z + \sum_{i \geq 2} \mu_i Z^i \in \Omega_0, \quad \mu_1 \notin 2U^*(pt),$$

such that: $E(D) + AM(D) + BN(D) = 0$ iff $M(Z), N(Z)$ belong to $(2 + J(Z))\Omega_*$ and $E(Z)$ to $T(Z)\Omega_*$ ($M(Z), N(Z), E(Z)$ are elements of Ω_*). We also recall the following notation: if $M(Z) = \sum_{i \geq r} a_i Z^i \in \Omega_{2n}$ with $a_r \neq 0$ then $\nu(M) = 4r$. Let W, V_1, V_2 be the $U_*(pt)$ -submodules of $\tilde{U}_*(B\Gamma)$ generated respectively by $\{W_{4n+3}\}_{n \geq 0}$, $\{u_{4n+1}\}_{n \geq 0}$, $\{v_{4n+1}\}_{n \geq 0}$.

THEOREM 2.4. (a) $\tilde{U}_*(B\Gamma) = W \oplus V_1 \oplus V_2$.

(b) In $\tilde{U}_{2p+1}(B\Gamma)$ we have $0 = a_0w_3 + a_1w_7 + \cdots + a_nw_{4n+3} = b_0u_1 + \cdots + b_mu_{4m+1}$ iff there are homogeneous polynomials $M(Z), M_1(Z)$ and homogeneous formal power series $N(Z), N_1(Z)$ of Ω_* satisfying: $b_mZ + b_{m-1}Z^2 + \cdots + b_0Z^{m+1} = M(Z)(2 + J(Z)) + N(Z)$, $a_nZ + a_{n-1}Z^2 + \cdots + a_0Z^{n+1} = M_1(Z)T(Z) + N_1(Z)$, $\nu(N) > 4(m+1)$, $\nu(N_1) > 4(n+1)$. Moreover: $b_0u_1 + \cdots + b_mu_{4m+1} = 0$ iff $b_0v_1 + \cdots + b_mv_{4m+1} = 0$.

Proof. (a) Suppose that $(a_0w_3 + \cdots + a_nw_{4n+3}) + (b_0u_1 + \cdots + b_mu_{4m+1}) + (c_0v_1 + \cdots + c_rv_{4r+1}) = 0$. Then a proof similar to that of Lemma 2.14 of [6] shows that $b_m = 2d_m$, $d_m \in U_*(pt)$. Consider $H(Z) = b_mZ + b_{m-1}Z^2 + \cdots + b_0Z^{m+1}$; we have: $H(Z) - d_mZ(2 + J(Z)) = b'_{m-1}Z^2 + \cdots + b'_0Z^{m+1} + F(Z)$, $\nu(F) > 4(m+1)$. Then $AH(D) = A[b'_{m-1}D^2 + \cdots + b'_0D^{m+1}] + AF(D)$ and by taking the cup product by w_{4m+7} we obtain $b_0u_1 + \cdots + b_mu_{4m+1} = b'_0u_1 + \cdots + b'_{m-1}u_{4(m-1)+1}$. As seen before, we have: $b'_{m-1} = 2d'_{m-1}$, $d'_{m-1} \in U_*(pt)$. We repeat the same process and after a finite number of operations we get $b_mZ + \cdots + b_0Z^{m+1} = M(Z)(2 + J(Z)) + N(Z)$, $M(Z)$ being a homogeneous polynomial and $N(Z)$ a homogeneous formal power series such that $\nu(N) > 4(m+1)$. Hence $b_0u_1 + \cdots + b_mu_{4m+1} = M(D)A(2 + J(D)) \cap w_{4m+7} = 0$. Similarly $c_0v_1 + \cdots + c_rv_{4r+1} = 0$ which ends the proof of part (a).

(b) Suppose that $a_0w_3 + \cdots + a_nw_{4n+3} = 0$. As in Proposition 2.6 of [6] we have $a_n = 8e_n$, $e_n \in U_*(pt)$. We form $a_nZ + \cdots + a_0Z^{n+1} - e_nT(Z) = a'_{n-1}Z^2 + \cdots + a'_0Z^{n+1} + F_1(Z)$, $\nu(F_1) > 4(n+1)$ and by taking the cup-product by w_{4n+7} we obtain: $a_0w_3 + \cdots + a_nw_{4n+3} = a'_0w_3 + a'_{n-1}w_{4(n-1)+3}$. As before, we have $a'_{n-1} = 8e'_{n-1}$, $e'_{n-1} \in U_*(pt)$. We repeat the same process with $a'_{n-1}Z^2 + a'_{n-2}Z^3 + \cdots + a'_0Z^{n+1}$ and after a finite number of operations we get: $a_nZ + \cdots + a_0Z^{n+1} = M_1(Z)T(Z) + N_1(Z)$, $\nu(N_1) > 4(n+1)$, $M_1(Z)$ being a homogeneous polynomial and $N_1(Z)$ a homogeneous formal power series. The proof of part (a) shows that $b_mZ + \cdots + b_0Z^{m+1} = M(Z)(2 + J(Z)) + N(Z)$, $\nu(N) > 4(m+1)$. The remaining part of (b) is evident. \square

(2) Orders of the Generators.

LEMMA 2.5. Suppose $t \in \tilde{U}_{2n+1}(B\Gamma)$, $t \neq 0$. If $a \in U_4(pt)$ is such that $a \notin 2U_4(pt)$ then $a \cdot t \neq 0$.

Proof. Since $t \neq 0$ there is an integer $q \geq 0$ such that $t \in J_{q,2n+1-q}$ and $t \notin J_{q-1,2n+1-(q-1)}$. We have either $q = 4s + 3$ or $q = 4s + 1$.

Suppose $q = 4s + 3$. We have the following commutative diagram:

$$\begin{array}{ccc}
 U_4(pt) \otimes J_{4s+3, 2n+1-(4s+3)} & \xrightarrow{\chi} & J_{4s+3, 2(n-2s+1)} \\
 1 \otimes h \downarrow & & \downarrow h \\
 U_4(pt) \otimes U_{2(n-2s-1)}(pt) \otimes \mathbb{Z}_8 & \xrightarrow{\times \otimes 1} & U_{2(n-2s+1)} \otimes \mathbb{Z}_8
 \end{array}$$

where h is the canonical map: $J_{**} \rightarrow E_{**}^\infty = H_*(B\Gamma, U_*(pt)) = U_*(pt) \otimes H_*(B\Gamma)$. It is enough to prove that in $U_*(pt) = \mathbb{Z}[x_1, x_2, \dots, x_4, \dots]$ if $a \in U_4(pt)$, $a \notin 2U_4(pt)$, $b \in U_{2k}(pt)$, $b \notin 8U_{2k}(pt)$ then $ab \notin 8U_{2(k+2)}(pt)$; we may suppose that a and b are monomials and then the assertion is clear. The case $q = 4s + 1$ is similar. \square

THEOREM 2.6. *We have $\text{ord } w_{4n+3} = 2^{2n+3}$.*

Proof. (a) $\text{ord } w_3 = 2^3$.

We have $0 = T(D) = 2^3D + H(D)D^2$ and $0 = T(D) \cap w_7 = 2^2w_3 + H(D) \cap (D^2 \cap w_7) = 2^3w_3$ because $D^2 \cap w_7 \in U_{-1}(B\Gamma) = 0$. Then by using the edge homomorphism $\mu': \tilde{U}_3(B\Gamma) \rightarrow \tilde{H}_3(B\Gamma) = \mathbb{Z}_8$ we see that $2^2w_3 \neq 0$. Hence $\text{ord } w_3 = 2^3$.

(b) *Suppose $\text{ord } w_{4i+3} = 2^{2i+3}$, $0 \leq i \leq n-1$.*

We have $0 = T(D) = 2^3D + 2\lambda_2D^2 + \lambda_3D^3 + \dots + \lambda_{n+1}D^{n+1} + H(D)D^{n+2}$, $\lambda_2 \in U^{-4}(pt) = U_4(pt)$, $\lambda_2 \notin 2U_4(pt)$. Take the cup-product by w_{4n+7} : $2^3w_{4n+3} + 2\lambda_2w_{4(n-1)+3} + \lambda_3w_{4(n-2)+3} + \dots + \lambda_{n+1}w_3 = 0$ and after multiplication by 2^{2n-1} we get: $2^{2n+2}w_{4n+3} + \lambda_22^{2n}w_{4(n-1)+3} = 0$; since $\text{ord } w_{4(n-1)+3} = 2^{2n+1}$ we have $2^{2n}w_{4(n-1)+3} \neq 0$ and by 2.5 $\lambda_22^{2n}w_{4(n-1)+3} \neq 0$ because $\lambda_2 \notin 2U_4(pt)$. Hence $2^{2n+2}w_{4n+3} \neq 0$. Now we have: $2^{2n+3}w_{4n+3} = -\lambda_22^{2n+1}w_{4(n-1)+3} = 0$. It follows that $\text{ord } w_{4n+3} = 2^{2n+3}$ which ends the proof of 2.6. \square

LEMMA 2.7. *If $G_n = U_{4n-2}(pt)w_3 + U_{4n}(pt)u_1 + U_{4n}(pt)v_1$, $G'_n = U_{4n}(pt)w_3 + U_{4n+2}(pt)u_1 + U_{4n+2}(pt)v_1$ then we have the exact sequences:*

$$\begin{aligned}
 0 \rightarrow G_n &\rightarrow \tilde{U}_{4n+1}(B\Gamma) \xrightarrow{\Delta} \tilde{U}_{4(n-1)+1}(B\Gamma) \rightarrow 0 \\
 0 \rightarrow G'_n &\rightarrow \tilde{U}_{4n+3}(B\Gamma) \xrightarrow{\Delta} \tilde{U}_{4(n-1)+3}(B\Gamma) \rightarrow 0
 \end{aligned}$$

Proof. We wish to show that the sequence:

$$0 \rightarrow G_n \rightarrow \tilde{U}_{4n+1}(B\Gamma) \xrightarrow{\Delta} \tilde{U}_{4(n-1)+1}(B\Gamma) \rightarrow 0$$

is exact. It follows by 2.3 that Δ is surjective and $G_n \subset \ker \Delta$. Suppose $0 = aw_3 + bu_1 + cv_1$, $a \in U_{4n-2}(pt)$, $b \in U_{4n}(pt)$, $c \in U_{4n}(pt)$. Then $a \cdot w_3 \in J_{3,4n-2}$ and since $bu_1 + cv_1 \in J_{1,4n}$ we have $a \cdot w_3 \in J_{2,4n-1} \supset J_{1,4n}$. If h denotes the quotient map: $J_{3,4n-2} \rightarrow J_{3,4n-2}/J_{2,4n-1} = H^3(B\Gamma, U_{4n-2}(pt)) = U_{4n-2}(pt)/8U_{4n-2}(pt)$, it follows that $h(aw_3) = 0$ and consequently $a = 2^3a'$. Hence $aw_3 = a'2^3w_3 = 0$ and then $bu_1 + cv_1 = 0$. Similarly we have $b = 2b'$, $c = 2c'$ which means that $0 = aw_3 + bu_1 + cv_1$ ($a \in U_{4n-2}(pt)$, $b \in U_{4n}(pt)$, $c \in U_{4n}(pt)$) if and only if $a = 2^3a'$, $b = 2b'$, $c = 2c'$. Hence $\text{ord } G_n = 2^k$, $k = 3 \text{Rank } U_{4n-2}(pt) + 2 \text{Rank } U_{4n}(pt)$. Now, we have $\text{ord } \ker \Delta = \text{ord } \tilde{U}_{4n+1}(B\Gamma)/\text{ord } \tilde{U}_{4(n-1)+1}(B\Gamma) = 2^k$ by 2.1. From $G_n \subset \ker \Delta$ and $\text{ord } G_n = \text{ord } \ker \Delta$ we see that the sequence $0 \rightarrow G_n \rightarrow \tilde{U}_{4n+1}(B\Gamma) \xrightarrow{\Delta} \tilde{U}_{4(n-1)+1}(B\Gamma) \rightarrow 0$ is exact. A similar proof shows that the sequence $0 \rightarrow G'_n \rightarrow \tilde{U}_{4n+3}(B\Gamma) \xrightarrow{\Delta} \tilde{U}_{4(n-1)+3}(B\Gamma) \rightarrow 0$ is exact. \square

THEOREM 2.8. *We have $\text{ord } u_{4n+1} = \text{ord } v_{4n+1} = 2^{n+1}$.*

Proof. If $n = 0$ the assertion is clear. Suppose $\text{ord } u_{4i+1} = 2^{i+1}$, $0 \leq i \leq n-1$. Then $\Delta(2^n u_{4n+1}) = 2^n u_{4(n-1)+1} = 0$ and since the sequence $0 \rightarrow G_n \rightarrow U_{4n+1}(B\Gamma) \xrightarrow{\Delta} U_{4(n-1)+1}(B\Gamma) \rightarrow 0$ is exact, (see 2.7), there are $a \in U_{4n-2}(pt)$, $b \in U_{4n}(pt)$, $c \in U_{4n}(pt)$ such that $2^n u_{4n+1} = aw_3 + bu_1 + cv_1$. It follows that $-bu_1 + 2^n \cdot u_{4n+1} = 0$ and $2^{n+1} \cdot u_{4n+1} = 0$; hence $\text{ord } u_{4n+1} \leq 2^{n+1}$. By Theorem 2.4 there are $M(Z), N(Z)$ in Ω_* such that: $2^n Z - bZ^{n+1} = M(Z)(2+J(Z)) + N(Z)$, $\nu(N) > 4(n+1)$. If $M(Z) = h_1 Z + h_2 Z^2 + \dots$, then we have:

$$\begin{aligned} 2^n Z - bZ^{n+1} &= (2 + \mu_1 Z + \mu_2 Z^2 + \dots)(h_1 Z + h_2 Z^2 + \dots) \\ &\quad + e_{n+2} Z^{n+2} + e_{n+3} Z^{n+3} + \dots, \quad \mu_1 \notin 2U_*(pt). \end{aligned}$$

A straightforward calculation shows that $2^{n-j}|h_j$ and $2^{n-j+1} \nmid h_j$, $1 \leq j \leq n$. We have: $-b = 2h_{n+1} + \mu_1 h_n + \mu_2 h_{n-1} + \dots + \mu_n h_1$; as $2|h_j$, $1 \leq j \leq n-1$, $2 \nmid h_n$, $2 \nmid \mu_1$ we have $2 \nmid b$. As a consequence we get $2^n u_{4n+1} \neq 0$ and $\text{ord } u_{4n+1} = 2^{n+1}$. Similarly $\text{ord } v_{4n+1} = 2^{n+1}$. \square

III. $\tilde{U}^*(B\Gamma_k)$, $k \geq 4$: generators, orders and relations. We have seen in [6], Section III, that there are elements $D_k \in \tilde{U}^4(B\Gamma_k)$, $B_k \in \tilde{U}^2(B\Gamma_k)$, $C_k \in \tilde{U}^2(B\Gamma_k)$ defined as Euler classes of irreducible unitary representations η_1, ξ_2, ξ_3 of Γ_k . Moreover in the same article

(Sec. III) we have determined three homogeneous formal power series $T_k(Z) \in \Omega_4$, $J_k(Z) \in \Omega_0$, $G_k(Z) \in \Omega_2$ such that $B_k(2 + J(D_k)) + G_k(D_k) = C_k(2 + J(D_k)) + G_k(D_k) = 0$ and there is $G'_k(Z) \in \Omega_2$ satisfying $G_k(Z) = (2 + J(Z))G'_k(Z)$. Then with $B'_k = B_k + G'_k(D_k)$, $C'_k = C_k + G'_k(D_k)$ and μ being the edge homomorphism: $U^2(B\Gamma_k) \rightarrow H^2(B\Gamma_k)$ we see that $\mu(B'_k) = \mu(B_k)$ and $\mu(C'_k) = \mu(C_k)$ are generators of the group $H^2(B\Gamma_k)$; $\mu(D_k)$ is obviously a generator of $H^4(B\Gamma_k)$. Moreover $B'_k(2 + J(D_k)) = C'_k(2 + J(D_k)) = 0$.

Now let $w'_{4n+3} \in \tilde{U}_{4n+3}(B\Gamma_k)$ be $[S^{4n+3}/\Gamma_k, q']$, q' being the inclusion $S^{4n+3}/\Gamma_k \subset B\Gamma_k$, $u'_{4n+1} = B'_k \cap w'_{4n+3} \in \tilde{U}_{4n+1}$, $v'_{4n+1} = C'_k \cap w'_{4n+3} \in \tilde{U}_{4n+1}(B\Gamma_k)$. Then we have the following theorems whose proofs are identical respectively to Theorem 2.2 and Theorem 2.4 and therefore will be omitted.

THEOREM 3.1. *The set $\{u'_{4n+1}, v'_{4n+1}, w'_{4n+3}\}_{n \geq 0}$ is a system of generators for the $U(pt)$ -module $\tilde{U}_*(B\Gamma_k)$.* \square

Now let W', V'_1, V'_2 be the $U_*(pt)$ -submodules of $\tilde{U}_*(B\Gamma_k)$ generated respectively by $\{w'_{4n+3}\}_{n \geq 0}$, $\{u'_{4n+1}\}_{n \geq 0}$, $\{v'_{4n+1}\}_{n \geq 0}$.

THEOREM 3.2. (a) $\tilde{U}_*(B\Gamma_k) = W' \oplus V'_1 \oplus V'_2$.

(b) In $\tilde{U}_{2p+1}(B\Gamma_k)$ we have $0 = a_0w'_3 + a_1w'_7 + \dots + a_nw'_{4n+3} = b_0u'_1 + \dots + b_mu'_{4m+1}$ iff there are homogeneous polynomials $M(Z), M_1(Z)$ and homogeneous formal power series $N(Z), N_1(Z)$ of Ω_* satisfying: $b_mZ + b_{m-1}Z^2 + \dots + b_0Z^{m+1} = M(Z)(2 + J(Z)) + N(Z)$, $a_nZ + a_{n-1}Z^2 + \dots + a_0Z^{n+1} = M_1(Z)T_k(Z) + N_1(Z)$, $\nu(N) > 4(m+1)$, $\nu(N_1) > 4(n+1)$. Moreover $b_0u'_1 + \dots + b_mu'_{4m+1} = 0$ iff $b_0v'_1 + \dots + b_mv'_{4m+1} = 0$. \square

There is a Smith homomorphism $\Delta: \tilde{U}_*(B\Gamma_k) \rightarrow \tilde{U}_*(B\Gamma_k)$ of degree -4 such that

$$\begin{aligned} \Delta(w'_{4n+3}) &= D_k \cap w'_{4n+3} = w'_{4(n-1)+3}, \\ \Delta(u'_{4n+1}) &= D_k \cap u'_{4n+1} = D_k \cap (B'_k \cap w'_{4n+3}) = B'_k \cap (D_k \cap w'_{4n+3}) \\ &= B'_k \cap w'_{4(n-1)+3} = u'_{4(n-1)+1}, \Delta(v'_{4n+1}) = v'_{4(n-1)+1}. \end{aligned}$$

If

$$\begin{aligned} F_n &= U_{4n}(pt)w'_3 + U_{4n+2}(pt)u'_1 + U_{4n+2}(pt)v'_1, \\ F'_n &= U_{4n-2}(pt)w'_3 + U_{4n}(pt)u'_1 + U_{4n}(pt)v'_1 \end{aligned}$$

then we have:

LEMMA 3.3. *The following sequences are exact:*

$$0 \rightarrow F_n \rightarrow U_{4n+3}(B\Gamma_k) \xrightarrow{\Delta} U_{4(n-1)+3}(B\Gamma_k) \rightarrow 0,$$

$$0 \rightarrow F'_n \rightarrow U_{4n+1}(B\Gamma_k) \xrightarrow{\Delta} U_{4(n-1)+1}(B\Gamma_k) \rightarrow 0.$$

Proof. The proof is similar to that of Lemma 2.7. □

It remains to calculate the orders of the generators.

THEOREM 3.4. *We have: $\text{ord } w'_{4n+3} = 2^{2n+k}$, $n \geq 0$.*

Proof. We have $0 = T_k(D_k) = 2^k D_k + H(D_k)D_k^2$ and then $0 = (2^k D_k + H(D_k)D_k^2) \cap w_7 = 2^k w_3$ because: $D_k^2 \cap w_7 \in \tilde{U}_{-1}(B\Gamma_k) = 0$. Now if μ' is the edge homomorphism: $U_3(B\Gamma_k) \rightarrow H_3(B\Gamma_k) = \mathbb{Z}_2 k$ then we have $\mu'(w'_3) = 1 \in \mathbb{Z}_2 k$ and consequently $2^{k-1} w_3 \neq 0$. Then $\text{ord } w_3 = 2^k$.

Suppose that $\text{ord } w'_{4i+3} = 2^{2i+k}$, $0 \leq i \leq n-1$. Then

$$\begin{aligned} 0 = T_k(D_k) \cap w'_{4n+7} &= 2^k w'_{4n+3} + 2^{k-2} \lambda'_2 w'_{4(n-1)+3} \\ &+ \cdots + 2^{k-i} \lambda'_i w'_{4(n-1)+3} + \cdots + 2\lambda'_{k-1} w'_{4(n-k+2)+3} \\ &+ \lambda'_k w'_{4(n-k+1)+3} + \cdots + \lambda'_m w'_{4(n-m+1)+3} + \cdots, \end{aligned}$$

the number of non-zero elements in this sum being finite. If $3 \leq i \leq k-1$ we have $2^{2n-1+k-i} w'_{4(n-i+1)+3} = 0$ because $\text{ord } w'_{4(n-i+1)+3} = 2^{2(n-i+1)+k}$ and $2(n-i+1)+k \leq 2n-1+k-i$ since $i \geq 3$. If $m \geq k$ (≥ 3) we have $2^{2n-1} w'_{4(n-m+1)+3} = 0$ because $\text{ord } w'_{4(n-m+1)+3} = 2^{2(n-m+1)+k}$ and $2(n-m+1)+k \leq 2n-1$ since $k \leq m \leq 2m-3$. It follows that $2^{2n-1+k} w'_{4n+3} + 2^{2n-3+k} \lambda'_2 w'_{4n-1} = 0$. Now $2^{2n-3+k} w'_{4n-1} \neq 0$ because $\text{ord } w'_{4n-1} = 2^{2n-2+k}$; since $\lambda'_2 \notin 2U^{-4}(pt)$ we have $2^{2n-3+k} \lambda'_2 w'_{4n-1} \neq 0$ (see 2.5). Hence

$$2^{2n-1+k} w'_{4n+3} \neq 0 \quad \text{and} \quad 2^{2n+k} w'_{4n+3} = -2^{2(n-1)+k} \lambda'_2 w'_{4n-1} = 0.$$

We have proved that $\text{ord } w_{4n+3} = 2^{2n+k}$. □

THEOREM 3.5. *We have: $\text{ord } u'_{4n+1} = \text{ord } v'_{4n+1} = 2^{n+1}$, $n \geq 0$, which are therefore independent of k .*

Proof. The proof of 3.5 is based on Theorem 3.2 and Lemma 3.3 and is exactly the same as the one of Theorem 2.8. □

REFERENCES

- [1] J. F. Adams, *Stable Homotopy and Generalized Homology*, University of Chicago Mathematics Lecture Notes, 1971.
- [2] J. M. Boardman, *Stable Homotopy Theory*, Mimeographed notes, Chapter V, VI, Warwick, 1966.
- [3] P. E. Conner and E. E. Floyd, *Differentiable Periodic Maps*, Springer, 1964.
- [4] —, *Periodic maps which preserve a complex structure*, Bull. Amer. Math. Soc., **70** (1964), 574–579.
- [5] —, *Torsion in SU-bordism*, Memoirs Amer. Math. Soc., **60** (1966).
- [6] A. Mesnaoui, *Unitary cobordism of classifying spaces of quaternion groups*, to be published.
- [7] D. Pitt, *On the complex bordism of the generalized quaternion groups*, J. London Math. Soc., **16** (1977), 142–148.
- [8] R. M. Switzer, *Algebraic Topology, Homotopy and Homology*, Vol. 2.2, Springer-Verlag, 1975.
- [9] R. H. Szczarba, *On tangent bundles of fibre spaces and quotient spaces*, Amer. J. Math., **86** (1964), 685–697.
- [10] G. Wilson, *K-theory invariants for unitary G-bordism*, Quart. J. Math., Oxford, **24** (1973), 499–526.

Received October 5, 1986 and, in revised form, August 15, 1988.

UNIVERSITÉ MOHAMMED V
B. P. 1014
RABAT, MOROCCO

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Marco Andreatta, Mauro Beltrametti and Andrew Sommese, Generic properties of the adjunction mapping for singular surfaces and applications	1
Chen-Lian Chuang and Pjek-Hwee Lee, On regular subdirect products of simple Artinian rings	17
Fernando Giménez and Vicente Miquel Molina, Volume estimates for real hypersurfaces of a Kaehler manifold with strictly positive holomorphic sectional and antiholomorphic Ricci curvatures	23
Richard J. Griego and Andrzej Korzeniowski, Asymptotics for certain Wiener integrals associated with higher order differential operators	41
Abdeslam Mesnaoui, Unitary bordism of classifying spaces of quaternion groups	49
Abdeslam Mesnaoui, Unitary cobordism of classifying spaces of quaternion groups	69
Jesper M. Møller, On equivariant function spaces	103
Bassam Nassrallah, A q-analogue of Appell's F_1 function, its integral representation and transformations	121
Peter A Ohring, Solvability of invariant differential operators on metabelian groups	135
Athanase Papadopoulos and R. C. Penner, Enumerating pseudo-Anosov foliations	159
Ti-Jun Xiao and Liang Jin, On complete second order linear differential equations in Banach spaces	175
Carl Widland and Robert F. Lax, Weierstrass points on Gorenstein curves	197