

Pacific Journal of Mathematics

**BANACH ALGEBRAS ASSOCIATED WITH SPHERICAL
REPRESENTATIONS OF THE FREE GROUP**

RYSZARD SZWARC

BANACH ALGEBRAS ASSOCIATED WITH SPHERICAL REPRESENTATIONS OF THE FREE GROUP

RYSZARD SZWARC

We prove that any spherical representation of the free group \mathbb{F} weakly contains the regular representation. Moreover C_π^* , the C^* -algebra associated with the spherical representation π , is a compact extension of the reduced C^* -algebra of \mathbb{F} . We also show that the standard projection onto radial functions admits extensions to C_π^* for a class of representations π of \mathbb{F} which includes spherical representations, as well as the regular representation and the universal representation.

Introduction. Let \mathbb{F}_r be a free group on r generators x_1, \dots, x_r . Let μ_1 be the finitely supported probability measure equidistributed on $\{x_1^{\pm 1}, x_2^{\pm 1}, \dots, x_r^{\pm 1}\}$. The operator of convolution by μ_1 is the analogue of the Laplace-Beltrami operator on Riemann rank one symmetric spaces. By [11] the l^1 -spectrum of μ_1 can be identified with the ellipse $E = \{z = x + iy: x^2 + (\frac{r}{r-1}y)^2 \leq 1\}$. Any point z of E corresponds in one-to-one fashion to a spherical function ϕ_z the eigenfunction of μ_1 with eigenvalue z . We refer to [11], [7] for this subject.

For real z spherical functions are positive definite and give rise to unitary representations of \mathbb{F}_r . Basing our argument on a particular realisation of these representations and on the simplicity of the reduced C^* -algebra of \mathbb{F}_r [10], we prove that all spherical representations weakly contain the regular representation. Moreover the C^* -algebras associated with spherical representations are compact extensions of $C_{\text{red}}^*(\mathbb{F}_r)$, the C^* -algebra associated with the regular representation.

Finally we consider the standard projection onto radial functions on \mathbb{F}_r and we prove that it is bounded on any C^* -algebra associated with spherical functions.

Acknowledgment. I am grateful to Uffe Haagerup for suggesting the problem and for helpful discussions.

Spherical representations. Let \mathbb{F}_r be a free group on r generators x_1, x_2, \dots, x_r , $r \geq 2$. Any element x of \mathbb{F}_r may be uniquely ex-

pressed as a reduced word in x_1, \dots, x_r and their inverses. The number of letters of this reduced word is called the length of x and is denoted by $|x|$.

A complex function on \mathbb{F}_r which values depend only on the length $|x|$ will be called radial. They are the analogues of bi- K -invariant functions on $\mathrm{SL}(2, \mathbb{R})$. The space $l_{\#}^1$ of absolutely summable radial functions forms a commutative Banach algebra with respect to convolution operation. This is due to the fact that if χ_n is defined as $\chi_n(x) = 1$ if $|x| = n$ and 0 otherwise, then:

$$(1) \quad \begin{aligned} \chi_n^* \chi_n &= \chi_{n+1} + (2r-1)\chi_{n-1}, & n \geq 2 \\ \chi_1^* \chi_1 &= \chi_2 + 2r\chi_0 \end{aligned}$$

(see [3], [4], [11]). In particular it means that $l_{\#}^1$ is generated by the function χ_1 . The Gelfand space of $l_{\#}^1$, when identified with the l^1 -spectrum of χ_1 , coincides ([11], Theorem 2.1) with the closed ellipse:

$$E = \left\{ z = x + iy \in \mathbb{C}: \left(\frac{x}{2r} \right)^2 + \left(\frac{y}{2r-2} \right)^2 \leq 1 \right\}.$$

Moreover multiplicative functionals on $l_{\#}^1$ are given by

$$(2) \quad l_{\#}^1 \ni f \mapsto \langle f, \bar{\phi}_z \rangle = \sum_{x \in \mathbb{F}_r} f(x) \phi_z(x),$$

where ϕ_z , $z \in E$, are bounded complex functions with properties:

- (i) $\phi_z(e) = 1$;
- (ii) $\chi_z^* \phi_z = z \phi_z$;
- (iii) ϕ_z is radial.

In analogy to $\mathrm{SL}(2, \mathbb{R})$ such functions are called spherical (see [3], [7]) and the properties (i), (ii), (iii) determine ϕ_z uniquely. Explicit formulas expressing ϕ_z can be found in [7], [11] and [13].

The theory developed in the papers [11], [7], [9] (see also [13]) gives that for any point z of the interior of E (including also two points $2r$ and $(-2r)$ which correspond to the only characters $x \mapsto 1$ and $x \mapsto (-1)^{|x|}$ on \mathbb{F}_r) the spherical function ϕ_z occurs as the matrix coefficient of a uniformly bounded representation of \mathbb{F}_r . The real segment $[-2r, 2r]$ corresponds to all unitary representations: in particular the segment $[-2\sqrt{2r-1}, 2\sqrt{2r-1}]$ (i.e. the l^2 -spectrum of convolution operator by χ_1) corresponds to the principal series while the two remaining parts $[-2r, -2\sqrt{2r-1})$ and $(2\sqrt{2r-1}, 2r]$ form the complementary series.

In the sequel we will need a realisation of spherical representations given in [13] because of its good additional features. In [13] the spherical functions are parametrized by the annulus $A = \{z \in \mathbb{C}: (2r-1)^{-1} \leq |z| \leq 1\}$ which can be mapped onto E by the function $\gamma(z) = (2r-1)z + \frac{1}{z}$. Taking it into account we reformulate below the main results of [13] with respect to the ellipse-parametrization.

THEOREM 1 ([12], Theorem 8 and remarks following it). *Let \mathbb{F}_r be a free group on r generators. There exists an analytic series of uniformly bounded representations π_z , $z \in E = \{x + iy: (\frac{x}{2r})^2 + (\frac{y}{2r-2})^2 \leq 1\}$, on a Hilbert space \mathcal{H} and a unit vector $\zeta \in \mathcal{H}$ such that:*

- (i) $\langle \pi_z(x)\zeta, \zeta \rangle = \phi_z(x)$, $x \in \mathbb{F}_r$;
- (ii) $\pi_z(x)^* = \pi_{\bar{z}}(x)^{-1}$, $x \in \mathbb{F}_r$;
- (iii) $\pi_z(x) - \pi_{z'}(x)$ has finite rank for any $z, z' \in E$ and $x \in \mathbb{F}_r$;
- (iv) Any representation π_z leaves no nontrivial subspace invariant and representations π_z are mutually inequivalent;
- (v) π_z is unitary if and only if $z \in \mathbb{R}$. Otherwise π_z cannot be made unitary by introducing equivalent inner product.

The series $[-2\sqrt{2r-1}, 2\sqrt{2r-1}]$ occurs in decomposition of the regular representation (see [11], Theorem 6.2). That is why it is called the principal series. Remaining parts of $[-2r, 2r]$ are called the complementary series.

The aim of the paper is to show that any representation of complementary series, as well as of principal series, weakly contains the regular representation. In some sense (see Lemma 2) we also prove it for nonunitary spherical representations. The crucial are Theorem 1 (iii) and the Powers' theorem which states that $C_{\text{red}}^*(\mathbb{F}_r)$, the reduced C^* -algebra of \mathbb{F}_r , is simple i.e. contains no nontrivial two-sided ideals.

Let λ denote the left regular representation of \mathbb{F}_r . By Powers' theorem any unitary representation weakly contained in λ is weakly equivalent to λ .

LEMMA 1. *Let (π, \mathcal{H}) be a unitary representation of \mathbb{F}_r weakly contained in the regular representation. Then for any compact operator K on \mathcal{H} and any $f \in l^1(\mathbb{F}_r)$ we have:*

$$\|\pi(f) + K\| \geq \|\pi(f)\|.$$

Proof. Let C_π^* be the C^* -algebra associated with π . Then by remarks preceding the lemma C_π^* is isometrically isomorphic to $C_{\text{red}}^*(\mathbb{F}_r)$.

Hence by [10] it is a simple C^* -algebra. Consider the homomorphism of C_π^* into the Calkin algebra $\mathcal{B}(\mathcal{H})/\mathcal{K}(\mathcal{H})$ (where $\mathcal{K}(\mathcal{H})$ denotes the set of all linear compact operators on \mathcal{H}) given on $l^1(\mathbb{F}_r)$ as

$$l^1(\mathbb{F}_r) \ni f \mapsto \pi(f) + \mathcal{K}(\mathcal{H}) \in \mathcal{B}(\mathcal{H})/\mathcal{K}(\mathcal{H}).$$

Because C_π^* has no ideals then this mapping is an isometry. Thus $\inf_{K \in \mathcal{K}(\mathcal{H})} \|\pi(f) + K\| = \|\pi(f)\|$ which proves the lemma.

LEMMA 2. *Let $z \in E$ and $f \in l^1(\mathbb{F}_r)$. Then $\|\pi_z(f)\| \geq \|\lambda(f)\|$. In particular any spherical representation π_z , $z \in E$ has trivial kernel in $l^1(\mathbb{F}_r)$.*

Proof. Let us apply Lemma 1 to a representation of the principal series e.g. π_0 . By Theorem 1 (iii) the operator $k = \pi_z(f) - \pi_0(f)$ is compact whatever f from $l^1(\mathbb{F}_r)$ is. Therefore

$$\|\pi_z(f)\| = \|\pi_0(f) + K\| \geq \|\pi_0(f)\| = \|\lambda(f)\|$$

because π_0 and λ are weakly equivalent.

Lemma 2 implies immediately:

THEOREM 2. *Any representation π_z of the complementary series of \mathbb{F}_r , except two characters $z = \pm 2r$, contains weakly the regular representation.*

For any $z \in E$ let C_{π_z} denote the completion of $l^1(\mathbb{F}_r)$ with respect to the norm $\|f\|_{C_{\pi_z}} = \|\pi_z(f)\|$. Then C_{π_z} is a Banach algebra which becomes a C^* -algebra whenever π_z is unitary i.e. $z \in [-2r, 2r]$.

THEOREM 3. *Let $z \in \int E$. Then the identity map $f \mapsto f$ on $l^1(\mathbb{F}_r)$ extends to an epimorphism from C_{π_z} onto $C_{\text{red}}^*(\mathbb{F}_r)$. Moreover the kernel of this map is isomorphic to the ideal of all compact operators on a Hilbert space.*

Proof. The first part follows from Lemma 2. Next observe that C_{π_z} is isometrically isomorphic to the norm closure of $\{\pi_z(f) : f \in l^1(\mathbb{F}_r)\}$ in $\mathcal{B}(\mathcal{H})$ because π_0 and λ are weakly equivalent. Thus we have to determine the kernel of the homomorphism between these two subalgebras of $\mathcal{B}(\mathcal{H})$ given by:

$$\pi_z(f) \longmapsto \pi_0(f), \quad f \in l^1(\mathbb{F}_r).$$

Let $\pi_z(f_n)$ be a Cauchy sequence in $B(\mathcal{H})$ such that $\pi_0(f_n) \xrightarrow{n} 0$. Hence $\pi_z(f_n) = \pi_0(f_n) + K_n$, where K_n , $n = 1, 2, \dots$, are compact operators (see Theorem 1 (iii)). This yields $\lim_n \pi_z(f_n) = \lim_n K_n$ is a compact operator. It means that the kernel is contained in $\mathcal{K}(\mathcal{H})$. On the other hand C_{π_z} contains $\mathcal{K}(\mathcal{H})$ by [13] (Remark following Proposition 2). Moreover the image of $\mathcal{K}(\mathcal{H})$ under this homomorphism must be trivial because it is a two-sided ideal in $C_{\pi_0}^*$.

REMARK. Theorem 2 implies that C_{π_z} are compact extensions of the reduced C^* -algebra of \mathbb{F}_r in the sense of [2] (see also [6]). All these extensions are *trivial* (see [2] for the definition) because they coincide with the trivial extension corresponding to a representation of the principal series.

PROPOSITION 1. *Let $z \in E$. Then the spectrum of the operator $\pi_z(\chi_1)$ coincides with $\{z\} \cup [-2\sqrt{2r-1}, 2\sqrt{2r-1}]$.*

Proof. By [13] (Theorem 5) the spectrum of $\pi_z(\chi_1)$ consists of $\{z\}$ and a subset of $[-2\sqrt{2r-1}, 2\sqrt{2r-1}]$ (recall that we use another parametrization). Thus we should prove only that the entire interval enters into the spectrum. However it follows from the fact that $\pi_z(\chi_1)$ is a finite dimensional perturbation of the selfadjoint operator $\pi_0(\chi_1)$ which spectrum coincides with the spectrum of $\lambda(\chi_1)$ in $C_{\text{red}}^*(\mathbb{F}_r)$ (because π_0 and λ are weakly equivalent) i.e. with the interval $[-2\sqrt{2r-1}, 2\sqrt{2r-1}]$.

Conditional expectation related to radial functions. As we have seen before the radial functions on \mathbb{F}_r form a commutative convolution algebra. There is a natural projection \mathcal{E} from the set of all complex function on \mathbb{F}_r onto radial functions:

$$\mathcal{E}f(x) = \frac{1}{\#\{x \in \mathbb{F}_r : |x| = n\}} \sum_{|x|=n} f(x).$$

The operator \mathcal{E} satisfies (cf. [11], [5])

- (i) $\mathcal{E}f = f$ if and only if f is radial;
- (ii) $\mathcal{E}(f * \mathcal{E}g) = \mathcal{E}(\mathcal{E}f * g) = \mathcal{E}(f * \mathcal{E}g)$;
- (iii) $\mathcal{E}f(e) = f(e)$.

These properties determine \mathcal{E} uniquely. By general theorems concerning weak conditional expectations \mathcal{E} extends to the contraction on the von Neumann algebra $\text{VN}(\mathbb{F}_r)$ of \mathbb{F}_r (see [5]).

Let π be a unitary representation of \mathbb{F}_r without kernel in $l^1(\mathbb{F}_r)$. There arises a question whether the operator \mathcal{E} can extend from $l^1(\mathbb{F}_r)$ to a contraction on $C_\pi^*(\mathbb{F}_r)$.

PROPOSITION 2. *Let π be a unitary representation of \mathbb{F}_r . Assume that π contains weakly all spherical representations π_t such that t belongs to the spectrum of the operator $\pi(\chi_1)$. Then \mathcal{E} extends by continuity to a contraction on $C_\pi^*(\mathbb{F}_r)$.*

Proof. It suffices to show that for any function f in $l^1(\mathbb{F}_r)$ there holds $\|\pi(\mathcal{E}f)\| \leq \|\pi(f)\|$. Denote by \mathcal{A} the unital C^* -algebra generated by $\pi(\chi_z)$. If $t \in \sigma(\pi(\chi_1))$ then by (2), the Gelfand-Naimark theorem and Theorem 2 (i) we have

$$\begin{aligned} \|\pi(\mathcal{E}f)\| &= \sup_{t \in \sigma(\pi(\chi_1))} |\langle \mathcal{E}f, \phi_t \rangle| = \sup_{t \in \sigma(\pi(\chi_1))} |\langle f, \phi_t \rangle| \\ &= \sup_{t \in \sigma(\pi(\chi_1))} |\langle \pi_t(f)\zeta, \zeta \rangle| \\ &\leq \sup_{t \in \sigma(\pi(\chi_1))} \|\pi_t(f)\| \leq \|\pi(f)\| \end{aligned}$$

(the second inequality holds because ϕ_t are radial while the last inequality follows from the assumptions).

The next theorem is already known for the regular (see [11], [5]) and the universal representation (see [8], Lemma B).

THEOREM 4. *Let π be one of the representations of the free group \mathbb{F}_r : regular, universal or spherical unitary representation. Then the projection \mathcal{E} onto radial functions can be extended from $l^1(\mathbb{F}_r)$ to the C^* -algebra C_π^* associated with π .*

Proof. It suffices to check that any of the mentioned representations satisfies the hypothesis of Proposition 2. Clearly it is valid for the regular and universal representations. By Powers' theorem it also holds for the principal series. Finally the case of complementary series follows from Proposition 1 and Theorem 2.

EXAMPLE. Theorem 3 does not hold for any unitary representation of \mathbb{F}_r which lifts to a faithful representation of $l^1(\mathbb{F}_r)$. In fact, let $\mathbb{F}_r = gp\{x_1, x_2, \dots, x_r\}$ and let π be a one-dimensional representation given by $\pi(x_1) = -1$, $\pi(x_2) = \dots = \pi(x_r) = 1$. Put $\Pi = \pi \oplus \lambda$ and let $f = \delta_{x_1} + \delta_{x_2} + \delta_{x_1^{-1}} + \delta_{x_2^{-1}}$. Thus $\Pi(f) = 0 \oplus \lambda(f)$ while

$$\Pi(\mathcal{E}f) = \frac{4(r-2)}{r} \oplus \frac{2}{r}\lambda(\chi_1).$$

Therefore by [1] (Theorem IV J) we have

$$\|\Pi(\mathcal{E}f)\| = \max \left\{ \frac{4(r-2)}{r}, \frac{4\sqrt{2r-1}}{r} \right\} \rightarrow 4, \quad \text{when } r \rightarrow \infty$$

and

$$\|\pi(f)\| = 2\sqrt{3} < 4.$$

REFERENCES

- [1] C. A. Akemann and P. A. Ostrand, *Computing norms in group C^* -algebras*, Amer. J. Math., **98** (1976), 1015-1047.
- [2] W. Arveson, *Notes on extensions of C^* -algebras*, Duke Math. J., **44** (1976), 329-355.
- [3] P. Cartier, *Fonctions harmoniques sur un arbre*, Symposia Math., **9** (1972), 203-270.
- [4] J. M. Cohen, *Operator norms on free groups*, Boll. Un. Math. Ital., **1-B** (1982), 1055-1065.
- [5] J. M. Cohen, and L. De Michele, *The radial Fourier-Stieltjes algebra of free groups*, Contemporary Math., **10** (1982), 33-40.
- [6] R. G. Douglas, *C^* -algebra extensions and K -homology*, Annals of Math. Studies 95, Princeton Univ. Press 1980.
- [7] A. Figà-Talamanca, and M. A. Picardello, *Spherical functions and harmonic analysis on free groups*, J. Funct. Anal., **47** (1982), 281-304.
- [8] U. Haagerup, *$M_0A(G)$ -functions which are not coefficients of uniformly bounded representations*, manuscript 1985.
- [9] A. M. Mantero and A. Zappa, *The Poisson transform on free groups and uniformly bounded representations*, J. Funct. Anal., **51** (1983), 372-400.
- [10] R. T. Powers, *Simplicity of the C^* -algebra associated with the free group on two generators*, Duke Math. J., **42** (1975), 151-156.
- [11] T. Pytlik, *Radial functions on free group and a decomposition of the regular representation into irreducible components*, J. Reine Angew. Math., **326** (1981), 124-135.
- [12] T. Pytlik, and R. Szwarc, *An analytic family of uniformly bounded representations of free groups*, Acta Math., **157** (1986), 287-309.
- [13] R. Szwarc, *An analytic series of irreducible representations of the free group*, Ann. Inst. Fourier (Grenoble), **38** (1988), 87-110.

Received June 8, 1988 and, in revised form November 5, 1988. This paper was written while the author held a C.N.R. fellowship at the University of Rome "La Sapienza".

WROCLAW UNIVERSITY
PL. GRUNWALDZKI 2/4
PL 50-384 WROCLAW, POLAND

AND

ISTITUTO "CASTELNUOVO"
UNIVERSITÀ "LA SAPIENZA"
PIAZZALE A. MORO 2
I-00185 ROMA, ITALY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Walter Bergweiler , On the fix-points of composite functions	1
Aldo Biancofiore, Maria Lucia Fania and Antonio Lanteri , Polarized surfaces with hyperelliptic sections	9
Ciprian Borcea , Deforming varieties of k -planes of projective complete intersections	25
Morton Brown , Fixed points for orientation preserving homeomorphisms of the plane which interchange two points	37
Hao Zhi Chuan , Note on the inequality of the arithmetic and geometric means	43
Paul Henry Edelman and Dennis E. White , Codes, transforms and the spectrum of the symmetric group	47
Pei-Kee Lin , The isometries of $H^\infty(E)$	69
James J. Moloney , Residue class domains of the ring of convergent sequences and of $C^\infty([0, 1], \mathbf{R})$	79
Zhong-Jin Ruan , The structure of pure completely bounded and completely positive multilinear operators	155
Wolfgang Ruess and William H. Summers , Weakly almost periodic semigroups of operators	175
Gideon Schwarz , A pretender to the title “canonical Moebius strip”	195
Ryszard Szwarc , Banach algebras associated with spherical representations of the free group	201