

Pacific Journal of Mathematics

SIMPLE PERIODIC MODULES OF TWISTED CHEVALLEY GROUPS

PETER FLEISCHMANN AND JENS CARSTEN JANTZEN

SIMPLE PERIODIC MODULES OF TWISTED CHEVALLEY GROUPS

PETER FLEISCHMANN AND JENS CARSTEN JANTZEN

Consider a finite twisted Chevalley group constructed over a field of prime characteristic and its representations over an algebraically closed field of the same characteristic. In this paper we classify all those irreducible representations that are periodic, i.e., that have a periodic projective resolution. There is always the Steinberg module that is both simple and projective. We show that there are further periodic simple modules only for groups of types 2A_2 and 2B_2 .

0. Introduction. Let K be an algebraically closed field of characteristic $p > 0$ and Γ a finite group with $p \nmid |\Gamma|$. A finite dimensional $K\Gamma$ -module M is called periodic, if it has a periodic minimal projective resolution. Of course projective modules are also periodic, but the non-projective ones are characterized by their complexity $c_\Gamma(M)$ being one. Here we understand complexity in the sense of Alperin [1].

In this article we look at the case where Γ is an “almost simple” twisted group of Lie type, defined over a finite field of the same characteristic p . We will give a classification of all periodic simple modules for these groups and the finite simple groups related to them.

It is a well-known fact that all finite groups of Lie type have a unique simple projective module, the Steinberg module St . We will prove in this paper that, if Γ is not of type 2A_2 or 2B_2 , then the only periodic simple $K\Gamma$ -module is St . With the classification for 2A_2 in [6], [7], and for 2B_2 in part 3 of this paper, we will achieve a full classification of simple periodic modules for twisted groups.

The corresponding classification for non-twisted Chevalley groups was recently accomplished in [9], [10]. Together with the initial classification for the type A_1 in [15], the final result is that for finite groups of Lie type there are simple periodic modules, other than St , only for groups of type A_1 , 2A_2 and 2B_2 .

To state our results more precisely, we have to introduce some notations which we take over from [5]. For conceptual reasons we prefer working with universal groups:

So let K be an algebraic closure of the prime field \mathbb{F}_p , let G be an almost simple, simply connected and connected affine algebraic

group over K , defined and split over \mathbb{F}_p . Let F be a Frobenius endomorphism of G , as in [5], T an F -stable maximal torus of G with character group $X(T)$ and $R \subset X(T)$ be the (indecomposable) root system of G . We choose a basis $\Pi = \{\alpha_1, \alpha_2, \dots, \alpha_l\}$ of R and assume F to induce a non-trivial automorphism τ of the Dynkin diagram of G . So either τ is of order 3 and R of type D_4 or τ is of order 2 and R of type A_l ($l > 1$), D_l ($l > 3$), E_6 , or B_2 , G_2 , F_4 (in which cases p must be 2, 3, 2 respectively).

Attached to F are natural numbers $i, e \in \mathbb{N}$ such that $F^i(t) = t^{p^e}$ for all $t \in T$ and the number $q := p^{e/i}$ is uniquely determined by G and F . The group G^F of F -fixed points is either a finite (universal) Steinberg group (usually denoted by ${}^3D_4(q^3)_{\text{s.c.}}$, ${}^2A_l(q^2)_{\text{s.c.}}$, ${}^2D_l(q^2)_{\text{s.c.}}$ or ${}^2E_6(q^2)_{\text{s.c.}}$) or a Ree or Suzuki group (usually denoted by ${}^2B_2(q^2)$, ${}^2F_4(q^2)$, ${}^2G_2(q^2)$ where $q^2 = 2^{2m+1}$ resp. $q^2 = 3^{2m+1}$ for some m).

Excluding the cases (a) ${}^2A_2(4)_{\text{s.c.}}$, (b) ${}^2B_2(2)_{\text{s.c.}}$, (c) ${}^2G_2(3)_{\text{s.c.}}$ and (d) ${}^2F_4(2)_{\text{s.c.}}$, their quotient by the center is simple (see [19], Theorem 34, pg. 188) and is usually denoted by ${}^3D_4(q^3)$, ${}^2A_l(q^2)$ etc. (or by ${}^3D_4(q)$, ${}^2A_l(q)$, according to the taste of the author.)

The groups in (a) and (b) are solvable, whereas in (c) and (d) their commutator subgroups have index 3 and 2 and are simple non-abelian finite groups ($\cong \text{PSL}_2(8)$ in (c), \cong Tits' simple group in (d)) ([19], pg. 188).

In this framework the main theorem of this article reads as follows:

0.1. THEOREM. *Let Γ be G^F , $G^F/Z(G^F)$ or $(G^F)'$, then all periodic simple $K\Gamma$ -modules are projective if and only if R is not of type A_2 or B_2 , in which case they are isomorphic to $\text{St}|_{G^F}$ or to a Clifford component of it, if $(G^F)' < G^F$.*

In §1 we will reduce the problem for the Steinberg groups to the case $G^F \cong {}^2A_3(q^2)_{\text{s.c.}} \cong \text{SU}_4(q^2)$, which will be dealt with in §2. Factoring out the center, which does not affect the Sylow- p -structure, leads to the result for the simple groups. Section 3 is devoted to the Ree and Suzuki groups.

1. The reduction to SU_4 . In this section we exclude the cases B_2 , G_2 and F_4 but keep the previous notations. So all roots in R have the same length. We assume moreover T to be defined and split over \mathbb{F}_p . So the usual Frobenius endomorphism $x \mapsto x^p$ on K gives rise to an endomorphism F_0 of G , acting on T via $F_0(t) = t^p$ for

all $t \in T$. There is an automorphism of G that has the same order as τ , commutes with F , stabilizes T and induces on $\Pi \subseteq X(T)$ the same map as τ . If this automorphism is also denoted by τ , then we can write $F = \tau \circ F_0^n = F_0^n \circ \tau$ for some integer $n > 0$; in this case $q = p^n$. We denote the fundamental dominant weights of R by λ_α and set $\rho := \sum \lambda_\alpha$ where the sum is over all simple roots. Let $X(T)^+$ be the set of dominant weights and set for all $n > 0$:

$$(1.1) \quad X_n(T) := \left\{ \lambda = \sum r_\alpha \lambda_\alpha \in X(T) \mid 0 \leq r_\alpha < p^n \text{ for all } \alpha \in \Pi \right\}.$$

For any $\lambda \in X(T)^+$ let $V(\lambda)$ be the Weyl module for G with highest weight λ and $L(\lambda)$ its unique simple quotient. One has $V((p^n - 1)\rho) = L((p^n - 1)\rho)$ for all n ; this module is often called the n th Steinberg module and denoted by St_n .

For any G -module M and any $r \in \mathbb{N}$ let $M^{(r)}$ denote the G -module got from M by twisting the G -action with F_0^r . If $\mu = \sum p^i \mu_i$ where i ranges from 0 to $n - 1$ and $\mu_i \in X_1(T)$, then

$$(1.2) \quad L(\mu) \cong L(\mu_0) \otimes L(\mu_1)^{(1)} \otimes \cdots \otimes L(\mu_{n-1})^{(n-1)}$$

by Steinberg's tensor product theorem [18].

According to another theorem of Steinberg [18] the $L(\lambda)$ with $\lambda \in X_n(T)$ remain irreducible when regarded as a representation of G^F , and each simple KG^F -module is isomorphic to exactly one $L(\lambda)$ with $\lambda \in X_n(T)$.

The n th Steinberg module $\text{St}_n = L((p^n - 1)\rho)$ is known to be projective as a KG^F -module. So in the case of the Steinberg groups our main theorem amounts to:

(1.3) **THEOREM.** *Suppose that G is a Steinberg group not of type A_2 . Then $L(\lambda)$ is not a periodic KG^F -module for any $\lambda \in X_n(T)$, $\lambda \neq (p^n - 1)\rho$.*

Using the corresponding results for non-twisted groups of type A_l , $l > 1$ (proved in [9]) we reduce the proof of this theorem to the case of 2A_3 (i.e. to the groups $\text{SU}_4(q^2)$) that will be dealt with in §2.

(1.4) **PROPOSITION.** *If Theorem 1.3 holds in the case 2A_3 , then it holds in all cases.*

Proof. Suppose that $L(\lambda)$ is periodic. Then for any subgroup H of G^F any direct summand M of $L(\lambda)$ as a KH -module is periodic. So we prove the proposition by constructing in each case some H and M such that M is not periodic for KH .

Here and in the following we will abbreviate a weight $\lambda = \sum r_i \lambda_{\alpha_i}$ by (r_1, r_2, \dots, r_l) , where $\Pi = \{\alpha_1, \dots, \alpha_l\}$.

Case 1. Type $A_l, l > 3$. Here the Dynkin diagram is:

$$\begin{array}{ccccccc} \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \cdots & \text{---} & \circ & \text{---} & \circ \\ \alpha_1 & & \alpha_2 & & \alpha_3 & & & & \alpha_{l-1} & & \alpha_l \end{array}$$

We consider $G^F \cong \mathrm{SU}_{l+1}(q^2)$ with respect to the hermitian form given by the unit matrix.

Then $c_1: g \mapsto \begin{pmatrix} g & 0 \\ 0 & 1 \end{pmatrix}$ and $c_2: g \mapsto \begin{pmatrix} 1 & 0 \\ 0 & g \end{pmatrix}$ define two canonical embeddings of $\mathrm{SU}_l(q^2) \rightarrow \mathrm{SU}_{l+1}(q^2)$. If L_J denotes the Levi subgroup for $J \subseteq \Pi$ in G , then the image $\mathrm{Im} c_1$ is contained in the derived group $L'_{\Pi \setminus \{\alpha_l\}} := L_1$ and $\mathrm{Im} c_2 \subseteq L'_{\Pi \setminus \{\alpha_1\}} := L_2$; hence we can apply a theorem of Smith [17]:

If $\lambda = (r_1, \dots, r_l) \in X_n(T)$ then $L(\lambda)|_{L_i} = L(\lambda_i) \oplus M_i$ where $L(\lambda_i)$ is an irreducible module for L_i with highest weight λ_i and M_i is some other L_i module. We get $\lambda_1 = (r_1, r_2, \dots, r_{l-1})$ and $\lambda_2 = (r_2, r_3, \dots, r_l)$. See also [16] for a proof of these facts.

Furthermore L_i is simply connected of type A_{l-1} . If $\lambda \neq (p^n - 1)\rho$, then we may assume without loss of generality that $\lambda_1 \neq (p^n - 1)(\rho - \lambda_{\alpha_l})$. Then $L(\lambda_1)$ is a simple periodic and non-projective $\mathrm{SU}_l(q^2)$ module in contradiction to the induction hypothesis.

Case 2. Type ${}^2D_l(q^2)_{\mathrm{s.c.}}$

$$\begin{array}{ccccccc} & & & & & & \circ \alpha_l \\ & & & & & & / \\ \circ & \text{---} & \circ & \text{---} & \circ & \cdots & \circ & \text{---} & \circ & & \\ \alpha_1 & & \alpha_2 & & & & & & \alpha_{l-2} & & \\ & & & & & & & & \backslash \\ & & & & & & & & \circ \alpha_{l-1} \end{array}$$

Let L_i be the derived group of the Levi subgroup corresponding to J_i , where $J_i = \{\alpha_i, \alpha_{i+1}\}$ for $i < l - 2$ and $J_{l-2} = \{\alpha_{l-2}, \alpha_{l-1}, \alpha_l\}$. Each L_i is F -stable and simply connected; one has $L_i^F \cong A_2(q)_{\mathrm{s.c.}}$ for $i < l - 2$ and $(L_{l-2})^F \cong {}^2A_3(q^2)_{\mathrm{s.c.}}$. If again $\lambda = (r_1, r_2, \dots, r_l)$, then by Smith's theorem $L(\lambda)|_{L_i^F} = L(\lambda_i) \oplus M_i$ as above with $\lambda_i = (r_i, r_{i+1})$ for $1 \leq i \leq l - 3$, and $\lambda_{l-2} = (r_{l-2}, r_{l-1}, r_l)$. Again, since $L(\lambda) \neq \mathrm{St}_n$, one of these $L(\lambda_i)$'s must be non-projective and hence non-periodic for L_i^F , by the corresponding result for $A_2(q)$ in [9] and the hypothesis.

Case 3. Type ${}^2E_6(q^2)_{\mathrm{s.c.}}$:

$$\begin{array}{ccccccc} & & & & \alpha_3 & & \\ & & & & | & & \\ \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ & \text{---} & \circ \\ \alpha_1 & & \alpha_2 & & & & \alpha_4 & & \alpha_5 \\ & & & & & & & & \\ & & & & & & & & \circ \\ & & & & & & & & \alpha_6 \end{array}$$

The same kind of argument applies here with $J_1 = \Pi \setminus \{\alpha_6\}$, $J_2 = \{\alpha_3, \alpha_6\}$ and $(L_{J_1})^F \cong {}^2A_5(q^2)$, $(L_{J_2})^F \cong A_2(q)$, using Case 1 and the result in [9].

Case 4. Type ${}^3D_4(q^3)_{\text{s.c.}}$:

$$\begin{array}{c}
 \circ \alpha_4 \\
 \diagup \\
 \circ \alpha_2 \\
 \diagdown \\
 \circ \alpha_3
 \end{array}
 \begin{array}{c}
 \circ \alpha_1 \\
 \text{---} \circ \alpha_2
 \end{array}, \quad R = \{\pm(\varepsilon_i \pm \varepsilon_j), i \neq j\}.$$

The Weyl group $W(G, T)$ acts on \mathbb{Z}^4 as permutations of the $\pm\varepsilon_i$'s with an even number of sign changes.

If $\alpha_1 := \varepsilon_1 - \varepsilon_2$, $\alpha_2 := \varepsilon_2 - \varepsilon_3$, $\alpha_3 := \varepsilon_3 - \varepsilon_4$, $\alpha_4 := \varepsilon_3 + \varepsilon_4$, then the τ -fixed points $\alpha_2, \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 = \varepsilon_1 + \varepsilon_3 =: \beta$, $\alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = \varepsilon_1 + \varepsilon_2$, form an A_2 -subsystem of R . Moreover W maps

$$(\alpha_2, \beta) \mapsto (\alpha_2, \alpha_1) \quad \text{under} \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & -3 & -2 & 4 \end{pmatrix}$$

$$(\alpha_1, \alpha_2) \mapsto (\alpha_2, \alpha_3) \quad \text{under} \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 4 & 1 \end{pmatrix}$$

$$(\alpha_1, \alpha_2) \mapsto (\alpha_2, \alpha_4) \quad \text{under} \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & -4 & -1 \end{pmatrix}$$

If U_α denotes the root subgroup of G , then U_{α_2}, U_β are F -stable and $U_{\alpha_2}^F, U_\beta^F \cong \mathbb{F}_q^+$. Let H be the subgroup $\langle U_{\pm\alpha_2}, U_{\pm\beta} \rangle \leq G$; then H is of type A_2 and H^F of type $A_2(q)$. For $i = 1, 3, 4$ let $J_i = \{\alpha_i, \alpha_2\}$; hence $L'_{J_i} =: L_i$ is of type A_2 but not F -stable. Let now $L(\lambda) \neq \text{St}_n$, with $\lambda = (r_1, r_2, r_3, r_4)$.

Then $L(\lambda)|_{L_i} = L(\lambda_i) \oplus M_i$, where $\lambda_i = (r_2, r_1)$ resp. (r_1, r_2) .

For each i there is some $w_i \in W$ such that $w_i(H) = w_i H w_i^{-1}$ is contained in L_i . Since $L(\lambda)|_{H^F}$ is periodic, also the restrictions $L(\lambda)|_{w_i(H^F)}$ must be periodic. This can be seen for instance looking at the rank variety in the sense of [3], [4] and the fact that rank varieties of conjugate groups are isomorphic. Since $w_i(H)$ is contained in L_i , also $L(\lambda_i)|_{w_i(H^F)}$ is periodic. But $w_i(H^F) \cong A_2(q)$ and $r_i < q$, so $L(\lambda_i)$ is also irreducible for $w_i(H^F)$, which forces r_i and r_2 to be $q - 1$ by the result in [9]. This gives a contradiction. \square

2. The Case SU_4 . Keep all assumptions and notations from §1 and assume that R is of type A_3 .

(2.1) LEMMA. Let $\mu = \sum_{l=0}^{n-1} p^l \mu_l \in X_n(T)$ with all $\mu_l \in X_1(T)$. If $L(\mu)$ is a periodic KG^F -module, then each μ_l has one of the following forms :

- (a) $(p-1, p-1, p-1) = (p-1)p$,
- (b) $(p-1, p-1, i)$ with $0 \leq i < p-1$,
- (b') $(i, p-1, p-1)$ with $0 \leq i < p-1$,
- (c) $(p-1, i, j)$ with $0 \leq i, j, i+j = p-2$,
- (c') $(i, j, p-1)$ with $0 \leq i, j, i+j = p-2$,
- (d) $(p-1, i, p-1)$ with $0 \leq i < p-1$,
- (e) $(i, p-1, j)$ with $0 \leq i, j < p-1$,
- (f) $(p-2-i, i, p-2-i)$ with $0 \leq i < p-1$.

Proof. Suppose $\mu = (m_1, m_2, m_3)$. The same argument as in the proof of (1.4), Case 1 shows that $L((m_1, m_2))$ and $L((m_2, m_3))$ are periodic as $KSU_3(q^2)$ -modules. The classification of these modules in [7] yields now the claim. \square

(2.2) LEMMA. Let $\lambda \in X_1(T)$ be of type (d), (e), (f) in Lemma (2.1). Then $\dim L(\lambda)$ is not divisible by p^3 .

Proof. The formulae for the characters of the $L(\lambda)$ in [12] §7 imply:

$$\begin{aligned} \dim L(p-1, i, p-1) &= \dim V(p-1, i, p-1) \\ &\quad - \dim V(p-(i+2), i, p-(i+2)) \\ &\quad \text{for } 0 \leq i < p-1; \end{aligned}$$

$$\begin{aligned} \dim L(i, p-1, j) &= \dim V(i, p-1, j) \\ &\quad - \dim V(p-(j+2), p-1, p-(i+2)) \\ &\quad + \dim V(p-(i+2), i+j+1-p, p-(j+2)) \\ &\quad \text{for } 0 \leq i, j < p-1, i+j > p-2; \end{aligned}$$

$$\dim L(i, p-1, j) = \dim V(i, p-1, j) \quad \text{for } 0 \leq i, j, i+j = p-2;$$

$$\begin{aligned} \dim L(i, p-1, j) &= \dim V(i, p-1, j) \\ &\quad - \dim V(j, p-(i+j+3), i) \\ &\quad \text{for } 0 \leq i, j, i+j < p-2; \end{aligned}$$

$$\begin{aligned} \dim L(p-2-i, i, p-2-i) &= \dim V(p-2-i, i, p-2-i) \\ &\quad \text{for } 0 \leq i < p-1. \end{aligned}$$

By Weyl's dimension formula

$$\dim V(r, s, t) = (r+1)(s+1)(t+1)(r+s+2)(s+t+2)(r+s+t+3)/12.$$

In each case one sees immediately that p^3 does not divide $\dim V(\mu)$ for any μ . In the case $\lambda = (p-1, i, p-1)$ one gets

$$\dim L(\lambda) \equiv 2 \dim V(\lambda) \not\equiv 0 \pmod{p^3} \quad \text{for } p \geq 5,$$

$$\dim L(\lambda) \equiv 2 \dim V(\lambda) \not\equiv 0 \pmod{p^2} \quad \text{for } p = 3,$$

$$\dim L(\lambda) = 14 \not\equiv 0 \pmod{8} \quad \text{for } p = 2.$$

The other cases are similar and are left to the reader. \square

Recall the definition of the rank variety $V_E M$ of a KE -module M for any elementary abelian p -group E as in [3] or [4]. So it is a homogeneous subvariety of the vector space K^m where m is the rank of E . It is called linear, if it is a linear subspace of K^m .

(2.3) LEMMA. *Let E be an elementary abelian p -subgroup of $G^F = \mathrm{SU}_4(q^2)$ and let $\mu \in X_1(T)$ be of type (b) or (c) in Lemma (2.1). Then*

$$V_E L(\mu) = V_E L((p-1, 0, p-2)).$$

Proof. Set $C^* = \{(r_1, r_2, r_3) \in X(T) \mid -1 \leq r_1, r_2, r_3, r_1 + r_2 + r_3 \leq p-3\}$. Then C^* is a fundamental domain for the "dot" operation (i.e., $w \cdot \mu = w(\mu + \rho) - \rho$) of W_p on X , where W_p is the affine Weyl group (as in [14], II, 6.1). So there is a unique $\mu' \in W_p \cdot \mu \cap C^*$ and a unique $\lambda' \in W_p \cdot \lambda \cap C^*$ where $\lambda = (p-1, 0, p-2)$; in fact $\lambda' = (-1, -1, 0)$.

We want to use the translation functors $T_{\lambda'}^{\mu'}$ and $T_{\mu'}^{\lambda'}$, cf. [14], II, 7.6. More precisely, we shall show that $T_{\lambda'}^{\mu'} L(\lambda) \cong L(\mu)$ and $T_{\mu'}^{\lambda'} L(\mu) \cong L(\lambda)$. As $T_{\lambda'}^{\mu'} L(\lambda)$ is a direct summand of $L(\lambda) \otimes M$ for a suitable finite dimensional G -module M , and $T_{\mu'}^{\lambda'} L(\mu)$ a direct summand of $L(\mu) \otimes M^*$, elementary properties of the rank variety (cf. [3]) yield $V_E L(\lambda) = V_E L(\mu)$.

If μ is of type (c) in Lemma (2.1), then λ and μ belong to the same facet for W_p . In this case $T_{\lambda'}^{\mu'} L(\lambda) \cong L(\mu)$ and $T_{\mu'}^{\lambda'} L(\mu) \cong L(\lambda)$ are special cases of [14], II, 7.15.

Suppose now that μ is of type b) in Lemma (2.1), i.e. $\mu = (p-1, p-1, i)$ for some i , $0 \leq i < p-1$. Then $\mu' = (-1, i, p-(i+2))$. One has $V(\lambda) = L(\lambda)$ and $V(\mu) = L(\mu)$ by [11], p. 120 or by [12].

We can compute the formal characters of $T_{\lambda'}^{\mu'} L(\lambda)$ and $T_{\mu'}^{\lambda'} L(\mu)$ using [14], II, 7.8. Let $s_i = s_{\alpha_i}$ be the reflection with respect to α_i (for $i = 1, 2, 3$) and let $s \in W_p$ be the affine reflection with $s(r, s, t) = (r+s+t-p, p-t, p-s)$ for all $(r, s, t) \in X(T)$. Then the stabilizer of μ' (under the dot action) in W_p consists of $\{1, s_1, s, s_1s, ss_1, s_1ss_1\}$, that of λ' of $\{1, s_1, s_2, s_1s_2, s_2s_1, s_1s_2s_1\}$. The intersection of these groups is $\{1, s_1\}$, a system of representatives modulo this intersection is $\{1, s_2, s_1s_2\}$ in $\text{Stab}(\lambda')$ and $\{1, s, s_1s\}$ in $\text{Stab}(\mu')$.

The element $w \in W_p$ with $w(r, s, t) = (p-r, r+s+t, p-t)$ has the property that $\mu = w \cdot \mu'$ and $\lambda = w \cdot \lambda'$. One has

$$\begin{aligned} ws_2 \cdot \mu' &= (p - (i+2), p-1, -1), \\ ws_1s_2 \cdot \mu' &= (p+i, p-(i+2), -1). \end{aligned}$$

These weights are not dominant, but become dominant after adding $\rho = (1, 1, 1)$. Therefore they contribute 0 to the sum in [14], II, 7.12, and we get $T_{\lambda'}^{\mu'} L(\lambda) = T_{\lambda'}^{\mu'} V(\lambda) \cong V(\mu) \cong L(\mu)$. Furthermore

$$\begin{aligned} ws \cdot \lambda' &= (2p-2, p-1, -1), \\ ws_1s \cdot \lambda' &= (0, 2p-2, -1). \end{aligned}$$

The same argument as above implies

$$T_{\mu'}^{\lambda'} L(\mu) = T_{\mu'}^{\lambda'} V(\mu) \cong V(\lambda) = L(\lambda). \quad \square$$

For $\alpha \in R$, let $U_{\alpha} = \{x_{\alpha}(t) | t \in K\}$ be the root subgroup of G with root homomorphism $x_{\alpha}: K \mapsto G$. We set:

$$H := U_{\alpha_1} \circ U_{\alpha_1+\alpha_2} \circ U_{\alpha_2+\alpha_3} \circ U_{\alpha_1+\alpha_2+\alpha_3} \cong (K^+)^4.$$

If $\{u_1, u_2, \dots, u_n\}$, $\{v_1, v_2, \dots, v_{2n}\}$ denote bases of \mathbb{F}_q and \mathbb{F}_{q^2} over \mathbb{F}_p respectively, then let $x_1(u_i) := x_{\alpha_2}(u_i)$,

$$x_2(v_j) := x_{\alpha_1+\alpha_2}(v_j)x_{\alpha_2+\alpha_3}(v_j^q), \quad x_3(u_k) := x_{\alpha_1+\alpha_2+\alpha_3}(u_k).$$

The group

$$E := H^F = \langle x_1(u_i)x_2(v_j)x_3(u_k) | 1 \leq i, k \leq n; 1 \leq j \leq 2n \rangle$$

is then an elementary abelian p -subgroup of rank $4n$ of $G^F = \text{SU}_4(q^2)$.

(2.4) LEMMA. *Let $\lambda \in X_1(T)$ be one of the weights in Lemma (2.1). Then*

$$\begin{aligned} \dim V_E L(\lambda) &\geq 4n - 2 \quad \text{for } \lambda \neq (p-1)\rho, \\ \dim V_E L((p-1)\rho) &\geq 4n - 4. \end{aligned}$$

Proof. Let $d := \dim V_E L(\lambda)$; then by [3], Proposition 5.1, there exists a “shifted subgroup” $E^\wedge \leq KE^*$, which is elementary abelian of order p^{4n-d} such that $L(\lambda)|_{E^\wedge}$ is free. In particular p^{4n-d} divides $\dim L(\lambda)$, which proves the claim for the weights of type (d), (e) and (f) by (2.2).

By (2.3) we are left with the weights $(p-1, 0, p-2)$ and $(p-1)\rho$; in both cases $L(\lambda) = V(\lambda)$. Now let $J := \{\alpha_1, \alpha_2\} \subseteq \Pi$; then H can also be described as the unipotent radical U_J of the standard parabolic subgroup $P_J = U_J \rtimes L_J \leq G$. If $\lambda = (r, s, t)$, then by Smith’s theorem $L(\lambda)|_{L_J} = L(\lambda)^{U_J} \oplus X$, where $L(\lambda)^{U_J}$ is the space of U_J -fixed points, which is also an irreducible module for $L'_J \cong A_1(K)_{\text{s.c.}} \times A_1(K)_{\text{s.c.}}$, with highest weight (r, t) , hence has dimension $(r+1)(t+1)$.

Since $E \subseteq U_J$ and E^\wedge is generated by elements of the form $1 + \sum a_i(x_i - 1)$ with $a_i \in K$, $x_i \in E$, it is clear that $L(\lambda)^{U_J} \subseteq L(\lambda)^{E^\wedge}$, which has dimension $\dim L(\lambda)/p^{4n-d}$ because $L(\lambda)$ is E^\wedge -free. So we get $p^{4n-d} \leq \dim L(\lambda)/(r+1)(t+1)$. Now we apply Weyl’s dimension formula and obtain $p^{4n-d} \leq p^2(p+1)/6$ for $\lambda = (p-1, 0, p-2)$ and $p^{4n-d} \leq p^4$ for $\lambda = (p-1)\rho$. \square

(2.5) PROPOSITION. *If M is a periodic simple $KSU_4(q^2)$ -module, then*

$$M \cong \text{St}_n.$$

Proof. By [3] 5.6, $V_E(X \otimes Y) = V_E(X) \cap V_E(Y)$ for KE -modules X and Y . It is easy to see that $\dim V_E(L(\mu)^{(n)}) = \dim V_E(L(\mu))$, so if $M (\not\cong \text{St}_n)$ is written in the form of (1.2), then at least for one μ_i we must have $\dim V_E(L(\mu_i)^{(i)}) \geq 4n - 2$ by (2.4). Since all rank varieties V_E are homogeneous affine subvarieties of K^{4n} the intersection formula for homogeneous varieties implies

$$\begin{aligned} \dim V_E M &= \dim \bigcap_{i=0}^{n-1} V_E(L(\mu_i)^{(i)}) \\ &\geq (n-1)(4n-4) + 4n - 2 - (n-1)4n \geq 2, \end{aligned}$$

which contradicts periodicity by [3], 7.6 and 8.1. \square

REMARK. This proposition together with (1.4) implies Theorem (1.3), and it is clear, since $G^F = (G^F)'$ if type (a) of the exceptions is excluded, how to get (0.1).

3. The Ree and Suzuki groups. In this section we deal with the cases, where R is of type G_2 , F_4 and B_2 , which were excluded earlier. Here twisted groups of Lie type can exist only if the characteristic of K is 3, 2 and 2 respectively, and τ always has order 2, interchanging the long and short roots.

PROPOSITION (3.1). *Let R be G_2 or F_4 , $\text{char } K = 3$ or 2 respectively and $\Gamma = G^F$, $G^F/Z(G^F)$, or $(G^F)'$; then every periodic simple $K\Gamma$ -module is projective, and either isomorphic to St_{G^F} or to a Clifford component of $\text{St}_{G^F}|_\Gamma$ if $\Gamma = (G^F)' \triangleleft G^F$ and $\Gamma \neq G^F$.*

Proof. We first prove this for $\Gamma = G^F$. By a result of Steinberg [18], §12, an analogue of (1.2) applies here, with $X_1(T)$ replaced by $X_1(T)' =: X'_1 = \{\mu = \sum r_\alpha \lambda_\alpha \in X_1(T) | \alpha \in \Pi \text{ and } r_\alpha = 0 \text{ if } \alpha \text{ is a long root}\}$.

Case 1. $G^F = {}^2G_2(q^2)_{\text{s.c.}}$ with $q^2 = 3^n$, $n = 2m + 1$ and $\text{char } K = 3$, $R = \{\pm\alpha, \pm(2\alpha + 3\beta), \pm(\alpha + 3\beta) \text{ (long roots)}, \pm\beta, \pm(\alpha + \beta), \pm(\alpha + 2\beta) \text{ (short roots)}\}$, and the Dynkin diagram is:

$$\alpha \Longleftrightarrow \beta$$

Here $X_1(T)' = \{(0, 0), (0, 1), (0, 2)\}$ and $\dim L(\lambda) = 1, 7, 27$ for $\lambda \in X_1(T)'$ respectively. See [18], §12.

Let $E = \{x(u, v) | u, v \in \mathbb{F}_{3^n}\}$ with

$$x(u, v) := x_{\alpha+3\beta}(u)x_{\alpha+\beta}(u^{3^m})x_{2\alpha+3\beta}(v)x_{\alpha+2\beta}(v^{3^m}).$$

Then E is elementary abelian of rank $2n$. With regard to the dimensions it is enough to look at $\lambda = (0, 2)$. Then $L(\lambda)$ has the following “monomial basis” with highest weight vector v_λ^+ :

$$\{x_{-(\alpha+\beta)}^{n_1} X_{-(\alpha+2\beta)}^{n_2} X_{-\beta}^{n_3} \circ v_\lambda^+ | 0 \leq n_i \leq 2\},$$

where $X_{-\gamma}$ are root vectors in $\text{Lie}(G)$.

If γ is any positive root, $x_\gamma: K \rightarrow G$ the corresponding root homomorphism and V_μ a weight subspace to the weight μ of some G module V , then one knows:

$$x_\gamma(t) \circ v_\mu \in \bigoplus_{i \in \mathbb{N}} V_{\mu+i\gamma}.$$

Since λ is the highest weight of $L(\lambda)$ and E consists of unipotent elements, E must fix the three basis vectors v_λ^+ , $X_{-\beta}v_\lambda^+$, $X_{-\beta}^2v_\lambda^+$. If $d := \dim V_E(L(0, 2))$, then as in the proof of (2.4), a suitable shifted elementary abelian $E^\wedge \leq (KE)^*$ with $|E^\wedge| = 3^{2n-d}$ acts freely on $L(0, 2)$ and we get $\dim L(0, 2)/|E^\wedge| = 3^{d-2n+3} \geq 3$ and hence $d \geq 2n - 2$. By elementary properties of rank varieties, $V_E(L(\lambda)) \cong K^{2n}$ for $\lambda = (0, 0)$ or $(0, 1)$. The analogue of (1.2) together with the intersection theorem for affine varieties yields $\dim V_E(M) \geq 2$ for all $M \neq \text{St}_{G^F}$.

Case 2. $G^F = {}^2F_4(q^2)$, $q^2 = 2^n$, $n = 2m + 1$ and $\text{char } K = 2$. Here $X_1(T)' = \{(0, 0, 0, 0), (0, 0, 0, 1), (0, 0, 1, 0), (0, 0, 1, 1)\}$ with dimensions for $L(\lambda)$: 1, 26, 246, 2^{12} respectively (see [20]). Let $\theta: K \rightarrow K$, $x \mapsto x^{2^m}$; we consider the following elements of R^+ ($\Pi = \{\alpha_1, \alpha_2, \alpha_3, \alpha_4\}$):

$$\begin{aligned} \gamma_1 &:= \alpha_1 + 2\alpha_2 + 3\alpha_3 + 2\alpha_4; & \gamma_2 &:= 2\alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4; \\ \delta_1 &:= \alpha_1 + 2\alpha_2 + 3\alpha_3 + \alpha_4; & \delta_2 &:= \alpha_1 + 3\alpha_2 + 4\alpha_3 + 2\alpha_4; \\ \varepsilon_1 &:= \alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4; & \varepsilon_2 &:= \alpha_1 + 2\alpha_2 + 4\alpha_3 + 2\alpha_4; \\ \varphi_1 &:= \alpha_1 + \alpha_2 + 2\alpha_3 + \alpha_4; & \varphi_2 &:= \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4; \\ \psi_1 &:= \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4; & \psi_2 &:= \alpha_1 + \alpha_2 + 2\alpha_3 + 2\alpha_4. \end{aligned}$$

Notice that $\gamma_1, \delta_1, \dots$ are short and $\gamma_2, \delta_2, \dots$ are long.

Let

$$\begin{aligned} x_1(u_1) &:= x_{\gamma_1}(u_1) \circ x_{\gamma_2}(u_1^{2\theta}); & x_2(u_2) &:= x_{\delta_1}(u_2) \circ x_{\delta_2}(u_2^{2\theta}); \\ x_3(u_3) &:= x_{\varepsilon_1}(u_3) \circ x_{\varepsilon_2}(u_3^\theta); & x_4(u_4) &:= x_{\varphi_1}(u_4) \circ x_{\varphi_2}(u_4^\theta); \\ x_5(u_5) &:= x_{\psi_1}(u_5) \circ x_{\psi_2}(u_5^{2\theta}); & \text{with } u_i &\in \mathbb{F}_{2^n}. \end{aligned}$$

Notice that these elements commute because of $\text{char } K = 2$, and hence the group $E := \langle x_i(u_i) | i = 1, 2, 3, 4, 5; u_i \in \mathbb{F}_{2^n} \rangle$ is an elementary abelian subgroup of G^F of rank $5n$.

For $L(0, 0, 1, 1)$ there is a monomial basis, consisting of elements of the form: $X_{-\omega_1}^{n_1} \circ \dots \circ X_{-\omega_{12}}^{n_{12}} v_\lambda^+$ with $n_i \in \{0, 1\}$, where $\{\omega_1, \omega_2, \dots, \omega_{12}\}$ is the set of short positive roots.

The 7 short roots $\alpha_3, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3, \alpha_4, \alpha_3 + \alpha_4, \alpha_2 + \alpha_3 + \alpha_4, \alpha_2 + 2\alpha_3 + \alpha_4$ have not full support in Π . Since the generators of E are products of x_γ 's where γ is a positive root with full support, an argument similar to that of Case 1 shows that basis vectors involving only monomials in those 7 roots have to be fixed by E and hence by any shifted subgroup E^\wedge of KE^* .

Since there are 2^7 such monomials, $\dim L(0, 0, 1, 1)^{E^\wedge} \geq 2^7$. If $d = \dim V_E(L(0, 0, 1, 1))$ then again we can choose E^\wedge to act freely on $L(0, 0, 1, 1)$ and to be of order 2^{5n-d} . So we get $\dim L(0, 0, 1, 1)/|E^\wedge| = 2^{12-5+d} \geq 2^7$; hence $d \geq 5n - 5$. Since all other $V_E(L(\lambda))$ for $\lambda \in X_1(T)'$ have dimension $\geq 5n - 1$ (4 does not divide $\dim L(\lambda)$ for $\lambda \neq (0, 0, 1, 1)$), the analogue of (1.2) yields:

$$\dim V_E(M) \geq 4 \quad \text{for all } M \neq \text{St}_{G^F}.$$

Proof for $G^F/Z(G^F)$ and $G^{F'}$. As $Z(G^F)$ is a p' -subgroup of G^F the same results hold for $\Gamma = G^F/Z(G^F)$. Now let $G^{F'} \neq G^F$; then $m = 0, n = 1$ and $q^2 = |G^F/G^{F'}| = p = 3$ in Case 1 and $= 2$ in Case 2.

By Clifford's theorem $\text{St}|_{G^{F'}}$ is completely reducible. Let V be a simple direct summand of $\text{St}|_{G^{F'}}$, with inertia group $I := I_{G^F}(V) = \{g \in G^F | V^g \cong V \text{ as } G^{F'} \text{ modules}\}$. Then $I = G^{F'}$. Otherwise by [8] §9, (9.9), $V = W|_{G^{F'}}$ for a simple G^F module W , which is isomorphic to St_{G^F} , since V is projective; but then W had to be a direct summand of the induced module V^{G^F} since W is projective, which contradicts the indecomposability of V^{G^F} given by Green's theorem [8] §16. So $\text{St}|_{G^F} \cong \bigoplus_{g \in G^F/G^{F'}} V^g$ and we claim that each periodic simple $G^{F'}$ module is isomorphic to one of these V^g 's.

Let S be an irreducible periodic module for $G^{F'}$. If $I = I_{G^F}(S) = G^F$ then again $S = W|_{G^{F'}}$ for a simple G^F module W . In Case 1, $E \subseteq G^{F'}$ and in Case 2 $E \cap G^{F'}$ has rank ≥ 4 ; hence 3 or 2^3 divides $\dim W$ respectively which forces W to be $\cong \text{St}_{G^F}$, leading to a contradiction. Hence $I = G^{F'}$ which implies that S^G is simple periodic for G^F . So $S^G \cong \text{St}_{G^F}$ and $S \cong V^g$ for a $g \in G^F$ by Frobenius reciprocity. In particular S is projective. \square

We conclude the treatment of twisted groups of Lie type, by computing the rank varieties of simple modules for $G^F = {}^2B_2(q^2)_{\text{s.c.}}$, which includes a classification of periodic simple modules, i.e. those whose variety has dimension 1.

Let $q^2 = 2^n, n = 2m + 1, \text{char } K = 2, \theta := (?)^2, \Pi = \{\alpha, \beta\}$. The Dynkin diagram is $\circ \text{---} \circ$. Let $P := \{x(t, u) | t, u \in \mathbb{F}_{2^{2m+1}}\} \in \text{Syl}_2(G^F)$, with $x(t, u) = x_\alpha(t)x_\beta(t^\theta)x_{\alpha+2\beta}(u)x_{\alpha+\beta}(t^{1+\theta} + u^\theta)$; then $|P| = 2^{4m+2}$ and $Z(P) = \{x(0, u)\}$ is maximal elementary abelian;

let $D := Z(P)$. Notice that there is only one conjugacy class of maximal elementary abelian 2-subgroups in G^F . Here $X_1(T)' = \{(0, 0), (0, 1)\}$ with dimensions 1 and 4 respectively.

(3.2) PROPOSITION. *Let $G^F = {}^2B_2(q^2)_{\text{s.c.}}$ and M be an irreducible KG^F -module; then $V_E(M)$ is linear and $\dim V_E(M) = n - s$, where s is the number of factors $\lambda_i = (0, 1)$ in the tensor product*

$$M = \bigotimes_{i=0}^{n-1} L(\lambda_i)^{(i)}, \quad \lambda_i \in X_1(T).$$

Proof. We need only to consider $L(0, 1)$ which has a basis $B = \{v^+, X_{-\beta}v^+, X_{-(\alpha+\beta)}v^+, X_{-\beta}X_{-(\alpha+\beta)}v^+\}$. A straightforward computation shows, that

$$\begin{aligned} x(0, u)v^+ &= v^+; & x(0, u)X_{-\beta}v^+ &= X_{-\beta}v^+; \\ x(0, u)X_{-(\alpha+\beta)}v^+ &= u^\theta v^+ + X_{-(\alpha+\beta)}v^+ \\ x(0, u)X_{-\beta}X_{-(\alpha+\beta)}v^+ &= uv^+ + u^\theta X_{-\beta}X_{-(\alpha+\beta)}v^+. \end{aligned}$$

Hence $x(0, u)$ is represented with respect to B by

$$\begin{bmatrix} 1 & & & \\ 0 & 1 & 0 & \\ u^\theta & 0 & 1 & \\ u & u^\theta & 0 & 1 \end{bmatrix}$$

Let now $\{u_i\}$ be an \mathbb{F}_2 -basis of \mathbb{F}_{2^n} , then $\{x(0, u_i) =: x_i\}$ is a basis of E . For $\mathbf{a} = (a_1, a_2, \dots, a_n) \in K^n$ we define $u_{\mathbf{a}} := 1 + \sum a_i(x_i - 1)$. Then we get, using Lemma 4.2 of [3]:

$$V_E(L(0, 1)) = \{\mathbf{a} \in K^n \mid \text{rk } u_{\mathbf{a}} - 1 < \tfrac{1}{2}4\} = \left\{ \mathbf{a} \in K^n \mid \sum a_i u_i^\theta = 0 \right\}.$$

The analogue of (1.2) now gives the result. \square

REMARK. It may be interesting to notice that non-projective simple periodic modules only occur in finite groups of Lie type, if the group has a split BN -pair of rank one. But 2G_2 shows, they need not necessarily occur in this case.

REFERENCES

- [1] J. L. Alperin, *Periodicity in groups*, Illinois J. Math., **21** (1977), 776–783.
- [2] J. L. Alperin and L. Evens, *Representations, resolutions and Quillen's dimension theorem*, J. Pure Appl. Algebra, **22** (1981), 1–9.

- [3] J. F. Carlson, *The varieties and the cohomology ring of a module*, J. Algebra, **85** (1983), 104–143.
- [4] —, *Module varieties and cohomology rings of finite groups*, Vorlesungen aus dem FB Mathematik der Universitaet Essen, **13** (1985).
- [5] R. W. Carter, *Finite Groups of Lie Type: Conjugacy Classes and Complex Characters*, New York etc. 1985 (Wiley).
- [6] P. Fleischmann, *Periodic simple modules for $SU_3(q^2)$ in the describing characteristic $p \neq 2$* , Math. Z., **198** (1988), 555–568.
- [7] —, *The complexities and rank varieties of the simple modules of ${}^2A_2(q^2)$ in the natural characteristic*, J. Algebra, **121** (1989), 399–408.
- [8] B. Huppert and N. Blackburn, *Finite Groups II*, Berlin etc. 1982 (Springer).
- [9] I. Janiszczak, *Irreducible periodic modules over $SL(m, q)$ in the describing characteristic*, Comm. in algebra, **15** (1987), 1375–1391.
- [10] I. Janiszczak and J. C. Jantzen, *Simple periodic modules over Chevalley groups*, to appear.
- [11] J. C. Jantzen, *Darstellungen halbeinfacher algebraischer Gruppen und zugeordnete kontravariante Formen*, Bonner math. Schriften, **63** (1973).
- [12] —, *Zur Charakterformel gewisser Darstellungen halbeinfacher Gruppen und Lie-Algebren*, Math. Z., **140** (1974), 127–149.
- [13] —, *Modular Representations of Reductive Groups*, in *Group Theory*, Beijing 1984, Springer Lecture Notes, **1185** (1986), 118–154.
- [14] —, *Representations of Algebraic Groups*, Orlando etc. 1987, (Academic Press).
- [15] A. V. Jeyakumar, *Periodic modules for the groups $SL(2, q)$* , Comm. in Algebra, **8** (1980), 1721–1735.
- [16] G. M. Seitz, *Representations and maximal subgroups of finite groups of Lie type*, Geometriae Dedicata, **25** (1988), 391–406.
- [17] S. D. Smith, *Irreducible modules and parabolic subgroups*, J. Algebra, **75** (1982), 286–289.
- [18] R. Steinberg, *Representations of algebraic groups*, Nagoya Math. J., **22** (1963), 33–56.
- [19] —, *Lectures on Chevalley Groups*, Yale Univ. (1968).
- [20] F. D. Veldkamp, *Representations of algebraic groups of type F_4 in characteristic 2*, J. Algebra, **16** (1970), 326–339.

Received July 25, 1988.

UNIVERSITAET-GHS-ESSEN
D-4300 ESSEN, W. GERMANY

AND

UNIVERSITAET
D-2000 HAMBURG 13, W. GERMANY

Current address for J. C. Jantzen: University of Oregon
Eugene, OR 97403

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1980 *Mathematics Subject Classification* (1985 *Revision*) scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (6 Vols., 12 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
Copyright © 1990 by Pacific Journal of Mathematics

Gustavo Corach, Horacio Porta and Lázaro Recht, Differential geometry of systems of projections in Banach algebras	209
Peter Fleischmann and Jens Carsten Jantzen, Simple periodic modules of twisted Chevalley groups	229
Niels Gronbaek, Amenability of discrete convolution algebras, the commutative case	243
Nguyễn H. V. Hung, The mod 2 equivariant cohomology algebras of configuration spaces	251
Wojciech Kucharz, Global almost analytic algebraicity of analytic sets	287
John Merrill, A class of consistent anti-Martin's axioms	301
Takafumi Murai, The power $3/2$ appearing in the estimate of analytic capacity	313
L. Panaitopol and Doru Stefanescu, On the generalized difference polynomials	341
Katsuhiro Shiohama, Takashi Shioya and Minoru Tanaka, Mass of rays on complete open surfaces	349
Gerhard Ströhmer, About compressible viscous fluid flow in a bounded region	359
A. Ülger, Arens regularity sometimes implies the RNP	377