

Pacific Journal of Mathematics

**AMENABILITY OF DISCRETE CONVOLUTION ALGEBRAS,
THE COMMUTATIVE CASE**

NIELS GRONBAEK

AMENABILITY OF DISCRETE CONVOLUTION ALGEBRAS, THE COMMUTATIVE CASE

NIELS GRØNBÆK

A Banach algebra \mathfrak{A} is called amenable if all bounded derivations into dual Banach \mathfrak{A} -modules are inner. Let S be a semigroup and let $l^1(S)$ be the corresponding discrete convolution algebra. This paper is on the theme: “On the hypothesis that $l^1(S)$ is amenable, what conclusions can be drawn about the (algebraic) structure of S ?” We give a complete characterization of commutative semigroups carrying amenable semigroup algebras. If S is commutative, then $l^1(S)$ is amenable if and only if S is a finite semilattice of groups, that is, there is a finite semilattice Y and disjoint commutative groups G_α ($\alpha \in Y$) such that $S = \bigcup_{\alpha \in Y} G_\alpha$ and $G_\alpha G_\beta \subseteq G_{\alpha\beta}$ ($\alpha, \beta \in Y$).

The theme above has previously been studied in [3] and [4]. In both papers it is apparent that the condition of amenability imposes strong algebraic constraints on the semigroup. In [3] a rather complete description of inverse semigroups carrying amenable semigroup algebras is given. Of particular interest for this paper is that a semilattice carries an amenable semigroup algebra if and only if it is finite [3, Theorem 10]. In [4] it is proved that, if a one-sided cancellative semigroup carries an amenable semigroup algebra, then it is a group. The result of this paper, that for a commutative semigroup S , the semigroup algebra $l^1(S)$ is amenable if and only if S is a finite lattice of groups, is proved by looking at the gross structure of S by means of the “principle of maximal homomorphic image of a given type”. Using the fact that homomorphic images of S carry amenable semigroup algebras when S does, we establish the necessity of the characterization by showing that each archimedean component of S is a group. This is obtained by applying the results from [3] and [4], mentioned above, to the maximal semilattice, the maximal cancellative, and the maximal separative homomorphic images of S . The sufficiency of the characterization is easily verified. Alternatively, it follows from [3, Theorem 8].

1. Preliminaries. We shall need some elementary semigroup theory. We prefer to keep our exposition self-contained, so although most of what follows can be found in standard texts on the subject,

we shall, with a few exceptions, give proofs in some detail. For a further discussion the reader is referred to [1]. Throughout S will denote a commutative semigroup, with the binary operation written multiplicatively.

1.1. DEFINITIONS. Consider the following conditions on S :

(A) Each element of S is an idempotent.

(B) For all $s, t \in S$ there is $n \in \mathbb{N}$ such that

$$s^n \in tS \quad \text{and} \quad t^n \in sS.$$

(C) $s^2 = t^2 = st \Rightarrow s = t$ ($s, t \in S$).

If S satisfies (A) we call S a *semilattice*.

If S satisfies (B) we call S *archimedean*.

If S satisfies (C) we call S *separative*.

An *ideal* in S is a subset I such that $SI \subseteq I$. A *prime ideal* in S is an ideal, whose complement is a subsemigroup of S .

A *congruence* on S is an equivalence relation which is compatible with the semigroup operation.

A congruence \sim on S will be called *separative* (cancellative, archimedean, etc.) if the semigroup S/\sim is separative (cancellative, archimedean, etc.).

1.2. DEFINITION. (Principle of maximal homomorphic image of a given type). Let \mathfrak{C} be a class of congruences on S , closed under intersections. Put $\rho_0 = \bigcap \{\rho \mid \rho \in \mathfrak{C}\}$. Then S/ρ_0 is the *maximal "type class \mathfrak{C} " homomorphic image of S* .

See also [1, p. 18] and [7, §1].

EXAMPLE. Let $\rho_0 = \bigcap \{\rho \mid s^2 \rho s \ (s \in S)\}$. Then S/ρ_0 is the maximal semilattice homomorphic image of S .

1.3. DEFINITION. Let $s \in S$ and choose $m \in \mathbb{N}$ smallest possible so that $s^m = s^{m+r}$ for some $r \in \mathbb{N}$. Then $\text{order}(s) = m$ and the smallest possible r is called $\text{period}(s)$. If no such $m \in \mathbb{N}$ can be found we put $\text{order}(s) = \infty$.

1.4. DEFINITION. Let S be a semigroup and suppose that there is a semilattice Y and disjoint subsemigroups S_α ($\alpha \in Y$) of S such that $S = \bigcup_{\alpha \in Y} S_\alpha$ and $S_\alpha S_\beta \subseteq S_{\alpha\beta}$ ($\alpha, \beta \in Y$). Then S is called a *semilattice of the subsemigroups S_α ($\alpha \in Y$)*.

The following lemma is the main structure theorem for commutative semigroups.

1.5. LEMMA. *Let S be a commutative semigroup and let Y be the maximal semilattice homomorphic image of S . Then there are disjoint archimedean subsemigroups S_α ($\alpha \in Y$) of S such that S is a semilattice of the semigroups S_α ($\alpha \in Y$). This decomposition of S into archimedean subsemigroups is unique up to isomorphism of Y , and S is separative if and only if each archimedean component S_α is cancellative.*

Proof. See [1, §4.3].

1.6. LEMMA. *On S define the relations:*

$$sct \Leftrightarrow \exists u \in S \ su = tu$$

and

$$s\sigma t \Leftrightarrow \exists n_0 \in \mathbb{N} \forall n \geq n_0 \ s^n = t^n.$$

Then c and σ are congruences and S/c is the maximal cancellative homomorphic image of S and S/σ is the maximal separative homomorphic image of S .

Proof. It is clear that both relations are congruences. Now suppose ρ is a cancellative congruence; that is, $su\rho tu \Rightarrow s\rho t$ ($s, t, u \in S$). Then clearly $sct \Rightarrow s\rho t$ ($s, t \in S$) so that $c \subseteq \rho$. Since c is cancellative we are done with the statements about c .

Now suppose that $s^2\sigma t^2\sigma st$; that is, there is $n_0 \in \mathbb{N}$ so that $s^{2n} = t^{2n} = s^n t^n$ for $n \geq n_0$. Then $s^{4n_0+1}t = ss^{2n_0} \cdot t^{2n_0} \cdot t = s^{2n_0+1}t^{2n_0+1} = s^{4n_0+2}$ so that for $n \geq 8n_0+2$ we have $s^n = t^n$. Hence $s\sigma t$, proving that σ is separative. Let ρ be a separative congruence. If $s\sigma t$, then there is $k \in \mathbb{N}$ so that $st^k = t^{k+1}$. In particular $st^k \rho t^{k+1}$. This gives

$$(st^{k-1})^2 = st^{k-2}st^k \rho st^{k-2}t^{k+1} = st^{k-1}t^k \rho t^{k+1}t^{k-1} = (t^k)^2.$$

With $x = st^{k-1}$ and $y = t^k$ we have $x^2\rho y^2\rho xy$ so that $x\rho y$, that is, $st^{k-1}\rho t^k$. Repeating as necessary, we get $st\rho t^2\rho s^2$, where the second relation follows from symmetry. Thus $s\rho t$, proving that $\sigma \subseteq \rho$. \square

1.7. LEMMA. $s^2\sigma s \Leftrightarrow \text{order}(s) < \infty$ and $\text{period}(s) = 1$. If e, f are idempotents in S , then $e\sigma f \Leftrightarrow e = f$.

Proof. Suppose $s^2\sigma s$. Then there is $n_0 \in \mathbb{N}$ so that $s^{2n} = s^n$ for $n \geq n_0$. If r is the period of s we have $2n \equiv n \pmod{r}$ for $n \geq n_0$ so that $r = 1$. The rest is obvious. \square

1.8. LEMMA. *S/σ is a group if and only if S is archimedean with unique idempotent.*

Proof. First suppose that S/σ is a group. From Lemma 1.7 it follows that S has a unique idempotent. Let $s, t \in S$. Since S/σ is a group there are $u, v \in S$ so that $su\sigma t$ and $tv\sigma s$. By definition of σ , s divides a power of t and t divides a power of s , that is, S is archimedean. Conversely, let $s \in S$ and let e denote the unique idempotent in S . Since S is archimedean there are $t, u \in S$ so that $st = e$ and $ue = s^{n_0}$ for some n_0 . We have $(es)^{n_0+p} = e^{n_0+p} s^{n_0} s^p = e^{n_0+p} ues^p = ues^p = s^{n_0+p}$ ($p \in \mathbb{N}$) so that $es\sigma s$. Clearly $st\sigma e$, so S/σ is a group. \square

2. The main theorem. For the remainder of this paper we shall assume that S is a commutative semigroup such that $l^1(S)$ is amenable. We shall make frequent use of the fact that, if T is a homomorphic image of S , then $l^1(T)$ is amenable, and if I is an ideal in S which is generated by an idempotent, then $l^1(I)$, being a closed $l^1(S)$ -ideal which is unital as a Banach algebra, is amenable [6, Proposition 5.1]. Thus, if $S = \bigcup_{\alpha \in Y} S_\alpha$ is the decomposition of S into its archimedean components, then the semilattice Y is finite, since $l^1(Y)$ is amenable ([3, Theorem 10]). We give Y the usual semilattice ordering $\alpha \leq \beta \Leftrightarrow \alpha\beta = \alpha$ ($\alpha, \beta \in Y$). Since Y is finite, Y has a minimal element, namely the product of all elements in Y .

It is convenient to start with the case where S is separative; that is, we are assuming that each archimedean component is cancellative.

2.1. LEMMA. *Let S and Y be as above and let α_0 be the minimal element of Y . Then S_{α_0} is a group.*

Proof. By [4, Theorem 2.3] S/c is a group. Let $s \in S_{\alpha_0}$. Then there is $t \in S$ so that for all $u \in S$ $stucu$, that is, for all $u \in S$ there is $v \in S$ so that $stuv = uv$. Since α_0 is minimal, $st \in S_{\alpha_0}$ and $uv \in S_{\alpha_0}$, so, using the cancellation law in S_{α_0} , we see that st is a neutral element in S_{α_0} . Consequently $l^1(S_{\alpha_0})$ can be identified canonically with an ideal generated by an idempotent in $l^1(S)$. It follows that $l^1(S_{\alpha_0})$ is amenable and therefore S_{α_0} is a group, again by [4, Theorem 2.3]. \square

2.2. LEMMA. *Let $l^1(S)$ be amenable and suppose that S is separative. Then S is a finite semilattice of groups.*

Proof. Let $S = \bigcup_{\alpha \in Y} S_\alpha$ be the decomposition of S into its archimedean components. Let $\beta \in Y$, and define $T = \bigcup_{\alpha \geq \beta} S_\alpha$. Then T is a subsemigroup of S and $S \setminus T$ is a (prime) ideal in S . Hence the canonical Banach space direct sum $l^1(S) = l^1(T) \oplus l^1(S \setminus T)$ is a semidirect product, so that $l^1(T)$ is amenable. Since β is minimal in $\{\alpha \in Y \mid \alpha \geq \beta\}$, Lemma 2.1 implies that S_β is a group. But β was arbitrary in Y . \square

We now turn to the general case.

2.3. LEMMA. *Suppose $l^1(S)$ is amenable. Then S is a finite semilattice of its archimedean components, $S = \bigcup_{\alpha \in Y} S_\alpha$. Each S_α has a unique idempotent e_α , and $e_\alpha S_\alpha$ is a group, isomorphic to the maximal separative homomorphic image of S_α .*

Proof. By Lemma 2.2 S/σ is a finite semilattice of groups, $S/\sigma = \bigcup_{\alpha \in Y} G_\alpha$. Let S_α be the preimage of G_α by the canonical map $S \rightarrow S/\sigma$. With slight abuse of notation we have $S_\alpha/\sigma = G_\alpha$, so that S_α is archimedean with unique idempotent, e_α say, by Lemma 1.8. It follows that $S = \bigcup_{\alpha \in Y} S_\alpha$ is the decomposition of S into its archimedean components. Now let $s \in S_\alpha$. Since G_α is a group, there is $t \in S_\alpha$ so that $st\sigma e_\alpha$, i.e. $(st)^n = e_\alpha$ for some $n \in \mathbb{N}$. Hence $e_\alpha s^{n-1} t^n$ is an inverse to $e_\alpha s$. Clearly the canonical map from $e_\alpha S_\alpha$ to G_α is surjective. Assume that $e_\alpha s \sigma e_\alpha$ for some $s \in S_\alpha$. Since $e_\alpha S_\alpha$ is a group it follows from Lemma 1.7 that $e_\alpha s = e_\alpha$, proving injectivity of the canonical map. \square

We shall finish the proof of the main theorem by proving that $e_\alpha S_\alpha = S_\alpha$ for each $\alpha \in Y$. This is done by exploiting that $l^1(S)$, being amenable, has a bounded approximate identity. First we need a definition.

2.4. DEFINITION. Let $s \in S$. Then we define

$$[ss^{-1}] = \{u \in S \mid us = s\}.$$

Since $l^1(S)$ has a bounded approximate identity $[ss^{-1}] \neq \emptyset$ for all $s \in S$ [4, Theorem 1.1].

2.5. LEMMA. *Let $S = \bigcup_{\alpha \in Y} S_\alpha$ be the decomposition of S into its archimedean components, as in Lemma 2.3, and let $s \in S_\alpha$. If $[ss^{-1}] \cap S_\alpha \neq \emptyset$, then $s \in e_\alpha S_\alpha$. If α is maximal in Y , then S_α is a group.*

Proof. Let $u \in [ss^{-1}] \cap S_\alpha$. Then $us \sigma e_\alpha s$. Since S_α/σ is a group we have $u \sigma e_\alpha$, i.e. $u^n = e_\alpha$ for some $n \in \mathbb{N}$. Hence $s = u^n s = e_\alpha s$. In general, if $s \in S_\alpha$ and $u \in [ss^{-1}] \cap S_\beta$, then $s = us \in S_\alpha \cap S_{\beta\alpha}$, so $\beta \geq \alpha$. Thus, when α is maximal in Y we have that $[ss^{-1}] \subseteq S_\alpha$ for all $s \in S_\alpha$. It follows that $e_\alpha S_\alpha = S_\alpha$, so that S_α is a group by Lemma 2.3. \square

2.6. LEMMA. *Let $s = \bigcup_{\alpha \in Y} S_\alpha$ be as in Lemma 2.3. Then $[ss^{-1}] \cap \{e_\alpha | \alpha \in Y\} \neq \emptyset$ for all $s \in S$. In particular $l^1(S)$ is unital.*

Proof. First note that, if $u \in [ss^{-1}]$, then $[uu^{-1}] \subseteq [ss^{-1}]$. Let $s \in S$ and let S_{α_0} be the archimedean component of s . Put $u_0 = s$ and choose successively $u_k \in [u_{k-1}u_{k-1}^{-1}]$. Let S_{α_k} be the archimedean component of u_k . As noted in the proof of Lemma 2.5 we have $\alpha_0 \leq \alpha_1 \leq \dots \leq \alpha_k \leq \dots$. Since $\text{card} Y < \infty$, we eventually have $S_{\alpha_k} = S_{\alpha_{k+1}}$, whence $[u_k u_k^{-1}] \cap S_{\alpha_k} \neq \emptyset$, so that $e_{\alpha_k} \in [u_k u_k^{-1}]$ by Lemma 2.5. As observed in the beginning of the proof $e_{\alpha_k} \in [ss^{-1}]$. From [5, Theorem 7.5] it follows that $l^1(S)$ has a unit. \square

We are now able to prove:

2.7. THEOREM. *Let S be a commutative semigroup. Then $l^1(S)$ is amenable if and only if S is a finite semilattice of commutative groups.*

Proof. The sufficiency has been noted in the introduction. Hence we assume that $l^1(S)$ is amenable. Let $s = \bigcup_{\alpha \in Y} S_\alpha$ be the decomposition as in Lemma 2.3. By Lemma 2.5 the theorem is true if $\text{card} Y = 1$. We proceed by induction on $n = \text{card} Y$. Assume that $n \geq 2$ and that the theorem is true for semigroups which are semilattices of archimedean semigroups with cardinality of the semilattice strictly less than n . Let α_0 be the minimal element in Y . Let $\beta \in Y \setminus \{\alpha_0\}$, and define $T_\beta = \bigcup_{\alpha \geq \beta} S_\alpha$. As in the proof of Lemma 2.2, we see that $l^1(T_\beta)$ is amenable. Thus, by the induction hypothesis, we have that S_α is a group for $\alpha \in Y \setminus \{\alpha_0\}$. We finish the induction step by proving that $S_{\alpha_0} = e_{\alpha_0} S_{\alpha_0}$. To this end, define a congruence \sim on S by

$$s \sim t \Leftrightarrow Ss = St \quad (s, t \in S).$$

Note that, if $s \sim t$, then $s \in St$, since $[ss^{-1}] \neq \emptyset$. Using that S_α is a group for $\alpha \neq \alpha_0$, we see that $S/\sim \cong \bigcup_{\alpha \neq \alpha_0} \{e_\alpha\} \cup S_{\alpha_0}/\sim$. Hence $l^1(S_{\alpha_0}/\sim)$ is (isomorphic to) a closed ideal of finite codimension in the

amenable Banach algebra $l^1(S/\sim)$, and therefore $l^1(S_{\alpha_0}/\sim)$ is itself amenable [2, Theorem 4.1]. From Lemma 2.5 we get that S_{α_0}/\sim is a group. In particular we have for all $s \in S_{ga_0}$ that $s \sim e_{\alpha_0}s$, so, by the note above, $S_{\alpha_0} \subseteq e_{\alpha_0}S_{\alpha_0}$. The induction step is hereby completed. \square

Acknowledgment. I wish to acknowledge a stimulating correspondence with Professor J. Duncan on the subject. I wish to thank Dr. K. B. Laursen for a careful reading of the manuscript.

REFERENCES

- [1] A. H. Clifford and G. B. Preston, *The algebraic theory of semigroups*, Math. Surveys no. 7, Amer. Math. Soc., Providence, Rhode Island 1967.
- [2] P. C. Curtis and R. J. Loy, *The structure of amenable Banach algebras*; (to appear in J. London Math. Soc.).
- [3] J. Duncan and I. Namioka, *Amenability of inverse semigroups and their semigroup algebras*, Proc. Royal Soc. Edinburgh, **80A** (1978), 309–321.
- [4] N. Grønbæk, *Amenability of weighted discrete convolution algebras on cancellative semigroups*, Proc. Royal Soc. Edinburgh, **110A** (1988), 351–360.
- [5] E. Hewitt and H. S. Zuckerman, *The l_1 -algebra of a commutative semigroup*, Trans. Amer. Math. Soc., **83** (1956), 70–97.
- [6] B. E. Johnson, *Cohomology in Banach algebras*; Mem. Amer. Math. Soc., **127** (1972).
- [7] T. Tamura, *The study of closets and free contents related to semilattice decomposition of semigroups*; in Semigroups, Proceedings of a symposium on semigroups held at Wayne State University, Detroit, Michigan, (1968); Academic Press, New York and London.

Received August 29, 1988.

KØBENHAVNS UNIVERSITETS MATEMATISKE INSTITUT
UNIVERSITETSPARKEN 5
2100 KØBENHAVN Ø, DENMARK

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1980 *Mathematics Subject Classification* (1985 *Revision*) scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (6 Vols., 12 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1990 by Pacific Journal of Mathematics

Gustavo Corach, Horacio Porta and Lázaro Recht, Differential geometry of systems of projections in Banach algebras	209
Peter Fleischmann and Jens Carsten Jantzen, Simple periodic modules of twisted Chevalley groups	229
Niels Gronbaek, Amenability of discrete convolution algebras, the commutative case	243
Nguyễn H. V. Hung, The mod 2 equivariant cohomology algebras of configuration spaces	251
Wojciech Kucharz, Global almost analytic algebraicity of analytic sets	287
John Merrill, A class of consistent anti-Martin's axioms	301
Takafumi Murai, The power $3/2$ appearing in the estimate of analytic capacity	313
L. Panaitopol and Doru Stefanescu, On the generalized difference polynomials	341
Katsuhiro Shiohama, Takashi Shioya and Minoru Tanaka, Mass of rays on complete open surfaces	349
Gerhard Ströhmer, About compressible viscous fluid flow in a bounded region	359
A. Ülger, Arens regularity sometimes implies the RNP	377