

Pacific Journal of Mathematics

ON THE GENERALIZED DIFFERENCE POLYNOMIALS

L. PANAITOPOL AND DORU STEFANESCU

ON THE GENERALIZED DIFFERENCE POLYNOMIALS

L. PANAITOPOL AND D. ŞTEFĂNESCU

We study some factorization properties of a family of polynomials which includes the generalized difference polynomials. We deduce new irreducibility criteria for polynomials in two variables with coefficients in an algebraically closed field. We also obtain new proofs for the irreducibility criteria of Ehrenfeucht and Angermüller.

Let k be a commutative algebraically closed field. A polynomial in two variables $P(X, Y) \in k[X, Y]$ is called a *difference polynomial* if $P(X, Y) = f(X) - g(Y)$, where $f, g \in k[X] \setminus k$.

A. Ehrenfeucht [6] and H. Tverberg [11] studied a case of irreducibility of the difference polynomials and A. Schinzel [10] established conditions for the factorization of the difference polynomials. L. A. Rubel and S. S. Abhyankar [2], L. A. Rubel, A. Schinzel and H. Tverberg [8] and G. Angermüller [3] studied some factorization and irreducibility conditions of the larger class of the *generalized difference polynomials*

$$(*) \quad Q(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i},$$

where $c \in k \setminus \{0\}$, $n \in \mathbb{N}^*$, $\deg P_n(X) = m \geq 1$ and $\deg P_i(X) < mi/n$ for every i , $1 \leq i \leq n - 1$.

In this paper we study factorization properties of polynomials of the form

$$(**) \quad F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i},$$

where $c \in k \setminus \{0\}$, $P_i(X) \in k[X]$, $n \geq 1$.

The family of the polynomials $(**)$ includes the class of the generalized difference polynomials $(*)$. We introduce a rational number $p_Y(F)$ associated with a polynomial $F(X, Y) \in k[X, Y]$ that satisfies $(**)$. We shall establish some properties of $p_Y(F)$ using a Newton polygon argument. We deduce irreducibility criteria for the polynomials $(**)$. We also obtain new proofs of the criteria of Ehrenfeucht and Angermüller.

DEFINITION. Let X, Y be two indeterminates over k and let

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y],$$

where $c \in k \setminus \{0\}$, $n \geq 1$, $P_i(X) \in k[X]$.

We call the *degree index* of the polynomial $F(X, Y)$ the rational number

$$p_Y(F) = \max \left\{ \frac{\deg P_i}{i}; 1 \leq i \leq n \right\}.$$

REMARKS. (i) $p_Y(F) = 0$ if and only if $F(X, Y) \in k[Y]$.

(ii) If

$$p_Y(F) = \frac{\deg P_n}{n} \quad \text{and} \quad p_Y(F) > \max \left\{ \frac{\deg P_i}{i}; 1 \leq i \leq n-1 \right\}$$

then $F(X, Y)$ is a generalized difference polynomial.

THEOREM 1. *Let*

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y],$$

where $c \in k \setminus \{0\}$, $P_i(X) \in k[X]$, $n \geq 1$. If $F = F_1 F_2$, with $F_1, F_2 \in k[X, Y] \setminus k$, then $p = \max(p_1, p_2)$, where $p = p_Y(F)$, $p_1 = p_Y(F_1)$, $p_2 = p_Y(F_2)$.

Proof. We shall prove that p can be obtained as a suitable number associated with the polynomial $G(X, Y) := Y^n F(X^{-1}, Y^{-1}) \in k((X))[Y]$. (G is a polynomial in Y with the coefficients meromorphic formal power series in X .)

Let

$$G(X, Y) = \sum_{i=0}^n H_i(X)Y^i \in k((X))[Y],$$

where $H_i(X) \in k((X))$. Let $r_i = \text{ord}_X H_i(X)$ and let

$$e(G) = \max \left(\frac{r_0 - r_1}{1}, \frac{r_0 - r_2}{2}, \dots, \frac{r_0 - r_n}{n} \right).$$

But

$$G(X, Y) = P_n \left(\frac{1}{X} \right) Y^n + P_{n-1} \left(\frac{1}{X} \right) Y^{n-1} + \dots + P_1 \left(\frac{1}{X} \right) Y + c.$$

Hence $r_0 = 0$ and $r_i = \text{ord}_X P_i(\frac{1}{X}) = -\text{deg}(P_i)$ for $i = 1, 2, \dots, n$. Therefore $r_0 - r_1 = 0 - (-\text{deg}(P_1)) = \text{deg}(P_1)$ and it follows that

$$(1) \quad e(G) = p_Y(F).$$

If the characteristic of the field k is zero or is positive and does not divide the degree n of the polynomial $G(X, Y)$ then $e(G)$ is the smallest exponent of a Puiseux series

$$y(X) = \sum_i c_i X^i \in \bigcup_{m=1}^{\infty} k((X^{1/m})) \quad \text{such that } G(X, y) = 0.$$

(Such a series exists because k is algebraically closed and the characteristic of k does not divide n .)

If the characteristic of k is positive and divides n then a root $y(X)$ of the equation $G(X, y) = 0$ is not necessary a Puiseux series, as it was remarked by C. Chevalley in [4], p. 64. In this case a root of the equation $G(X, y) = 0$ is a general power series $y(X) = \sum_{i \in S(f)} c_i X^i \in k((T^{\mathbb{Q}}))$, where the support $S(f)$ is a well ordered subset of \mathbb{Q} . Let i_0 be the smallest of the exponents of $y(X)$. Then i_0 can be obtained with a Newton polygon argument (cf. [7] pp. 42–48) as

$$i_0 = \max \left(\frac{r_0 - r_1}{1}, \frac{r_0 - r_2}{2}, \dots, \frac{r_0 - r_n}{n} \right).$$

Indeed, it suffices to remark that the determination of i_0 above does not depend on the characteristic. Therefore $i_0 = e(G)$.

Let $G_1(X, Y), G_2(X, Y) \in k((X))[Y]$ corresponding to the polynomials F_1 and F_2 respectively. From a result of G. Dumas relative to the Newton polygon of the product of two polynomials ([5], pp. 216–217) it follows that $e(G) = \max(e(G_1), e(G_2))$. From (1) we deduce that $p = \max(p_1, p_2)$.

PROPOSITION 2. *Let*

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y],$$

where $c \in k^*, n \geq 1$ and let $m = \text{deg } P_n(X)$. Let us suppose that $p_Y(F) = m/n$. If $F = F_1 F_2$, with $F_1, F_2 \in k[X, Y] \setminus k$ then $p_Y(F) = p_Y(F_1) = p_Y(F_2)$.

Proof. Let

$$F_i(X, Y) = c_i Y^{n_i} + \sum_{j=1}^{n_i} P_{ij}(X) Y^{n_i-j},$$

where $c_i \in k^*$, $P_{ij}(X) \in k[X]$ and let $m_i = \deg P_{in_i}$ ($i = 1, 2$). Then $n_1 + n_2 = n$, $m_1 + m_2 = m$.

From Theorem 1 it follows that

$$\frac{m_1}{n_1} \leq \frac{m}{n} \quad \text{and} \quad \frac{m_2}{n_2} \leq \frac{m}{n}.$$

But

$$\frac{m}{n} = \frac{m_1 + m_2}{n_1 + n_2}.$$

We deduce that $m_2 n_1 \geq m_1 n_2 \geq m_2 n_1$, and hence $m_2 n_1 = m_1 n_2$. Therefore

$$\frac{m}{m_2} = \frac{m_1 + m_2}{m_2} = \frac{n_1 + n_2}{n_2} = \frac{n}{n_2}.$$

It follows that $m/n = m_2/n_2$, and hence $p_Y(F_2) = p_Y(F)$. In the same way we deduce that $p_Y(F_1) = p_Y(F)$.

COROLLARY 3. *Let*

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y],$$

where $c \in k^*$, $P_i(X) \in k[X]$, $n \geq 1$ and let $m = \deg P_n(X)$. If $p_Y(F) = m/n$ and $(m, n) = 1$ then the polynomial $F(X, Y)$ is irreducible in $k[X, Y]$.

Proof. Let us suppose that there are $F_1, F_2 \in k[X, Y] \setminus k$ such that $F = F_1 F_2$. We suppose F_1, F_2 expressed as in the proof of the former proposition.

Because $p_Y(F) = p_Y(F_1) = p_Y(F_2) = m/n$ there is $i \in \{1, 2, \dots, n_1\}$ such that

$$\frac{\deg(P_{1i})}{i} = \frac{m}{n}.$$

Therefore $im = n \cdot \deg(P_{1i})$. Since $(m, n) = 1$ there is $s \in \mathbb{N}^*$ such that $i = sn$. It follows that $i = n_1 = n$ and $\deg(P_{1i}) = \deg(P_{1n_1}) = \deg(P_{1n}) = m$. Therefore $F_2(X, Y) \in k[X]$ and we deduce that $P_n(X) = P_{1n}(X)F_2(X, Y)$. We conclude that $F_2(X, Y) \in k$, a contradiction. It follows that $F(X, Y)$ is irreducible in $k[X, Y]$.

REMARKS. (i) If the characteristic of k does not divide n one can prove Corollary 3 using the Newton-Puiseux expansion theorem [1], 5.14.

(ii) The class of the polynomials

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y], \quad c \in k^*,$$

$$P_i(X) \in k[X], \quad n \geq 1$$

such that

$$p_Y(F) = \frac{\deg(P_n)}{n}$$

includes the family of the generalized difference polynomials. Therefore Corollary 3 establishes an irreducibility criterion for the generalized difference polynomials.

LEMMA 4. *Let*

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, y], \quad c \in k^*, \quad P_i(X) \in k[X]$$

and let $f \in [X] \setminus k, g \in k[Y] \setminus k$. Then

$$p_Y(F(f, g)) = \frac{\deg(f)}{\deg(g)} \cdot p_Y(F).$$

Proof. Let $u = \deg(f), v = \deg(g)$ and $H(X, Y) = F(f(X), g(Y))$. Then

$$H(X, Y) = c[g(Y)]^n + P_1(f(X))[g(Y)]^{n-1} + \dots + P_{n-1}(f(X))g(Y) + P_n(f(X)).$$

Because

$$\frac{\deg[P_i(f(X))]}{iv} \geq \frac{\deg[P_i(f(X))]}{s} \quad \text{for } iv \leq s \leq nv$$

it follows that

$$p_Y(G) = \max \left\{ \frac{\deg P_i(f)}{iv}; 1 \leq i \leq n \right\}.$$

But $\deg P_i(f) = u \cdot \deg(P_i)$. Therefore

$$p_Y(H) = \frac{u}{v} \cdot p_Y(F) = \frac{\deg(f)}{\deg(g)} \cdot p_Y(F).$$

COROLLARY 5. *Let*

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y], \quad c \in k^*,$$

$$P_i(X) \in k[X], \quad n \geq 1$$

such that $m = \deg(P_n) \geq 1$ and $p_Y(F) = m/n$. If $f \in k[X] \setminus k$ and $g \in k[X] \setminus k$ are such that $(m \cdot \deg(f), n \cdot \deg(g)) = 1$ then the polynomial $F(f, g)$ is irreducible in $k[X, Y]$.

Proof. Let $H(X, Y) = F(f, g)$. From the above lemma it follows that $p_Y(G) = m \cdot \deg(f)/n \cdot \deg(g)$. From Corollary 3 it follows that $F(f, g)$ is irreducible in $k[X, Y]$.

REMARK. Corollary 5 was obtained by G. Angermüller in [3] with different methods in the special case $F(X, Y)$ is a generalized difference polynomial.

THEOREM 6. *Let*

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y], \quad c \in k^*,$$

$$P_i(X) \in k[X], \quad n \geq 1$$

and let $a = \deg_X F(X, Y)$. If $p_Y(F) = a/b$, $(a, b) = 1$, then the polynomial $F(X, Y)$ is irreducible in $k[X, Y]$ or it has a factor from $k[Y]$.

Proof. Let us suppose that there are $F_1, F_2 \in k[X, Y] \setminus k$ such that $F = F_1 F_2$. Let

$$F_i(X, Y) = c_i Y^{n_i} + \sum_{j=1}^{n_i} P_{ij}(X) Y^{n_i-j} \in k[X, Y], \quad c_i \in k^*,$$

$$P_{ij}(X) \in k[X] \quad (i = 1, 2).$$

From Theorem 1 it follows that we may suppose that $p_Y(F_1) = p_Y(F)$. Hence there is $j \in \{1, 2, \dots, n_1\}$ such that $\deg(P_{1j})/j = a/b$, i.e. $ai = b \cdot \deg(P_{1j})$.

Since $(a, b) = 1$ it follows a divides $\deg(P_{1j})$. But $0 \leq \deg(P_{1j}) \leq \deg_X(F) = a$. Therefore $\deg(P_{1j}) = 0$ or a .

If $\deg(P_{1j}) = 0$ then $p_Y(F_1) = 0$; hence $p_Y(F) = 0$ and it follows that $F(X, Y) \in k[Y]$.

If $\deg(P_{1j}) = a$ then $\deg_X(F_2) = 0$. Therefore F_2 is a polynomial from $k[Y]$.

It follows that $F(X, Y)$ is irreducible or has a factor from $k[Y]$.

REMARKS. (i) If the polynomial $F(X, Y)$ has a factor from $k[Y]$ then this factor is the greatest common divisor of the polynomials

$Q_i \in k[Y]$ such that

$$F(X, Y) = \sum_{i=0}^a Q_i(Y)X^{a-i}.$$

(ii) If $p_Y(F) = m/n$, where $m = \deg(P_n)$, then $m/n \geq \deg(P_i)/i$ for every $i = 1, 2, \dots, n$. Hence $m \geq (n/i) \cdot \deg P_i \geq \deg P_i$ and it follows that $m = a = \deg_X(F)$. Therefore the class of the generalized difference polynomials is contained in the family of the polynomials satisfying the assumptions from Theorem 6.

COROLLARY 7. *Let*

$$F(X, Y) = cY^n + \sum_{i=1}^n P_i(X)Y^{n-i} \in k[X, Y],$$

where $c \in k^*$, $P_i(X) \in k[X]$, $n \geq 1$, let $a = \deg_X F(X, Y)$ and $f \in k[X] \setminus k$, $g \in k[X] \setminus k$. If $p_Y(F) = a/b$ and $(a \cdot \deg(f), b \cdot \deg(g)) = 1$ then $F(f, g)$ is irreducible in $k[X, Y]$ or it has a factor from $k[Y]$.

Proof. Let $u = \deg(f)$, $v = \deg(g)$ and $H(X, Y) = F(f(X), g(Y))$. From Lemma 4 it follows that $p_Y(H) = ua/vb$. Since $\deg_X(H) = \deg(f) \cdot \deg_X(F) = ua$ and $(ua, vb) = 1$ we conclude by Theorem 6.

REFERENCES

- [1] S. S. Abhyankar, *Expansion Techniques in Algebraic Geometry*, Tata Institute, Bombay (1977).
- [2] S. S. Abhyankar and L. A. Rubel, *Every difference polynomial has a connected zero-set*, J. Indian Math. Soc., **43** (1979), 69–78.
- [3] G. Angermüller, *A generalization of Ehrenfeucht's irreducibility criterion* preprint, Univ. Erlangen, (1985), 1–8.
- [4] C. Chevalley, *Algebraic Functions of One Variable*, Amer. Math. Soc. Math. Surv., **6** (1951).
- [5] G. Dumas, *Irréductibilité des polynômes à coefficients rationnels*, J. de Math. (6^e série), **2** (1906), 191–258.
- [6] A. Ehrenfeucht, *Kryterium absolutnej nierozkładalności wielomianów*, Prace Mat., **2** (1958), 167–169.
- [7] K. Hensel and G. Landsberg, *Theorie der algebraischen Funktionen einer Variablen*, Teubner, Leipzig (1902).
- [8] L. A. Rubel, A. Schinzel, and H. Tverberg, *On difference polynomials and hereditary irreducible polynomials*, J. Number Theory, **12** (1980), 230–235.
- [9] A. Schinzel, *Reducibility of polynomials in several variables*, Bull. Polon. Ac. Sc-Ser. Math., **11** (1963), 633–638.
- [10] —, *Reducibility of polynomials of the form $f(X) - g(Y)$* , Colloq. Math., **18** (1967), 213–218.

- [11] H. Tverberg, *A remark on Ehrenfeucht's criterion for irreducibility of polynomials*, Prace Mat., **18** (1964), 117–118.

Received July 28, 1988.

UNIVERSITY OF BUCHAREST
STR. ACADEMIEI
14 BUCHAREST, ROMANIA

AND

UNIVERSITY OF BUCHAREST
P.O. Box 52-11
BUCHAREST-MAGURELE, ROMANIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

ROBION KIRBY
University of California
Berkeley, CA 94720

C. C. MOORE
University of California
Berkeley, CA 94720

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1980 *Mathematics Subject Classification* (1985 Revision) scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is issued monthly as of January 1966. Regular subscription rate: \$190.00 a year (6 Vols., 12 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1990 by Pacific Journal of Mathematics

Gustavo Corach, Horacio Porta and Lázaro Recht, Differential geometry of systems of projections in Banach algebras	209
Peter Fleischmann and Jens Carsten Jantzen, Simple periodic modules of twisted Chevalley groups	229
Niels Gronbaek, Amenability of discrete convolution algebras, the commutative case	243
Nguyễn H. V. Hung, The mod 2 equivariant cohomology algebras of configuration spaces	251
Wojciech Kucharz, Global almost analytic algebraicity of analytic sets	287
John Merrill, A class of consistent anti-Martin's axioms	301
Takafumi Murai, The power $3/2$ appearing in the estimate of analytic capacity	313
L. Panaitopol and Doru Stefanescu, On the generalized difference polynomials	341
Katsuhiro Shiohama, Takashi Shioya and Minoru Tanaka, Mass of rays on complete open surfaces	349
Gerhard Ströhmer, About compressible viscous fluid flow in a bounded region	359
A. Ülger, Arens regularity sometimes implies the RNP	377