

# Pacific Journal of Mathematics

## **REAL ANALYTIC REGULARITY OF THE SZEGŐ PROJECTION ON CIRCULAR DOMAINS**

SO-CHIN CHEN

# REAL ANALYTIC REGULARITY OF THE SZEGÖ PROJECTION ON CIRCULAR DOMAINS

SO-CHIN CHEN

**In this note we show that  $D \subseteq \mathbb{C}^n$ ,  $n \geq 2$ , is a smooth bounded pseudoconvex domain with real analytic defining function  $r(z)$  such that  $\sum_{k=1}^n z_k(\partial r/\partial z_k) \neq 0$  holds near some  $x_0 \in bD$ , then if  $g \in C^\omega(bD)$ , we have that the Szegő projection of  $g$ ,  $Sg$ , is real analytic near  $x_0$ . In particular if  $D$  is a smooth bounded complete Reinhardt (or Reinhardt) pseudoconvex domain with real analytic boundary, then the Szegő projection  $S$  preserves real analyticity globally.**

**I. Introduction.** Let  $D \subseteq \mathbb{C}^n$ ,  $n \geq 2$ , be a smooth bounded pseudoconvex domain. Denote by  $L^2(bD)$  the space of square-integrable functions on the boundary and by  $H^2(bD)$  the closed subspace of  $L^2(bD)$  whose Poisson integrals are holomorphic in  $D$ . Then we define the Szegő projection  $S$  to be the orthogonal projection from  $L^2(bD)$  onto  $H^2(bD)$ . It is represented by integration against the Szegő kernel function  $S(\omega, z)$ , i.e., for  $f \in L^2(bD)$ , we have

$$Sf(\omega) = \int_{bD} S(\omega, z)f(z) d\sigma_z.$$

The smooth regularity of the Szegő projection for a large class of domains has been established, for instance, see [1] [2], [3], [4], [8], [9], [10], [12]. Hence in this paper we are going to study the real analytic regularity of the Szegő projection on circular domains. The problems can be formulated as follows.

1. (Global version): Suppose that the boundary of  $D$  is real analytic, then does the Szegő projection  $S$  map analytic functions to analytic functions, i.e.,  $S: C^\omega(bD) \rightarrow C^\omega(bD)$ ?
2. (Local version): Suppose that the boundary of  $D$  is real analytic near some point  $x_0 \in bD$ , then does the Szegő projection  $S$  preserve real analyticity near  $x_0$ ?

These problems are quite open. The only results we know so far are due to D. Tartakoff [13]. He showed the following.

**THEOREM (Tartakoff).** *On a real, real analytic CR-manifold of dimension  $2n - 1$  whose Levi form is non-degenerate and which satisfies  $Y(q)$ , then  $\square_b$  is locally real analytic hypoelliptic on  $(p, q)$ -forms.*

For global result in this case see also Tartakoff [14].

The condition  $Y(q)$  means that the Levi form has  $\max(q + 1, n - q)$  eigenvalues of the same sign or  $\min(q + 1, n - q)$  pairs of eigenvalues of opposite sign at each point. Therefore if one has a smooth bounded strictly pseudoconvex domain  $D$  in  $\mathbb{C}^n$ ,  $n \geq 3$ , with real analytic boundary, then one can apply the following formula

$$(1.1) \quad S = \text{Id} - \bar{\partial}_b^* N_b \bar{\partial}_b,$$

where  $N_b$  is the boundary Neumann operator, to show that the Szegő projection  $S$  preserves real analyticity locally (hence globally too). However, we must point out here that this theorem does not apply to the domains in  $\mathbb{C}^2$ , because condition  $Y(1)$  is violated on such domains.

Very recently M. Derridj and D. Tartakoff [7] showed that if the defining function near  $0 \in bD$  can be expressed as

$$(1.2) \quad \text{Im } \omega = h(|z|^2)$$

with  $h$  real analytic and  $h(0) = 0$ , then again a local theorem holds near  $0 \in bD$ .

In this paper we prove the following main results.

**THEOREM 1.** *Let  $D \subseteq \mathbb{C}^n$ ,  $n \geq 2$ , be a smooth bounded pseudoconvex circular domain with real analytic defining function  $r(z)$ . Suppose that  $\sum_{k=1}^n z_k(\partial r / \partial z_k) \neq 0$  holds near some  $x_0 \in bD$  and that  $f$  is globally real analytic, i.e.,  $f \in C^\omega(bD)$ . Then  $Sf$  is real analytic near  $x_0$ .*

We remark here that (i) Theorem 1 is also true in dimension two, (ii) it is not quite a local theorem, because we need  $f$  to be globally real analytic. It follows from Theorem 1 that we have

**COROLLARY 2.** *Let  $D \subseteq \mathbb{C}^n$ ,  $n \geq 2$ , be a smooth bounded pseudoconvex circular domain with real analytic defining function  $r(z)$ . If  $\sum_{k=1}^n z_k(\partial r / \partial z_k) \neq 0$  holds for all  $z \in bD$ , then the Szegő projection preserves real analyticity globally.*

Since the transversal condition always holds on a complete Reinhardt domain, see Chen [6], we have

**COROLLARY 3.** *Let  $D \subseteq \mathbb{C}^n$ ,  $n \geq 2$ , be a smooth bounded complete Reinhardt pseudoconvex domain with real analytic boundary. Then the Szegő projection preserves real analyticity globally.*

Next if we drop the transversal condition, we can show

**THEOREM 4.** *Let  $D \subseteq \mathbb{C}^n$ ,  $n \geq 2$ , be a smooth bounded Reinhardt pseudoconvex domain with real analytic boundary. Then the Szegő projection preserves real analyticity globally.*

The author would like to thank Mei-chi Shaw for bringing this problem to his attention.

**II. Proof of the main results.** A domain  $D$  in  $\mathbb{C}^n$  is called circular if  $(z_1, \dots, z_n) \in D$  implies  $(e^{i\theta} z_1, \dots, e^{i\theta} z_n) \in D$  for all  $\theta \in \mathbb{R}$ , and  $D$  is called Reinhardt (or multi-circular) if  $(z_1, \dots, z_n) \in D$  implies  $(e^{i\theta_1} z_1, \dots, e^{i\theta_n} z_n) \in D$  for  $\theta_1, \dots, \theta_n \in \mathbb{R}$ .  $D$  is called complete Reinhardt if  $(z_1, \dots, z_n) \in D$  implies  $(\omega_1, \dots, \omega_n) \in D$  with  $|\omega_j| \leq |z_j|$  for all  $j = 1, \dots, n$ .

First we recall that the Szegő kernel function  $S(\omega, z)$  can be represented as follows. Let  $\{\varphi_j\}_{j=1}^\infty$  be an orthonormal basis for  $H^2(bD)$ . Here we have identified each element  $f \in H^2(bD)$  with its Poisson integral. Then we have

$$(2.1) \quad S(\omega, z) = \sum_{j=1}^\infty \varphi_j(\omega) \overline{\varphi_j(z)}, \quad \forall \omega, z \in D,$$

and the expression is independent of the choice of the basis.

Next we prove some basic facts on circular domains. Define an  $S^1$ -action on  $D$  as follows,

$$\begin{aligned} \pi: S^1 \times \overline{D} &\rightarrow \overline{D}, \\ (e^{i\theta}, z) &\mapsto \omega = e^{i\theta} \cdot z = (e^{i\theta} z_1, \dots, e^{i\theta} z_n). \end{aligned}$$

Then for each fixed  $\theta$ ,  $\pi_b = \pi|_{bD}$  is a CR-diffeomorphism of  $bD$ .

**LEMMA 2.2.** *Let  $D \subseteq \mathbb{C}^n$ ,  $n \geq 2$ , be a smooth bounded circular domain. Then for each fixed  $\theta$  we have  $\pi_b^* d\sigma_\omega = d\sigma_z$ , where  $d\sigma$  denotes the surface element on  $bD$ .*

*Proof.* Let  $r(\omega)$  be the defining function for  $D_\omega$ . Here we use subscripts to emphasize the domain we consider. Then  $r \circ \pi(z)$  will be a defining function for  $D_z$ . By using the  $*$ -operator (cf. [11]), we have

$$\pi_b^* d\sigma_\omega = \pi_b^* \left( \frac{2}{\|dr\|} i_\omega^* (*\partial r) \right),$$

where  $i: D \hookrightarrow \mathbb{C}^n$  is the inclusion map. Then by direct computation, we get

$$\begin{aligned} \pi_b^* d\sigma_\omega &= \frac{2}{\|d(r \circ \pi)\|} \pi_b^* i_\omega^* \left( \sum_{k=1}^n \frac{1}{2^{n-1} \cdot i^n} \frac{\partial r}{\partial \omega_k}(\omega) d\omega_k \bigwedge_{j \neq k} (d\bar{\omega}_j \wedge d\omega_j) \right) \\ &= \frac{2}{\|d(r \circ \pi)\|} i_z^* \cdot \pi^* \left( \sum_{k=1}^n \frac{1}{2^{n-1} \cdot i^n} \frac{\partial r}{\partial \omega_k}(\omega) d\omega_k \bigwedge_{j \neq k} (d\bar{\omega}_j \wedge d\omega_j) \right) \\ &= \frac{2}{\|d(r \circ \pi)\|} i_z^* (*\partial(r \circ \pi)) \\ &= d\sigma_z. \end{aligned}$$

This completes the proof of the lemma.

From Lemma 2.2 and the representation of the Szegő kernel function we obtain immediately the following transformation law for the Szegő kernel on circular domains.

**LEMMA 2.3.** *Under the hypotheses of Lemma 2.2, we have  $S(\omega, z) = S(\pi(\omega), \pi(z))$ .*

Now we define the crucial vector field  $T$  on  $bD$  as follows. Let  $z$  be a point near the boundary. Define

$$\begin{aligned} \Lambda_z: S^1 &\rightarrow \bar{D} \\ e^{i\theta} &\mapsto e^{i\theta} \cdot z = (e^{i\theta} z_1, \dots, e^{i\theta} z_n), \end{aligned}$$

and denote by  $\Lambda_{z,*}$  the differential mapping of  $\Lambda_z$ . Then define the vector field  $T$  at  $z$  by

$$\begin{aligned} T(z) &= \frac{1}{2} \Lambda_{z,*} \left( \frac{\partial}{\partial \theta} \Big|_{\theta=0} \right) = \frac{i}{2} \sum_{j=1}^n z_j \frac{\partial}{\partial z_j} - \frac{i}{2} \sum_{j=1}^n \bar{z}_j \frac{\partial}{\partial \bar{z}_j} \\ &= \frac{i}{2} X - \frac{i}{2} \bar{X}, \end{aligned}$$

where  $X = \sum_{j=1}^n z_j (\partial / \partial z_j)$ .

The vector field  $T$  defined in this way has many nice properties. First of all, if  $(\partial r/\partial z_n)(x_0) \neq 0$ , then

$$L_k = \frac{\partial r}{\partial z_k} \frac{\partial}{\partial z_n} - \frac{\partial r}{\partial z_n} \frac{\partial}{\partial z_k}, \quad k = 1, \dots, n - 1,$$

forms a local basis for  $T^{1,0}(bD)$ . Hence by assumption  $Xr \neq 0$  near  $x_0$ ,  $L_1, \dots, L_{n-1}, \bar{L}_1, \dots, \bar{L}_{n-1}$  and  $T$  form a basis for complex tangent space on  $bD$  near  $x_0$ , and it is shown in [6] that we have

$$(2.4) \quad [T, L_k] = -iL_k, \quad [T, \bar{L}_k] = i\bar{L}_k, \quad \text{for } k = 1, \dots, n - 1.$$

The next lemma shows that  $T$  commutes with the Szegő projection  $S$  on such domains.

**LEMMA 2.5.** *Under the hypotheses of Theorem 1, we have  $TSu = STu$  for all  $u \in C^\infty(bD)$ .*

*Proof.* Since  $Su \in C^\infty(bD)$ , one can apply  $T$  to  $Su$ . Also by the construction of  $T$ , we get by writing  $z = e^{it} \cdot \eta$ ,

$$\begin{aligned} 2TSu(\omega) &= \frac{d}{dt} \int_{bD} S(e^{it} \cdot \omega, z)u(z) d\sigma_z|_{t=0} \\ &= \frac{d}{dt} \int_{bD} S(e^{it} \cdot \omega, e^{it} \cdot \eta)u(e^{it} \cdot \eta) d\sigma_\eta|_{t=0} \\ &= \frac{d}{dt} \int_{bD} S(\omega, \eta)u(e^{it} \cdot \eta) d\sigma_\eta|_{t=0} \\ &= 2STu(\omega). \end{aligned}$$

This completes the proof of the lemma.

Now we begin to prove Theorem 1. So we assume that  $f \in C^\omega(bD)$ . It follows from the smooth regularity of the Szegő projection that we have  $Sf \in C^\infty(bD)$ . Denote by  $L$  (or  $\bar{L}$ ) any of the  $L_i$ 's (or  $\bar{L}_i$ 's) with  $i \leq n - 1$ , and by  $\overset{(-)}{L}$  any of  $L_i$ 's or  $\bar{L}_i$ 's with  $i \leq n - 1$ . Also denote by  $Z$  either  $\overset{(-)}{L}$  or  $T$ . Denote by  $a_{(1)}$  any of the finite collection of analytic functions that occur in commutators of  $L, \bar{L}$  and  $T$  and integration by parts. Let  $a_{(j+1)} = a_{(1)} \cdot a_{(j)}$  or  $a_{(j+1)} = Za_{(j)}$ . Hence there exists a constant  $R > 0$  such that for all  $k \in \mathbb{N}$ , we have

$$(2.6) \quad |a_{(k)}| \leq RR^k k!.$$

The proof for Theorem 1 will be complete if one can show

$$(2.7) \quad \|Z_1 \cdots Z_q S f\|_{L^2(U)} \leq M M^q q!,$$

for all  $q \in \mathbb{N}$  and some open neighborhood  $U$  of  $x_0$  and some  $M > 0$  uniformly in  $q$ . We may assume that  $Xr \neq 0$  holds on some neighborhood of  $\bar{U}$ .

Let  $\varphi$  be a cut-off function with  $\varphi \equiv 1$  in some neighborhood of  $x_0$  and  $\text{supp } \varphi$  contained in  $U$ . Denote by  $\varphi'$  any first derivative of  $\varphi$ . Then by Lemma 2.5 one can estimate the pure terms quite easily,

$$(2.8) \quad \|\varphi T^p S f\| = \|\varphi S T^p f\| \leq \|T^p f\| \leq R_f R_f^p p!,$$

for some  $R_f > 0$ . To estimate the mixed terms we use  $\text{Op}(k, q)$  to denote any differential operator of order  $q$  formed out of the  $\overset{(-)}{L}$  and  $T$  in all order with precisely  $k \overset{(-)}{L}$ 's. Thus if  $k = 0$ , by (2.8), we have

$$\|\varphi \text{Op}(0, q) S f\| = \|\varphi T^q S f\| \leq R_f R_f^q q!.$$

Hence what remains in this note is to estimate the term  $\text{Op}(s, q) S f$  with  $s \geq 1$ . We will use underline to mean at most such terms are being considered and  $c$  is a constant depending only on  $n$ . First we write

$$(2.9) \quad \begin{aligned} \text{Op}(s, p) &= T^r \overset{(-)}{L} \text{Op}(s-1, p-r-1) \\ &= \overset{(-)}{L} T^r \text{Op}(s-1, p-r-1) \\ &\quad + \sum_{j=1}^r (\pm) i T^{r-j} \overset{(-)}{L} T^{j-1} \text{Op}(s-1, p-r-1) \\ &= \overset{(-)}{L} \text{Op}(s-1, p-1) + \sum_{j=1}^r (\pm) i \text{Op}(s, p-1). \end{aligned}$$

Define

$$\begin{aligned} I_p(\varphi) &= \|\varphi \text{Op}(s, p) S f\| + \|\varphi T \text{Op}(s-1, p-1) S f\| \\ &\quad + \sum_{i=1}^{n-1} (\|\varphi L_i \text{Op}(s-1, p-1) S f\| \\ &\quad \quad \quad + \|\varphi \bar{L}_i \text{Op}(s-1, p-1) S f\|). \end{aligned}$$

By (2.9) we see that

$$\begin{aligned} I_p(\varphi) &\leq 2 \sum_{i=1}^{n-1} (\|\varphi L_i \text{Op}(s-1, p-1) S f\| + \|\varphi \bar{L}_i \text{Op}(s-1, p-1) S f\|) \\ &\quad + \|\varphi T \text{Op}(s-1, p-1) S f\| + (p-1) \|\varphi \text{Op}(s, p-1) S f\|. \end{aligned}$$

We estimate the first term as follows:

$$\begin{aligned}
 & \|\varphi L \text{Op}(s-1, p-1)Sf\|^2 \\
 &= (\varphi L \text{Op}(s-1, p-1)Sf, \varphi L \text{Op}(s-1, p-1)Sf) \\
 &= \|\varphi \bar{L} \text{Op}(s-1, p-1)Sf\|^2 \\
 &\quad + (\varphi \text{Op}(s-1, p-1)Sf, -\varphi[\bar{L}, L] \text{Op}(s-1, p-1)Sf) \\
 &\quad + 2(\varphi' \text{Op}(s-1, p-1)Sf, \varphi L \text{Op}(s-1, p-1)Sf) \\
 &\quad + 2(\varphi' \text{Op}(s-1, p-1)Sf, \varphi \bar{L} \text{Op}(s-1, p-1)Sf) \\
 &\quad + (a_{(1)}\varphi \text{Op}(s-1, p-1)Sf, \varphi L \text{Op}(s-1, p-1)Sf) \\
 &\quad + (a_{(1)}\varphi \text{Op}(s-1, p-1)Sf, \varphi \bar{L} \text{Op}(s-1, p-1)Sf).
 \end{aligned}$$

It shows that for  $C_1 > 0$  we have

$$\begin{aligned}
 & \|\varphi L \text{Op}(s-1, p-1)Sf\| \\
 & \leq \|\varphi \bar{L} \text{Op}(s-1, p-1)Sf\| \\
 & \quad + C_1(\|\varphi \text{Op}(s-1, p-1)Sf\| + 4\|\varphi' \text{Op}(s-1, p-1)Sf\| \\
 & \quad \quad \quad + 2\|a_{(1)}\varphi \text{Op}(s-1, p-1)Sf\|) \\
 & \quad + \frac{1}{C_1}(3\|\varphi L \text{Op}(s-1, p-1)Sf\| + 3\|\varphi \bar{L} \text{Op}(s-1, p-1)Sf\| \\
 & \quad \quad \quad + \|\varphi[\bar{L}, L] \text{Op}(s-1, p-1)Sf\|).
 \end{aligned}$$

LEMMA 2.11. (i)  $[\bar{L}, L] = \underline{(2n-1)a_{(1)}Z}$ .  
 (ii)

$$\begin{aligned}
 \bar{L} \text{Op}(s-1, p-1) &= \text{Op}(s-1, p-1)\bar{L} \\
 &\quad + \sum_{j=1}^{p-1} \underline{(2(2n-1))^j \binom{p-1}{j}} a_{(j)} Z_j Z_{j+1} \cdots Z_{p-1},
 \end{aligned}$$

where  $Z_j$  denotes any  $Z$ . Hence one may choose  $c = 2(2n-1)$ .

*Proof.* (i) is trivial. For (ii) we have

$$\begin{aligned}
 [\bar{L}, \text{Op}(s-1, p-1)] &= [\bar{L}, Z_1 \cdots Z_{p-1}] \\
 &= \sum_{j=1}^{p-1} \underline{\binom{p-1}{j}} [[\cdots[\bar{L}, Z_1]\cdots], Z_j] Z_{j+1} \cdots Z_{p-1}.
 \end{aligned}$$

Hence the conclusion follows immediately.



Since  $Sf$  is annihilated by  $\bar{L}$ , we obtain

$$\begin{aligned}
 (2.12) \quad I_p(\varphi) &\leq 2 \sum_{i=1}^{n-1} \{2\|\varphi \bar{L}_i \text{Op}(s-1, p-1)Sf\| \\
 &\quad + C_1(\|\varphi \text{Op}(s-1, p-1)Sf\| \\
 &\quad \quad + 4\|\varphi' \text{Op}(s-1, p-1)Sf\| \\
 &\quad \quad + 2R^2\|\varphi \text{Op}(s-1, p-1)Sf\|) \\
 &\quad + \frac{1}{C_1}(3\|\varphi L_i \text{Op}(s-1, p-1)Sf\| \\
 &\quad \quad + 3\|\varphi \bar{L}_i \text{Op}(s-1, p-1)Sf\| \\
 &\quad \quad + cR^2\|\varphi Z \text{Op}(s-1, p-1)Sf\|) \} \\
 &\quad + \|\varphi T \text{Op}(s-1, p-1)Sf\| \\
 &\quad + (p-1)\|\varphi \text{Op}(s, p-1)Sf\|.
 \end{aligned}$$

Choose  $C_1 = \max(24, 8ncR^2)$ ; then we get

$$\begin{aligned}
 (2.13) \quad I_p(\varphi) &\leq 8 \sum_{i=1}^{n-1} \sum_{j=1}^{p-1} \underline{c^j \binom{p-1}{j}} \|a_{(j)}\varphi Z_j Z_{j+1} \cdots Z_{p-1} Sf\| \\
 &\quad + 4(n-1)C_1(\|\varphi \text{Op}(s-1, p-1)Sf\| \\
 &\quad \quad + 4\|\varphi' \text{Op}(s-1, p-1)Sf\| \\
 &\quad \quad + 2R^2\|\varphi \text{Op}(s-1, p-1)Sf\|) \\
 &\quad + 2\|\varphi T \text{Op}(s-1, p-1)Sf\| \\
 &\quad + 2(p-1)\|\varphi \text{Op}(s, p-1)Sf\| \\
 &\leq 8(n-1) \sum_{j=1}^{p-1} \underline{c^j \binom{p-1}{j}} \cdot RR^j j! I_{p-1}(\varphi) \\
 &\quad + 16(n-1)C_1 I_{p-1}(\varphi') \\
 &\quad + 4(n-1)C_1(1 + 2R^2)I_{p-1}(\varphi) \\
 &\quad + 2\|\varphi \text{Op}(s-1, p)Sf\| + 2(p-1)I_{p-1}(\varphi).
 \end{aligned}$$

Since the vector field  $\overset{(-)}{L}$  in general is defined only locally, we need a special cut-off function which is of compact support, but behaves like an analytic function up to certain order.

**LEMMA (Ehrenpreis).** *Let  $x_0 \in bD$  and  $U_1, U_2$  be two neighborhoods of  $x_0$  with  $U_1 \Subset U_2 \Subset U$ ; then there exists a constant  $M > 0$  such that for any integer  $k$  one can find  $\varphi_k \in C_0^\infty(U_2)$  with*

$0 \leq \varphi_k \leq 1$ ,  $\varphi_k \equiv 1$  on  $U_1$  and satisfying

$$(2.14) \quad |D^\alpha \varphi_k| \leq M(Mk)^{|\alpha|}, \quad \text{for } |\alpha| \leq k + 1.$$

Now one can replace  $\varphi$  by  $\varphi_p^{(b)}$ , some derivative of  $\varphi_p$  of order  $\leq b$ , and  $p$  by  $p - b$  in (2.13). Then we obtain

$$(2.15) \quad \begin{aligned} I_{p-b}(\varphi_p^{(b)}) &\leq 8(n-1) \sum_{j=1}^{p-b-1} \frac{c^j (p-b-1)}{j} \cdot RR^j j! I_{p-b-j}(\varphi_p^{(b)}) \\ &\quad + 16(n-1)C_1 I_{p-b-1}(\varphi_p^{(b+1)}) \\ &\quad + 4(n-1)C_1(1+2R^2)I_{p-b-1}(\varphi_p^{(b)}) \\ &\quad + 2\|\varphi_p^{(b)} \text{Op}(s-1, p-b)Sf\| \\ &\quad + 2(p-b-1)I_{p-b-1}(\varphi_p^{(b)}). \end{aligned}$$

There are five terms in (2.15). Inductively we will show that for  $p \geq b \geq 0$ , we have

$$(2.16) \quad I_{p-b}(\varphi_p^{(b)}) \leq R_1(R_2p)^b(R_3p)^{p-b}R_4^s,$$

where  $R_1, R_2, R_3$  and  $R_4$  are some constants, i.e., we will show that each term can be bounded by  $\frac{1}{5}R_1(R_2p)^b(R_3p)^{p-b}R_4^s$ . The initial step  $s = 0$  or  $p = b$  is easy to check. Hence we assume that (2.16) is true for  $b > b_0$  or  $s < s_0$ . Then we prove the case  $b = b_0$  and  $s = s_0$ .

$$\begin{aligned} \text{Term 1} &\leq 8(n-1) \sum_{j=1}^{p-b_0-1} \frac{(p-b_0-1)!}{(p-b_0-j-1)!j!} \\ &\quad \cdot R(cR)^j j! R_1(R_2p)^{b_0}(R_3p)^{p-b_0-j} \cdot R_4^{s_0} \\ &= R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0} \\ &\quad \cdot \sum_{j=1}^{p-b_0-1} \frac{(p-b_0-1)!}{(p-b_0-j-1)!p^j} \cdot \frac{8(n-1)R(cR)^j}{R_3^j} \\ &\leq \frac{1}{5}R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0}, \end{aligned}$$

provided  $R_3$  is chosen large enough.

$$\begin{aligned} \text{Term 2} &\leq 16(n-1)C_1R_1(R_2p)^{b_0+1} \cdot (R_3p)^{p-b_0-1} \cdot R_4^{s_0} \\ &= R_1(R_2p)^{b_0} \cdot (R_3p)^{p-b_0} \cdot R_4^{s_0} \cdot \left( \frac{16(n-1)C_1R_2}{R_3} \right) \\ &\leq \frac{1}{5}R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0}, \end{aligned}$$

provided  $R_3 \geq 80(n-1)C_1R_2$ .

$$\begin{aligned} \text{Term 3} &\leq 4(n-1)C_1(1+2R^2) \cdot R_1(R_2p)^{b_0}(R_3p)^{p-b_0-1} \cdot R_4^{s_0} \\ &= R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0} \cdot \left( \frac{4(n-1)C_1(1+2R^2)}{R_3p} \right) \\ &\leq \frac{1}{5}R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0}, \end{aligned}$$

provided  $R_3 \geq 20(n-1)C_1(1+2R^2)$ .

$$\begin{aligned} \text{Term 4} &\leq 2R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0-1} \\ &\leq \frac{1}{5}R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0}, \end{aligned}$$

provided  $R_4 \geq 10$ .

$$\begin{aligned} \text{Term 5} &\leq 2(p-b_0-1)R_1(R_2p)^{b_0}(R_3p)^{p-b_0-1} \cdot R_4^{s_0} \\ &= R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0} \left( \frac{2(p-b_0-1)}{R_3p} \right) \\ &\leq \frac{1}{5}R_1(R_2p)^{b_0}(R_3p)^{p-b_0} \cdot R_4^{s_0}, \end{aligned}$$

provided  $R_3 \geq 10$ . This completes the proof of (2.16). In particular, we have shown that for  $k \leq p$ ,

$$\begin{aligned} (2.17) \quad \|\varphi_p \text{Op}(k, p)Sf\| &\leq I_p(\varphi_p) \leq R_1(R_3p)^p R_4^k \\ &\leq R_1(R_3R_4p)^p \\ &\leq MM^p p!, \end{aligned}$$

for some large constant  $M > 0$  depending only on  $R_1, R_3$  and  $R_4$ . This also completes the proof of Theorem 1.

To prove Theorem 4 one can apply the above techniques almost verbally with slight modification. The key point is that for every point  $x_0 \in bD$  we have to choose a vector field like  $T$  for Theorem 1 which is transversal to  $T^{1,0}(bD) \oplus T^{0,1}(bD)$  locally near  $x_0$  and commutes nicely with  $\overset{(-)}{L}$  and the Szegő projection. This can be done easily. By rotational symmetry of the domain one can choose a direction, say  $n$ , such that  $(z_n(\partial r/\partial z_n))(x_0) \neq 0$  holds in some neighborhood of  $x_0$ . Then define  $S^1$ -action on  $\bar{D}$  as follows.

$$\begin{aligned} \pi: S^1 \times \bar{D} &\rightarrow \bar{D} \\ (e^{i\theta}, z) &\mapsto e^{i\theta} \cdot z = (z_1, \dots, z_{n-1}, e^{i\theta} z_n). \end{aligned}$$

Such an action will generate a vector field  $T_n = iz_n(\partial/\partial z_n) - i\bar{z}_n(\partial/\partial \bar{z}_n)$  with the desired properties. So we are done. For more details in this case see [5].

## REFERENCES

- [1] H. P. Boas, *Regularity of the Szegő projection in weakly pseudoconvex domains*, Indiana Univ. Math. J., **34** (1985), 217–223.
- [2] ———, *The Szegő projection: Sobolev estimates in regular domains*, Trans. Amer. Math. Soc., **300** (1987), 109–132.
- [3] H. P. Boas, S. C. Chen and E. J. Straube, *Exact regularity of the Bergman and Szegő projections on domains with partially transverse symmetries*, Manuscripta Math., **62** (1988), 467–475.
- [4] L. Boutet de Monvel and J. Sjöstrand, *Sur la singularité des noyaux de Bergman et de Szegő*, Astérisque, **34-35** (1976), 123–164.
- [5] S. C. Chen, *Global real analyticity of solutions to the  $\bar{\partial}$ -Neumann problem on Reinhardt domains*, Indiana Univ. Math. J., **37**, No. 2 (1988), 421–430.
- [6] ———, *Global analytic hypoellipticity of the  $\bar{\partial}$ -Neumann problem on circular domains*, Invent. Math., **92** (1988), 173–185.
- [7] M. Derridj and D. S. Tartakoff, *Local analyticity for  $\square_b$  and the  $\bar{\partial}$ -Neumann problem at certain weakly pseudoconvex points*, Comm. Partial Differential Equations, **13** (12) (1988), 1521–1600.
- [8] N. Kerzman and E. M. Stein, *The Szegő kernel in terms of Cauchy-Fantappiè kernels*, Duke Math. J., **45** (1978), 197–224.
- [9] J. J. Kohn, *Boundaries of Complex Manifolds*, in Proc. Conference on Complex Analysis (Minneapolis), Springer-Verlag, New York, 1965.
- [10] D. H. Phong and E. M. Stein, *Estimates for the Bergman and Szegő projections on strongly pseudo-convex domains*, Duke Math. J., **44** (1977), 695–704.
- [11] R. M. Range, *Holomorphic Functions and Integral Representations in Several Complex Variables*, Springer-Verlag, New York, 1986.
- [12] E. J. Straube, *Exact regularity of Bergman, Szegő and Sobolev space projections in non-pseudoconvex domains*, Math. Zeit., **192** (1986), 117–128.
- [13] D. S. Tartakoff, *On the local real analyticity of solutions to  $\square_b$  and the  $\bar{\partial}$ -Neumann problem*, Acta Math., **145** (1980), 177–204.
- [14] ———, *On the global real analyticity of solutions to  $\square_b$* , Comm. Partial Differential Equations, **1** (1976), 283–311.

Received April 10, 1989. Partially supported by the NSF.

SUNY  
ALBANY, NY 12222



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093

R. FINN  
Stanford University  
Stanford, CA 94305

HERMANN FLASCHKA  
University of Arizona  
Tucson, AZ 85721

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305

C. C. MOORE  
University of California  
Berkeley, CA 94720

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH  
(1906-1982)

B. H. NEUMANN

F. WOLF  
(1904-1989)

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1991 by Pacific Journal of Mathematics

<b>Yuri A. Abramovich</b> , Operators preserving disjointness on rearrangement invariant spaces .....	201
<b>Andrew French Acker and Kirk Lancaster</b> , Existence and geometry of a free boundary problem for the heat equation .....	207
<b>So-Chin Chen</b> , Real analytic regularity of the Szegő projection on circular domains .....	225
<b>Chen-Lian Chuang</b> , An independence property of central polynomials .....	237
<b>Peter Larkin Duren and M. Schiffer</b> , Robin functions and energy functionals of multiply connected domains .....	251
<b>Johan Henricus Bernardus Kemperman</b> , Sets of uniqueness and systems of inequalities having a unique solution .....	275
<b>Ka-Lam Kueh</b> , Fourier coefficients of nonholomorphic modular forms and sums of Kloosterman sums .....	303
<b>Gerard J. Murphy</b> , Ordered groups and crossed products of $C^*$ -algebras ...	319
<b>You-Qiang Wang</b> , The $p$ -parts of Brauer character degrees in $p$ -solvable groups .....	351
<b>Hidenobu Yoshida</b> , Harmonic majorization of a subharmonic function on a cone or on a cylinder .....	369