

Pacific Journal of Mathematics

THE PROPER FORCING AXIOM AND STATIONARY SET REFLECTION

ROBERT EMILE BEAUDOIN

THE PROPER FORCING AXIOM AND STATIONARY SET REFLECTION

ROBERT E. BEAUDOIN

Our main result is that the proper forcing axiom (PFA) is equiconsistent with “PFA + there is a nonreflecting stationary subset of ω_2 .” More generally we show for any cardinals $n < m \leq \aleph_2$ that if $\text{PFA}^+(n)$ is consistent with ZFC then so is “ $\text{PFA}^+(n)$ + there are m mutually nonreflecting stationary subsets of ω_2 .” As corollaries we can show that if $n < m \leq \aleph_1$ then $\text{PFA}^+(n)$ (if consistent) does not imply $\text{PFA}^+(m)$, and that PFA (if consistent) does not imply Martin’s maximum.

1. Introduction. Recently much attention has been given to various strengthenings of Martin’s axiom for \aleph_1 . Following [FMS] let us denote by $\text{MA}(\Gamma)$, where Γ is a class of partial orders, the statement:

If $P \in \Gamma$ and Δ is a family of at most \aleph_1 dense subsets of P , then there is a Δ -generic filter on P .

Thus letting Γ be the class of all partial orders having the c.c.c., $\text{MA}(\Gamma)$ becomes Martin’s axiom (for \aleph_1 dense sets). Taking Γ to be the class of proper partial orders, $\text{MA}(\Gamma)$ becomes the proper forcing axiom (PFA). Taking Γ to be the class of all orders P so that forcing the P preserves the stationarity of subsets of ω_1 , $\text{MA}(\Gamma)$ becomes Martin’s maximum (MM), discussed in [FMS].

One may fortify these axioms by demanding that the filter obtained not only be generic, but also respect the stationarity of a collection of subsets (in the generic extension) of ω_1 . That is, one may consider the axioms

$\text{MA}^+(\Gamma, \kappa)$: If $P \in \Gamma$, Δ is a family of at most \aleph_1 dense subsets of P , and $\{\mathbf{S}_\alpha : \alpha < \kappa\}$ is a family of terms, each forced by every condition in P to denote a stationary subset of ω_1 , then there is a Δ -generic filter G on P so that for every $\alpha < \kappa$, $\mathbf{S}_\alpha(G)$ is stationary in ω_1 .

(Here $\mathbf{S}_\alpha(G) = \{\beta < \omega_1 : \exists p \in G \ p \Vdash \beta \in \mathbf{S}_\alpha\}$, the interpretation of the term \mathbf{S}_α by the filter G .) If Γ is the class of proper

partial orders $\text{MA}^+(\Gamma, \kappa)$ is commonly called $\text{PFA}^+(\kappa)$. The usual consistency proof of PFA in fact establishes consistency of $\text{PFA}^+(\aleph_1)$ with ZFC (relative to the consistency of $\text{ZFC} +$ “There is a supercompact cardinal”); it is not hard to see that $\text{PFA}^+(\aleph_2)$ is inconsistent with ZFC. Baumgartner conjectured in [Ba] that $\text{PFA}^+(1)$ is actually stronger than PFA; on the other hand, he also proved that if Γ is the class of partial orders having the c.c.c., then $\text{MA}(\Gamma)$ (i.e., ordinary MA_{\aleph_1}) and $\text{MA}^+(\Gamma, n)$ are equivalent for all finite n . Shelah [Sh, Remark 6A3] has proved that $\text{MM}^+(n)$, if consistent, does not imply $\text{PFA}^+(m)$ ($n < m \leq \aleph_2$, both cardinals), more than establishing Baumgartner’s conjecture. Simpler proofs were given by Veličković [V]. Our arguments, announced at the Southeastern Logic Symposium (Gainesville, February 1985), only show that $\text{PFA}^+(n)$, if consistent, does not imply $\text{PFA}^+(m)$, but are considerably simpler still. Our methods also establish that PFA does not (if consistent) imply MM (as, of course, do Shelah’s and Veličković’s).

The wedge we will drive between PFA and $\text{PFA}^+(1)$ is a nonreflecting stationary set. A stationary set S on a cardinal κ reflects if there is a limit $\alpha < \kappa$ with $\text{cf } \alpha > \omega$ such that $S \cap \alpha$ is stationary in α ; we shall say S reflects at α . Let us call a family F of stationary sets on κ *mutually reflecting* if for some limit $\alpha < \kappa$ every $S \in F$ reflects at α ; otherwise we call F *mutually nonreflecting*. For brevity’s sake, we adopt Shelah’s notation for sets of ordinals of specified cofinality: $S_k^j = \{\alpha < \omega_j : \text{cf } \alpha = \omega_k\}$. Our main result (Theorem 2.6) is that for any $n < m$, $\text{PFA}^+(n)$ (if consistent with ZFC) is consistent with the existence of a family of m mutually nonreflecting stationary subsets of S_0^2 . But (trivially generalizing a theorem from [Ba] for one stationary set), letting Γ be the class of all countably closed partial orders, $\text{MA}^+(\Gamma, m)$ implies that every family of m stationary subsets of ω_2 is mutually reflecting. So $\text{PFA}^+(n)$ is weaker than $\text{PFA}^+(m)$. In independent (unpublished) work along the same lines, Magidor constructed, by forcing over a model with a supercompact cardinal, a model of set theory in which PFA holds and there is a nonreflecting stationary subset of ω_2 (so $\text{PFA}^+(1)$ fails). His argument is essentially the same as ours.

As another application of Theorem 2.6, we can derive (modulo consistency of PFA) the following theorem from our doctoral dissertation [Be] (unpublished): There is a model of ZFC in which there are no ω_2 -Aronszajn threes but there is a nonreflecting stationary subset of S_0^2 . This follows from Theorem 2.6 (with $n = 0$ and $m = 1$) and

Baumgartner's result (see [To] for a proof) that PFA implies there are no ω_2 -Aronszajn trees. In [Be] we obtained such a model by forcing over a model with a weakly compact cardinal; the proof of Theorem 2.6 is an elaboration of this earlier construction.

Finally, let us mention an open question. In [Sh], Shelah has shown that MM implies $\text{MA}^+(\Gamma, 1)$, for Γ the class of countably closed partial orders. We do not know if MM implies $\text{MA}^+(\Gamma, m)$ for m between 2 and \aleph_1 (and the same Γ). Question: For which $m \leq \aleph_2$ does MM imply that every family of m stationary subsets of S_0^2 is mutually reflecting? We cannot rule out the possibility that MM is consistent with the existence of a pair of mutually nonreflecting stationary subsets of S_0^2 .

2. Adjoining mutually nonreflecting stationary sets. Our main goal is to show, for any cardinals n and m with $n < m \leq \aleph_2$, that $\text{PFA}^+(n)$ is equiconsistent with $\text{PFA}^+(n) +$ "There is a mutually nonreflecting family of m stationary subset of ω_2 ". Starting from a model of $\text{PFA}^+(n)$ we will force to add such a family. Let P be the set of all functions p from m into $\mathcal{P}(\omega_2)$, such that there is a $\beta < \omega_2$ so that for every $i < m$ $p(i) \subseteq \beta \cap S_0^2$, for every $\alpha \in S_1^2$, $p(i) \cap \alpha$ is nonstationary in α , and for $j \neq i$ $p(i) \cap p(j) = \emptyset$. For each $p \in P$ let $\text{lh}(p) = \inf\{\beta : \forall i < m \ p(i) \subseteq \beta\}$, and call this ordinal the length of p . Endow P with a partial ordering: $p \leq q$ iff for every $i < m$, $p(i) \cap \text{lh}(q) = q(i)$.

LEMMA 2.1. *P is countably complete and ω_2 -Baire, and any V -generic filter on P is equiconstructible over V with an enumeration E of a family of m pairwise disjoint, mutually nonreflecting stationary subsets of S_0^2 , whose union is costationary in S_0^2 .*

Proof. If $\langle p_k : k < \omega \rangle$ is a decreasing sequence in P then defining $p(i) = \bigcup\{p_k(i) : k < \omega\}$ for each $i < m$ clearly makes p the infimum of $\{p_n : n < \omega\}$ in P . So P is countably complete.

Next, suppose that $\langle D_\sigma : \sigma < \omega_1 \rangle$ is a sequence of dense open subsets of P , and that $p \in P$ is arbitrary. We shall find $q \leq p$ such that $q \in \bigcap\{D_\sigma : \sigma < \omega_1\}$. Construct $\langle (p_\sigma, \alpha_\sigma) : \sigma < \omega_1 \rangle$ by induction on σ such that:

- (1) $p_0 = p$ and $\alpha_0 = \text{lh}(p) + 1$.
- (2) $p_{\sigma+1} \leq p_\sigma$, $p_{\sigma+1} \in D_\sigma$, $\text{lh}(p_{\sigma+1}) > \alpha_\sigma$, and $\alpha_{\sigma+1} = \text{lh}(p_{\sigma+1}) + 1$.

- (3) If σ is limit then $\alpha_\sigma = \sup\{\alpha_\tau : \tau < \sigma\}$, for some $i < m$ $p_\sigma(i) = \bigcup\{p_\tau(i) : \tau < \sigma\} \cup \{\alpha_\sigma + \omega\}$, and for the remaining $j \neq i$ $p_\sigma(j) = \bigcup\{p_\tau(j) : \tau < \sigma\}$.

So for limit σ p_σ is a condition of length $\alpha_\sigma + \omega$ extending each p_τ , such that $\alpha_\sigma \notin \bigcup\{p_\sigma(i) : i < m\}$. Note for any σ that $\alpha_{\sigma+1}$, being a successor ordinal, is not in $\bigcup\{p_{\sigma+2}(i) : i < m\}$, whence for any σ and τ , $\alpha_\sigma \notin \bigcup\{p_\tau : i < m\}$. For each $i < m$ let $q(i) = \bigcup\{p_\sigma(i) : i < m\}$, and let $\alpha = \sup\{\alpha_\sigma : \sigma < \omega_1\}$. Then $\{\alpha_\sigma : \sigma < \omega_1\}$ is club in α and disjoint from $\bigcup\{q(i) : i < m\}$, and it follows easily that q is a condition in P . Clearly $q \leq p_\sigma$ for each σ , so $q \leq p$ belongs to each D_σ , which shows that P is ω_2 -Baire.

Now suppose F is a V -generic filter on P , and define a function E with domain m by setting $E(i) = \bigcup\{p(i) : p \in F\}$. As $F = \{p \in P : \forall i < m p(i) = E(i) \cap \text{lh}(p)\}$, F and E are equiconstructible. Clearly E enumerates a family of m pairwise disjoint sets, and for each $\alpha < \omega_2$ and $i < m$ $E(i) \cap \alpha$ is nonstationary in α . We claim that each $E(i)$ is stationary in ω_2 . Supposing $p \in P$ and $p \Vdash$ “ \mathbf{C} is club in ω_2 ,” it is enough to find $q \leq p$ and $\alpha \in q(i)$ such that $q \Vdash$ “ $\alpha \in \mathbf{C}$.”

Fix λ regular and so large that $P \in H(\lambda)$, and let $<^*$ well-order $H(\lambda)$. Choose a countable elementary substructure N of $(H(\lambda), \varepsilon, <^*)$ such that $p, P, \mathbf{C} \in N$. (The name \mathbf{C} may be identified with $\{(r, \beta) : r \Vdash \beta \in \mathbf{C}\}$.) Let $\alpha = \sup(N \cap \omega_2)$. Choose a descending sequence $\langle p_k : k < \omega \rangle$ of conditions in $P \cap N$ so that $p_0 = p$ and for every $D \in N$, if $D \subseteq P$ is dense below p then $p_k \in D$ for some k . Let $p' = \inf\{p_k : k < \omega\}$. Simple dense-set arguments show $\text{lh}(p') = \alpha$ and $p' \Vdash$ “ α is a limit point of \mathbf{C} .” Thus $p' \Vdash$ “ $\alpha \in \mathbf{C}$.” Define q by setting $q(i) = p'(i) \cup \{\alpha\}$, and setting $q(j) = p(j)$ for $j \neq i$. Clearly q is a condition such that $q \leq p$, $\alpha \in q(i)$, and $q \Vdash$ “ $\alpha \in \mathbf{C}$,” as desired.

Finally we must show that $S_0^2 - \bigcup\{E(i) : i < m\}$ is stationary. It suffices, given $p \in P$ such that $p \Vdash$ “ \mathbf{C} is club in ω_2 ,” to find $q \leq p$ and $\alpha \in S_0^2$ such that $q \Vdash$ “ $\alpha \in \mathbf{C}$ ” and $\alpha < \text{lh}(q)$, yet $\alpha \notin \bigcup\{q(i) : i < m\}$. As above we can find p' and α so that $p' \leq p$, $\text{lh}(p') = \alpha \in S_0^2$, and $p' \Vdash$ “ $\alpha \in \mathbf{C}$.” Now choose some $i < m$, let $q(i) = p'(i) \cup \{\alpha + \omega\}$, and for $j \neq i$ let $q(j) = p'(j)$. Clearly $q \in P$ is as desired, and so the lemma is proved. \square

REMARK. If $m = 1$ then forcing with P adjoins a single nonreflecting stationary subset of S_0^2 which is also costationary in S_0^2 .

Now our main result is easy to state: If V is a model of set theory plus $\text{PFA}^+(n)$ then any generic extension of V by forcing with P is also a model of $\text{PFA}^+(n)$. Hence $\text{PFA}^+(n)$ and $\text{PFA}^+(n) +$ “there are m mutually nonreflecting stationary subsets of ω_2 ” are equiconsistent (over ZFC). For use in the proof, we establish the following notation and terminology. E will always be a family constructed as in Lemma 2.1 from a V -generic filter on P , and \mathbf{E} will be the canonical P -term for such a family. Thus any generic extension of V by P has the form $V[E]$. (E and the generic filter are equiconstructible.) Note that forcing with P preserves \aleph_1 and \aleph_2 , so n and m are not collapsed. Note also that if $p \in P$, $\alpha \in S_0^2$, and for some $i < m$ either $p \Vdash “\alpha \in \mathbf{E}(i)”$ or $p \Vdash “\alpha \notin \mathbf{E}(i)”$ then $\alpha < \text{lh}(p)$.

We shall call $(Q, \Delta, \langle S_i : i < n \rangle)$ an *obnoxious triple* if Q is a proper partial order, Δ a family of at most \aleph_1 dense subsets of Q , and each S_i a Q -term for a stationary subset of ω_1 , such that there is no Δ -generic filter G on Q for which $S_i(G)$ is stationary for every $i < n$. Call Q *obnoxious* if there are a Δ and $\langle S_i : i < n \rangle$ so that $(Q, \Delta, \langle S_i : i < n \rangle)$ is an obnoxious triple. So $\text{PFA}^+(n)$ states that there is no obnoxious Q .

LEMMA 2.2. *If there is an obnoxious Q then there is an obnoxious triple $(Q', \Delta, \langle S_i : i < n \rangle)$ so that Q' collapses \aleph_2 to cardinality \aleph_1 and each $D \in \Delta$ is dense and open in Q' .*

Proof. Suppose $(Q, \{F_\sigma : \sigma < \omega_1\}, \langle S_i : i < n \rangle)$ is an obnoxious triple. If Q does not collapse \aleph_2 let C be the Levy partial order for adding a function from ω_1 onto ω_2 with countable conditions, defined in the extension by Q , and if Q does collapse \aleph_2 let C be the trivial order. Since C is countably closed, $Q * C$ is proper. Let $D_\sigma = \{(q, c) \in Q * C : \exists q' \in F_\sigma q \leq q'\}$. Clearly D_σ is dense open and if G were a $\{D_\sigma : \sigma < \omega_1\}$ -generic filter on $Q * C$ then $\{q : \exists c (q, c) \in G\}$ would be $\{F_\sigma : \sigma < \omega_1\}$ -generic on Q . So $(Q * C, \{D_\sigma : \sigma < \omega_1\}, \langle S_i : i < n \rangle)$ is an obnoxious triple. \square

The following is Lemma 8.2 of [Ba].

LEMMA 2.3. *If α is an ordinal of uncountable cofinality and $S \subseteq \{\beta < \alpha : \text{cf } \beta = \omega\}$, then S is stationary in α iff $\{x \in [\alpha]^{\leq \aleph_0} : \sup(x) \in S\}$ is stationary in $[\alpha]^{\leq \aleph_0}$.*

Using Lemma 2.3 it is easy to show that proper partial orders preserve stationarity of sets of ordinals of countable cofinality. (This is well known, but as we have no reference we include the simple proof.)

LEMMA 2.4. *If Q is a proper partial order, α is regular and uncountable, and $S \subseteq \{\beta < \alpha : \text{cf } \beta = \omega\}$ is stationary, then \Vdash_Q “ S is stationary in α .” (Note that Q may collapse α , though in the extension by Q α must remain uncountable.)*

Proof. By Lemma 2.3 $\{x \in [\alpha]^{\leq \aleph_0} : \text{sup}(x) \in S\}$ is stationary, and as Q is proper this set remains stationary in the extension by Q . But then applying Lemma 2.3 again in V^Q shows that S remains stationary there. \square

Now suppose E is a mutually nonreflecting family of stationary sets added to V by forcing with P , Q is a proper partial order in $V[E]$ that collapses \aleph_2 , and G is $V[E]$ -generic on Q . In $V[E, G]$ ω_2^V has a club subset of order type ω_1 ; let f be the increasing enumeration of such a club. After forcing with Q each $E(i)$ remains stationary in ω_2 by Lemma 2.4, so $f^{-1''}E(i)$ is stationary in ω_1 ; let R_i be the usual partial order for shooting a club through the complement of $f^{-1''}E(i)$. That is, R_i is the set of all continuous, strictly increasing functions r whose domains are countable successor ordinals and whose ranges are subsets of $\omega_1 - f^{-1''}E(i)$. Since R_i adjoins a club subset C of ω_1 such that $f''C$ is disjoint from $E(i)$ and $f''C$ is club in ω_2 , R_i is not proper. Central to the arguments of this section is the following lemma.

LEMMA 2.5. *Let E , Q , f , i and R_i be as above. Let $\mathbf{f} \in V$ be any $P * Q$ -term for f , such that every condition forces \mathbf{f} to denote the increasing enumeration of a club in ω_2 of type ω_1 . Then:*

- (a) *The iteration $P * Q * R_i$ is a proper partial order.*
- (b) *For all sufficiently large regular λ there is a D club in $[H(\lambda)]^{\leq \aleph_0}$ such that if $N \in D$, $(p, q, r) \in P * Q * R_i \cap N$, $\alpha = N \cap \omega_1$, and $\zeta = \text{sup}(N \cap \omega_2)$, then there is an extension (p', q', r') of (p, q, r) for which $\text{lh}(p') \leq \zeta + \omega + 1$ and $(p', q') \Vdash \mathbf{f}(\alpha) = \zeta$.”*

Proof. (a) Fix λ regular and so large that \mathbf{f} and the power set of $P * Q * R_i$ both belong to $H(\lambda)$. Let $N \prec H(\lambda)$ be countable with $P * Q * R_i$, $\mathbf{f} \in N$, and let $(p, q, r) \in P * Q * R \cap N$ be arbitrary; it

suffices to find an $(N, P * Q * R_i)$ -generic extension of (p, q, r) . Let $\alpha = N \cap \omega_1$ and $\zeta = \sup(N \cap \omega_2)$. Of course $\alpha \in \omega_1$ and $\zeta \in S_0^2$.

Begin by choosing $p' \leq p$ an (N, P) -generic condition as follows. Let $\langle D_k : k < \omega \rangle$ enumerate the dense subset of P contained in N , and define a descending sequence $\langle p_k : k < \omega \rangle$ of conditions in P such that $p_0 = p$ and for each k , $p_{k+1} \in D_k \cap N$. Let $p^* = \inf\{p_k : k < \omega\}$. Then $\text{lh}(p^*) = \zeta$. Let $p'(i) = p^*(i) \cup \{\zeta + \omega\}$, and for $j \neq i$ let $p'(j) = p^*(j)$. Then $p' \in P$ is (N, P) -generic, $p' \leq p$, $\text{lh}(p') = \zeta + \omega + 1$, and $p' \Vdash \text{“}\zeta \notin E(i)\text{”}$.

Since $\Vdash \text{“}Q \text{ is proper”}$, there is a term q' for a condition in Q such that $p' \Vdash \text{“}q' \leq q \text{ and } q' \text{ is } (N[E], Q)\text{-generic.} \text{”}$ Then (p', q') is an $(N, P * Q)$ -generic extension of (p, q) . It follows that $(p', q') \Vdash \text{“}N[E, G] \cap \omega_1 = \alpha \text{ and } \sup(N[E, G] \cap \omega_2^V) = \zeta \text{”}$. (Here of course G is a term denoting the $V[E]$ -generic filter on Q .) Work for the moment in an extension $V[E, G]$, where E is constructed from a V -generic filter on P containing the condition p' , G is $V[E]$ -generic on Q , and $q' \in G$. Then $N[E, G]$ is a countable elementary substructure of $H(\lambda)^{V[E, G]}$ containing R_i and r , with $N[E, G] \cap \omega_1 = \alpha$ and $\sup(N[E, G] \cap \omega_2) = \zeta \notin E(i)$. As $\mathbf{f} \in N$, $f \in N[E, G]$. It follows by elementarity that $\sup f''(\alpha) = \zeta$, and by continuity $f(\alpha) = \zeta$.

Let $\langle F_k : k < \omega \rangle$ enumerate the dense subsets of R_i belonging to $N[E, G]$. Using the elementarity of $N[E, G]$ we can choose a descending sequence $\langle r_k : k < \omega \rangle$ of conditions in R_i , so that $r_0 = r$ and $r_{k+1} \in F_k \cap N[E, G]$. Let $r^* = \bigcup\{r_k : k < \omega\} \cup \{(\alpha, \alpha)\}$. Clearly r^* is increasing and continuous, and as $f(\alpha) = \zeta \notin E(i)$, r^* belongs to R_i . Clearly r^* extends each r_k , and so r^* is an $(N[E, G], R_i)$ -generic extension of r .

Returning to V , we can find a term r' denoting r^* . Since E and G were arbitrary such that $q' \in G$ and E is the union of a generic filter containing p' , we can choose r' so that $(p', q') \Vdash \text{“}r' \text{ is } (N[E, G], R_i)\text{-generic.} \text{”}$ It follows that (p', q', r') is an $(N, P * Q * R_i)$ -generic extension of (p, q, r) , which completes the proof that $P * Q * R_i$ is proper.

(b) Let λ be as in the proof of (a) and let D be the set of all countable elementary substructures of $H(\lambda)$ having F and $P * Q * R_i$ as elements. Given $N \in D$ and $(p, q, r) \in P * Q * R_i \cap N$, let $\alpha = N \cap \omega_1$ and $\zeta = \sup(N \cap \omega_2)$, and choose (p', q', r') extending (p, q, r) as in part (a). Clearly $\text{lh}(p') = \zeta + \omega + 1$. In the proof of part (a) we saw that for any generic extension $V[E, G]$ such that p' belongs to the generic filter on P and q' belongs to G , $V[E, G] \models \text{“}f(\alpha) = \zeta \text{”}$. Hence $(p', q') \models \text{“}f(\alpha) = \zeta \text{”}$. \square

Now we can prove the main result.

THEOREM 2.6. *If $\text{PFA}^+(n)$ holds in V then $\models p$ “ $\text{PFA}^+(n)$.” Thus if $\text{PFA}^+(n)$ is consistent with ZFC then so is “ $\text{PFA}^+(n)$ + there is a family of m mutually nonreflecting stationary subsets of ω_2 .”*

Proof. Supposing the contrary, there must be a P -term Q and a condition p_0 , so that $p_0 \Vdash$ “ Q is an obnoxious partial order.” Since $\{p \in P : p \leq p_0\}$ is isomorphic to P , we may assume p_0 is the trivial condition. By Lemma 2.2 we may assume that forcing with Q collapses \aleph_2 to cardinality \aleph_1 , and find P -terms $\langle \mathbf{D}_\sigma : \sigma < \omega_1 \rangle$ and $P * Q$ -terms $\langle \mathbf{S}_i : i < n \rangle$ such that

$$\Vdash \text{“} (Q, \langle \mathbf{D}_\sigma : \sigma < \omega_1 \rangle, \langle \mathbf{S}_i : i < n \rangle \text{ is an obnoxious triple, ”}$$

and \Vdash “ \mathbf{D}_σ is dense open” for each σ . As above, let \mathbf{E} be a P -term for the mutually nonreflecting stationary family added by P , and let \mathbf{f} be a $P * Q$ -term, forced by every condition to denote the increasing enumeration of a club in ω_2^V of order type ω_1 . We may define the partial orders R_i as in the discussion preceding Lemma 2.5, and by that lemma each $P * Q * R_i$ is proper.

In any extension by $P * Q$, \mathbf{E} is a family of m pairwise disjoint sets. We claim that there is an $i^* < m$ such that $S_j - f^{-1''}\mathbf{E}(i)$ is stationary for every $j < n$. Otherwise, as $n < m$, there would be $i_0 < i_1 < m$ and $j < n$ such that both $S_j - f^{-1''}\mathbf{E}(i_0)$ and $S_j - f^{-1''}\mathbf{E}(i_1)$ were nonstationary, and so the union of these two sets would be nonstationary, but would also contain S_j , a contradiction.

We claim further that forcing with R_{i^*} preserves the stationarity of each S_j . For in the extension by $P * Q$, $S_j - f^{-1''}\mathbf{E}(i^*)$ is stationary. So if $r \Vdash$ “ \mathbf{B} is club in ω_1 ,” then we can find a countable $N \prec H(\omega_2)$ (taken in the extension by $P * Q$) containing r , R_{i^*} , and \mathbf{B} , such that $\alpha \in S_j$ but $f(\alpha) \notin \mathbf{E}(i^*)$, where $\alpha = N \cap \omega_1$. Choose $\langle r_k : k < \omega \rangle$ a descending sequence in $R_{i^*} \cap N$ such that $r_0 = r$, and for every dense subset D of R_{i^*} in N there is a k with $r_k \in D$. Let $r^* = \bigcup \{r_k : k < \omega\} \cup \{(\alpha, \alpha)\}$. Then $r^* \leq r$ and $r^* \Vdash$ “ $\alpha \in \mathbf{B}$.” So the set of conditions forcing S_j to remain stationary is dense in R_{i^*} .

In V we can find (p_0, q_0) forcing each $S_j - f^{-1''}\mathbf{E}(i^*)$ to be nonstationary. Replacing $P * Q$ with $\{(p, q) : (p, q) \leq (p_0, q_0)\}$, we may assume (p_0, q_0) is trivial. Thus each term \mathbf{S}_j is forced to denote a stationary set by every condition in $P * Q * R_{i^*}$.

Let A be the set of all conditions $(p, q, r) \in P * Q * R_{i^*}$ so that for some $\alpha < \omega_1$ and some $\zeta < \omega_2$, $\text{lh}(p) \leq \zeta + \omega + 1$ and $(p, q) \Vdash \text{“f}(\alpha) = \zeta\text{.”}$ Given any condition (p, q, r) we can find λ and D as in Lemma 2.5(b), and $N \in D$ such that $(p, q, r) \in N$. Letting $\alpha = N \cap \omega_1$ and $\zeta = \sup(N \cap \omega_2)$, we see by Lemma 2.5(b) that (p, q, r) has an extension in A . So A is dense in $P * Q * R_{i^*}$. Therefore A (with the ordering inherited from $P * Q * R_{i^*}$) is a proper partial order.

Let \mathbf{C} be the canonical term for the club adjoined by R_{i^*} . Pick λ regular and so large that $P * Q * R_{i^*}$, $\langle \mathbf{D}_\sigma : \sigma < \omega_1 \rangle$, $\langle \mathbf{S}_i : i < n \rangle$, \mathbf{f} , and \mathbf{C} all belong to $H(\lambda)$. Let $M \prec H(\lambda)$ contain all these sets, contain all countable ordinals, and have cardinality \aleph_1 . Using $\text{PFA}^+(n)$, find an M -generic filter H_0 on A such that for each $j < n$, $\mathbf{S}_j(H_0)$ is stationary. Let H be the upward closure of H_0 in $P * Q * R_{i^*}$. Then H is an M -generic filter on $P * Q * R_{i^*}$, each $\mathbf{S}_j(H)$ is stationary, and every element of H has an extension in $H \cap A$.

For $i < m$ let $p^*(i) = \bigcup \{p(i) : \exists q, r (p, q, r) \in H\}$. Let $\beta^* = \sup\{\text{lh}(p) : \exists q, r (p, q, r) \in H\}$. Clearly $p^*(i) \subseteq S_0^2 \cap \beta^*$ and for each $\beta \neq \beta^*$ there is an $i < m$ so that $p^*(i) \cap \beta$ is nonstationary in β . We shall show that $p^*(i^*) \cap \beta^*$ is nonstationary in β^* , from which it follows that p^* is in fact a condition in P . To this end, let f^* be the set of all pairs (α, ζ) so that there is a condition $(p, q, r) \in H$ such that $(p, q) \Vdash \text{“f}(\alpha) = \zeta\text{,”}$ and let C^* be the set of all $\alpha < \omega_1$ so that there is a condition $(p, q, r) \in H$ such that $(p, q, r) \Vdash \text{“}\alpha \in \mathbf{C}\text{.”}$ Since H is pairwise compatible f^* is an increasing function, and since H is M -generic f^* is continuous with domain ω_1 and C^* is club in ω_1 . Let $\zeta^* = \sup f^{*-1} \omega_1$. As $\Vdash \text{“}\mathbf{f}''\mathbf{C} \cap \mathbf{E}(i^*) = \emptyset\text{,”}$ $f^{*''}C^*$ is club in ζ^* and disjoint from $p^*(i^*)$.

We claim that $\zeta^* = \beta^*$. Assume the contrary. If $\zeta^* > \beta^*$ choose $\alpha \in C^*$ large enough that for every $(p, q, r) \in H$, $\text{lh}(p) < f^*(\alpha)$. By the definition of f^* and C^* (and pairwise compatibility of H) there are $(p, q, r) \in H$ and $\zeta < \omega_2$ so that $f^*(\alpha) = \zeta$ and $(p, q, r) \Vdash \text{“}\alpha \in \mathbf{C}$ and $\mathbf{f}(\alpha) = \zeta\text{.”}$ Now $\Vdash \text{“}\mathbf{f}''\mathbf{C} \cap \mathbf{E}(i^*) = \emptyset\text{,”}$ so $(p, q, r) \Vdash \text{“}\zeta \notin \mathbf{E}(i^*)\text{.”}$ Thus $\zeta \leq \text{lh}(p)$. But $\text{lh}(p) < f^*(\alpha) = \zeta$, a contradiction. On the other hand, if $\zeta^* < \beta^*$ choose $(p, q, r) \in H$, $i < m$, and $\eta \in p(i)$ with $\zeta^* \leq \eta$. Extending if need be, we may assume that $(p, q, r) \in A$. So there are $\alpha < \omega_1$ and $\zeta < \omega_2$ such that $\text{lh}(p) \leq \zeta + \omega + 1$ and $(p, q) \Vdash \text{“f}(\alpha) = \zeta\text{.”}$ Clearly $f^*(\alpha) = \zeta$, and as f^* is increasing, $\zeta + \omega + 1 \leq f^*(\alpha + \omega + 1) < \zeta^*$. Hence $\zeta^* \leq \eta \leq \text{lh}(p) < \zeta^*$,

another contradiction. Thus the claim is proved. Therefore p^* is a condition in P . Choose a V -generic filter on P containing p^* , and let E be the mutually nonreflecting family added by this filter. As E and the generic filter are equiconstructible, we may speak of the interpretation τ^E of an arbitrary P -term τ by E . For each $\sigma < \omega_1$, let $D_\sigma = \mathbf{D}_\sigma^E$, and for $j < n$ let $\mathbf{S}'_j = \mathbf{S}_j^E$. By assumption $(Q^E, \langle D_\sigma : \sigma < \omega_1 \rangle, \langle \mathbf{S}'_j : j < n \rangle)$ is an obnoxious triple in $V[E]$.

Let $G = \{q^E : \exists p, r (p, q, r) \in H\}$. It is not hard to show G is a filter on Q^E . For each $\sigma < \omega_1$, $\{(p, q, r) : p \Vdash "q \in \mathbf{D}_\sigma"\}$ is dense in $P * Q * R_i$ and belongs to M . So, as H is M -generic G is $\langle D_\sigma : \sigma < \omega_1 \rangle$ -generic. Now suppose $\alpha \in \mathbf{S}_j(H)$. Then for some $(p, q, r) \in H$, $(p, q) \Vdash "\alpha \in \mathbf{S}_j."$ Since $p \geq p^*$, p belongs to the V -generic filter we have chosen on P , and so $V[E] \models q^E \Vdash "\alpha \in \mathbf{S}'_j."$ Hence $\alpha \in \mathbf{S}'_j(G)$. Thus $\mathbf{S}_j(H) \subseteq \mathbf{S}'_j(G)$, and so $\mathbf{S}'_j(G)$ is stationary.

In $V[E]$ we have found G , a $\langle D_\sigma : \sigma < \omega_1 \rangle$ -generic filter on Q^E , with $\mathbf{S}'_j(G)$ stationary for each $j < n$. But this is a contradiction since $(Q^E, \langle D_\sigma : \sigma < \omega_1 \rangle, \langle \mathbf{S}'_j : j < n \rangle)$ is obnoxious. The theorem is therefore proved. \square

REMARK. A similar argument can be given with ω_2 replaced by any regular cardinal greater than ω_1 . Hence if $n < m$, $V \models \text{PFA}^+(n)$, and $\kappa > \omega_1$ is regular, then there is a countably closed, κ -Baire partial order P such that $V^P \models \text{PFA}^+(n)$ and in V^P there is a mutually nonreflecting family of m pairwise disjoint stationary subsets of $\{\alpha < \kappa : \text{cf } \alpha = \omega\}$.

3. Excluding mutually nonreflection stationary sets. In [Ba] it is shown that $\text{PFA}^+(1)$ implies that for every $\kappa \geq \omega_2$ with $\text{cf } \kappa > \omega_1$, every stationary subset of $\{\alpha < \kappa : \text{cf } \alpha = \omega\}$ reflects. Thus Theorem 2.6 has the immediate corollary that PFA (if consistent with ZFC) does not imply $\text{PFA}^+(1)$. It is simple to generalize this argument to show, for any $n < m$, that $\text{PFA}^+(n)$, if consistent, does not imply $\text{PFA}^+(m)$. We include the proof for the sake of completeness. Let Γ be the class of all countably closed partial orders.

THEOREM 3.1. *$\text{MA}^+(\Gamma, m)$ implies, for every $\kappa \geq \omega_2$ with $\text{cf } \kappa > \omega_1$, that every family of m stationary subsets of $\{\alpha < \kappa : \text{cf } \alpha = \omega\}$ is mutually reflecting.*

Proof. Suppose $\langle E_i : i < m \rangle$ is a family of stationary subsets of $\{\alpha < \kappa : \text{cf } \alpha = \omega\}$. Let P be the Levy order for collapsing κ

to cardinality \aleph_1 with countable conditions, and let f be a P -term for a continuous, increasing function from ω_1 into κ with range cofinal in κ . As P is countably closed, $\Vdash_P \forall \alpha < \omega_1 \exists g \in V. \alpha \in g$. So $D_\alpha = \{p \in P : \exists g \in V. p \Vdash \alpha \in g\}$ is dense for each α . For each $i < m$ let A_i be a P -term such that $\Vdash_P A_i = \{\alpha < \omega_1 : f(\alpha) \in E_i\}$. Since P is proper, E_i is stationary in the extension by P (by Lemma 2.4), and so A_i is forced to denote a stationary set. Applying $\text{MA}^+(\Gamma, m)$, there is a $\langle D_\alpha : \alpha < \omega_1 \rangle$ -generic filter G such that each $A_i(G)$ is stationary.

Letting $f^* = \{(\alpha, \zeta) : \exists p \in G. p \Vdash f(\alpha) = \zeta\}$, f^* is an increasing, continuous function from ω_1 into κ . Put $\eta = \sup \text{ran}(f^*)$. Note that $\text{ran}(f^*)$ is club in η . Thus for any club C in η , $f^{*-1} \cap C$ is club in ω_1 . Choosing $\alpha \in f^{*-1} \cap C \cap A_i(G)$, we have $f^*(\alpha) \in C \cap E_i$. So each $E_i \cap \eta$ is stationary, and $\langle E_i : i < m \rangle$ reflects mutually at η . \square

MM can be substituted for $\text{MA}^+(\Gamma, 1)$ in the hypothesis, as is shown by Theorem 9 of [FMS]: MM implies, for every regular $\kappa \geq \omega_2$, that every stationary subset of $\{\alpha < \kappa : \text{cf } \alpha = \omega\}$ contains a closed set of order type ω_1 (and so reflects). (It is not hard to prove the conclusion for every $\kappa \geq \omega_2$ with $\text{cf } \kappa > \omega_1$, given this theorem for regular κ .)

COROLLARY 3.2. (a) For any $n < m < \aleph_1$, $\text{PFA}^+(n)$ does not imply $\text{PFA}^+(m)$ (unless of course $\text{PFA}^+(n)$ is inconsistent).

(b) PFA does not imply MM (unless PFA is inconsistent).

Proof. (a) By Theorem 2.6, if $\text{PFA}^+(n)$ is consistent then there is a model of $\text{PFA}^+(n)$ in which there is a family of m mutually nonreflecting stationary subsets of ω_2 . Were $\text{PFA}^+(n)$ to imply $\text{PFA}^+(m)$, in this model there could be, by Theorem 3.1, no such family.

(b) Again, by theorem 2.6 (take $n = 0$ and $m = 1$), if PFA is consistent then it has a model in which there is a nonreflecting stationary subset of ω_2 . But if PFA implied MM then, by Theorem 9 of [FMS], there could be no such model. \square

REFERENCES

- [Ba] J. Baumgartner, *Applications of the proper forcing axiom*, in the *Handbook of Set-theoretic Topology* (K. Kunen and J. Vaughan, editors), North-Holland, Amsterdam, (1984), 913–959.
- [Be] R. Beaudoin, *On Uncountable Trees and Linear Orders*, doctoral dissertation, Dartmouth College (unpublished), 1984.

- [FMS] M. Foreman, M. Magidor, and S. Shelah, *Martin's maximum, saturated ideals, and non-regular ultrafilters*, part 1, *Ann. of Math.*, **127** (1988), 1–47.
- [Sh] S. Shelah, *The semi-proper forcing axiom implies Martin's maximum but not PFA^+* , *J. Symbolic Logic*, **52** (1987), 360–367.
- [To] S. Todorčević, *Trees and linearly ordered sets*, in the *Handbook of Set-theoretic Topology* (K. Kunen and J. Vaughan, editors), North-Holland, Amsterdam, (1984), 235–293.
- [V] B. Veličković, *Forcing axioms and stationary sets*, to appear.

Received August 29, 1989.

AUBURN UNIVERSITY
AUBURN UNIVERSITY, AL 36849-5310

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

R. FINN
Stanford University
Stanford, CA 94305

HERMANN FLASCHKA
University of Arizona
Tucson, AZ 85721

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906–1982)

B. H. NEUMANN

F. WOLF
(1904–1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

Takao Akahori and Harunori Ameku , On the Romanov kernel and Kuranishi's L^2 -estimate for $\bar{\partial}_b$ over a ball in the strongly pseudo convex boundary	1
Robert Emile Beaudoin , The proper forcing axiom and stationary set reflection	13
Hans Engler , A matrix Volterra integrodifferential equation occurring in polymer rheology	25
Zhong Ge , On a constrained variational problem and the spaces of horizontal paths	61
Yutaka Hemmi , Higher homotopy commutativity of H -spaces and the mod p torus theorem	95
G. D. Johnson , An intrinsic characterization of a class of minimal surfaces in constant curvature manifolds	113
Min Ho Lee , Conjugates of equivariant holomorphic maps of symmetric domains	127
Jacek Nikiel, H. Murat Tuncali and Edward D. Tymchatyn , On the rim-structure of continuous images of ordered compacta	145
Tara Lynn Smith , Generalized Clifford-Littlewood-Eckmann groups	157
Tara Lynn Smith , Generalized Clifford-Littlewood-Eckmann groups II: Linear representations and applications	185