

Pacific Journal of Mathematics

**SINGULARITY OF THE RADIAL SUBALGEBRA OF $\mathcal{L}(F_N)$
AND THE PUKÁNSZKY INVARIANT**

FLORIN G. RADULESCU

SINGULARITY OF THE RADIAL SUBALGEBRA OF $\mathcal{L}(F_N)$ AND THE PUKÁNSZKY INVARIANT

FLORIN RĂDULESCU

Let $\mathcal{L}(F_N)$ be the von Neumann algebra of the free group with N generators x_1, \dots, x_N , $N \geq 2$ and let A be the abelian von Neumann subalgebra generated by $x_1 + x_1^{-1} + \dots + x_N + x_N^{-1}$ acting as a left convolutor on $l^2(F_N)$. The radial algebra A appeared in the harmonic analysis of the free group as a maximal abelian subalgebra of $\mathcal{L}(F_N)$, the von Neumann algebra of the free group. The aim of this paper is to prove that A is singular (which means that there are no unitaries u in $\mathcal{L}(F_N)$ excepting those coming from A such that $u^*Au \subseteq A$). This is done by showing that the Pukánszky invariant of A is infinite, where the Pukánszky invariant of A is the type of the commutant of the algebra \mathcal{A} in $B(l^2(F_N))$ generated by A and $x_1 + x_1^{-1} + \dots + x_N + x_N^{-1}$ regarded also as a right convolutor on $l^2(F_N)$.

1. Introduction. Let M be a type II_1 factor with trace τ , $\tau(1) = 1$ and $A \subseteq M$ a maximal abelian von Neumann subalgebra (briefly a M.A.S.A.). Following J. Dixmier [1], let $N_M(A) = \{u \in M \mid u \text{ unitary, } uAu^* = A\}$ be the normalizer of A in M and $B = N_M(A)''$ the von Neumann subalgebra generated by $N_M(A)$ in M . According to the size of B in M , A is called singular if $B = A$ and A is called regular (or Cartan) if $B = M$. While examples of regular M.A.S.A.'s are readily available by the classical group measure space construction from a free action of a discrete group on a measure space, examples of singular M.A.S.A.'s are more difficult to obtain (see, e.g., [1], [6], [9], [10], [5]).

The aim of this paper is to show that in the von Neumann algebra $M = \mathcal{L}(F_N)$ of the free group with N generators X_1, X_2, \dots, X_N , the radial algebra (i.e. the abelian von Neumann subalgebra generated by $X_1 + \dots + X_N + X_1^{-1} + \dots + X_N^{-1}$) is singular. This algebra has been studied intensively in [2], [3], [7] because of its connections with the problem of computing spectra of convolutors and with representation theory of F_N . In particular in [7] it is shown that the radial algebra is a M.A.S.A. in $\mathcal{L}(F_N)$.

To prove our result we need in fact to prove more than the singularity of A . In order to express this we recall some definitions.

Let $\|X\|_2 = \tau(X^*X)^{1/2}$ be the Hilbert norm given by τ on M , let $L^2(M, \tau)$ be the completion of M with respect to this norm so that M acts (in the standard way) on $L^2(M, \tau)$. Let also $J: L^2(M, \tau) \mapsto L^2(M, \tau)$ be the canonical conjugation (given by $Jx = x^*$ for x in M) and let $\mathcal{A} = (A \vee JAJ)''$ be the (abelian) von Neumann subalgebra generated in $B(L^2(M, \tau))$ by A and JAJ . Since any automorphism of M is unitarily implemented on $L^2(M, \tau)$, it follows that the type of the algebra \mathcal{A}' is an invariant for A . This invariant was considered by Ambrose-Singer and also by Pukánszky in [6]. Moreover the latter showed that in the hyperfinite factor there are singular M.A.S.A.'s, A_n such that the corresponding \mathcal{A}_n 's are of the homogeneous type I_n on $I_{B(L^2(M, \tau))} - p_1$ (where p_1 is the cyclic projection onto $\overline{A}^{\|\cdot\|_2} \subseteq L^2(M, \tau)$).

The link between this invariant and the classification of M.A.S.A.'s recalled at the beginning is given by a result of S. Popa ([4]); \mathcal{A} is maximal abelian whenever A is a Cartan M.A.S.A. and (consequently) if \mathcal{A} is of the homogeneous type I_n ($n \geq 2$) on $I_{B(L^2(M, \tau))} - p_1$, then A is singular.

We prove that for the radial algebra this invariant is infinite (for each N) and therefore A is singular.

In an earlier version of this paper the proofs were very complicated. I am greatly indebted to Florin Boca who simplified the proofs by noticing the relations in Lemma 1. Also the actual proof of Lemma 5 is due to him.

2. Singularity of the radial algebra. Let $N \geq 2$ be an integer and F_N be the free group with N generators X_1, X_2, \dots, X_N . Let $M = \mathcal{L}(F_N)$ be the associated von Neuman algebra (which is the weakly closed subalgebra of $B(l^2(F_N))$ generated by the left convolution operators on $l^2(F_N)$). It is well known that $\mathcal{L}(F_N)$ is a type II_1 factor that acts standardly on $l^2(F_N)$, (see [8]) and that with this identification the norm $\|\cdot\|_\tau$ coincides with the usual norm $\|\cdot\|_2$ on $l^2(F_N)$.

By $M_0 = \mathbb{C}[F_N]$ we denote the group ring of F_N over \mathbb{C} , viewed as a subalgebra of M (thus M_0 is the linear space of all finite sums $\sum_{w \in F_N} \lambda_w \cdot w$ (where λ_w are complex numbers) endowed with the usual product structure). Since the void word \emptyset (the unity of F_N) coincides with the unity of $\mathcal{L}(F_N)$, we shall simply write 1 instead of λ_\emptyset . By means of the identification of M with a subspace of $L^2(M, \tau)$, M_0 corresponds to the subspace of finite support sequences in $l^2(F_N)$.

Recall that the canonical length function $|\cdot|$ on F_N is defined by $|w| = |\alpha_1| + |\alpha_2| + \cdots + |\alpha_p|$ if $w \in F_N$ has the reduced form $X_{i_1}^{\alpha_1} \cdots X_{i_p}^{\alpha_p}$ ($i_j \neq i_{j+1}$).

For any integer $n \geq 0$ we denote by $M_0^n \subseteq M_0$ the linear span of all words in F_N of length n and by q_n we denote the projection of $l^2(F_N)$ onto M_0^n ; χ_n will be the vector in M_0^n defined by

$$\chi_n = \sum_{|w|=n} w.$$

In [2], [3], [7] it is proved that

$$A = \overline{\text{Sp}\{\chi_n | n \geq 0\}}^w \subseteq M$$

(the closure being taken with respect to the weak topology on M) is an abelian von Neumann subalgebra of $\mathcal{L}(F_N)$ (called the radial algebra); this in fact a consequence of the following recurrence relations:

$$(1) \quad \begin{aligned} \chi_1 \cdot \chi_1 &= \chi_2 + 2N, \\ \chi_1 \cdot \chi_n &= \chi_n \chi_1 = \chi_{n+1} + (2N - 1)\chi_{n-1}, \quad n \geq 1. \end{aligned}$$

Moreover, by [7], A is a M.A.S.A. in M .

Let as before $\mathcal{A} = (A \vee JAJ)''$ be the (abelian) von Neumann subalgebra of $B(l^2(F_N))$ generated by A and JAJ and for any ξ in $l^2(F_N)$ let p_ξ denote the cyclic projection of $l^2(F_N)$ onto $\overline{\text{Sp}\mathcal{A}\xi}^{\|\cdot\|_2} = \overline{\text{Sp}A\xi A}^{\|\cdot\|_2}$. Our aim is to show that \mathcal{A}' is of the homogeneous type I_∞ on $I - p_1$ (where $1 \in M \subseteq L^2(M, \tau)$) and to do this we will construct an infinite family of vectors $\{\xi_n\}_n$ in $l^2(F_N)$ such that their corresponding cyclic projections $\{p_{\xi_n}\} \in \mathcal{A}'$ are mutually orthogonal and of central support $I - p_1$ in \mathcal{A}' .

In proving this we will need also to consider for each integer $n \geq 0$ the projections P_n onto $\overline{\text{Sp}\{AWA | w \in F_N, |w| \leq n\}}^{\|\cdot\|_2}$ and the linear subspaces $S_n \subseteq M_0^n$ spanned by $\{q_n(\chi_1 w), q_n(w \chi_1), w \in F_N, |w| \leq n - 1\}$ (so that $S_0 = (0)$). We will later identify S_n with the range of the projection $P_{n-1} \wedge q_n$. Note that with the notation before $p_1 = P_0$.

To describe the structure of the cyclic projections associated with an arbitrary vector ξ in M_0^l , $l \geq 1$, we will also use the following notation: for all r, s integers let

$$\xi_{r,s} = q_{r+s+1}(\chi_r \xi \chi_s) \quad \text{if } r, s \geq 0$$

and if r or s are strictly negative, let $\xi_{r,s} = 0$.

Our first purpose is to show that for every γ in $M_0^l \ominus S_l$, $l \geq 1$ the projection p_γ commutes with all q_n , $n \geq 0$ and to describe the range of $p_\gamma \wedge q_n$. To do this we need first the following lemma which is analogous to the recurrence relations (1).

LEMMA 1. (a) If γ is in M_0^l , $l \geq 1$, then

$$\begin{aligned}\chi_1 \gamma_{r,s} &= \gamma_{r+1,s} + (2N-1)\gamma_{r-1,s} \quad \text{for } r \geq 1, s \geq 0, \\ \gamma_{r,s} \chi_1 &= \gamma_{r,s+1} + (2N-1)\gamma_{r,s-1} \quad \text{for } r \geq 0, s \geq 1.\end{aligned}$$

(b) If γ is in $M_0^l \ominus S_l$ for some $l \geq 2$ then the relations in (a) are also true for $r, s \geq 0$, i.e. we have

$$\chi_1 \gamma_{0,s} = \gamma_{1,s}; \quad \gamma_{s,0} \chi_1 = \gamma_{s,1}, \quad s \geq 0.$$

(c) If $\gamma = \sum_{|X|=1} c_X \cdot X$ belongs to $M_0^1 \ominus S_1$ and $\varepsilon \in \{\pm 1\}$ is such that $c_{(X^{-1})} = \varepsilon c_X$ for all X in F_N with $|X| = 1$ then

$$\begin{aligned}\chi_1 \gamma_{0,s} &= \gamma_{1,s} - \varepsilon \gamma_{0,s-1}, \quad s \geq 0, \\ \gamma_{s,0} \chi_1 &= \gamma_{s,1} - \varepsilon \gamma_{s-1,0}, \quad s \geq 0.\end{aligned}$$

(d) Finally if γ belongs to $M_0^l \ominus S_l$, $l \geq 1$, then

$$\gamma \chi_s = \gamma_{0,s} - \gamma_{0,s-2}, \quad s \geq 0.$$

Proof. (a) is proved exactly as the relations (1) are proved in [2] and (b), (c) are proved by similar arguments. For example, if γ is as in the statement of (c) and $s \geq 1$ an integer, then

$$\begin{aligned}\chi_1 \gamma_{0,s} &= \gamma_{1,s} + \sum_{|w|=s} \left(\sum_{X \neq (f(w))^{-1}} c_X \right) \cdot w \\ &= \gamma_{1,s} - \sum_{|w|=s} (c_{(f(w))^{-1}}) \cdot w = \gamma_{1,s} - \varepsilon \sum_{|w|=s} (c_{f(w)}) w \\ &= \gamma_{1,s} - \varepsilon \gamma_{0,s-1},\end{aligned}$$

where by $f(w)$ we denoted the first letter of w and where we used the equality $\sum_{|X|=1} c_X = 0$ which follows from the fact that γ being orthogonal to S_1 it is also orthogonal on χ_1 .

Finally (d) is proved by induction from the preceding relations.

The next lemma shows that for γ in $M_0^l \ominus S_l$, p_γ commutes with q_n and range $p_\gamma \wedge q_n = \text{Sp}\{\gamma_{r,s}; r+s=n-1\}$ for all $n \geq 1$.

LEMMA 2. Suppose γ is in $M_l^0 \oplus S_l$, $l \geq 2$, ε in $\{\pm 1\}$ and β in $M_0^1 \oplus S_1$, $\beta = \sum c_X$ such that $c_X = \varepsilon c_{(X^{-1})}$ for $|X| = 1$. Then for all $n, m \geq 0$

- (a) $\chi_n \gamma \chi_m = \gamma_{n,m} - (\gamma_{n,m-2} + \gamma_{n-2,m}) + \gamma_{n-2,m-2}$,
 (b)

$$\chi_n \beta \chi_m = \beta_{n,m} - (\beta_{n-2,m} + \beta_{n,m-2} + \varepsilon \beta_{n-1,m-1}) + \sum_{k \geq 2} (-\varepsilon)^k (\varepsilon \beta_{n-k-1,m-k+1} + \varepsilon \beta_{n-k+1,m-k-1} + 2\beta_{n-k,m-k}),$$

(c) $\gamma_{n,m} = \sum_{r \leq n, s \leq m} \chi_r \gamma \chi_s$ where (r, s) runs over all possible values such that r, s have the same parity as n, m respectively.

(d) $\beta_{n,m} = \sum_{r \leq n, s \leq m} (\varepsilon)^{n-r} \chi_r \beta \chi_s$ where (r, s) runs over all possible values such that $r - s$ has the same parity as $n - m$.

In particular $\text{Sp}((\chi_n \gamma \chi_m)_{n,m \geq 0}) = \text{Sp}((\gamma_{n,m})_{n,m \geq 0})$ and similarly for β .

Proof. We will prove only (b), (d) since (a), (c) can be proved in a similar (but easier) way.

To prove (b) note that the case $n = 0$ follows straight from the preceding lemma (point (d)) and hence we can proceed with the proof by induction according to n . Assume that we have already carried the induction up to n and hence we have to compute $\chi_{n+1} \beta \chi_m$. For the sake of simplicity we assume $n \geq 2$ and by the use of the second relation from (1) (when $n = 1$ we have to use the first) we get for any $m \geq 0$

$$\begin{aligned} \chi_{n+1} \beta \chi_m &= (\chi_1 \cdot \chi_n - (2N - 1)\chi_{n-1}) \beta \chi_m \\ &= \chi_1 (\chi_n \beta \chi_m) - (2N - 1)\chi_{n-1} \beta \chi_m. \end{aligned}$$

In order to describe the proof we introduce the following linear operators L and M on $\text{Sp}(\beta_{n,m})$ defined by $L(\beta_{n,m}) = \beta_{n+1,m}$ and

$$M(\beta_{n,m}) = \begin{cases} (2N - 1)\beta_{n-1,m} & \text{if } n \geq 1, \\ (-\varepsilon)\beta_{0,m-1} & \text{if } n = 0, \end{cases}$$

so that by Lemma 1.c

$$\chi_{n+1} \beta \chi_m = (L + M)(\chi_n \beta \chi_m) - (2N - 1)\chi_{n-1} \beta \chi_m.$$

One easily observes that $L(\chi_n \beta \chi_m)$ contains all the required terms in the expansion of $\chi_{n+1} \beta \chi_m$ except those of the form $\beta_{0,r}$. But on the other hand $M(\chi_n \beta \chi_m)$ gives $(2N - 1)\chi_{n-1} \beta \chi_m$ (by the induction

hypothesis) plus the above-mentioned missing terms. This completes the proof of (b). Similarly (d) is proved by using

$$\beta_{n+1,m} = \chi_1 \beta_{n,m} - (2N - 1) \beta_{n-1,m}, \quad n \geq 1.$$

The next lemma shows in particular that whenever γ, γ' are orthogonal vectors in $M_0^l \ominus S_l$ the associated cyclic projections are orthogonal.

LEMMA 3. *Let γ, γ' be vectors in $M_0^l \ominus S_l, l \geq 2$. Corresponding to $\varepsilon, \varepsilon' \in \{\pm 1\}$ as in the statement of the preceding lemma, let β, β' be in $M_0^1 \ominus S_1$. Then for any $n, n', m, m' \geq 0$*

(a) $\langle \gamma_{n,m}, \gamma_{n',m'} \rangle_2 = \delta_{n,n'} \delta_{m,m'} (2N - 1)^{n+m} \langle \gamma, \gamma' \rangle_2$.

(b)

$$\begin{aligned} \langle \beta_{n,m}, \beta_{n',m'} \rangle_2 &= \delta_{\varepsilon,\varepsilon'} \delta_{n+m,n'+m'} (2N - 1)^{n+m} \\ &\quad \cdot (-\varepsilon(2N - 1))^{-|n-n'|} \langle \beta, \beta' \rangle_2, \end{aligned}$$

(where by $\langle \cdot, \cdot \rangle_2$ we denote the scalar product on $l^2(F_N)$ and δ_{ij} is the Kronecker symbol: δ_{ij} is nonzero only if $i = j$ and in this case $\delta_{ii} = 1$).

Proof. It is obvious that both sides in (a) and (b) vanish if $n + m \neq n' + m'$. Hence we will assume that $n + m = n' + m' = k$. Let first ξ, ξ' be elements in $M_0^l \ominus S_l, l \geq 1$; using (1) and the fact that $q_l(\chi_n \cdot \xi_{n',m'} \cdot \chi_{m-2}) = 0$ we deduce

$$q_l(\chi_n \xi_{n',m'} \chi_m) = q_l(\chi_n \cdot \xi_{n',m'} \cdot \chi_1 \cdot \chi_{m-1})$$

whenever $m \geq 1$ (and a similar relation to the left of ξ). Using this and Lemma 1 (and since $q_l(\chi_n \xi_{n',m'+1} \chi_{m-1}) = 0$) we obtain

$$q_l(\chi_n \xi_{n',m'} \chi_m) = (2N - 1) q_l(\chi_n \xi_{n',m'-1} \chi_{m-1}), \quad \text{for } m, m' \geq 1,$$

$$q_l(\xi_{s,0} \cdot \chi_s) = \begin{cases} 0 & \text{if } \xi = \gamma \text{ for } s \geq 1, \\ (-\varepsilon) q_l(\xi_{s-1,0} \chi_{s-1}) & \text{if } \xi = \beta, \text{ for } s \geq 1. \end{cases}$$

(and similar relations to the left of ξ). Assuming $n \geq n'$ (so $m \leq m'$) we obtain by induction

$$q_l(\chi_n \gamma_{n',m'} \chi_m) = \delta_{n,n'} (2N - 1)^{n+m'} \gamma,$$

$$q_l(\chi_n \beta_{n',m'} \chi_m) = (2N - 1)^{n+m'} (-\varepsilon)^{|n-n'|} \beta.$$

The proof is now accomplished by noticing that

$$\begin{aligned} \langle \xi'_{n',m'}, \xi_{n,m} \rangle_2 &= \langle \xi'_{n',m'}, \chi_n \xi \chi_m \rangle_2 \\ &= \langle \chi_n \xi'_{n',m'} \chi_m, \xi \rangle_2 = \langle q_l(\chi_n \xi'_{n',m'} \chi_m), \xi \rangle_2. \end{aligned}$$

We can now show that the space S_l coincides with the range of the projection $P_{l-1} \wedge q_l$ and deduce that P_{l-1} and q_l are commuting projections. As a corollary we will obtain that the cyclic projections corresponding to two orthogonal vectors in $\text{Sp}(M_0^l \ominus S_l \mid l \geq 0)$ are orthogonal:

LEMMA 4. (a) *The projections P_{l-1} and q_l commute and S_l is the range of $P_{l-1}q_l = P_{l-1} \wedge q_l$ (so that $M_0^l \ominus S_l$ is the range of $q_l \ominus P_{l-1}$) for all $l \geq 0$.*

(b) *If ξ_i belongs to $\text{Sp}(M_0^l \ominus S_l \mid l \geq 0)$, $i = 1, 2$, and $\langle \xi_1, \xi_2 \rangle_2 = 0$ then p_{ξ_1} and p_{ξ_2} are orthogonal.*

(c) *If $p = 2N - 1$, $\alpha_l = \dim(q_l \ominus P_{l-1})$, $l \geq 0$, then $\alpha_0 = 1$, $\alpha_1 = p$, $\alpha_2 = p^2 - p - 1$ and $\alpha_l = p^{l-3}(p-1)^2(p+1)$ for $l \geq 3$. In particular $q_l \ominus P_{l-1}$ is nonnull for all l .*

Proof. (b) is an easy consequence of (a) and of the preceding lemma. (a) is obvious for $l = 0$ since $P_0 = 0$ and in general it will be proved by induction according to l . Assuming that we have carried the induction up to l , we prove (a) for $(l+1)$ instead of l . First we prove that for each w in F_N , $|w| \leq l$,

$$(3) \quad q_{l+1}(\chi_p w \chi_q) \in S_{l+1} \quad \text{for all } p, q \geq 0.$$

By the induction hypothesis it is sufficient to show that for any $k \leq l$ and any γ in $M_0^k \ominus S_k$,

$$q_{l+1}(\chi_p \gamma \chi_q) \text{ belongs } S_{l+1} \quad \text{for all } p, q \geq 0.$$

This means (by Lemma 2) that we have to show that $\{\gamma_{p,q}\}_{-p+q=l+1-k}$ is contained in S_{l+1} . But this follows from the fact that whenever $p+q = l-k$, $q_{l+1}(\chi_1 \gamma_{p,q}) = \gamma_{p+1,q}$ and $q_{l+1}(\gamma_{p,q} \chi_1) = \gamma_{p,q+1}$ (by Lemma 1). Hence (3) is true and it follows that

$$\text{range } q_{l+1}P_l \subseteq S_{l+1} \subseteq \text{range } P_l \wedge q_{l+1}$$

(the second inclusion being obvious). Hence P_l commutes with q_{l+1} and S_{l+1} equals the range of $q_{l+1}P_l$.

To prove (c) note that by Lemma 2 for any γ in $M_0^l \ominus S_l$, $l \geq 1$, p_γ commutes with all q_n and $\dim(p_\gamma \wedge q_n) = n - l + 1$ for $n \geq l$ (and zero for $n < l$). Hence the following formula holds for $l \geq 1$:

$$\alpha_l = p^{l-1}(p+1) - (1 + l\alpha_1 + (l-1)\alpha_2 + \cdots + 2\alpha_{l-1})$$

and by an elementary induction argument we get (c).

The following computational lemma will be used only in the next lemma. Probably it is known but for the sake of completeness we include its proof here.

LEMMA 5. *Let a be a real number with $|a| < 1$. There are strictly positive numbers B, C depending on a such that for any integer $k \geq 0$ and any $\lambda_0, \dots, \lambda_k$ in \mathbb{C}*

$$B \left(\sum |\lambda_i|^2 \right) \leq \sum_{i,j} \lambda_i \bar{\lambda}_j a^{|i-j|} \leq C \left(\sum_i |\lambda_i|^2 \right).$$

Proof. Let A be the linear operator on \mathbb{C}^{k+1} given by the matrix $a_{ij} = a^{|i-j|}$, $i, j = 0, 1, \dots, k$, and let N_0 be the nilpotent operator given by $n_{ij} = \delta_{i,j+1}$ and D the diagonal operator with only nonnull entries $d_{00} = d_{kk} = 1$. Then

$$A = I + a(N_0 + N_0^*) + \dots + a^k(N_0^k + (N_0^*)^k)$$

and an elementary computation shows that A is invertible and

$$A^{-1} = (1 - a^2)^{-1}((1 + a^2)I - a(N_0 + N_0^*) - a^2D).$$

Hence

$$\begin{aligned} \|A\| &\leq (1 + |a|)(1 - |a|)^{-1} = C, \\ \|A^{-1}\| &\leq (1 - a^2)^{-1}(1 + 2|a| + 2a^2) = B^{-1} \end{aligned}$$

and the lemma follows now from the inequality

$$\|A^{-1}\|^{-1} \|\xi\|^2 \leq \langle A\xi, \xi \rangle \leq \|A\| \|\xi\|^2, \quad \xi \in \mathbb{C}^{k+1}$$

which holds true since A is positive definite.

Finally start with nonnull vectors β in $M_0^1 \ominus S_1$ and γ in $M_0^l \ominus S_l$, $l \geq 2$. To prove that the cyclic projections p_β and p_γ have the same central support in \mathcal{A}' it is clearly sufficient to show that the linear mapping T_0 defined by the requirement $T_0(\chi_n \beta \chi_m) = \chi_n \gamma \chi_m$ is well defined and extends to a bounded invertible operator from $\overline{\text{Sp}(\chi_n \beta \chi_m)}^{\|\cdot\|_2}$ onto $\overline{\text{Sp}(\chi_n \gamma \chi_m)}^{\|\cdot\|_2}$. This is done in the following lemma.

LEMMA 6. *Let β and γ be as in the statement of Lemma 2. Then the linear mapping T_0 on $\text{Sp}\{\chi_n \beta \chi_m\}$ into $\text{Sp}\{\chi_n \gamma \chi_m\}$ defined by $T_0(\chi_n \beta \chi_m) = \chi_n \gamma \chi_m$ is well defined and extends to a bounded linear*

operator from $\overline{\text{Sp}(\chi_n \beta \chi_m)}^{\|\cdot\|_2}$ onto $\overline{\text{Sp}(\chi_n \gamma \chi_m)}^{\|\cdot\|_2}$. In particular p_β and p_γ have the same central support in \mathcal{A}' .

Proof. Let $T'_0: \text{Sp}(\chi_n \beta \chi_m) \mapsto \text{Sp}(\chi_n \gamma \chi_m)$ be the linear map defined by requiring $T'_0 \beta_{n,m} = \gamma_{n,m}$, and let $S: \text{Sp}\{\chi_n \gamma \chi_m\} \rightarrow \{\text{Sp} \chi_n \gamma \chi_m\}$ be defined by $S \gamma_{n,m} = \gamma_{n-1,m-1}$. By the preceding lemma and by Lemma 3, T'_0 extends to a bounded invertible operator (also denoted by T'_0) from $\overline{\text{Sp}(\chi_n \beta \chi_m)}^{\|\cdot\|_2}$ into $\overline{\text{Sp}\{\chi_n \gamma \chi_m\}}^{\|\cdot\|_2}$. Also by Lemma 3, we have $\|S\| \leq (2N-1)^{-2}$ so that $I_{\overline{\text{Sp}\{\chi_n \gamma \chi_m\}}^{\|\cdot\|_2}} + S$ is also bounded and invertible. Hence the linear map (which is the composite map $T'_0(I+S)$)

$$\beta_{n,m} \mapsto \gamma_{n,m} + \varepsilon \gamma_{n-1,m-1}$$

extends to a bounded invertible operator from $\overline{\text{Sp}\{\chi_n \beta \chi_m\}}^{\|\cdot\|_2}$ into $\overline{\text{Sp}\{\chi_n \gamma \chi_m\}}^{\|\cdot\|_2}$ which by Lemma 2(b) and (d) coincides with T_0 . This completes the proof of the lemma.

We can now state and prove our main result.

THEOREM 7. *Let $A = \{X_1 + \cdots + X_N + X_1^{-1} + \cdots + X_N^{-1}\}''$ be the radial algebra in $\mathcal{L}(F_N)$. Let $\mathcal{A} = (A \vee JAJ)''$ be the (abelian) von Neumann algebra generated in $B(l^2(F_N))$ by A and JAJ . Then $\mathcal{A}' \subseteq B(l^2(F_N))$ is of the homogeneous type I_∞ on $I_{B(l^2(F_N))} - p_1$ (where p_1 is the cyclic projection from $l^2(F_N)$ onto $\overline{A}^{\|\cdot\|_2}$). In particular (by [4]) A is singular.*

Proof. For each $l \geq 0$ take a basis $\{\xi_{i,l}\}_i$ of $M_0^l \ominus S_l$ with $\xi_{0,0} = 1 \in M_0^0$, and when $l = 1$ we have to choose the vectors $\xi_{i,1}$ to be like in Lemma 2 for some $\varepsilon = \pm 1$ (which is always possible). By Lemma 6 $\{p_{\xi_{i,l}}\}_{i,l}$ have all the same central support in \mathcal{A}' . By Lemma 4, $\sum_{l \geq 1} p_{\xi_{i,l}} = I - p_1$ and hence, since $p_1 \in \mathcal{A}$ (by Lemma 3.1 in [4]), it follows that the central support of $p_{\xi_{i,l}}$ in \mathcal{A}' is $l - p_1$ for each $l \geq 1$. Since by Lemma 4 the family $\{p_{\xi_{i,l}}\}_{l \geq 1}$ is infinite, the theorem is proved.

REFERENCES

- [1] J. Dixmier, *Sous-anneaux abeliens maximaux dans les facteurs de type fini*, Ann. Math., **59** (1954), 279–286.
- [2] J. M. Cohen, *Operator norms on free groups*, Boll. Un. Mat. Ital., A1, 1982.

- [3] A. Figà-Talamanca and M. Picardello, *Harmonic Analysis on Free Groups*, Lecture Notes in Pure and Appl. Math., Vol. 87, Dekker, New York, 1983.
- [4] S. Popa, *Notes on Cartan subalgebras in type II_1 factors*, Math. Scand., **57** (1985), 171–188.
- [5] —, *Singular maximal Abelian *-subalgebras in continuous von Neumann algebras*, J. Funct. Anal., **50**, No. 2 (1985), 151–165.
- [6] L. Pukánszky, *On maximal abelian subalgebras of factors of type II_1* , Canad. J. Math., **12** (1960), 289–296.
- [7] T. Pytlík, *Radial functions on free groups and a decomposition of the regular representation into irreducible components*, J. Reine Angew. Math., **326** (1981), 124–135.
- [8] S. Strătilă, *Modular Theory in Operator Algebras*, Editura Academiei, Bucharest, Abacus Press, 1981.
- [9] M. Takesaki, *On the unitary equivalence among the components of decompositions of representations of involutive Banach algebras and the associated diagonal algebras*, Tohoku Math. J., **15** (1963), 365–393.
- [10] R. I. Tauer, *Maximal abelian subalgebras in finite factors of type II* , Trans. Amer. Math. Soc., **114** (1965), 281–308.

Received January 5, 1990 and in revised form October 1, 1990.

NATIONAL INSTITUTE FOR SCIENTIFIC AND TECHNICAL CREATION

Bd. PĂCII 220

79622 BUCHAREST, ROMANIA

Current address: University of California
Los Angeles, CA 90024-1555

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721

R. FINN
Stanford University
Stanford, CA 94305

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH
(1906-1982)

B. H. NEUMANN

F. WOLF
(1904-1989)

K. YOSHIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1991 by Pacific Journal of Mathematics

Michael G. Eastwood and A. M. Pilato , On the density of twistor elementary states	201
Brian E. Forrest , Arens regularity and discrete groups	217
Yu Li Fu , On Lipschitz stability for F.D.E	229
Douglas Austin Hensley , The largest digit in the continued fraction expansion of a rational number	237
Uwe Kaiser , Link homotopy in \mathbb{R}^3 and S^3	257
Ronald Leslie Lipsman , The Penney-Fujiwara Plancherel formula for abelian symmetric spaces and completely solvable homogeneous spaces	265
Florin G. Radulescu , Singularity of the radial subalgebra of $\mathcal{L}(F_N)$ and the Pukánszky invariant	297
Albert Jeu-Liang Sheu , The structure of twisted $SU(3)$ groups	307
Morwen Thistlethwaite , On the algebraic part of an alternating link	317
Thomas (Toma) V. Tonev , Multi-tuple hulls	335
Arno van den Essen , A note on Meisters and Olech's proof of the global asymptotic stability Jacobian conjecture	351
Hendrik J. van Maldeghem , A characterization of the finite Moufang hexagons by generalized homologies	357
Bun Wong , A note on homotopy complex surfaces with negative tangent bundles	369
Chung-Tao Yang , Any Blaschke manifold of the homotopy type of CP^n has the right volume	379