

# Pacific Journal of Mathematics

## **LUSTERNIK-SCHNIRELMANN INVARIANTS IN PROPER HOMOTOPY THEORY**

R. AYALA, ELADIO DOMÍNGUEZ MURILLO,  
ALBERTO MÁRQUEZ PÉREZ AND A. QUINTERO

## LUSTERNIK-SCHNIRELMANN INVARIANTS IN PROPER HOMOTOPY THEORY

R. AYALA, E. DOMÍNGUEZ, A. MÁRQUEZ, AND A. QUINTERO

**We introduce and study proper homotopy invariants of the Lusternik-Schnirelmann type,  $\text{p-cat}(-)$ ,  $\text{p-Cat}(-)$ , and  $\text{cat } \varepsilon(-)$  in the category of  $T_2$ -locally compact spaces and proper maps. As an application,  $\mathbb{R}^n$  ( $n \neq 3$ ) is characterized as (i) the unique open manifold  $X$  with  $\text{p-Cat}(X) = 2$ , or (ii) the unique open manifold with one strong end and  $\text{p-cat}(x) = 2$ .**

**Introduction.** The category  $\text{cat}(X)$  of a space  $X$  in the sense of Lusternik and Schnirelmann (L-S category) is the smallest number  $k$  such that there exists an open covering  $\{X_1, \dots, X_k\}$  of  $X$  for which each inclusion  $X_j \subseteq X$  is nullhomotopic in  $X$ . This concept was introduced by the quoted authors in their studies on calculus of variations [16] and they used it as a lower bound for the number of critical points of a differentiable real function on a manifold. The basic work on the homotopical significance of  $\text{cat}$  is due to Borsuk (see [5]). Borsuk's work was continued by Fox [10].

Here we present the definition and the basic properties of a new numerical topological invariant for  $T_2$ -locally compact spaces which agrees with the notion of L-S category for  $T_2$ -compact spaces. This invariant, denoted  $\text{p-cat}(X)$ , is called the proper L-S category of  $X$  and turns out to be a proper homotopy invariant of  $X$ . Hence,  $\text{p-cat}(X)$  is a finer invariant than  $\text{cat}(X)$ .

In [10] several generalizations of L-S category are suggested. More explicitly, a general notion of L-S  $\mathcal{A}$ -category with respect to a class  $\mathcal{A}$  of spaces has been developed by Puppe and Clapp in [6]. Our work shares some common points with [6] but does not fit into the notion of L-S  $\mathcal{A}$ -category since we entirely deal with proper maps instead of ordinary continuous maps.

Another generalization of L-S category has been given in [1], where L-S category for pro-objects in  $\text{pro-Top}$  is defined. This idea is related to proper L-S category by the Edwards-Hastings embedding (see [8]) which provides a close link between proper homotopy theory and homotopy in  $\text{pro-Top}$ .

We shall work entirely in the category  $\mathfrak{P}_\infty$  of non-compact  $T_2$ -locally compact  $\sigma$ -compact spaces and proper maps. Notice that any  $T_2$ -compact space  $X$  can be regarded in  $\mathfrak{P}_\infty$  as the wedge  $X \vee J$ , where  $J$  will stand for the half-line  $[0, \infty)$ .

We recall that a proper map (p-map) is a continuous map  $f: X \rightarrow Y$  such that  $f^{-1}(K)$  is compact for each compact  $K \subseteq Y$ . Proper homotopy (p-homotopy), proper deformation (p-deformation), etc. can be defined in the natural way. The symbols “ $\simeq$ ”, “ $\simeq_p$ ” and “ $\cong$ ” stand for homotopy equivalence, p-homotopy equivalence and homeomorphism respectively.

Finally we also recall the notion of end in proper homotopy. A Freudenthal end of  $X \in \mathfrak{P}_\infty$  is an element of the inverse limit  $\mathcal{F}(X) = \varprojlim \pi_0(X - K)$  where  $K$  ranges over the family of compact subsets of  $X$  and  $\pi_0$  stands for the set of connected components. The topology of  $X$  can be enlarged to a topology on  $X \cup \mathcal{F}(X)$  in such a way that  $\mathcal{F}(X)$  turns out to be homeomorphic to a closed set of the Cantor set (see [11] for details). It is easy to check that any p-map  $f: X \rightarrow Y$  induces a continuous map  $f_*: \mathcal{F}(X) \rightarrow \mathcal{F}(Y)$  such that  $f_*$  is a homeomorphism when  $f$  is a p-homotopy equivalence.

**1. Basic properties.** In [8] the following lemma is proven,

1.1. LEMMA ([8; 6.3.5]). *Any space  $X$  in  $\mathfrak{P}_\infty$  admits a p-map  $r: X \rightarrow J$  unique up to p-homotopy.*

1.2. DEFINITION. A closed subset  $C \subseteq X$  is said to be properly deformable to  $J$  in  $X$  if there exists a diagram in  $\mathfrak{P}_\infty$

$$\begin{array}{ccc} C & \hookrightarrow & X \\ r \downarrow & & \uparrow \alpha \\ J & \longleftarrow & \end{array}$$

commutative up to p-homotopy. Notice that we may use  $r$  as the restriction of a p-map  $X \rightarrow J$  given by Lemma 1.1.

1.3. DEFINITION. Given a space  $X$  in  $\mathfrak{P}_\infty$ ,  $A \subseteq X$  is said to be properly categorical (p-categorical) in  $X$  if there is a closed neighbourhood of  $A$  properly deformable to  $J$  in  $X$ .

An open covering  $\{U_\alpha\}$  of  $X$  is said to be p-categorical if each  $U_\alpha$  is p-categorical in  $X$  (i.e.  $\overline{U}_\alpha$  is properly deformable to  $J$  in  $X$ ).

The p-category of  $X$ ,  $\text{p-cat}(X)$ , is the least number  $n$  such that  $X$  admits a p-categorical open covering with  $n$  elements. If no finite p-categorical covering exists then  $\text{p-cat}(X) = \infty$ .

LEMMA 1.4 [4; 3.4]. *Let  $P$  be a locally compact metrizable space and  $Q$  a locally compact ANR. Suppose  $X$  is a closed subset of  $P$  and  $f, g: X \rightarrow Q$  are  $p$ -homotopic maps. If  $\bar{f}, \bar{g}$  are extensions of  $f$  and  $g$  respectively, there exists a closed neighbourhood  $U$  of  $X$  such that  $\bar{f}|_U$  and  $\bar{g}|_U$  are  $p$ -homotopic.*

Hence, if we deal with locally finite polyhedra we have

1.5. PROPOSITION. *Given a locally finite polyhedron  $P$ ,  $\text{p-cat}(P)$  is the smallest number  $n$  such that  $P$  can be covered by  $n$   $p$ -categorical subpolyhedra.*

*Proof.* Given a  $p$ -categorical open covering  $\{W_i\}$  let  $H^i: \bar{W}_i \times I \rightarrow P$  be a  $p$ -deformation of  $\bar{W}_i$ . Since  $P$  is ANR by Lemma 1.4 there is a closed neighbourhood  $\Omega_i$  of  $\bar{W}_i$  and a  $p$ -extension  $\bar{H}^i: \Omega_i \times I \rightarrow P$ . Now by [23; 3.5] we may take a closed subpolyhedron  $P_i$  such that  $\bar{W}_i \subseteq P_i \subseteq \text{int } \Omega_i$  and thus  $\{P_i\}$  is a covering of  $P$  with  $P_i$   $p$ -categorical in  $P$ . Conversely, if  $P$  is covered by  $m$   $p$ -categorical subpolyhedra we may use regular neighbourhoods to obtain a  $p$ -categorical open covering with  $m$  elements.

The following properties of  $\text{p-cat}(-)$  are straightforwardly checked:

1.6. PROPOSITION. (i)  $\text{cat}(X) \leq \text{p-cat}(X)$ . *If  $X$  is compact the equality holds.*

(ii)  $\text{p-cat}(X) = 1$  *if and only if  $X \simeq_p J$ .*

(iii) *Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow X$  be  $p$ -maps such that  $fg \simeq_p \text{id}_Y$ . Then  $\text{p-cat}(Y) \leq \text{p-cat}(X)$ . In particular,  $\text{p-cat}(-)$  is a  $p$ -homotopy invariant.*

(iv) *If  $A$  is a  $p$ -retract of  $X$  then  $\text{p-cat}(A) \leq \text{p-cat}(X)$ .*

(v) *If  $A$  and  $B$  are closed and  $X = \text{int } A \cup \text{int } B$ , then  $\text{p-cat}(X) \leq \text{p-cat}(A) + \text{p-cat}(B)$ .*

The next proposition gives us an elementary relation between the set of Freudenthal ends of  $X$  and  $\text{p-cat}(X)$ .

1.7. PROPOSITION. *Given a space  $X$  in  $\mathfrak{B}_\infty$  we have  $\text{card}(\mathcal{F}(X)) \leq \text{p-cat}(X)$ .*

*Proof.* If  $\text{p-cat}(X) = \infty$  there is nothing to prove. Otherwise, if  $C_1, C_2, \dots, C_m$  is a  $p$ -categorical closed covering of  $X$ , the natural inclusions  $K_j: C_j \rightarrow X$  induce continuous maps  $k_j: \mathcal{F}(C_j) \rightarrow$

$\mathcal{F}(X)$ . Moreover, it is easy to check that  $\mathcal{F}(X) = \bigcup \text{Im } k_{j*}$  and since  $C_j$  is  $\mathfrak{p}$ -deformable to  $J$  we get that each  $\text{Im } k_{j*}$  is an one-point set. Hence  $\text{card}(\mathcal{F}(X)) \leq \mathfrak{p}\text{-cat}(X)$ .

1.8. EXAMPLES. (a)  $\mathfrak{p}\text{-cat}(\mathbb{R}^n) = 2 > 1 = \text{cat}(\mathbb{R}^n)$ .

(b) For any  $T_2$ -compact space  $X$ ,

$$\text{cat}(X) = \mathfrak{p}\text{-cat}(X \times J) = \mathfrak{p}\text{-cat}(X \vee J).$$

(c) If  $\tilde{S}^n$  is the space obtained from  $J$  by attaching one copy of  $S^n$  at each natural number  $n \in J$  we have  $\mathfrak{p}\text{-cat}(\tilde{S}^n) = 2$ .

(d) If  $X$  is a space in  $\mathfrak{P}_\infty$  and  $r: X \rightarrow J$  is a  $\mathfrak{p}$ -map we can define (up to  $\mathfrak{p}$ -homotopy) the proper cone  $C_p X$  of  $X$  as the push-out of the diagram  $X \times I \xleftarrow{i_0} X \xrightarrow{r} J$  and the proper mapping cone  $C_p f$  of a  $\mathfrak{p}$ -map  $f$  is defined in the natural way (see [2] for details). If  $f = r$ ,  $C_p r$  turns to be the proper suspension  $\sum_p X$  of  $X$ . As in the ordinary case it follows from 1.6(iii) and (iv)

$$\mathfrak{p}\text{-cat}(C_p f) \leq \mathfrak{p}\text{-cat}(Y) + 1.$$

In particular  $\mathfrak{p}\text{-cat}(\sum_p X) \leq 2$ .

(e) As a consequence of (d),  $\mathfrak{p}\text{-cat}(X) \leq \dim X + 1$ , if  $X$  is a locally finite CW-complex with only one Freudenthal end.

(f) The notion of categorical sequence due to Fox (see [10]) can be translated into proper terms. Namely, given a space  $X$  in  $\mathfrak{P}_\infty$  a sequence of open sets  $V_1 \subseteq \dots \subseteq V_n = X$  for which each difference  $V_i - V_{i-1}$  is  $\mathfrak{p}$ -categorical ( $V_0 = \emptyset$ ), is called a  $\mathfrak{p}$ -categorical sequence. It is easy to check that  $\mathfrak{p}\text{-cat } X \leq n$  if and only if  $X$  admits a  $\mathfrak{p}$ -categorical sequence of length  $n$ . By using this result, one shows the inequality  $\mathfrak{p}\text{-cat}(X \times Y) \leq \mathfrak{p}\text{-cat}(X) + \mathfrak{p}\text{-cat}(Y) - 1$ .

(g) The inequality  $\max\{\text{cat } X, \text{cat } Y\} \leq \text{cat}(X \times Y)$  holds in ordinary L-S category but not in proper L-S category as the following example shows. Let  $X = \mathbb{R}^2$  and  $Y = J$ ; then  $\mathfrak{p}\text{-cat}(X \times Y) = 1 < \mathfrak{p}\text{-cat}(\mathbb{R}^2)$ .

The following definition provides a new proper L-S invariant which is the translation of Ganea's strong L-S category into proper homotopy (see [15]).

DEFINITION 1.9. given a space  $X$  in  $\mathfrak{P}_\infty$ , the strong L-S category of  $X$  is the smallest integer  $\mathfrak{p}\text{-Cat}(X)$  such that there exists a space  $Y$  in  $\mathfrak{P}_\infty$   $\mathfrak{p}$ -homotopically equivalent to  $X$  which can be covered

by  $\mathbf{p}\text{-Cat}(X)$  closed sets each with the same  $\mathbf{p}$ -homotopy type as  $J$ . Obviously  $\mathbf{p}\text{-cat}(X) \leq \mathbf{p}\text{-Cat}(X)$ .

The relation between  $\mathbf{p}\text{-Cat}(X)$  and the 1-LC at  $\infty$  condition is given in the following theorem. We firstly recall that a space  $X$  in  $\mathfrak{P}_\infty$  is 1-LC at  $\infty$  if given a sequence of compact subsets  $\{L_j\}$  with  $L_j \subseteq \text{int } L_{j+1}$  and  $X = \bigcup L_j$ , the inverse sequence

$$(1.9.1) \quad \pi_1(X - L_1, \alpha(t_1)) \xleftarrow{\theta_1} \pi_1(X - L_2, \alpha(t_2)) \xleftarrow{\theta_2} \dots$$

is trivial, where  $\alpha: J \rightarrow X$  is a proper map with  $\alpha([t_i, \infty)) \subseteq X - L_i$  ( $i \geq 1$ ) and  $\theta_i$  is the inclusion induced homomorphism followed by the change of basepoint isomorphism given by  $\alpha|[t_i, t_{i+1}]$ .

**THEOREM 1.10.** *Let  $X$  be a space in  $\mathfrak{P}_\infty$  with one Freudenthal end. If  $\mathbf{p}\text{-Cat}(X) = 2$ ,  $X$  is 1-LC at  $\infty$  if and only if the inverse sequence*

$$(1.10.1) \quad H_1(X - L_1) \leftarrow H_1(X - L_2) \leftarrow \dots \leftarrow H_1(X - L_n) \leftarrow \dots$$

is trivial.

*This theorem is an immediate corollary of*

**PROPOSITION 1.11.** *Let  $X$  be as in Theorem 1.10. Assume that  $X$  can be covered by two one-ended  $\mathbf{p}$ -categorical closed subsets  $U$  and  $V$ . Then  $X$  is 1-LC at  $\infty$  if and only if the inverse sequence (1.10.1) is trivial.*

*Proof.* Let  $\{U, V\}$  be a  $\mathbf{p}$ -categorical open covering with  $U$  and  $V$  one-ended. We claim that  $U \cap V$  also is one-ended or equivalently that the inverse sequence

$$(1.11.1) \quad \tilde{H}_0(U \cap V - L_1) \leftarrow \dots \leftarrow \tilde{H}_0(U \cap V - L_n) \leftarrow \dots$$

is trivial. In the commutative diagram

$$\begin{array}{ccccc} H_1(X - L_n) & \longleftarrow & H_1(X - L_{n_1}) & & \\ \downarrow & & \downarrow & & \\ \tilde{H}_0(U \cap V - L_n) & \xleftarrow{(1)} & \tilde{H}_0(U \cap V - L_{n_1}) & \xleftarrow{(2)} & \tilde{H}_0(U \cap V - L_{n_2}) \\ & & \downarrow & & \downarrow \\ & & \tilde{H}_0(U - L_{n_1}) \oplus \tilde{H}_0(V - L_{n_1}) & \longleftarrow & \tilde{H}_0(U - L_{n_2}) \oplus \tilde{H}_0(V - L_{n_2}) \end{array}$$

where the vertical arrows are provided by the respective Mayer-Vietoris sequences. We may choose  $n_2 > n_1 > n$  such that the upper

and lower horizontal arrows are trivial. This implies that the composition of the morphisms (1) and (2) is trivial and so the inverse sequence (1.11.1) is trivial.

We have the following facts:

(i) Since  $U$  and  $V$  are  $\mathfrak{p}$ -categorical, we may choose  $n_3 > n$  such that any loop either in  $U - L_{n_3}$  or in  $V - L_{n_3}$  is null-homotopic in  $X - L_n$ .

(ii) On the other hand, since  $U \cap V$  is one-ended there is  $n_4 > n_3$  such that all the connected components of  $U \cap V - L_{n_4}$  are included in the same connected component of  $U \cap V - L_{n_3}$ .

Given any loop  $f: (I, \{0, 1\}) \rightarrow (X - L_{n_4}, *)$ , by Lebesgue’s Lemma we may find a partition  $0 = t_0 < t_1 < \dots < t_n = 1$  such that  $f|_{[t_i, t_{i-1}]}$  is included either in  $U - L_{n_4}$  or in  $V - L_{n_4}$ . Therefore, by (ii),  $f$  is a loop in  $X - L_{n_3}$  that can be expressed as a product of loops either in  $U - L_{n_3}$  or in  $V - L_{n_3}$  and each factor is null-homotopic in  $X - L_n$  by (i). Thus the morphism

$$\pi_1(X - L_{n_4}) \rightarrow \pi_1(X - L_n)$$

is trivial and  $X$  is 1-LC at  $\infty$ .

**COROLLARY 1.12.** *Let  $X$  be a homologically trivial open  $n$ -manifold ( $n \geq 3$ ) with  $\mathfrak{p}\text{-Cat}(X) = 2$ . Then  $X$  is 1-LC at  $\infty$ .*

*Proof.* As  $n \geq 3$ , Poincaré-Lefschetz Duality arguments (see [7; 3.2]) show that the inverse sequence (1.10.1) is semistable (i.e. satisfies the Mittag-Leffler condition) and its inverse limit is trivial. Then by [18; II.6.2.2] the inverse sequence (1.10.1) is trivial. Now the result follows from Theorem 1.11.

**2. Proper L-S category and L-S category in pro-Top.** We recall that  $\text{cat}(f) \leq n$  for a continuous map  $f$  from  $X$  to  $Y$ , if there is an open covering  $\{V_1, \dots, V_n\}$  of  $X$  such that  $f|_{V_j}$  is homotopically trivial for  $1 \leq j \leq n$ .

In [1] a notion of L-S category for inverse systems is defined. Namely, given an inverse systems  $\chi = \{X_\alpha; p_{\alpha\beta}\}$ , the L-S category of  $\chi$ ,  $\text{cat}(\chi)$ , is  $\leq n$  if for each  $\alpha$  there exists  $\beta > \alpha$  such that  $\text{cat}(p_{\alpha\beta}) \leq n$ .

If  $\text{pro-Top}$  is the category of topological inverse systems and promorphisms, Edwards and Hastings (see [8]) have proven that there exists a natural functor  $\varepsilon: \mathfrak{P}_\infty \rightarrow \text{pro-Top}$ ,  $\varepsilon(X)$  being the inverse

system  $\{U_1 \leftarrow U_2 \leftarrow \dots\}$  where  $U_k = X - C_k$  and  $C_k$  is an increasing sequence of compacts with  $C_k \subseteq \text{int } C_{k+1}$  and  $X = \bigcup C_k$ . A relation between  $\text{p-cat}(X)$  and  $\text{cat}(\varepsilon(X))$  is given by

2.1. THEOREM. *If  $X$  is a space in  $\mathfrak{P}_\infty$ ,  $\text{p-cat}(X) \geq \text{cat}(\varepsilon(X))$ .*

*Proof.* If  $\{W_1, \dots, W_n\}$  is a  $\text{p}$ -categorical open covering of  $X$ , we consider the covering  $\{W_j \cap U_k\}$  of  $U_k$ . Let  $r_j: \overline{W}_j \rightarrow J$  and  $\alpha_j: J \rightarrow X$  be  $\text{p}$ -maps such that there are  $\text{p}$ -homotopies  $H^j: \overline{W}_j \times I \rightarrow X$  between  $\alpha_j \circ r_j$  and the inclusion  $\overline{W}_j \subseteq X$ . For each  $k$  there exists  $s(k) \geq k$  such that  $H^j(W_j \cap U_{s(k)} \times I) \subseteq U_k$ . But since  $J$  is contractible it follows that  $U_{s(k)}$  is null-homotopic in each  $W_j \cap U_{s(k)}$ . So  $\text{cat}(\varepsilon(X)) \leq \text{p-cat}(X)$ .

2.2. REMARKS. (a) We now consider the category  $(\text{pro-Top}, \text{Top})$  whose objects are arrows  $f: \chi \rightarrow A$  where  $\chi$  is an object in  $\text{pro-Top}$  and  $A$  is a space regarded as the constant inverse system. Morphisms are pairs  $(\phi, h)$ : where  $\phi: \chi \rightarrow \xi$  is a pro-morphism and  $h: A \rightarrow B$  is a continuous map compatible with  $\phi$  via the bonding maps. Edwards and Hastings proved that  $\varepsilon: \mathfrak{P}_\infty \rightarrow (\text{pro-Top}, \text{Top})$  given by  $\varepsilon(X) = X \leftarrow U_1 \leftarrow U_2 \dots$  is a full embedding (see [8]). We can also prove that  $\text{cat}(\varepsilon(X)) \leq \text{p-cat}(X)$  in this case.

(b) There exist spaces with  $\text{cat}(\varepsilon(X)) < \text{p-cat}(X)$ . Indeed, let  $X$  be the punctured torus. Then  $\text{p-cat}(X) = 3$  according to Corollary 3.3 but obviously  $\text{cat}(\varepsilon(X)) = \text{cat}(S^1) = 2$ .

2.3. PROPOSITION. *If  $M$  is a compact connected triangulable manifold with boundary and  $W = M - \partial M$  then  $\text{cat}(\partial M) \leq \text{p-cat}(W) \leq \text{cat}(\partial M) + \text{cat}(M)$ . Furthermore, if  $M$  is contractible then  $\text{p-cat}(W) = \text{cat}(\partial M)$ .*

*Proof.* The inverse system  $\varepsilon(W)$  in  $(\text{pro-Top}, \text{Top})$  is

$$W \leftarrow \partial M \times [0, \infty) \leftarrow \partial M \times [1, \infty) \dots$$

Now we have  $\text{cat}[i: \partial M \times [0, \infty) \rightarrow W] \leq \text{cat}(\partial M \times [0, \infty)) = \text{cat}(\partial M)$  and  $\text{cat}(\varepsilon(X)) = \max\{\text{cat}(i), \text{cat}(\partial M)\} = \text{cat}(\partial M)$ . So, by Remark 2.2(a)  $\text{cat}(\partial M) \leq \text{p-cat}(W)$ .

On the other hand if we take a copy of  $M$ ,  $M' \subseteq W$  then  $\text{p-cat}(W) \leq \text{p-cat}(M' \vee J) + \text{p-cat}(\partial M' \times J) = \text{cat}(M) + \text{cat}(\partial M)$  by 1.6(viii) and 1.8(b).



If we assume that  $M$  is contractible, we get  $\text{cat}(\partial M) \leq \text{p-cat}(W) \leq \text{cat}(\partial M) + 1$ . But actually  $\text{p-cat}(W) \leq \text{cat}(\partial M)$ . Indeed, if  $\{W_j\}_{j \leq n}$  is a categorical covering of  $\partial M$ , we may assume that  $\overline{W}_1$  is a subpolyhedron (see Proposition 1.5). By [22; 6.30] we extend the proper deformation of  $\overline{W}_1 \times J$  into  $\{*\} \times J$  to a proper deformation of  $\overline{W}_1 \times J \cup M$ . Thus  $\{\overline{W} \times J \cup M, W_2 \times J, \dots, W_n \times J\}$  is a p-categorical covering of  $W$ .

The invariant  $\text{cat } \varepsilon(X)$  can be used to give some results on the behaviour of the inverse sequence

$$(2.3.1) \quad \pi_1(X - K_1) \leftarrow \pi_1(X - K_2) \leftarrow \dots \leftarrow \pi_1(X - K_n) \leftarrow \dots$$

where  $\{K_i\}$  is a sequence of compact subsets with  $K_i \subseteq \text{int } K_{i+1}$ , and  $X = \bigcup K_i$ .

**2.4. THEOREM.** *Let  $X$  be a locally finite polyhedron with one Freudenthal end and  $\text{cat } \varepsilon(X) = 2$ . Assume that the inverse sequence of abelian groups*

$$H_1(X - K_1) \leftarrow H_1(X - K_2) \leftarrow \dots \leftarrow H_1(X - K_n) \leftarrow \dots$$

*is trivial. Then the inverse sequence (2.3.1) is pro-isomorphic in  $\text{pro-}\mathcal{G}$  to an inverse sequence of finitely generated groups.*

**THEOREM 2.5.** *Under the hypotheses of Theorem 2.4,  $X$  is 1-LC at  $\infty$  if and only if the inverse sequence (2.3.1) is semistable, i.e.  $\lim^1 \pi_1(X - K_j) = *$ .*

*Proof of Theorem 2.4.* We may choose the sequence  $\{K_j\}$  with  $U_j = \overline{X - K_j}$  subpolyhedron of  $X$ . Up to pro-isomorphism we may replace  $\pi_1(X - K_j)$  by  $G_j = \pi_1(U_j)$  in (2.3.1). By using the 1-skeleton of  $X$ , it is easy to check that there is a commutative diagram

$$(2.4.1) \quad \begin{array}{ccccccc} F(L_1) & \longleftarrow & F(L_2) & \longleftarrow & F(L_3) & \longleftarrow & \dots & \longleftarrow & F(L_n) & \longleftarrow & \dots \\ \downarrow & & \downarrow & & \downarrow & & & & \downarrow & & \\ G_1 & \longleftarrow & G_2 & \longleftarrow & G_3 & \longleftarrow & \dots & \longleftarrow & G_n & \longleftarrow & \dots \end{array}$$

where  $F(L_n)$  denotes the corresponding free group of basis  $L_n$ , the differences  $L_n - L_{n+1}$  are finite, and the bonding morphisms are the natural inclusions.

Since the inverse sequence

$$G_1^{ab} \leftarrow G_2^{ab} \leftarrow G_3^{ab} \leftarrow \dots \leftarrow G_n^{ab} \leftarrow \dots$$

is trivial, by abelianizing (2.4.1) we readily check that  $G_n^{ab}$  is finitely generated for  $n \geq 1$ .

Now, since

$$\text{cat } \varepsilon(X) = \sup\{\text{cat}[i_n : U_n \rightarrow U_{n-1}]\} = 2$$

we get from [15; 7.3] that for any  $n \geq 2$ , there is  $\mu_n$  such that

$$\begin{array}{ccc} U_n & \xrightarrow{\mu_n} & U_{n-1} \vee U_{n-1} \\ \downarrow i_n & & \downarrow l_{n-1} \\ U_{n-1} & \xrightarrow{\Delta} & U_{n-1} \times U_{n-1} \end{array}$$

is commutative up to homotopy. This diagram induces a commutative diagram

$$(2.4.2) \quad \begin{array}{ccc} G_n & \xrightarrow{\mu_{n*}} & G_{n-1} * G_{n-1} \\ \downarrow i_{n*} & & \downarrow l_{n-1*} \\ G_{n-1} & \xrightarrow{\Delta} & G_{n-1} \times G_{n-1} \end{array}$$

where  $l_{n*}$  induces the natural epimorphism  $a * b \rightarrow (a, b)$ .

From the Kurosh Subgroup Theorem (see [17]) the group  $l_{n-1*}^{-1} \Delta i_{n*}(G_n)$  is a free group. Therefore,  $\mu_{n*} : G_n \rightarrow \text{Im } \mu_{n*}$  is an epimorphism onto a free group and so  $(\text{Im } \mu_{n*})^{ab}$  is a finitely generated free abelian group. Hence,  $\text{Im } \mu_{n*}$  is a finitely generated free group.

On the other hand,  $\Delta$  is injective and so the commutativity of (2.4.2) yields a natural epimorphism  $\text{Im } \mu_{n*} \twoheadrightarrow \text{Im } i_{n*}$ . Thus,  $\text{Im } i_{n*}$  is a finitely generated group.

It is a well-known fact that (2.3.1) is pro-isomorphic to the inverse sequence

$$(2.4.3) \quad \text{Im } i_{1*} \leftarrow \text{Im } i_{2*} \leftarrow \dots \leftarrow \text{Im } i_{n*} \leftarrow \dots$$

*Proof of Theorem 2.5.* If in addition  $\lim^1 G_j = *$ , we know that the inverse sequence (2.3.1) satisfies the Mittag-Leffler condition (see [18; p. 174]) and we may assume that the bonding morphisms in (2.4.3) are onto. We shall denote  $\text{Im } i_{n*}$  by  $H_n$ .

Since the sequences (2.3.1) and (2.4.3) are pro-isomorphic, the abelianization of (2.4.3) is trivial. Furthermore, as the bonding morphisms are onto we get that  $H_n$  is its own commutator subgroup. Thus,  $H_n^{\text{ab}} = 0$ .

On the other hand, diagram (2.4.2) induces a commutative diagram

$$(2.4.4) \quad \begin{array}{ccc} H_n & \xrightarrow{\mu'} & H_{n-2} * H_{n-2} \\ \downarrow (1) & & \downarrow i_{n-2*} \\ H_{n-2} & \xrightarrow{\Delta} & H_{n-2} \times H_{n-2} \end{array}$$

where (1) is the natural bonding morphism and  $\mu' = (i_{n-1} * i_{n-1}) \circ \mu : H_n \rightarrow H_{n-2} * H_{n-2}$ .

As in the proof of Theorem 2.4,  $\text{Im } \mu'$  is a free group and the epimorphism  $H_n^{\text{ab}} = 0 \rightarrow (\text{Im } \mu')^{\text{ab}}$  implies that  $\text{Im } \mu'$  is trivial. Since  $\Delta$  is injective, the morphism on the left side of (2.4.4) is trivial. This completes the proof.

**COROLLARY 2.6.** *Let  $X$  be a homologically trivial open  $n$ -manifold ( $n \geq 3$ ) with only one strong end and  $\text{cat}(\varepsilon(X)) = 2$ . Then  $X$  is 1-LC at  $\infty$ .*

We recall that the strong ends of  $X$  are the elements in the set of  $p$ -homotopy classes  $[J; X]_p$ . It is a well-known fact that  $[J; X]_p \cong \lim^1 \pi_1(X - K_j)$ . Now the proof of Corollary 2.6 is similar to the proof of Corollary 1.12.

**REMARK 2.7.** Notice that Theorem 2.1 and Theorem 2.4 imply that the contractible open 3-manifolds described in [20; Theorem 1] have  $p$ -cat  $\geq 3$ . These manifolds have infinitely many strong ends.

Indeed, the inverse sequence of  $\pi_1$ 's of those manifolds are of the type

$$\mathcal{S} = \left\{ \star_0^\infty G_i \leftarrow \star_1^\infty G_i \leftarrow \dots \leftarrow \star_n^\infty G_i \dots \right\}$$

where “ $\star$ ” stands for the free product of groups and the bonding morphisms are the natural inclusions (see [24] and [20]). The sequence  $\mathcal{S}$  is not pro-isomorphic to a sequence of finitely generated groups since the existence of such an isomorphism would yield an inclusion  $\star_n^\infty G_i \subseteq F \subseteq \star_0^\infty G_i$  for some finitely generated group  $F$ , if  $n \geq n_0$  for some  $n_0$ .

**REMARK 2.8.** The following simple example shows that Theorem 2.4 does not imply the condition 1-LC at  $\infty$ .

Let  $F(x, y)$  be the free group with generators  $x$  and  $y$ , and let  $f: S^1 \vee S^1 \rightarrow S^1 \vee S^1$  be a continuous realization of the morphism  $\phi: F(x, y) \rightarrow F(x, y)$  given by  $\phi(x) = [x, y]$  and  $\phi(y) = [x^{-1}, y]$ , with  $[x, y] = xyx^{-1}y^{-1}$ .

We now define  $T$  as the telescope

$$T = X \times [0, 1] \cup_f X \times [1, 2] \cup_f \dots \cup_f X \times [n, n + 1] \dots$$

with  $X = S^1 \vee S^1$ . It is clear that  $T$  is under the hypotheses of Theorem 2.4 but the inverse sequence (2.3.1) agrees with

$$F(x, y) \xleftarrow{\phi} F(x, y) \xleftarrow{\phi} F(x, y) \xleftarrow{\phi} \dots$$

whose  $\lim^1$  is not trivial.

**3. Cohomology and p-category.** Several p-homotopy invariant cohomologies have appeared in the literature. We next give some examples.

3.1. EXAMPLES. (a) Various cohomologies with compact supports (singular, Čech type, Alexander-Spanier, etc.). See [19] for details.

(b) Cohomology of the end of  $X$ ,  $H_e^*(X)$ . Namely,  $H_e^*(X)$  is defined as the cohomology of the cochain complex  $C^*(X)/C_c^*(X)$  where  $C_c^*(X)$  stands for the subgroup of singular cochains with compact supports. Alternatively,  $H_e^*(X) = \varinjlim \{H^*(U_j)\}$  where  $\{U_j\}$  is a system of  $\infty$ -neighbourhoods of  $X$ . See [13] for details.

(c) Pro-cohomology of  $X$  with coefficients in the inverse system  $\mathcal{S}$ . Given  $\{U_j\}$  as above,  $\mathcal{H}^*(X; \mathcal{S})$  is the cohomology of the cochain complex  $C^*(X; \mathcal{S}) = (\text{pro-}\mathcal{A}b, \mathcal{A}b)(\mathcal{E}_*(X); \mathcal{S})$ , where  $\mathcal{E}_*(X)$  denotes the inverse system  $\{C_*(X) \leftarrow C_*(U_1) \leftarrow \dots\}$ . Here  $\mathcal{A}b$  stands for the category of abelian groups and  $(\text{pro-}\mathcal{A}b, \mathcal{A}b)$  is constructed from  $\mathcal{A}b$  as  $(\text{pro-}\mathcal{T}op, \mathcal{T}op)$  is constructed from  $\mathcal{T}op$  in §3. See [14] for details.

If  $\mathfrak{H}^*$  is any of the above cohomologies, relative groups  $\mathfrak{H}^*(X, A)$  are easily defined and restricted morphisms  $\rho: \mathfrak{H}^*(X, A) \rightarrow \mathfrak{H}^*(X, B)$  are natural when  $B \subseteq A$ . If  $A$  is closed there is an exact sequence

$$\dots \rightarrow \mathfrak{H}^n(X, A) \xrightarrow{\rho} \mathfrak{H}^n(X) \xrightarrow{i^*} \mathfrak{H}^n(A) \xrightarrow{\delta} \mathfrak{H}^{n+1}(X, A) \rightarrow \dots$$

where  $i: A \subseteq X$  is the inclusion. Also cup-products can be defined for  $\mathfrak{H}^*$ .

As in the ordinary case, p-invariant cohomologies provide a lower bound for proper L-S category. Namely,

**3.2. PROPOSITION.** *Given  $X$  in  $\mathfrak{P}_\infty$ , let  $l(\mathfrak{H}^*(X))$  be  $\sup\{n \in \mathbb{N}; \exists \alpha_1, \alpha_2, \dots, \alpha_n \in \mathfrak{H}^*(X) \text{ with } \alpha_1 \cup \alpha_2 \cup \dots \cup \alpha_n \neq 0\}$ . Then  $\text{p-cat}(X) \geq l(\mathfrak{H}^*(X)) + 1$ .*

*Proof.* Let  $A_1, A_2, \dots, A_n$  be a p-categorical open covering of  $X$ . Since  $i^*: \mathfrak{H}^*(X) \rightarrow \mathfrak{H}^*(\overline{A}_k)$  is trivial for each  $k \leq n$ , given  $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathfrak{H}^*(X)$ , there are elements  $\alpha'_k \in \mathfrak{H}^*(X, A_k)$  with  $\rho(\alpha'_k) = \alpha_k$  ( $k \leq n$ ) in the diagram

$$\begin{array}{ccc}
 & \xrightarrow{\rho'} & \mathfrak{H}^*(X, A_k) \\
 \downarrow & \nearrow \rho' & \downarrow \rho \\
 \mathfrak{H}^*(X, \overline{A}_k) & \xrightarrow{\rho''} & \mathfrak{H}^*(X) \xrightarrow{i^*=0} \mathfrak{H}^*(\overline{A}_k)
 \end{array}$$

Now the commutative diagram

$$\begin{array}{ccc}
 \mathfrak{H}^*(X, A_1) \otimes \mathfrak{H}^*(X, A_2) \otimes \dots \otimes \mathfrak{H}^*(X, A_n) & \xrightarrow{\cup} & \mathfrak{H}^*(X, A_1 \cup A_2 \cup \dots \cup A_n) = 0 \\
 \downarrow \rho & & \downarrow \rho \\
 \mathfrak{H}^*(X) \otimes \mathfrak{H}^*(X) \otimes \dots \otimes \mathfrak{H}^*(X) & \xrightarrow{\cup} & \mathfrak{H}^*(X)
 \end{array}$$

yields  $\alpha_1 \cup \alpha_2 \cup \dots \cup \alpha_n = 0$ .

**3.3. COROLLARY.** *If  $W$  is a non-compact surface with only one Freudenthal end then*

$$\text{p-cat}(W) = \begin{cases} 2 & \text{if } W \cong \mathbb{R}^2, \\ 3 & \text{otherwise.} \end{cases}$$

*Proof.* If  $W \not\cong \mathbb{R}^2$  we can find elements  $\alpha, \beta \in H_c^1(W; \mathbb{Z}_2)$  with  $\alpha \cup \beta \neq 0$  and then  $\text{p-cat}(W) \geq 3$ . By 1.8(e) we conclude  $\text{p-cat}(W) = 3$ .

Also we may use the cohomology with compact supports in the following

**3.4. EXAMPLE.** There are contractible spaces with only one Freudenthal end and infinite proper L-S category. Let  $T^n$  be the  $n$ -dimensional torus ( $n \geq 0, T^0 = *$ ) and  $\pi_n: T^n \rightarrow T^{n-1}$  ( $n \geq 1$ ) the natural projection. Then  $X = \bigcup X_n$ , where  $X_n$  is the mapping cylinder of  $\pi_n$ , is a contractible space and  $H_c^1(X; \mathbb{Z})$  is the exterior algebra  $\Lambda(x_1, \dots, x_n, \dots)$  with infinitely many generators. Then  $\text{p-cat}(X) = \infty$ .

Now we characterize  $\mathbb{R}^n$  among all the manifolds by using proper L-S category. Namely,

**THEOREM 3.5.** *Let  $X$  be a connected open manifold ( $n \geq 1$ ). Then  $X \simeq_p \mathbb{R}^n$  if and only if one of the following statements holds:*

- (i)  $\mathbf{p}\text{-Cat}(X) = 2$
- (ii)  $\mathbf{p}\text{-cat}(X) = 2$  and  $X$  has only one strong end.

*Proof.* Since  $\text{cat}(X) \leq \mathbf{p}\text{-cat}(X) = 2$ , it follows that  $\pi_1(X)$  is a free group (see [15, §4]) and so  $X$  is orientable.

For any  $\mathbb{Z}_p$  with  $p$  prime we have the isomorphisms

$$H_c^q(X; \mathbb{Z}_p) \stackrel{D}{\cong} H_{n-q}(X; \mathbb{Z}_p) \stackrel{\alpha}{\cong} H^{n-q}(X; \mathbb{Z}_p)$$

given by the Poincaré Duality (see [19; 11.2]) and the ordinary algebraic duality respectively. Then, if  $\varepsilon \neq 0 \in H_c^q(X; \mathbb{Z}_p)$  and  $\mu_X \in H_n^\Pi(X; \mathbb{Z}_p)$  is the fundamental class of  $X$  in the homology of “infinite chains”, we get

$$0 = \mu_X \cap (\varepsilon \cup \alpha(D(\varepsilon))) = D(\varepsilon) \cap \alpha(D(\varepsilon)) = 1$$

where “ $\cap$ ” and “ $\cup$ ” are the natural cap and cup products  $H_p^\Pi \otimes H_c^r \rightarrow H_{p-r}$  and  $H_c^q \otimes H^r \rightarrow H_c^{q+r}$  respectively (see [19; 10.4] for details).

Therefore,  $H_{n-q}(X; \mathbb{Z}_p) = H_c^q(X; \mathbb{Z}_p) = 0$  for all  $q$  and the Universal Coefficient Theorem proves that  $X$  is homologically trivial. In particular  $\pi_1(X) = 1$  and  $X$  is contractible by the Whitehead Theorem. Then  $X = \mathbb{R}^n$  if  $n = 1, 2$ .

On the other hand, if  $n \geq 3$  we have the Poincaré Duality  $H_q^\Pi(X; \mathbb{Z}) \cong H^{n-q}(X; \mathbb{Z}) = 0$  (see [19; 11.4]). Since  $X$  is properly 1-connected at infinity by Corollary 1.12 or Corollary 2.6, a convenient proper Hurewicz Theorem (see [3; 3.4]) yields that  $H_n^\Pi(X; \mathbb{Z}) \cong [\mathbb{R}^n; X]_p$  and the fundamental class  $\mu_X$  can be represented by a  $p$ -map  $f: \mathbb{R}^n \rightarrow X$ . It is easily checked that  $f_*: H_*^\Pi(\mathbb{R}^n) \rightarrow H_*^\Pi(X)$  is an isomorphism and again by [19; 11.4],  $f^*: H_c^*(X) \rightarrow H_c^*(\mathbb{R}^n)$  is an isomorphism. Now,  $f$  is a  $p$ -homotopy equivalence by [9; 4.9].

By using the results of Siebenmann [21] and Freedman [12] we get

**THEOREM 3.6.** *If  $X$  is a connected open  $n$ -manifold ( $n \neq 3$ ), then  $X$  is homeomorphic to  $\mathbb{R}^n$  if and only if 3.5(i) or 3.5(ii) holds.*

**REMARK 3.7.** We do not know whether Theorem 3.5(ii) is true without the condition on strong ends.

Given an embedding  $i: J \rightarrow X$ , we can define a new L-S type proper invariant  $\text{hp-cat}(X, i)$  as follows (cf. [15; §4 and §5]).

We define the  $n$ th canonical proper fat wedge of  $(X, i)$  as the union

$$T_p^n(X, i) = \bigcup \{A_k, i \leq k \leq n\}$$

where  $A_k$  is the product of  $(n - 1)$  copies of  $X$  with  $J$ , where  $J$  is placed in the  $k$ th factor. Then  $\text{hp-cat}(X, i) \leq n$  if there is a diagram in  $\mathfrak{B}_\infty$

$$\begin{array}{ccc} X & \xrightarrow{\Delta} & \Pi^n X \\ \downarrow n & & \uparrow \varphi(i) \\ T_p^n(X, i) & \xrightarrow{\quad} & \end{array}$$

commutative up to the  $p$ -homotopy. Here  $\Delta$  is the diagonal map and  $\varphi(i)$  is the natural embedding induced by  $i$ .

In contrast with Theorem 3.6, the following result holds.

**3.7. PROPOSITION.** *For any contractible open manifold  $W$ , and any embedding  $i: J \rightarrow W$ ,  $\text{hp-cat}(W, i) = 2$ .*

*Proof.* It is clear that  $\text{hp-cat}(W) \geq 2$  and  $\text{hp-cat}(\mathbb{R}) = 2$ . For  $\dim(W) = n \geq 2$ , it is known that  $W \times W$  is 1-connected at infinity (see [13; 1.8]). And by [12] and [21]  $W \times W \cong^h \mathbb{R}^{2n}$ . Finally we apply the Proper Cellular Approximation Theorem (see [9]) to deform  $h \circ \Delta: W \rightarrow \mathbb{R}^{2n}$  onto the  $n$ -skeleton of  $\mathbb{R}^{2n}$  which may be regarded as  $J$  since  $\mathbb{R}^{2n}$  is properly  $(2n - 2)$ -connected. So,  $\text{hp-cat}(W) \leq 2$ .

**Acknowledgments.** The authors wish to thank the referee for his/her helpful comments.

#### REFERENCES

- [1] V. V. Agaronjan and J. M. Smirnov, *The Ljusternik-Šnirel'man category for procategories*, Soviet Math. Dokl., **20** (1973), 516–518.
- [2] R. Ayala, E. Domínguez and A. Quintero, *A theoretical framework for proper homotopy theory*, Math. Proc. Camb. Philos. Soc., **107** (1990), 475–482.
- [3] ———, *Hurewicz Theorem for homology at  $\infty$* , preprint.
- [4] B. J. Ball and R. B. Sher, *A theory of proper shape for locally compact metric spaces*, Fund. Math., **86** (1974), 162–194.
- [5] K. Borsuk, *Über den Lusternik-Schnirelmannschen Begriff der Kategorie*, Fund. Math., **26** (1936), 123–136. Abstracted: *Sur la notion de la catégorie de M. L. Lusternik et Schnirelmann*, Comptes Rend. Acad. Sci. Paris, **198** (1934), 1731–1732.
- [6] M. Clapp and D. Puppe, *Invariants of the Lusternik-Schnirelmann type and the topology of critical sets*, Trans. Amer. Math. Soc., **298** (1986), 603–620.

- [7] A. Z. Dymov, *On the behaviour at infinity of the fundamental group of a homologically trivial manifold*, Math. USSR Izv., **11** (1977), 529–550.
- [8] D. A. Edwards and H. M. Hastings, *Čech and Steenrod Homotopy Theories with Applications to Geometric Topology*, Lecture Notes 542, Springer, 1976.
- [9] F. T. Farrell, L. R. Taylor, and J. B. Wagoner, *The Whitehead theorem in proper category*, Compositio Math., **27** (1973), 1–23.
- [10] R. H. Fox, *On the Lusternik-Schnirelmann category*, Ann. of Math., **42** (1941), 333–370.
- [11] H. Freudenthal, *Über die enden topologischer Räume und Gruppen*, Math. Zeit., **33** (1931), 692–713.
- [12] M. H. Freedman, *The topology of 4-manifolds*, J. Differential Geom., **17** (1982), 375–453.
- [13] L. Glaser, *Geometrical Combinatorial Topology*. Vol. II, Van Nostrand, 1972.
- [14] L. J. Hernández, *About the extension problem for proper maps*, Topology Appl., **25** (1987), 51–64.
- [15] I. M. James, *On category in the sense of Lusternik-Schnirelmann*, Topology, **17** (1987), 51–64.
- [16] L. Lusternik and L. Schnirelmann, *Méthodes Topologiques dans les problèmes variationnels*, Hermann, 1934.
- [17] W. Magnus, A. Karrass, and D. Solitar, *Combinatorial Group Theory*, Wiley, 1966.
- [18] S. Mardesic and J. Segal, *Shape Theory*, North-Holland, 1982.
- [19] W. S. Massey, *Homology and Cohomology Theory*, M. Dekker, 1978.
- [20] D. R. McMillan, *Some contractible open 3-manifolds*, Trans. Amer. Math. Soc., **102** (1962), 373–382.
- [21] L. Siebenmann, *On detecting Euclidean Space homotopically among topological manifolds*, Inv. Math., **6** (1968), 245–261.
- [22] R. M. Switzer, *Algebraic Topology-Homotopy and Homology*, Springer, 1975.
- [23] J. H. C. Whitehead, *Simplicial spaces, nuclei and  $m$ -groups*, Proc. London Math. Soc., **45** (1939), 243–327.
- [24] J. H. C. Whitehead and M. H. A. Newman, *On the group of certain linkage*, Quart J. Math., **8** (1937), 14–21.

Received August 20, 1990 and in revised form June 3, 1991.

R. AYALA, A. MÁRQUEZ, A. QUINTERO  
DPTO. ALGEBRA, COMPUTACIÓN, GEOMETRÍA Y TOPOLOGÍA  
FACULTAD DE MATEMÁTICAS  
APDO. 1160  
41080-SEVILLA, SPAIN

AND

D. DOMÍNGUEZ  
DPTO. MATEMÁTICAS  
FACULTAD DE CIENCIAS  
CIUDAD UNIVERSITARIA  
50009-ZARAGOZA, SPAIN





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555-05

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112

F. MICHAEL CHRIST  
University of California  
Los Angeles, CA 90024-1555

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093

NICHOLAS ERCOLANI  
University of Arizona  
Tucson, AZ 85721

R. FINN  
Stanford University  
Stanford, CA 94305

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305

C. C. MOORE  
University of California  
Berkeley, CA 94720

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH  
(1906-1982)

B. H. NEUMANN

F. WOLF  
(1904-1989)

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1992 by Pacific Journal of Mathematics

<b>R. Ayala, Eladio Domínguez Murillo, Alberto Márquez Pérez and A. Quintero, Lusternik-Schnirelmann invariants in proper homotopy theory</b> .....	201
<b>Hari Bercovici and Dan-Virgil Voiculescu, Lévy-Hinčin type theorems for multiplicative and additive free convolution</b> .....	217
<b>L. J. Bunce and Cho-Ho Chu, Compact operations, multipliers and Radon-Nikodým property in <math>JB^*</math>-triples</b> .....	249
<b>Marius Dadarlat, Gabriel Nagy, András Némethi and Cornel Pasnicu, Reduction of topological stable rank in inductive limits of <math>C^*</math>-algebras</b> .....	267
<b>François Dumas and Robert Vidal, Dérivations, et hautes dérivations, dans certains corps gauches de series de Laurent</b> .....	277
<b>Mourad Ismail and Xin Li, On sieved orthogonal polynomials. IX: Orthogonality on the unit circle</b> .....	289
<b>X. T. Liang and Y. W. Lu, A Phragmén-Lindelöf theorem</b> .....	299
<b>Mark Stephen Reeder, On certain Iwahori invariants in the unramified principal series</b> .....	313
<b>Shohei Tanaka, On the representation of the determinant of Harish-Chandra's <math>C</math>-function of <math>SL(n, \mathbb{R})</math></b> .....	343
<b>Fritz von Haeseler and Guentcho Svetoslavov Skordev, Borsuk-Ulam theorem, fixed point index and chain approximations for maps with multiplicity</b> .....	369