

# Pacific Journal of Mathematics

## **ALMOST $s$ -TANGENT MANIFOLDS OF HIGHER ORDER**

MANUEL (RODRIGUEZ) DE LEÓN, J. A. OUBIÑA,  
P. R. RODRIGUES AND MODESTO R. SALGADO

## ALMOST $s$ -TANGENT MANIFOLDS OF HIGHER ORDER

M. DE LEÓN, J. A. OUBIÑA, P. R. RODRIGUES, AND M. SALGADO

**We introduce the notion of almost  $s$ -tangent structures of higher order by abstracting the geometric structure of the space of  $k$ -jets  $J^k(\mathbb{R}, M)$ . These structures are a natural extension of almost tangent structures of higher order.**

**1. Introduction.** Almost tangent structures on even-dimensional manifolds were introduced by Clark and Bruckheimer [1] and Eliopoulos [6] around 1960 and have been investigated by many authors (see [15] and references therein). As it is well known the tangent bundle  $TM$  of a manifold  $M$  carries a canonical almost tangent structure (hence the name). This almost tangent structure plays an important role in the Lagrangian formulation of particle dynamics [15]. Crampin [2] showed that integrable almost tangent structures are relevant to study the inverse problem of Lagrangian dynamics and Cantrijn, Cariñena, Crampin and Ibort [3] proposed a geometric method of reduction of degenerate Lagrangian systems in this framework.

The notion of almost tangent structure of higher order is due to Eliopoulos [7]. An almost tangent structure of order  $k$  on a  $((k+1)n)$ -dimensional manifold is defined by abstracting the geometric structure of the tangent bundle of order  $k$  of an  $n$ -dimensional manifold. Tangent bundles of higher order are the natural framework to develop the Lagrangian dynamics of higher order (see [14, 4]). Recently, de León, Giraldo and Rodrigues [12, 17] have obtained results similar to those of Cantrijn et al. in order to reduce degenerate Lagrangian systems of higher order in the framework of higher order integrable almost tangent structures.

In [20] Oubiña extended the notion of almost tangent structures to odd-dimensional manifolds and introduced a new type of geometric structures, the so called almost  $s$ -tangent structures, their model being the stable tangent bundle  $J^1(\mathbb{R}, M) \equiv \mathbb{R} \times TM$ . These structures are involved in the study of the non-autonomous Lagrangian systems, and the inverse problem of non-autonomous Lagrangian dynamics can be reformulated in terms of almost  $s$ -tangent structures [13].

In the present paper we extend the notion of almost  $s$ -tangent structure to higher orders. The geometrical model is now the space of  $k$ -jets  $J^k(\mathbb{R}, M)$ , which can be identified to  $\mathbb{R} \times T^k M$ . In §2, we introduce the notion of almost  $s$ -tangent structure  $(J, \omega, \xi)$  of order  $k$  on a manifold  $V$  of dimension  $(k+1)n+1$ . In §3,  $(J, \omega, \xi)$  is interpreted as a  $\mathbf{G}$ -structure, for a certain Lie subgroup  $\mathbf{G}$  of  $\text{Gl}((k+1)n+1, \mathbb{R})$ , and it is proved that  $(J, \omega, \xi)$  is integrable if and only if the Nijenhuis tensor of  $J$  vanishes and  $\omega$  is closed. Moreover, we prove that the existence of an almost  $s$ -tangent structure of order  $k$  on a manifold  $V$  is equivalent to a reduction of the structure group of  $TV$  to  $O(n) \times \cdots \times O(n) \times 1$ . Section 4 is devoted to showing that some special submanifolds of an almost tangent manifold of higher order inherit an almost  $s$ -tangent structure of the same order.

Since  $J^{2k-1}(\mathbb{R}, M)$  is the evolution space of a non-autonomous Lagrangian system of order  $k$ , we feel that almost  $s$ -tangent structures of higher order might be relevant to give a geometric procedure in order to reduce degenerate non-autonomous Lagrangian systems of higher order. They could also be useful in discussing the inverse problem for higher order autonomous systems (see [10, 12, 16]).

**2. Almost  $s$ -tangent structures of higher order.** Throughout this paper it is assumed that all differential structures are of  $C^\infty$ -class.

Let  $M$  be a manifold of dimension  $n$  and  $T^k M$  be the tangent bundle of order  $k$  of  $M$ . Then  $T^k M$  carries a canonical integrable almost tangent structure  $F$  of order  $k$  (see [14]), which is actually the  $(k-1)$ -lift of the identity tensor field of  $M$  to  $T^k M$  in the sense of Morimoto [19]. Let  $(z^i)$  be local coordinates for  $M$  and  $(z_0^i, z_1^i, \dots, z_k^i)$  the induced coordinates of  $T^k M$ . Then the components of  $F$  in every such coordinate system are

$$\begin{pmatrix} 0 & 0 \\ I_{kn} & 0 \end{pmatrix}.$$

This notion can be extended to the space of jets of order  $k$ , i.e.  $J^k(\mathbb{R}, M)$ . In fact, the  $((k+1)n+1)$ -dimensional manifold  $J^k(\mathbb{R}, M)$  can be identified with  $\mathbb{R} \times T^k M$  in a very natural way. Then we can define a canonical tensor field on  $J^k(\mathbb{R}, M)$  given by  $J = F + (\partial/\partial t) \otimes dt$ , where  $t$  is the global coordinate function for  $\mathbb{R}$ . Hence  $J$  has rank  $kn+1$  and satisfies  $J^{k+1} = (\partial/\partial t) \otimes dt$ .

This suggests the following

**DEFINITION.** Let  $V$  be a differentiable manifold of dimension  $(k+1)n+1$ . A triple  $(J, \omega, \xi)$ , where  $J$  is a tensor field of type

(1, 1),  $\omega$  is a 1-form and  $\xi$  is a vector field on  $V$  such that

$$\begin{aligned} (1) \quad & \omega(\xi) = 1, \\ (2) \quad & J^{k+1} = \omega \otimes \xi, \\ (3) \quad & \text{rank } J = kn + 1, \end{aligned}$$

will be called an *almost  $s$ -tangent structure of order  $k$*  and the manifold  $V$  an *almost  $s$ -tangent manifold of order  $k$* .

The following proposition follows from (1) and (2).

**PROPOSITION 2.1.** *We have  $J\xi = \lambda\xi$  and  $\omega J = \lambda\omega$ , where  $\lambda^{k+1} = 1$ .*

*Proof.* Clearly  $J\xi = J(J^{k+1}\xi) = J^{k+1}(J\xi) = \omega(J\xi)\xi$  and so  $J\xi = \lambda\xi$ , being  $\lambda^{k+1} = 1$ . Also  $\omega(JX)\xi = J^{k+1}(JX) = J(J^{k+1}X) = \omega(X)J\xi = \lambda\omega(X)\xi$ , that is  $\omega J = \lambda\omega$ .  $\square$

Let  $Q$  be the  $((k+1)n)$ -dimensional distribution defined by the condition  $\omega = 0$ , i.e.  $Q = \ker \omega$ . By Proposition 2.1 we have  $JQ \subset Q$  and hence  $J|_Q^{k+1} = 0$ . Since  $\text{rank}(J|_Q) = kn$ ,  $J$  acts on  $Q$  as an almost tangent structure operator of order  $k$ . Also we have

$$\ker J \subset \ker J^2 \subset \dots \subset \ker J^k \subset \ker J^{k+1} = Q$$

and

$$\dim \ker J^r = rn, \quad 1 \leq r \leq k+1.$$

**REMARK.** It is to be noted that, by virtue of well known properties of nilpotent operators on vector spaces, conditions (1) and (2) in the definition imply that  $\text{rank } J \leq kn+1$ . Thus the condition (3) requires  $\text{rank } J$  to be maximal.

**EXAMPLE 2.1.** The space  $J^k(\mathbb{R}, M)$  of jets of order  $k$  on a differentiable manifold  $M$  is an almost  $s$ -tangent manifold of order  $k$ .

**EXAMPLE 2.2.** (A principal 1-bundle over an almost tangent manifold of order  $k$ .) Let  $V(M, G)$  be a principal bundle over a manifold  $M$ ,  $G$  being a connected Lie group of dimension 1 with Lie algebra  $\mathfrak{g}$ . Let  $F$  be an almost tangent structure of order  $k$  on  $M$ . If  $a \in \mathfrak{g} \cong \mathbb{R}$ , we denote by  $a^*$  the fundamental vector field on  $V$  corresponding to  $a$ . In particular, if  $e$  is the unit vector of  $\mathfrak{g} \cong \mathbb{R}$ , we shall denote  $\xi = e^*$ . Let  $\omega$  be a connection form on  $V$  and, for every vector field

$X$  on  $M$ , denote by  $X^H$  the horizontal lift of  $X$  with respect to  $\omega$ . We define a tensor field  $J$  of type  $(1, 1)$  on  $V$  by putting

$$JX^H = (FX)^H, \quad Ja^* = a\xi$$

for all vector fields  $X$  on  $M$  and for all  $a \in \mathfrak{g} \equiv \mathbb{R}$ . Then  $(J, \omega, \xi)$  is an almost  $s$ -tangent structure of order  $k$  on  $V$  and the distribution  $Q$  on  $V$ , defined by  $\omega = 0$ , is the horizontal distribution of the connection.

**3. The structure group.** Next we shall describe an almost  $s$ -tangent structure of order  $k$  as a  $G$ -structure.

Let  $(J, \omega, \xi)$  be an almost  $s$ -tangent structure of order  $k$  on a manifold  $V$  of dimension  $(k+1)n+1$ . Let  $P$  be the one-dimensional distribution determined by  $\xi$ . We have  $T_xV = Q_x \oplus P_x$ , for each  $x \in V$ . If  $S_x$  is a complementary subspace of  $\ker J_x^k$  in  $Q_x$  then we have  $T_xV = S_x \oplus \ker J_x^k \oplus P_x$ . Thus, if  $\{X_1, \dots, X_n\}$  is a basis for  $S_x$  then  $\{X_i, JX_i, \dots, J^kX_i, \xi_x\}$  is a basis for  $T_xV$  called an *adapted frame*. If  $S'_x$  is another complementary subspace of  $\ker J_x^k$  in  $Q_x$  and  $\{X'_i, JX'_i, \dots, J^kX'_i, \xi_x\}$  another adapted frame, then we have

$$\begin{aligned} X'_i &= A_i^j X_j + (A_1)_i^j JX_j + \dots + (A_k)_i^j J^k X_j, \\ JX'_i &= A_i^j JX_j + (A_1)_i^j J^2 X_j + \dots + (A_{k-1})_i^j J^{k-1} X_j, \\ &\dots \dots \\ J^k X'_i &= A_i^j J^k X_j, \\ \xi_x &= \xi_x. \end{aligned}$$

Therefore two adapted frames are related by a matrix  $\mathbf{A} \in \text{Gl}((k+1)n+1, \mathbb{R})$  of the form

$$\mathbf{A} = \begin{pmatrix} A & 0 & 0 & \dots & 0 & 0 & 0 \\ A_1 & A & 0 & \dots & 0 & 0 & 0 \\ A_2 & A_1 & A & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots & \vdots \\ A_{k-1} & A_{k-2} & A_{k-3} & \dots & A & 0 & 0 \\ A_k & A_{k-1} & A_{k-2} & \dots & A_1 & A & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 \end{pmatrix}.$$

Let  $\mathbf{G}$  be the set of such matrices. Then  $\mathbf{G}$  is a closed subgroup of  $\text{Gl}((k+1)n+1, \mathbb{R})$  and therefore a Lie subgroup of  $\text{Gl}((k+1)n+1, \mathbb{R})$ . Let  $B_{\mathbf{G}}$  be the set of adapted frames at all points of  $V$ . It can be easily proved that  $B_{\mathbf{G}}$  defines a  $\mathbf{G}$ -structure on  $V$ .

Notice that, with respect to an adapted frame,  $J$  is represented by the matrix

$$J_0 = \begin{pmatrix} 0 & 0 & 0 \\ I_{kn} & 0 & 0 \\ 0 & 0 & \lambda \end{pmatrix}$$

where  $\lambda^{k+1} = 1$ ,  $J\xi = \lambda\xi$ . In fact, the group  $\mathbf{G}$  can be described as the invariance group of the matrix  $J_0$ , i.e.  $\mathbf{A} \in \mathbf{G}$  if and only if  $\mathbf{A}J_0\mathbf{A}^{-1} = J_0$ .

Suppose now given a  $\mathbf{G}$ -structure  $B$  on  $V$ . Then we may define a tensor field  $J$  of type  $(1, 1)$  on  $V$  as follows:

$$J_x(X) = p \circ J_0 \circ p^{-1}(X),$$

where  $X \in T_xV$ ,  $x \in V$  and  $p \in B$  is a linear frame at  $x$ ,  $p: \mathbb{R}^{(k+1)n+1} \rightarrow T_xV$ . From the definition of  $\mathbf{G}$ ,  $J_x(X)$  is independent of the choice of  $p$ . In other words,  $J_x$  is defined as the linear endomorphism of  $T_xV$  which has at  $x$  the matrix representation  $J_0$  with respect to one of the linear frames determined at  $x$  by  $B$ , and hence with respect to any other. Also, we define a vector field  $\xi$  and a 1-form  $\omega$  on  $V$  as follows. If  $\{e_1, \dots, e_{(k+1)n+1}\}$  is the canonical basis for  $\mathbb{R}^{(k+1)n+1}$ , then  $\xi_x = p(e_{(k+1)n+1}) = p(0, \dots, 0, 1)$ , and  $\omega_x(p(e_\alpha)) = 0$ , for  $1 \leq \alpha \leq (k+1)n$  and  $\omega_x(p(e_{(k+1)n+1})) = 1$ . One can easily check that  $(J, \omega, \xi)$  defines an almost  $s$ -tangent structure of order  $k$  on  $V$ .

Summing up, we have proved the following

**PROPOSITION 3.1.** *A  $((k+1)n+1)$ -dimensional manifold admits an almost  $s$ -tangent structure of order  $k$  if and only if the structure group of its tangent bundle is reducible to  $\mathbf{G}$ .*

**COROLLARY 3.1.** *Every almost  $s$ -tangent manifold of odd order is orientable.*

The integrability of  $(J, \omega, \xi)$  as a  $\mathbf{G}$ -structure means that around each point of  $V$  there exists a coordinate system in which  $J$  is represented by the constant matrix  $J_0$ , being  $\lambda^{k+1} = 1$ ,  $J\xi = \lambda\xi$ . In other words, around each point of  $V$  there is a coordinate system  $(z_0^i, z_1^i, \dots, z_k^i, t)$  such that

$$(4) \quad J \left( \frac{\partial}{\partial z_r^i} \right) = \frac{\partial}{\partial z_{r+1}^i}, \quad 0 \leq r \leq k-1, \quad J \left( \frac{\partial}{\partial z_k^i} \right) = 0,$$

$$J \left( \frac{\partial}{\partial t} \right) = \lambda \left( \frac{\partial}{\partial t} \right), \quad \xi = \frac{\partial}{\partial t}, \quad \omega = dt.$$

Let us remark that the canonical almost  $s$ -tangent structure of order  $k$  on  $J^k(\mathbb{R}, M)$  is integrable.

**PROPOSITION 3.2.** *An almost  $s$ -tangent structure  $(J, \omega, \xi)$  of order  $k$  is integrable if and only if  $N_J = 0$  and  $d\omega = 0$ , where  $N_J$  denotes the Nijenhuis tensor of  $J$ .*

*Proof.* Clearly, if  $(J, \omega, \xi)$  is integrable then  $N_J = 0$  and  $d\omega = 0$ . Conversely, suppose that  $d\omega = 0$  and  $N_J = 0$ . Hence,  $Q$  and of course  $P$  are integrable distributions. Then, around each point of  $V$ , there exists a cubic coordinate neighbourhood  $U$ , with local coordinates  $(\bar{z}_r^i, \bar{t})$ ,  $-\varepsilon < \bar{t} < \varepsilon$ , such that

$$Q = \left\langle \frac{\partial}{\partial \bar{z}_r^i} \right\rangle, \quad P = \left\langle \frac{\partial}{\partial \bar{t}} \right\rangle.$$

Then  $\xi = f(\partial/\partial \bar{t})$  and  $\omega = g d\bar{t}$ , with  $g = 1/f$ . Since  $d\omega = 0$ , we deduce that  $\partial g/\partial \bar{z}_r^i = 0$  and so  $g = g(\bar{t})$ . Now, we introduce a new coordinate system  $(z_r^i, t')$  where  $z_r^i = \bar{z}_r^i$ ,  $t' = h(\bar{t})$ , with  $h(\bar{t})$  a primitive function of  $g(\bar{t})$ . With respect to  $(z_r^i, t')$ , we have

$$Q = \left\langle \frac{\partial}{\partial z_r^i} \right\rangle, \quad \xi = \frac{\partial}{\partial t'}, \quad \omega = dt'.$$

Since  $N_J = 0$  and  $Q$  is integrable, we know that  $J|_Q$  is an integrable almost tangent structure of order  $k$  on the integral manifolds of  $Q$  ([11]). Consider the submanifold  $W$  of  $U$  defined by  $t' = 0$ . Then, there exists a coordinate open  $U'$  in  $W$ , with local coordinates  $(\tilde{z}_r^i)$  such that

$$J \left( \frac{\partial}{\partial \tilde{z}_r^i} \right) = \frac{\partial}{\partial \tilde{z}_{r+1}^i}, \quad 0 \leq r \leq k-1, \quad J \left( \frac{\partial}{\partial \tilde{z}_k^i} \right) = 0.$$

Consider the open coordinate  $U' \times (-\varepsilon, \varepsilon)$  in  $V$ , with local coordinates  $(z_r^i = \tilde{z}_r^i, t = t')$ . We set

$$J \left( \frac{\partial}{\partial z_r^i} \right) = \sum_{\substack{0 \leq p \leq k \\ 1 \leq l \leq n}} (A_r^i)_l^p \frac{\partial}{\partial z_p^l} + B_r^i \frac{\partial}{\partial t}.$$

Since  $N_J = 0$  and  $d\omega = 0$  we deduce that  $L_\xi J = 0$ . Then

$$\frac{\partial (A_r^i)_l^p}{\partial t} = 0, \quad \frac{\partial B_r^i}{\partial t} = 0,$$

and, consequently,

$$\begin{aligned} (A_r^i)^p(z_s^j, t) &= (A_r^i)^p(z_s^j, 0) = \delta_j^i \delta_s^{r+1}, & 0 \leq i \leq k-1, \\ (A_k^i)^p(z_s^j, t) &= (A_k^i)^p(z_s^j, 0) = 0, \\ (B_r^i)(z_s^j, t) &= (B_r^i)(z_s^j, 0) = 0, & 0 \leq r \leq k. \end{aligned}$$

Hence  $J$ ,  $\omega$  and  $\xi$  are locally expressed by (4).  $\square$

Now, let  $h$  be an arbitrary Riemannian metric on  $V$  and let  $S$  be the subbundle of  $Q$  orthogonal to  $\ker J^k$  with respect to  $h$ . Then  $J_S^p: S \rightarrow Q$  is injective and  $J^p S \cap J^q S = 0$ , for  $0 \leq p, q \leq k$ ,  $p \neq q$ , where  $J^0$  is the identity tensor field. Also we have a direct sum decomposition

$$J^r S_x \oplus J^{r+1} S_x \oplus \cdots \oplus J^k S_x = (\ker J^{k-r+1})_x, \quad 0 \leq r \leq k,$$

at each point  $x$  of  $V$  and, therefore,

$$(5) \quad TV = \underbrace{S \oplus JS \oplus \cdots \oplus J^k S}_Q \oplus P.$$

Let  $\rho^r: TV \rightarrow J^r S$  be the canonical projection. We define a Riemannian metric on  $V$  by

$$g_x(X, Y) = \sum_{r=0}^k h(J^{k-r} \rho^r(X), J^{k-r} \rho^r(Y)) + \omega(X)\omega(Y),$$

for all  $X, Y \in T_x(V)$ ,  $x \in V$ . It is clear that  $g(X, \xi) = \omega(X)$ , so that  $Q$  and  $P$  are orthogonal with respect to  $g$ ; moreover,  $g_x(X, Y) = 0$  if  $X \in J^p S$ ,  $Y \in J^q S$ ,  $0 \leq p < q \leq k$ , and  $g_x(X, Y) = g_x(J^r X J^r Y)$  for  $X, Y \in S_x$ ,  $1 \leq r \leq k$ . Hence the decomposition (5) is orthogonal with respect to  $g$  and we have

**PROPOSITION 3.3.** *Let  $V(J, \omega, \xi)$  be an almost  $s$ -tangent manifold of order  $k$ ,  $P$  the one-dimensional distribution determined by  $\xi$  and  $Q = \ker \omega$ . Then there exists a subbundle  $S$  of  $Q$  and a Riemannian metric  $g$  on  $V$  such that  $S, JS, \dots, J^k S$  and  $P$  are mutually orthogonal and*

- (i)  $g(X, \xi) = \omega(X)$  for all vector fields  $X$  on  $V$ ,
- (ii)  $g_x(X, Y) = g_x(J^r X, J^r Y)$  for all  $X, Y \in S_x$ ,  $x \in V$ ,  $1 \leq r \leq k$ .



We shall say that a metric  $g$  with the properties stated in this proposition is a *compatible metric*.

Now we can obtain a reduction of the structural group  $G$  in Proposition 3.1.

**PROPOSITION 3.4.** *A  $((k + 1)n + 1)$ -dimensional differentiable manifold  $V$  admits an almost  $s$ -tangent structure of order  $k$  if and only if the structure group of its tangent bundle is reducible to  $O(n) \overset{\times}{\cong} \overset{\dots}{\dots} \overset{\times}{\cong} O(n) \times 1$ , where  $O(n) \overset{\times}{\cong} O(n)$  denotes the group of diagonal products of  $O(n)$ .*

*Proof.* Suppose  $(J, \omega, \xi)$  is an almost  $s$ -tangent structure of order  $k$  on  $V$  and let  $g$  be a compatible metric. We can choose a local orthonormal basis  $\{X_\alpha\}_{\alpha=1, \dots, n}$  for  $S$  in a coordinate neighbourhood  $U$ ; then  $\{J^r X_\alpha\}_{\alpha=1, \dots, n}$  is an orthonormal basis for  $J^r S$  on  $U$ ,  $1 \leq r \leq k$ . We call the local orthonormal basis

$$\{X_\alpha, JX_\alpha, \dots, J^k X_\alpha, \xi\}_{\alpha=1, \dots, n}$$

a  $J$ -basis. With respect to this frame, the matrix representing  $J$  is

$$J_0 = \begin{pmatrix} 0 & 0 & 0 \\ I_{kn} & 0 & 0 \\ 0 & 0 & \lambda \end{pmatrix}$$

where  $\lambda^{k+1} = 1$ ,  $J\xi = \lambda\xi$ . With respect to another  $J$ -basis it is easily verified that the transformation matrix will be of the form

$$\begin{pmatrix} A & 0 & 0 \\ & k & \\ & \ddots & \\ 0 & A & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

where the  $n \times n$  matrix  $A$  is orthogonal. The converse is trivial.  $\square$

**EXAMPLE 3.1.** (*The Riemannian Heisenberg manifold  $\Gamma \backslash H_{k+1}$* .) The real Heisenberg group  $H_{k+1}$  is the Lie subgroup of  $Gl(k + 2, \mathbb{R})$  consisting of all matrices of the form

$$a = \begin{pmatrix} 1 & X & z \\ 0 & I_{k+1} & {}^t Y \\ 0 & 0 & 1 \end{pmatrix}$$

where  $X = (x_0, \dots, x_k)$ ,  $Y = (y_0, \dots, y_k) \in \mathbb{R}^{k+1}$  and  $z \in \mathbb{R}$ .  $H_{k+1}$  is a connected simply connected Lie group of dimension  $2(k+1) + 1$  (see [9]).

A global system of coordinates  $(x^i, y^i, z)$ ,  $1 \leq i \leq k+1$ , on  $H_{k+1}$  is defined by

$$x^i(a) = x_i, \quad y^i(a) = y_i, \quad z(a) = z, \quad 0 \leq i \leq k.$$

A basis for the left invariant 1-forms on  $H_{k+1}$  is given by

$$dx^i, \quad dy^i, \quad \gamma = dz - \sum_{i=0}^k x^i dy^i, \quad 0 \leq i \leq k,$$

and its dual basis of left invariant vector fields is

$$\left\{ \frac{\partial}{\partial x^i}, \frac{\partial}{\partial y^i} + x^i \frac{\partial}{\partial z}, \frac{\partial}{\partial z}; 0 \leq i \leq k \right\}.$$

If we put

$$\hat{J} = \sum_{i=0}^{k-1} \left\{ dx^i \otimes \frac{\partial}{\partial x^{i+1}} + dy^i \otimes \left( \frac{\partial}{\partial y^{i+1}} + x^{i+1} \frac{\partial}{\partial z} \right) \right\} + \gamma \otimes \frac{\partial}{\partial z}$$

then  $(\hat{J}, \gamma, \partial/\partial z)$  is an almost  $s$ -tangent structure of order  $k$  on  $H_{k+1}$  and the left invariant metric defined by

$$\langle \cdot, \cdot \rangle = \sum_{i=0}^k \{ dx^i \otimes dx^i + dy^i \otimes dy^i \} + \gamma \otimes \gamma$$

is a compatible metric.

Let  $\Gamma$  be the subgroup of matrices of  $H_{k+1}$  with integer entries and consider the space of right cosets  $\Gamma \backslash H_{k+1}$ . If  $g$  is the unique Riemannian metric on  $\Gamma \backslash H_{k+1}$  for which the canonical projection  $\pi$  from  $(H_{k+1}, \langle \cdot, \cdot \rangle)$  to  $\Gamma \backslash H_{k+1}$  is a Riemannian covering, the pair  $(\Gamma \backslash H_{k+1}, g)$  is called a Riemannian Heisenberg manifold (see [8]). Since the tensor fields that we have considered on  $H_{k+1}$  are left invariant they descend to  $\Gamma \backslash H_{k+1}$ . So we obtain an almost  $s$ -tangent structure  $(J, \omega, \xi)$  of order  $k$  on  $\Gamma \backslash H_{k+1}$ , with

$$J \circ \pi_* = \pi_* \circ \hat{J}, \quad \pi^* \omega = \gamma, \quad \xi = \pi_* \left( \frac{\partial}{\partial z} \right).$$

Moreover  $g$  is a compatible metric. Consider the subbundle  $S$  of  $Q = \ker \omega$  that is orthogonal to  $\ker J^k$  with respect to the metric  $g$ . Then each distribution  $J'S$  is generated by the vector fields

$X_r = \pi_*(\partial/\partial x^r)$  and  $Y_r = \pi_*(\partial/\partial y^r + x^r\partial/\partial z)$ . Since  $[X_r, Y_r] = \xi$  for  $0 \leq r \leq k$  and the other bracket products are all zero, then  $N_J(X_r, Y_r) \neq 0$  and the structure  $(J, \omega, \xi)$  is not integrable. Notice that the distribution  $Q$  is not integrable either.

**4. Submanifolds of almost tangent manifolds of order  $k$ .** Let  $W$  be a differentiable manifold of dimension  $(k+1)(n+1)$  with an almost tangent structure  $F$  of order  $k$ , that is  $F^{k+1} = 0$ ,  $\text{rank } F = k(n+1)$ . Let  $V$  be a submanifold of  $W$  of codimension  $k$  and  $\iota: V \rightarrow W$  the imbedding map. Suppose that there exists a vector field  $\xi$  on  $V$  such that the vector fields  $C = F\iota_*\xi, FC, \dots, F^{k-1}C$  along  $V$  are linearly independent at each point and nowhere tangent to  $V$ . Then the vector field  $C - \iota_*\xi$  along  $V$  is also nowhere tangent to  $V$  and  $C - \iota_*\xi, FC, \dots, F^{k-1}C$  are linearly independent at each point of  $V$ .

Now, we define a tensor field  $J$  of type  $(1, 1)$  and  $k$  differential 1-forms  $\alpha_1, \dots, \alpha_k$  on  $V$  by

$$(6) \quad F\iota_*X = \iota_*JX + \alpha_1(X)(C - \iota_*\xi) + \alpha_2(X)FC + \dots + \alpha_k(X)F^{k-1}C.$$

Applying  $F$  to both sides of (6)  $k$  times we have

$$\begin{aligned} 0 &= \iota_*(J^{k+1}(X) - \alpha_1(J^k X)\xi) + (\alpha_1(J^k X) - \alpha_1(J^{k-1} X))C \\ &\quad + (\alpha_2(J^k X) + \alpha_1(J^{k-1} X) - \alpha_1(J^{k-2} X))FC \\ &\quad + \dots + (\alpha_k(J^k X) + \alpha_{k-1}(J^{k-1} X) + \dots + \alpha_1(JX) - \alpha_1(X))F^{k-1}C. \end{aligned}$$

In particular, we have

$$J^{k+1}(X) = \alpha_1(J^{k-1} X)\xi, \quad \alpha_1(J^k X) = \alpha_1(J^{k-1} X).$$

Putting  $\omega = \alpha_1 \circ J^{k-1}$ , we get  $J^{k+1} = \omega \otimes \xi$ . Now, setting  $X = \xi$  in (6) we obtain

$$C = \iota_*(J\xi - \alpha_1(\xi)\xi) + \alpha_1(\xi)C + \alpha_2(\xi)FC + \dots + \alpha_k(\xi)F^{k-1}C.$$

Thus  $\alpha_1(\xi) = 1$ ,  $J\xi = \xi$  and  $\omega(\xi) = \alpha_1(J^{k+1}\xi) = 1$ . Since  $\text{rank } F = k(n+1)$ , then  $\text{rank } J = kn+1$ . Hence, we have proved the following

**PROPOSITION 4.1.** *Let  $W$  be a  $(k+1)(n+1)$ -dimensional differentiable manifold with an almost tangent structure  $F$  of order  $k$  and let  $(V, \iota)$  a submanifold of  $W$  of codimension  $k$ . If there exists a vector field  $\xi$  on  $V$  such that  $C = F\iota_*\xi, FC, \dots, F^{k-1}C$  are linearly independent at each point and nowhere tangent to  $V$ , then  $V$  admits an almost  $s$ -tangent structure of order  $k$ .*

EXAMPLE 4.1 ([20]). The tangent sphere bundle of a Riemannian manifold  $M$  inherits an almost  $s$ -tangent structure of order 1 from the canonical almost tangent structure of the tangent bundle  $TM$ .

EXAMPLE 4.2. (*A submanifold of  $T^2M$ .*) Let  $M$  be an  $(n + 1)$ -dimensional Riemannian manifold with metric  $g$ ,  $T^2M$  its tangent bundle of order 2 and  $TM \oplus TM$  the Whitney sum of the tangent bundle of  $M$  with itself. Let  $\{U, x_i\}$  be a coordinate neighbourhood of  $M$  and  $(x^i, v^i, w^i)$  the corresponding coordinates of  $TM \oplus TM$ , where  $(v^i, w^i)$  are the fibre coordinates. On the other hand, let  $(x^i, y^i, z^i)$  be the induced coordinates in  $T^2U$ , that is, if  $j^2\sigma$  is the 2-jet at  $0 \in \mathbb{R}$  of a differentiable curve  $\sigma: \mathbb{R} \rightarrow M$  then

$$x^i(j^2\sigma) = x^i(\sigma(0)), \quad y^i(j^2\sigma) = \frac{d(x^i \circ \sigma)}{dt}(0),$$

$$z^i(j^2\sigma) = \frac{1}{2} \frac{d^2(x^i \circ \sigma)}{dt^2}(0).$$

If  $\nabla$  is the Riemannian connection of  $g$ , there exists a diffeomorphism ([5], [18])  $\varphi: T^2M \rightarrow TM \oplus TM$  given by

$$\varphi(j^2\sigma) = (\dot{\sigma}(0), (\nabla_{\dot{\sigma}(0)}\dot{\sigma})(0))$$

where  $\dot{\sigma}$  is the tangent vector field to  $\sigma$ . In coordinates

$$T^2U \xrightarrow{\varphi} TU \oplus TU$$

$$(x^i, y^i, z^i) \mapsto (x^i, y^i, 2z^i + y^j y^k \Gamma_{jk}^i),$$

$\Gamma_{jk}^i$  are the local components of the Riemannian connection.

Let  $F$  be the standard almost tangent structure of order 2 on  $T^2M$ , given by

$$F\left(\frac{\partial}{\partial x^i}\right) = \frac{\partial}{\partial y^i}, \quad F\left(\frac{\partial}{\partial y^i}\right) = \frac{\partial}{\partial z^i}, \quad F\left(\frac{\partial}{\partial z^i}\right) = 0.$$

By means of the diffeomorphism  $\varphi$ , the tensor field  $F$  defines an almost tangent structure  $\bar{F}$  of order 2 on  $TM \oplus TM$ , which is given by

$$\bar{F}\left(\frac{\partial}{\partial x^i}\right) = \frac{\partial}{\partial v^i} + 2v^k \Gamma_{ik}^j \frac{\partial}{\partial w^j},$$

$$\bar{F}\left(\frac{\partial}{\partial v^i}\right) = 2\frac{\partial}{\partial w^i}, \quad \bar{F}\left(\frac{\partial}{\partial w^i}\right) = 0.$$

Let us consider the tangent sphere bundle  $V_1M$  of  $M$ , and let  $V_2M$  be the subbundle of the fibre product  $V_1M \oplus V_1M$ , whose fibre at each

point  $x \in M$  is

$$(V_2M)_x = \{(u, v) \in (V_1M)_x \oplus (V_1M)_x \mid g(u, v) = 0\}.$$

We consider  $\bar{V} = V_1M \oplus V_1M - V_2M$ , which is the submanifold of codimension 2 of  $TM \oplus TM$  defined by the conditions

$$g_{ij}v^i v^j = g_{ij}w^i w^j = 1, \quad g_{ij}v^i w^j \neq 0.$$

Let  $\iota: \bar{V} \rightarrow TM \oplus TM$  be the imbedding map. We define a vector field  $\bar{\xi}$  on  $\bar{V}$  by

$$\iota_*\bar{\xi} = v^i \left( \frac{\partial}{\partial x^i} \right)^H$$

where  $(\partial/\partial x^i)^H$  is the horizontal lift of  $\partial/\partial x^i$  to  $TM \oplus TM$  with respect to the Riemannian connection of  $M$ ; that is

$$\left( \frac{\partial}{\partial x^i} \right)^H = \frac{\partial}{\partial x^i} - \Gamma_{ik}^j v^k \frac{\partial}{\partial v^j} - \Gamma_{ik}^j w^k \frac{\partial}{\partial w^j}.$$

The vector fields

$$C = \bar{F}\iota_*\bar{\xi} = v^i \frac{\partial}{\partial v^i} \quad \text{and} \quad \bar{F}C = 2v^i \frac{\partial}{\partial w^i}$$

are nowhere tangent to  $\bar{V}$  and they are linearly independent at each point of  $\bar{V}$ . By Proposition 4.1,  $\bar{V}$  inherits an almost tangent structure  $(\bar{J}, \bar{\omega}, \bar{\xi})$  of order 2, which is given by

$$\bar{F}\iota_*X = \iota_*\bar{J}X + \alpha_1(X)(C - \iota_*\bar{\xi}) + \alpha_2(X)\bar{F}C, \quad \bar{\omega}(X) = \alpha_1(\bar{J}X).$$

Hence  $(J = \varphi^{-1}\bar{J}\varphi_*, \omega = \varphi^*\bar{\omega}, \xi = \varphi^{-1}\bar{\xi})$  is an almost  $s$ -tangent structure of order 2 on the submanifold  $V = \varphi^{-1}(\bar{V})$  of  $T^2M$ .

### REFERENCES

- [1] R. S. Clark and M. R. Bruckheimer, *Sur les structures presque tangentés*, C. R. Acad. Sci. Paris, **251** (1960), 627–629.
- [2] M. Crampin, *On the differential geometry of the Euler-Lagrange equations, and the inverse problem of Lagrangian dynamics*, J. Phys. A: Math. Gen., **14** (1981), 2567–2575.
- [3] F. Cantrijn, J. F. Cariñena, M. Crampin and L. A. Ibort, *Reduction of degenerate Lagrangian systems*, J. Geom. Phys., **3** (3) (1986), 353–400.
- [4] M. Crampin, W. Sarlet and F. Cantrijn, *Higher-order differential equations and higher-order Lagrangian dynamics*, Math. Proc. Cambridge Philos. Soc., **99** (1986), 565–587.
- [5] C. T. J. Dodson and M. S. Radiiovici, *Tangent and frame bundles of order two*. I, An. Ştiinţ. Univ. “Al. I. Cuza” Iaşi Sect. I a Mat., **28** (1982), 63–71.

- [6] H. Eliopoulos, *Structures presque-tangentes sur les variétés différentiables*, C. R. Acad. Sci. Paris, **255** (1962), 1563–1565.
- [7] —, *Structures  $r$ -tangentes sur les variétés différentiables*, C. R. Acad. Sci. Paris, **263** (1966), 413–416.
- [8] C. S. Gordon and E. N. Wilson, *The spectrum of the laplacian on Riemannian Heisenberg manifolds*, Michigan Math. J., **33** (1986), 253–271.
- [9] Y. Haraguchi, *Sur une généralisation des structures de contact*, Thèse, Univ. du Haute Alsace, Mulhouse, 1981.
- [10] L. A. Iborn and C. López-Lacasta, *On the existence of local and global Lagrangians for ordinary differential equations*, J. Phys. A: Math. Gen., **23** (1990), 4779–4792.
- [11] J. Lehmann-Lejeune, *Sur l'intégrabilité de certaines  $G$ -structures*, C. R. Acad. Sci. Paris, **258** (1964), 5326–5329.
- [12] M. de León, A. Giraldo and P. R. Rodrigues, *Sur la réduction de systèmes lagrangiens dégénérés d'ordre supérieur*, C. R. Acad. Sci. Paris Sér. I Math., **309** (1989), 363–366.
- [13] M. de León, M. H. Mello and P. R. Rodrigues, *Reduction of degenerate non-autonomous Lagrangian systems*, preprint UM-CSIC, 1991.
- [14] M. de León and P. R. Rodrigues, *Generalized Classical Mechanics and Field Theory*, North-Holland Math. Ser. 112, Amsterdam, 1985.
- [15] —, *Method of Differential Geometry in Analytical Mechanics and Field Theory*, North-Holland Math. Ser. 158, Amsterdam, 1989.
- [16] —, *Regular Lagrangian dynamical systems of higher order*, preprint UM-CSIC, 1989.
- [17] —, *Lagrangiens dégénérés d'ordre supérieur et réduction des systèmes hamiltoniens associés*, preprint UM-CSIC, 1990.
- [18] M. de León and E. Vázquez, *On the geometry of the tangent bundle of order 2*, An. Univ. București, **34** (1985), 40–48.
- [19] A. Morimoto, *Liftings of tensor fields and connections to tangent bundles of higher order*, Nagoya Math. J., **40** (1970), 99–120.
- [20] J. A. Oubiña, *Almost  $s$ -tangent structures*, Geom. Dedicata, **14** (1983), 395–403.

Received February 12, 1991 and in revised form June 5, 1991. Partilly supported by DGICYT-Spain, proyecto PB88-0012, Xunta de Galicia, Spain, Proxecto XUGA8050189 and CNPq-Brazil, Proc. 301115/79.

UNIDAD DE MATEMÁTICAS  
 CONSEJO SUPERIOR DE INVESTIGACIONES CIENTÍFICAS  
 SERRANO 123, 28006 MADRID, SPAIN

DEPARTAMENTO DE XEOMETRÍA E TOPOLOXÍA  
 FACULTADE DE MATEMÁTICAS  
 UNIVERSIDADE DE SANTIAGO DE COMPOSTELA, SPAIN

AND

DEPARTAMENTO DE GEOMETRIA  
 INSTITUTO DE MATEMATICA  
 UNIVERSIDADE FEDERAL FLUMINENSE  
 24.210 NITEROI, RJ, BRAZIL



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112

F. MICHAEL CHRIST  
University of California  
Los Angeles, CA 90024-1555

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093

NICHOLAS ERCOLANI  
University of Arizona  
Tucson, AZ 85721

R. FINN  
Stanford University  
Stanford, CA 94305

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305

C. C. MOORE  
University of California  
Berkeley, CA 94720

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH  
(1906-1982)

B. H. NEUMANN

F. WOLF  
(1904-1989)

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555-05.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1992 by Pacific Journal of Mathematics



<b>Manuel (Rodriguez) de León, J. A. Oubiña, P. R. Rodrigues and Modesto R. Salgado</b> , Almost $s$ -tangent manifolds of higher order . . . . .	201
<b>Martin Engman</b> , New spectral characterization theorems for $S^2$ . . . . .	215
<b>Yuval Zvi Flicker</b> , The adjoint representation $L$ -function for $GL(n)$ . . . . .	231
<b>Enrique Alberto Gonzalez-Velasco and Lee Kenneth Jones</b> , On the range of an unbounded partly atomic vector-valued measure . . . . .	245
<b>Takayuki Hibi</b> , Face number inequalities for matroid complexes and Cohen-Macaulay types of Stanley-Reisner rings of distributive lattices . . . . .	253
<b>Hervé Jacquet and Stephen James Rallis</b> , Kloosterman integrals for skew symmetric matrices . . . . .	265
<b>Shulim Kaliman</b> , Two remarks on polynomials in two variables . . . . .	285
<b>Kirk Lancaster</b> , Qualitative behavior of solutions of elliptic free boundary problems . . . . .	297
<b>Feng Luo</b> , Actions of finite groups on knot complements . . . . .	317
<b>James Joseph Madden and Charles Madison Stanton</b> , One-dimensional Nash groups . . . . .	331
<b>Christopher K. McCord</b> , Estimating Nielsen numbers on infrасolvmanifolds . . . . .	345
<b>Gordan Savin</b> , On the tensor product of theta representations of $GL_3$ . . . . .	369
<b>Gerold Wagner</b> , On means of distances on the surface of a sphere. II. (Upper bounds) . . . . .	381