

*Pacific  
Journal of  
Mathematics*

**ORIENTATION AND STRING STRUCTURES ON LOOP SPACE**

DENNIS MCLAUGHLIN

# ORIENTATION AND STRING STRUCTURES ON LOOP SPACE

DENNIS A. McLAUGHLIN

**This paper deals rigorously with the notion of a string structure and the topological obstruction to its existence. The question of orienting loop space is discussed and shown to be directly analogous to orientation of finite dimensional manifolds. Finally, equivariant string structures are considered.**

**1. Introduction.** Physicists working on the grand unification program have recently been led to consider particles, not as points on some manifold  $M$ , but rather as loops on  $M$ . This novel idea has resulted in efforts to formulate a theory of spinors on  $LM$ , the free loop space of  $M$  [3]. The theory is called string theory.

Since  $LM$  is an infinite dimensional manifold, placing this on a rigorous mathematical footing is a real challenge. The first problem is to define the Dirac operator and involves constructing a spinor bundle on which it acts. E. Witten and T. P. Killingback have made considerable progress in this regard.

Witten in [19], [20] and Atiyah in [2] argued that  $LM$  should be considered orientable exactly when  $M$  is a spin manifold. Killingback [10] looked at bundles on  $LM$  whose structural groups are loop groups. He defined a string structure as a lifting of the structural group to a central extension of the loop group by a circle. The candidate for the spinor bundle on  $LM$  is then a certain infinite dimensional vector bundle associated with the string structure. Just as in finite dimensions, there is a topological obstruction to defining this bundle. Killingback argued that it is essentially the first Pontrjagin class of  $M$ . In this paper, we clarify these results and prove them rigorously.

In §2, we examine the orientability of loop space. Suppose that  $P \rightarrow M$  is an  $SO(n)$ -bundle. By taking free loops, we obtain an  $LSO(n)$ -bundle  $LP \rightarrow LM$  in a natural way. Assuming that  $M$  is simply connected, we show that it is possible to reduce the structural group of  $LP \rightarrow LM$  to the connected component of the identity if and only if  $P \rightarrow M$  admits a spin structure. The condition that  $M$  be simply connected is reasonable, since it is equivalent to  $LM$

being connected. We will see that  $LSO(n)$  is the structural group of the tangent bundle to loop space. Combining this result with the observation of Atiyah and Witten, we find that orientability of loop space is exactly analogous to orientation in finite dimensions.

Now suppose that  $P \rightarrow M$  has a spin structure  $Q \rightarrow M$ . A string structure for this bundle is a lifting of the structural group of  $LQ \rightarrow LM$  to a central extension of  $LSpin(n)$  by the circle. In §3, we prove (compare [10])

**THEOREM.** *Suppose that  $M$  is simply connected and  $n \geq 5$ . Then  $LQ \rightarrow LM$  has a string structure if  $\frac{1}{2}p_1(P) = 0$ , where  $p_1(P)$  is the first Pontrjagin class of  $P \rightarrow M$ . The converse is also true if we further assume that  $\pi_2(M) = 0$ .*

In §4, we consider the case where a compact, abelian group  $G$  acts on  $M$ . It is easy to see that there exists a  $G$ -equivariant spin structure if and only if the equivariant Stiefel-Whitney class  $w_2(P)_G$  vanishes. As noted in [19], the analogous result for string theory would be that a  $G$ -equivariant string structure exists if  $\frac{1}{2}p_1(P)_G$  is zero. We will prove this statement and show also that the converse is, in general, false. Finally, we show that the spinor bundle on  $LM$  is equivariant under the action of rotating loops (at least if  $\pi_2(M) = 0$ ). The novelty here is that this action is not induced from one on  $M$ .

I thank J.-L. Brylinski for drawing my attention to the problem of defining the spinor bundle on  $LM$  and for many useful comments. I also thank T. Goodwillie for pointing out some corrections.

**2. Orientability of loop space.** Let  $M$  be a compact, connected, orientable, Riemannian manifold of dimension  $n$ .  $LM$  will denote the space of smooth loops on  $M$ . It is an infinite dimensional, paracompact manifold modelled on the topological vector space  $L\mathbb{R}^n$  (with the topology of uniform convergence of the functions and all their derivatives) [14, Chapter 3]. If  $M$  happens to be a Lie group, then  $LM$  is an infinite dimensional Lie group—a loop group.

A tangent vector to a loop  $\gamma$  in  $M$  is an “infinitesimal deformation” of  $\gamma$  and therefore can be regarded as a map  $v: \theta \rightarrow T_{\gamma(\theta)}M$ . Thus, we see that the space  $T_\gamma LM$  of all such vectors is precisely the space of sections of the pullback bundle  $\gamma^*TM \rightarrow S^1$ . But this is a trivial bundle since  $M$  is orientable. Therefore, we can identify  $T_\gamma LM$  with  $L\mathbb{R}^n$ . In this way, we see that the tangent bundle to loop space is an infinite dimensional, locally trivial vector bundle with fiber  $L\mathbb{R}^n$ . It is

associated to the  $L\text{SO}(n)$ -bundle  $LFM \rightarrow LM$ , obtained in a natural way by taking free loops on the frame bundle  $FM \rightarrow M$ , (compare [10]). Thus,  $LFM \rightarrow LM$  plays the role of the frame bundle on loop space.

However, as topological spaces,  $L\text{SO}(n) \cong \Omega\text{SO}(n) \times \text{SO}(n)$ , where  $\Omega\text{SO}(n)$  denotes the pointed loop space. It follows that  $L\text{SO}(n)$  has two connected components,  $n \geq 3$ . The question arises whether the structural group of  $LFM \rightarrow LM$  can be reduced to  $L^0\text{SO}(n)$ , the connected component of the identity. A complete answer is given by

**PROPOSITION 2.1.** *Suppose that  $M$  is simply connected and that  $P \rightarrow M$  is an  $\text{SO}(n)$ -bundle,  $n \geq 4$ . The following are equivalent:*

- (1) *The structural group of  $LP \rightarrow LM$  is reducible to  $L^0\text{SO}(n)$ .*
- (2)  *$P \rightarrow M$  has a spin structure.*
- (3) *The structural group of  $LP \rightarrow LM$  can be lifted to  $L\text{Spin}(n)$ .*

*Proof.* (1)  $\Leftrightarrow$  (2). First we characterize the obstruction to reducing the structural group of  $LP \rightarrow LM$  to  $L^0\text{SO}(n)$ . There are classifying spaces  $B\text{L}\text{SO}(n)$  and  $BL^0\text{SO}(n)$ , for  $L\text{SO}(n)$  and  $L^0\text{SO}(n)$ -bundles respectively. Moreover, the inclusion map  $i: L^0\text{SO}(n) \rightarrow L\text{SO}(n)$  induces a double covering  $B(i): BL^0\text{SO}(n) \rightarrow B\text{L}\text{SO}(n)$  [4]. Reducing the structural group of  $LP \rightarrow LM$  corresponds to lifting its classifying map in the following diagram.

$$\begin{array}{ccc}
 & & BL^0\text{SO}(n) \\
 & & \downarrow B(i) \\
 LM & \longrightarrow & B\text{L}\text{SO}(n)
 \end{array}$$

Since  $BL^0\text{SO}(n)$  is connected, the double covering  $B(i)$  is non-trivial. There is a unique, well-defined obstruction to finding a section of  $B(i)$  and it lies in  $H^1(B\text{L}\text{SO}(n); \mathbb{Z}_2)$ . From the Hurewicz Theorem, we see that this group is  $\mathbb{Z}_2$ . Therefore, the non-zero element must generate the obstruction. By functoriality, we conclude that the obstruction to reducing the structural group of  $LP \rightarrow LM$  is the pull-back of this element by  $f$ . We denote it by  $\lambda(P)$ .

The evaluation map  $ev: LM \times S^1 \rightarrow M$  defined by evaluating a loop against an angle, induces a map  $ev^*: H^2(M) \rightarrow H^2(LM \times S^1)$ . Composing this with “integration over  $S^1$ ” (denoted  $\int_{S^1}$ ), we obtain a map  $\int_{S^1} \circ ev^*: H^2(M) \rightarrow H^1(LM)$ . To complete the proof, we

show that this composition carries the second Stiefel-Whitney class of  $P \rightarrow M$  isomorphically to  $\lambda(P)$ .

Taking free loops on the universal  $SO(n)$ -bundle  $ESO(n) \rightarrow BSO(n)$  yields a bundle with contractible total space, on which  $LSO(n)$  acts freely. This means that  $LESO(n) \rightarrow LBSO(n)$  is actually a model for the universal  $LSO(n)$ -bundle. Furthermore, we see that by “looping” the classifying map of  $P \rightarrow M$ , we obtain the classifying map of  $LP \rightarrow LM$ . But,  $w_2(P)$  is the pullback of the generator of  $H^2(BSO(n); \mathbb{Z}_2)$  under the classifying map of  $P \rightarrow M$ . Therefore, the assertion will follow, if we show that  $\int_{S^1} \circ ev^*$  is an isomorphism when  $M$  is simply connected.

We work on the level of homology. By the Hurewicz Theorem,  $H_2(M)$  is generated by some map  $f: S^2 \rightarrow M$ . But,  $S^2$  can be covered by loops which meet only at one point, and the parameter space for this set of loops is a copy of  $S^1$ . Using this, we obtain a map  $g: S^1 \rightarrow LM$  and a corresponding element  $[g]$  in  $H_1(LM)$ . The map  $g$  has the property that its evaluation against the circle is exactly  $f$ . Thus,  $ev_*$  maps  $[g] \otimes [id] \in H_2(LM \times S^1)$  to  $[f]$ , the generator of  $H_2(M)$ . Conversely, starting with a map  $g: S^1 \rightarrow LM$ , we can realize  $S^1$  as the parameter space for such loops and then evaluate to produce  $f: S^2 \rightarrow M$ . We conclude that  $\int_{S^1} \circ ev^*$  is an isomorphism if  $\pi_1(M) = 0$ .

(2)  $\Leftrightarrow$  (3). There is another evaluation map  $ev_0: LM \rightarrow M$  which maps a loop to its basepoint. It induces a bundle morphism

$$\begin{array}{ccc} LP & \longrightarrow & P \\ \downarrow & & \downarrow \\ LM & \longrightarrow & M \end{array}$$

which in turn induces a morphism of the associated spectral sequences. This leads to the following commutative diagram.

$$\begin{array}{ccccccc} \dots & \longrightarrow & H^1(P; \mathbb{Z}_2) & \longrightarrow & H^1(SO(n); \mathbb{Z}_2) & \longrightarrow & H^2(M; \mathbb{Z}_2) \dots \\ & & \downarrow & & \downarrow & & \downarrow \\ \dots & \longrightarrow & H^1(LP; \mathbb{Z}_2) & \longrightarrow & E^{0,1} & \longrightarrow & E^{2,0} \dots \end{array}$$

Since  $H^1(LSO(n); \mathbb{Z}_2) \cong \mathbb{Z}_2$ , it follows that  $E^{0,1} \cong \mathbb{Z}_2$ . Lifting the structural group of  $LP \rightarrow LM$  to  $LSpin(n)$ , corresponds to finding an element of  $H^1(LP; \mathbb{Z}_2)$  which restricts to the generator of  $E^{0,1}$ . By exactness, the image of this element in  $E^{2,0}$  is the obstruction. We denote it by  $\nu(P)$ .

We know from [1] that  $w_2(P)$  is the image of the generator of  $H^1(\mathrm{SO}(n); \mathbb{Z}_2)$  in  $H^2(M; \mathbb{Z}_2)$ . Moreover,  $ev_0$  has an obvious section obtained by regarding  $M$  as the constant loops in  $LM$ . Therefore,  $ev_0^*$  is injective, and we conclude that it carries  $w_2(P)$  isomorphically to  $\nu(P)$ , completing the proof.

Witten [20] and Atiyah [2] both argued that  $LM$  should be considered orientable exactly when  $M$  is spin. Combining this with the above proposition, we see that orientation of loop space is directly analogous to orientation of finite dimensional manifolds.

**COROLLARY 2.2.**  *$LM$  is orientable if and only if the structural group of the tangent bundle can be reduced to the connected component of the identity.*

**3. String structures on loop space.** To define a spin structure in finite dimensions, one lifts the structural group of the frame bundle  $FM \rightarrow M$  to  $\mathrm{Spin}(n)$ , the universal cover of  $\mathrm{SO}(n)$  for  $n \geq 3$ . The corresponding notion for loop space is called a string structure, which we now describe.

Let  $Q \rightarrow M$  be a spin structure for the  $\mathrm{SO}(n)$ -bundle  $P \rightarrow M$ . Following Killingback in [10], we define a string structure to be a lifting of the structural group of  $LQ \rightarrow LM$  to  $L\widehat{\mathrm{Spin}}(n)$ , a non-trivial, central extension of  $L\mathrm{Spin}(n)$  by  $S^1$ . We will be especially interested in  $L\widehat{\mathrm{Spin}}(n)$ , the universal such extension. For the existence of such extensions and the fact that they are completely determined by their topological class as  $S^1$ -bundles, see [14, Chapter 4].

The motivation for this definition is the following: In trying to formally define the Dirac operator on loop space, one is led to an infinite dimensional Clifford algebra. The spinor representation of this algebra is then a representation of some extension of  $L\mathrm{SO}(n)$ . The novelty here is that it is an extension by a circle, rather than a discrete group as one might expect [14, Chapter 12]. It follows that the spinor representation is also a representation of some extension of  $L\mathrm{Spin}(n)$  by the circle. We will return to this later. We want to prove (compare [10])

**THEOREM 3.1.** *Let  $P \rightarrow M$  be an  $\mathrm{SO}(n)$ -bundle over a simply connected manifold,  $n \geq 5$ . Suppose that  $Q \rightarrow M$  is a spin structure for this bundle. Then  $LQ \rightarrow LM$  has a string structure if  $\frac{1}{2}p_1(P)$  vanishes, where  $p_1(P)$  denotes the first Pontrjagin class of  $P \rightarrow M$ . The converse is also true, if we further assume that  $M$  is doubly connected.*

First, we explain what we mean by  $\frac{1}{2}p_1(P)$ .

**LEMMA 2.2.** *A spin structure  $Q \rightarrow M$  for the  $SO(n)$ -bundle  $P \rightarrow M$  determines a class in  $H^4(M; \mathbb{Z})$ , which when multiplied by 2 is exactly  $p_1(P)$ . It is the pullback of the generator of  $H^4(B \operatorname{Spin}(n); \mathbb{Z})$  by the classifying map of  $Q \rightarrow M$ .*

*Proof.* The projection  $\pi: \operatorname{Spin}(n) \rightarrow \operatorname{SO}(n)$  induces a fibration  $B(\pi): B \operatorname{Spin}(n) \rightarrow B \operatorname{SO}(n)$  with fiber  $B\mathbb{Z}_2$  [4]. The classifying maps of  $P \rightarrow M$  and  $Q \rightarrow M$  induce the following commutative diagram.

$$\begin{array}{ccc} H^4(B \operatorname{SO}(n); \mathbb{Z}) & \longrightarrow & H^4(M; \mathbb{Z}) \\ \downarrow B(\pi)^* & & \downarrow \operatorname{id} \\ H^4(B \operatorname{Spin}(n); \mathbb{Z}) & \longrightarrow & H^4(M; \mathbb{Z}) \end{array}$$

We will show that the image of  $B(\pi)^*$  has index 2 and that both groups in the left-hand column are  $\mathbb{Z}$ . Since  $p_1(P)$  is the image of the generator of  $H^4(B \operatorname{SO}(n); \mathbb{Z})$  under the classifying map of  $P \rightarrow M$ , the result will follow.

The integral cohomology of  $B \operatorname{SO}(n)$  can be computed using the following facts [4], [13]: The only torsion in  $H^*(B \operatorname{SO}; \mathbb{Z})$  is 2-torsion and  $H^*(B \operatorname{SO}; \mathbb{Z}_2)$  is a polynomial algebra generated by the Stiefel-Whitney classes  $w_2, w_3, \dots$ . Moreover,  $H^*(B \operatorname{SO}; \mathbb{Q})$  is a polynomial algebra generated by elements in degree  $4i$ , the Pontrjagin classes. We deduce that  $H^i(B \operatorname{SO}(n); \mathbb{Z})$  is zero in degrees 1 and 2,  $\mathbb{Z}_2$  in degrees 3 and 5, and  $\mathbb{Z}$  in degree 4. On the other hand, it follows from the Hurewicz Theorem that  $H^4(B \operatorname{Spin}(n); \mathbb{Z}) \cong \mathbb{Z}$ .

It is well known [7] that  $H^*(B\mathbb{Z}_2; \mathbb{Z}) \cong \mathbb{Z}_2[a]/(2a)$ . Therefore, in the spectral sequence associated to  $B(\pi)$ ,  $d_3: E^{0,4} \rightarrow E^{3,2}$  maps  $a^2$ , the generator of  $H^4(B\mathbb{Z}_2; \mathbb{Z})$ , to  $2a \cdot da$  and so must be zero. The only other relevant differential is  $d_3: E^{2,2} \rightarrow E^{5,0}$ . This is actually an isomorphism. To see this, note that both groups are  $\mathbb{Z}_2$ . If  $d_3$  were the zero map,  $E^{5,0}$  would survive in the spectral sequence giving something non-zero in  $H^5(B \operatorname{Spin}(n))$ . But  $B \operatorname{Spin}(n)$  is 3-connected so the Hurewicz homomorphism is surjective in degree 5. Since  $\pi_5(B \operatorname{Spin}(n)) \cong \pi_4(\operatorname{SO}(n))$  is at most 2-torsion [17, Part 2], we must have  $H^5(B \operatorname{spin}(n); \mathbb{Z}) = 0$ —a contradiction.

These facts yield a short exact sequence

$$0 \rightarrow H^4(B \operatorname{SO}(n); \mathbb{Z}) \rightarrow H^4(B \operatorname{Spin}(n); \mathbb{Z}) \rightarrow \mathbb{Z}_2 \rightarrow 0,$$

and the lemma follows.

Now we discuss the obstruction to defining a string structure. Since  $L\text{Spin}(n)$  is connected and simply connected, the spectral sequence of the bundle  $LQ \rightarrow LM$  degenerates in low degrees and we obtain an exact sequence

$$\begin{aligned} 0 \rightarrow H^2(LM; \mathbb{Z}) &\rightarrow H^2(LQ; \mathbb{Z}) \\ &\rightarrow H^2(L\text{Spin}(n); \mathbb{Z}) \rightarrow H^3(LM; \mathbb{Z}) \rightarrow \dots \end{aligned}$$

Lifting the structural group of  $LQ \rightarrow LM$  to  $L\widehat{\text{Spin}}(n)$  corresponds to finding a circle bundle  $\widehat{LQ}$  over  $LQ$  which restricts to the bundle  $L\widehat{\text{Spin}}(n) \rightarrow L\text{Spin}(n)$  in each fiber. Circle bundles over  $LQ$  and  $L\text{Spin}(n)$  are classified by elements of  $H^2(LQ; \mathbb{Z})$  and  $H^2(L\text{Spin}(n); \mathbb{Z})$ . The latter group is  $\mathbb{Z}$  and the bundle  $L\widehat{\text{Spin}}(n) \rightarrow L\text{Spin}(n)$  corresponds to the generator. We conclude that the existence of a string structure corresponding to  $L\widehat{\text{Spin}}(n)$  is equivalent to the existence of an element of  $H^2(LQ; \mathbb{Z})$ , which restricts to the generator of  $H^2(L\text{Spin}(n); \mathbb{Z})$ . From the exact sequence above, the image of the generator in  $H^3(LM; \mathbb{Z})$  is the obstruction to defining this string structure. We denote it by  $\mu(Q)$ . It is the pullback of the generator of  $H^3(LB\text{Spin}(n); \mathbb{Z})$  by the classifying map of  $LQ \rightarrow LM$ . The obstruction to defining a string structure corresponding to any other central extension, is just some multiple of  $\mu$ . Note also, that inequivalent string structures are classified by elements of  $H^2(LM; \mathbb{Z})$ .

We prove Theorem 3.1 by showing that  $\int_{S^1} \circ ev^*$  maps  $\frac{1}{2}p_1(P)$  to  $\mu$  and does so injectively when  $M$  is 2-connected. By functoriality, the first assertion will follow from the fact that

$$\int_{S^1} \circ ev^*: H^4(B\text{Spin}(n); \mathbb{Z}) \rightarrow H^3(LB\text{Spin}(n); \mathbb{Z})$$

is an isomorphism for  $n \geq 5$ . To see this, we use a similar argument to the one given in Proposition 2.1. By the Hurewicz Theorem,  $H_4(B\text{Spin}(n))$  is generated by a map  $f: S^4 \rightarrow B\text{Spin}(n)$ . If we cover  $S^4$  by loops meeting at only one point, the parameter space for such loops is  $S^3$ . Using this, we produce a map  $g: S^3 \rightarrow L\text{Spin}(n)$  which “evaluates” to  $f$ . Proceeding exactly as before yields the result. In fact this argument shows that  $\int_{S^1} \circ ev^*$  is injective on those classes which are cohomologous to maps of spheres into our manifold. If  $M$  is assumed to be 2-connected, this will certainly be the case, since then the Hurewicz homomorphism is an isomorphism in degree 3 and surjective in degree 4.

REMARK. The case  $n = 4$  must be treated separately, since  $\mathrm{SO}(4)$  is no longer simple. But  $\pi_i(B\mathrm{Spin}(4)) = 0$  for  $i = 1, 2, 3$ , so that  $\int_{S^1} \circ ev^*$  is still an isomorphism. But now,

$$H^4(B\mathrm{Spin}(n); \mathbb{Z}) \cong H^3(LB\mathrm{Spin}(n); \mathbb{Z}) \cong \mathbb{Z} \oplus \mathbb{Z}.$$

As before, this result allows us to study the obstruction to lifting the structural group of  $LQ \rightarrow LM$  to a central extension of  $L\mathrm{Spin}(4)$  by  $S^1$ . However, the universal central extension is now an extension by a 2-torus, not a circle, since  $\mathrm{SO}(4)$  has two simple factors [14, Chapter 4].

The spinor bundle on loop space is a certain infinite dimensional vector bundle associated with a string structure. To define it, we need the appropriate representation of  $L\widetilde{\mathrm{Spin}}(n)$ . Let  $H$  be the Hilbert space  $L^2(S^1; \mathbb{R}^{2n})$  and  $e_1, \dots, e_{2n}$  the standard basis for  $\mathbb{R}^{2n}$ . If  $\varepsilon_k = \frac{1}{2}(e_{2k-1} + ie_{2k})$ , the elements  $\varepsilon_k z^m$  and  $\bar{\varepsilon}_k z^m$  form a basis for the complexification  $H_{\mathbb{C}}$ . Let  $W$  denote the subspace of  $H_{\mathbb{C}}$  spanned by  $\varepsilon_k z^r$  and  $\bar{\varepsilon}_l z^s$ , for  $r \geq 0$  and  $s \geq 1$ . Then it is shown in [14, Chapter 12], that  $L^0\mathrm{SO}(n)$  (and hence  $L\mathrm{Spin}(n)$ ) acts projectively on the Hilbert space completion of  $\bigwedge(W)$  and on the completions of  $\bigwedge^{\mathrm{even}}(W)$ ,  $\bigwedge^{\mathrm{odd}}(W)$  by two inequivalent, irreducible representations. Therefore, they are actual representations of some central extension  $L\widetilde{\mathrm{Spin}}(n)$  by  $S^1$ . If  $LQ \rightarrow LM$  admits a string structure, we can form the associated bundles  $\widetilde{LQ} \times_{L\widetilde{\mathrm{Spin}}(n)} \bigwedge^{\pm}(W) \rightarrow LM$ . Sections of these bundles are called strings.

REMARK. One would now like to define the Dirac operator acting on these bundles. This is discussed in [15] and involves difficult analytical problems. However, it is possible to give such a definition in a neighbourhood of the constant loops. This is done in [18].

**4. Equivariant string structures.** Suppose that a compact, abelian group  $G$  acts on  $M$  and let  $EG \rightarrow BG$  denote the universal  $G$ -bundle. We can then form the space  $EG \times_G M$ , the quotient of  $EG \times M$  by the diagonal action of  $G$ . This space is fibered over  $BG$  with fiber  $M$  and its cohomology  $H_G^*(M)$  is, by definition, the equivariant cohomology of  $M$ . If  $P \rightarrow M$  is a  $G$ -equivariant  $\mathrm{SO}(n)$ -bundle, then this construction yields an  $\mathrm{SO}(n)$ -bundle  $EG \times_G P \rightarrow EG \times_G M$ . There is a commutative diagram of classifying maps,

$$\begin{array}{ccc}
 M & \longrightarrow & BSO(n) \\
 \downarrow i & & \downarrow \text{id} \\
 EG \times_G M & \longrightarrow & BSO(n)
 \end{array}$$

where  $i$  is the inclusion of the fiber. If, in addition,  $P \rightarrow M$  has a  $G$ -equivariant spin structure  $Q \rightarrow M$ , then there is a similar diagram involving  $B\text{Spin}(n)$  and the classifying map of  $Q \rightarrow M$ . We conclude that the bundle  $EG \times_G Q \rightarrow EG \times_G M$  is a spin structure for  $EG \times_G P \rightarrow EG \times_G M$ . Therefore, the second Stiefel-Whitney class of the latter bundle must be zero. This class is denoted by  $w_2(P)_G$ .

Conversely, suppose that  $P \rightarrow M$  is  $G$ -equivariant and that  $w_2(P)_G = 0$ . Then, the bundle  $EG \times_G P \rightarrow EG \times_G M$  admits a spin structure  $\tilde{Q} \rightarrow EG \times_G M$ . But  $\tilde{Q} \rightarrow EG \times_G P$  is a double cover which by [11] corresponds to some  $G$ -equivariant double cover  $Q \rightarrow P$ . Note that the main result of [11] applied here because by [9]  $P$  has the homotopy type of a  $G$ -CW complex. Since  $Q \rightarrow P$  is the pullback of  $\tilde{Q} \rightarrow EG \times_G P$  by the inclusion  $P \hookrightarrow EG \times_G P$ , we conclude that  $Q \rightarrow M$  is a  $G$ -equivariant spin structure for  $P \rightarrow M$ .

As noted in [19], the analogous result for string theory would be that a  $G$ -equivariant string structure exists if and only if the equivariant class  $\frac{1}{2}p_1(P)_G$  vanishes—the “rigidity condition” (see also [6]). In fact, this is not completely true, as we will now see. First observe that the action of  $G$  on  $M$  induces an action on  $LM$ . If  $Q \rightarrow M$  is  $G$ -equivariant, then  $LQ \rightarrow LM$  is an equivariant  $L\text{Spin}(n)$ -bundle under this induced action. As above, we obtain a commutative diagram,

$$\begin{array}{ccc}
 LM & \longrightarrow & BL\text{Spin}(n) \\
 \downarrow & & \downarrow \text{id} \\
 EG \times_G LM & \longrightarrow & BL\text{Spin}(n).
 \end{array}$$

Suppose now that  $LQ \rightarrow LM$  has a  $G$ -equivariant string structure  $\underline{LQ} \rightarrow LM$ . Then arguing as before, we can conclude that  $EG \times_G \underline{LQ} \rightarrow EG \times_G LM$  is a string structure for the bundle  $EG \times_G LQ \rightarrow EG \times_G LM$ . Thus, the obstruction  $\underline{\mu}_G$  to lifting the structural group of  $EG \times_G LQ \rightarrow EG \times_G LM$  to  $L\widetilde{\text{Spin}}(n)$  must vanish. The reasoning used in §3 shows that  $\underline{\mu}_G$  lies in  $H_G^3(LM; \mathbb{Z})$ , and is the pullback of the generator of  $H^3(BL\text{Spin}(n); \mathbb{Z})$  by the classifying map of  $EG \times_G LQ \rightarrow EG \times_G LM$ .

Conversely, suppose that  $\mu_G = 0$ . Then the structural group of  $EG \times_G LQ \rightarrow EG \times_G LM$  can be lifted to  $L\widetilde{\text{Spin}}(n)$ , yielding a new bundle  $\widetilde{W} \rightarrow EG \times_G LM$ . The map  $\widetilde{W} \rightarrow EG \times_G LQ$  has the structure of an  $S^1$ -bundle. According to [12], the space  $LQ$  has the homotopy type of a locally finite CW complex. Therefore, we can apply the main result of [8] to obtain a corresponding  $G$ -equivariant circle bundle  $W \rightarrow LQ$ . As before, we conclude that  $W \rightarrow LM$  is a  $G$ -equivariant string structure for  $LQ \rightarrow LM$ .

The evaluation map  $ev: LM \times S^1 \rightarrow M$  is  $G$ -equivariant (with the trivial action on  $S^1$ ). Therefore it induces a map

$$ev_G: (EG \times_G LM) \times S^1 \rightarrow EG \times_G M.$$

Composing with integration over  $S^1$ , we obtain a map from  $H_G^4(M; \mathbb{Z})$  to  $H_G^3(LM; \mathbb{Z})$ , and a commutative diagram:

$$\begin{array}{ccc} H_G^4(M; \mathbb{Z}) & \xrightarrow{\int_{S^1} \circ ev_G^*} & H_G^3(LM; \mathbb{Z}) \\ \uparrow & & \uparrow \\ H^4(B \text{Spin}(n); \mathbb{Z}) & \xrightarrow{\int_{S^1} \circ ev^*} & H^3(LB \text{Spin}(n); \mathbb{Z}). \end{array}$$

The vertical maps are induced from the classifying maps of  $EG \times_G Q \rightarrow EG \times_G M$  and  $EG \times_G LQ \rightarrow EG \times_G LM$ . By definition,  $\frac{1}{2}p_1(P)_G$  is the pullback of the generator of  $H^4(B \text{Spin}(n); \mathbb{Z})$  by the classifying map of  $EG \times_G Q \rightarrow EG \times_G M$ . From the diagram, we see that it is mapped to  $\mu_G$ . From this, we conclude that if  $\frac{1}{2}p_1(P)_G$  vanishes then  $LQ \rightarrow LM$  has a  $G$ -equivariant string structure. Conversely, suppose that there exists an equivariant string structure, so that  $\mu_G$  is zero. To conclude that  $\frac{1}{2}p_1(P)_G$  is zero, we need to know whether  $\int_{S^1} \circ ev_G^*$  is injective. There is a commutative diagram,

$$\begin{array}{ccc} H_G^4(M; \mathbb{Z}) & \xrightarrow{\int_{S^1} \circ ev_G^*} & H_G^3(LM; \mathbb{Z}) \\ \downarrow & & \downarrow \\ H^4(M; \mathbb{Z}) & \xrightarrow{\int_{S^1} \circ ev^*} & H^3(LM; \mathbb{Z}) \end{array}$$

where the vertical arrows are induced from the homotopy inclusion of the fiber. Thus, we see that injectivity will depend on the cohomology

of  $BG$  and the injectivity of  $\int_{S^1} \circ ev^*$ . In the important case where  $G = S^1$ , the map is not generally injective. For instance, if  $M$  is 3-connected, then the spectral sequence associated to  $ES^1 \times_{S^1} M \rightarrow BS^1$  yields an exact sequence

$$0 \rightarrow \mathbb{Z} \rightarrow H_{S^1}^4(M; \mathbb{Z}) \rightarrow H^4(M; \mathbb{Z}) \rightarrow 0.$$

Summarizing, we have proved

**PROPOSITION 4.1.** *Let  $G$  be a compact, abelian group and  $P \rightarrow M$  a  $G$ -equivariant  $SO(n)$ -bundle.*

(1)  *$P \rightarrow M$  admits a  $G$ -equivariant spin structure if and only if  $w_2(P)_G$  vanishes.*

(2) *Suppose that  $n \geq 5$  and that  $w_2(P)_G = 0$ . Then there exists a  $G$ -equivariant string structure if  $\frac{1}{2}p_1(P)_G$  vanishes. The converse is not true in general.*

The proposition does not generalize to non-compact groups, since the result of [11] does not apply in this case. If  $G = \mathbb{Z}$ , the above argument does show that the existence of a  $\mathbb{Z}$ -equivariant spin structure forces  $w_2(P)_\mathbb{Z}$  to vanish. Moreover, if we assume  $M$  is 2-connected, then  $\int_{S^1} \circ ev_\mathbb{Z}^*$  is injective, since  $B\mathbb{Z} = S^1$ . Therefore, the existence of a  $\mathbb{Z}$ -equivariant string structure also forces  $\frac{1}{2}p_1(P)_\mathbb{Z}$  to be zero.

Closely related to this is the question of whether a given diffeomorphism  $f$  preserves spin and string structures. The map  $f$  generates a  $\mathbb{Z}$ -action on  $M$ . The space  $E\mathbb{Z} \times_\mathbb{Z} M$  can then be identified with the “mapping cylinder”

$$M_f = \frac{M \times I}{(x, 0) \sim (f(x), 1)}.$$

The bundle  $E\mathbb{Z} \times_\mathbb{Z} P \rightarrow E\mathbb{Z} \times_\mathbb{Z} M$  is just  $P_f \rightarrow M_f$ . If  $P \rightarrow M$  is  $\mathbb{Z}$ -equivariant, there is a diagram of bundle morphisms,

$$\begin{array}{ccccc} SO(n) & \xlongequal{\quad} & SO(n) & \longrightarrow & pt. \\ \downarrow & & \downarrow & & \downarrow \\ P & \longrightarrow & P_f & \longrightarrow & S^1 \\ \downarrow & & \downarrow & & \parallel \\ M & \longrightarrow & M_f & \longrightarrow & S^1. \end{array}$$

The induced morphisms of spectral sequences yield a commutative array of cohomology groups in low degrees. In the Gysin sequence associated to  $P_f \rightarrow S^1$ , the arrow  $H^1(P) \rightarrow H^1(P)$  can be identified with  $f^* - \text{id}^*$ . Thus, we can conclude that there is a spin structure invariant under  $f$  if and only if  $w_2(P_f) = 0$ .

Suppose now that such a bundle  $Q \rightarrow M$  exists. The induced action of  $\mathbb{Z}$  on  $LM$  is generated by a map  $F$ , related in the obvious manner to  $f$ . Arguing in a similar fashion, we see that  $LQ \rightarrow LM$  has a string structure invariant under  $F$  if and only if  $(LQ)_F \rightarrow (LM)_F$  admits a string structure. Of course the latter bundle is  $E\mathbb{Z} \times_{\mathbb{Z}} LQ \rightarrow E\mathbb{Z} \times_{\mathbb{Z}} LM$ . Assuming that  $M$  is 2-connected, the proof of Proposition 4.1 shows that  $\int_{S^1} \circ ev_{\mathbb{Z}}^*$  is injective. We conclude that there is an invariant string structure if and only if  $\frac{1}{2}p_1(P)_{\mathbb{Z}} = \frac{1}{2}p_1(P_f) = 0$ . This was noted in [10].

The most natural action on  $LM$  is that of rotating loops. It differs from the above situation in that it is not inherited from an action on  $M$ . We now ask whether a given string structure  $\widetilde{LQ} \rightarrow LM$  is equivariant under this action. This is equivalent to requiring that the  $S^1$ -action on  $LQ$  lifts to the  $S^1$ -bundle  $\widetilde{LQ} \rightarrow LQ$ . Again, we apply the main result of [8]. We conclude that the  $S^1$ -action lifts if and only if there is a commutative diagram of classifying maps,

$$\begin{array}{ccc} LQ & \longrightarrow & BS^1 \\ \downarrow i & & \downarrow \text{id} \\ ES^1 \times_{S^1} LQ & \longrightarrow & BS^1. \end{array}$$

But,  $\widetilde{LQ} \rightarrow LQ$  corresponds to some element  $\mu$  in  $H^2(LQ; \mathbb{Z})$ . Therefore, such a diagram exists if and only if  $\mu$  lies in the image of  $i^*: H^2_{S^1}(LQ; \mathbb{Z}) \rightarrow H^2(LQ; \mathbb{Z})$ . From the spectral sequence associated to  $ES^1 \times_{S^1} LQ \rightarrow BS^1$ , we see that the obstruction to this diagram is  $d_2(\mu) \in H^2(BS^1; \mathbb{Z}) \otimes H^1(LQ; \mathbb{Z})$ . This will be zero, for instance, when  $\pi_2(M)$  is pure torsion, since then  $H^1(LM; \mathbb{Z}) = 0$ , forcing  $H^1(LQ; \mathbb{Z}) = 0$ . At least in this case, we can state

**PROPOSITION 4.2.** *Suppose  $M$  is simply connected and that  $\pi_2(M)$  is torsion. Then every string structure for the  $SO(n)$ -bundle  $P \rightarrow M$  is necessarily equivariant under the action of rotating loops.*

**CONCLUDING REMARKS.** Witten has shown in [19] that the natural action of  $S^1$  on  $LM$  carries deep topological information. Suppose

the spin manifold  $M$  admits an  $S^1$ -action and  $p_1(M)$  is torsion. Then, by formally applying the Atiyah-Bott-Lefschetz fixed point formula, he showed that the index of the Dirac operator on  $LM$  is a modular form.

The group,  $\text{Diff}^+(S^1)$ , of orientation-preserving diffeomorphisms, also acts on loop space, in a more refined way than the circle. Moreover, this action lifts to a projective (infinitesimal) action on the bundle of spinors, and gives some geometrical insight into elliptic cohomology [5].

Finally, we could have defined the notion of a string structure for any compact, semi-simple Lie group  $G$  and its universal cover  $\tilde{G}$ , rather than  $\text{SO}(n)$  and  $\text{Spin}(n)$ . Exactly the same methods apply to study this situation.

#### REFERENCES

- [1] M. F. Atiyah, *Riemann surfaces and spin structures*, Ann. Sci. École. Norm. Sup., **4** (1971), 47–62.
- [2] —, *Circular symmetry and stationary-phase approximation*, Astérisque No. 131, **1** (1985), 43–59.
- [3] O. Alvarez, T. P. Killingback, M. Margano and P. Windey, *String theory and loop space index theorems*, Comm. Math. Phys., **111** (1) (1987), 1–10.
- [4] A. Borel, *Topics in the Homology Theory of Fiber Bundles*, Lecture Notes in Math., vol. **36**, Springer-Verlag, 1967.
- [5] J.-L. Brylinski, *Representations of loop groups, Dirac operators on loop space, and modular forms*, preprint (1988).
- [6] R. Bott and C. Taubes, *On the rigidity theorems of Witten*, J. Amer. Math. Soc., **2** (1989), 137–186.
- [7] R. Bott and L. Tu, *Differential Forms in Algebraic Topology*, Springer Verlag, 1982.
- [8] A. Hattori and T. Yoshida, *Lifting compact group actions in fiber bundles*, Japanese J. Math. (N.S.), no. 1, **2** (1976), 13–25.
- [9] S. Illman, *The equivariant triangulation theorem for actions of compact Lie groups*, Math. Ann., **262** (1983), 487–501.
- [10] T. P. Killingback, *World sheet anomalies and loop geometry*, Nucl. Phys., B-Part. Phys., **288** (1987), 578–588.
- [11] R. Lashof, J. P. May and G. B. Segal, *Equivariant bundles with abelian structural group*, Homotopy Theory, Proc. Conf. Evanston, Contemp. Math., **19**, 167–176.
- [12] J. Milnor, *On spaces having the homotopy type of a CW complex*, Trans. Amer. Math. Soc., **90** (1959), 272–280.
- [13] J. Milnor and J. Stasheff, *Characteristic Classes*, Princeton University Press, 1974.
- [14] A. Pressley and G. Segal, *Loop Groups*, Oxford University Press, 1986.
- [15] G. Segal, *Elliptic cohomology*, Séminaire Bourbaki, 40e année, no. 695, 1987–88.
- [16] P. Shanahan, *An introduction to the Atiyah-Singer index theorem*, SLN 638, 1976.

- [17] N. Steenrod, *The Topology of Fiber Bundles*, Princeton University Press, 1951.
- [18] C. H. Taubes,  *$S^1$ -actions and elliptic genera*, preprint (1987).
- [19] E. Witten, *The Index of the Dirac Operator on Loop Space*, Elliptic curves and modular forms in algebraic topology, Lecture Notes in Math., vol. 1326, Springer Verlag, 1988.
- [20] E. Witten, *Global anomalies in string theory*, Symposium on anomalies, geometry, topology, ed. Bardeen and White, World Scientific, 1985.

Received March 14, 1990.

PRINCETON UNIVERSITY  
PRINCETON, NJ 08544

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555  
vsv@math.ucla.edu

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112  
clemens@math.utah.edu

F. MICHAEL CHRIST  
University of California  
Los Angeles, CA 90024-1555  
christ@math.ucla.edu

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093  
tenright@ucsd.edu

NICHOLAS ERCOLANI  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

R. FINN  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720  
vfr@math.berkeley.edu

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

C. C. MOORE  
University of California  
Berkeley, CA 94720

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106  
mgscharl@henri.ucsb.edu

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## ASSOCIATE EDITORS

R. ARENS

E. F. BECKENBACH  
(1906–1982)

B. H. NEUMANN

F. WOLF  
(1904–1989)

K. YOSHIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

# PACIFIC JOURNAL OF MATHEMATICS

Volume 155    No. 1    September 1992

---

Characterization of modular correspondences by geometric properties	1
ALLAN RUSSELL ADLER	
Representations of convex nondentable sets	29
SPIROS ARGYROS and IRENE DELIYANNI	
Isomorphisms of spaces of continuous affine functions	71
CHO-HO CHU and HENRY BRUCE COHEN	
Universal classes of Orlicz function spaces	87
FRANCISCO LUIS HERNÁNDEZ RODRÍGUEZ and CESAR RUIZ	
Asymptotic behavior of the curvature of the Bergman metric of the thin domains	99
KANG-TAE KIM	
Quadratic central polynomials with derivation and involution	111
CHARLES PHILIP LANSKI	
Nonsplit ring spectra and products of $\beta$ -elements in the stable homotopy of Moore spaces	129
JIN KUN LIN	
Orientation and string structures on loop space	143
DENNIS MCLAUGHLIN	
Homomorphisms of Bunce-Deddens algebras	157
CORNEL PASNICU	
Certain $C^*$ -algebras with real rank zero and their corona and multiplier algebras. Part I	169
SHUANG ZHANG	
Correction to: "On the density of twistor elementary states"	199
MICHAEL G. EASTWOOD and A. M. PILATO	