

*Pacific
Journal of
Mathematics*

**STUDYING LINKS VIA CLOSED BRAIDS. VI. A
NONFINITENESS THEOREM**

JOAN BIRMAN AND WILLIAM W. MENASCO

STUDYING LINKS VIA CLOSED BRAIDS VI: A NON-FINITENESS THEOREM

JOAN S. BIRMAN AND WILLIAM W. MENASCO

Exchange moves were introduced in an earlier paper by the same authors. They take one closed n -braid representative of a link to another, and can lead to examples where there are infinitely many conjugacy classes of n -braids representing a single link type.


THEOREM 1. *If a link type has infinitely many conjugacy classes of closed n -braid representatives, then $n \geq 4$ and the infinitely many classes divide into finitely many equivalence classes under the equivalence relation generated by exchange moves.*

This theorem is the last of the preliminary steps in the authors' program for the development of a calculus on links in S^3 .

THEOREM 2. *Choose integers $n, g \geq 1$. Then there are at most finitely many link types with braid index n and genus g .*

Introduction. This paper is the sixth in a series in which the authors study the closed braid representatives of an oriented link type \mathcal{L} in oriented 3-space. The earlier papers in the series are [B-M,I]-[B-M,V]. An overall view of the program may be found in [B-M]. The long-range goal of the program is to classify link types, up to isotopy in oriented 3-space, using techniques based upon the theory of braids. This paper is the last of the preliminary steps on the way to so doing.

Let \mathcal{L} be an oriented link in oriented 3-space, and let \mathbf{L} be a closed n -braid representative of \mathcal{L} , with braid axis \mathbf{A} . If the isotopy class of \mathbf{L} in $S^3 - \mathbf{A}$ has a representative which has the very special form illustrated in Figure 1 (see next page), then \mathbf{L} is said to admit an *exchange move*, as illustrated in Figure 1. (The example shown there is a 4-braid; however if each strand is replaced by some number of parallel strands, it can be reinterpreted as an n -braid, for any n .) Exchange moves take closed n -braids to closed n -braids, in general changing the conjugacy class. Figure 2 (see next page) shows how n -braids which admit exchange moves may be modified to produce infinitely many closed n -braid representatives of \mathcal{L} . In effect, the exchange move allows one to replace the sub-braid \mathbf{X}



Exchange move

FIGURE 1




FIGURE 2

by a very special conjugate of itself, leaving Y invariant. For generic braids X and Y the infinitely many n -braids so produced will be in distinct conjugacy classes, because the isotopy which is shown in general not realizable as an isotopy in the complement of A . See [VB] for a proof of this assertion, for specific choices of X and Y . To understand how this phenomenon can lead to serious complications in the n -braid representations of knots and links, recall that it was shown in [B-M,V] that there is a 4-braid representative of the unknot which admits an exchange move, and that by modifying it as in Figure 2 one obtains infinitely many distinct conjugacy classes of 4-braid representatives of the unknot. But then, any closed braid representative of any link may be connect-summed with this particular closed braid representative of the unknot, to produce infinitely many similar examples for every link type.

The main result in this paper is that in fact exchange moves are the *only* way to produce infinitely many distinct conjugacy classes of closed n -braid representatives of a link type.

THEOREM 1. *Assume that \mathcal{L} admits infinitely many conjugacy classes of n -braid representatives. Then $n \geq 4$, and the infinitely many conjugacy classes divide into finitely many equivalence classes under the equivalence relation generated by exchange moves.*

As a corollary to Theorem 1, we will be able to say something about the exponent sum of a minimum-string braid representative of a link. Our corollary relates to

The Jones Conjecture. Let \mathcal{L} be a link of braid index n , and let \mathbf{L} and \mathbf{L}' be any two n -braid representatives. Express \mathbf{L} and \mathbf{L}' in any way as words \mathbf{W} and \mathbf{W}' in the elementary braids $(\sigma_i)^{\pm 1}$. Then the exponent sums of \mathbf{W} and \mathbf{W}' coincide.

By the work of Vaughan Jones [J] this conjecture is known to be true for $n = 3$ and 4. Our contribution is a weak version of the Jones Conjecture, which follows directly from Theorem 1.

COROLLARY. *Let \mathcal{L} be a link of braid index n . Then, among the n -braid representatives of \mathcal{L} , at most finitely many exponent sums can occur.*

The referee has observed that there is another result which follows from the proof of Theorem 1:

THEOREM 2. *Choose integers $n, g \geq 1$. Then there are at most finitely many link types with braid index n and genus g .*

Most of this paper will be devoted to the proof of Theorem 1. At the very end, after we have completed our proof of Theorem 1, we will prove the Corollary and Theorem 2.

Acknowledgment. We thank the referee for the detailed attention which he or she gave to our work, and for pointing out Theorem 2 to us.

Proof of Theorem 1. By hypothesis we are given a link type \mathcal{L} which has infinitely many conjugacy classes $\{[\mathbf{L}_i]; i = 1, 2, 3, \dots\}$ of n -braid representatives. We will first show that all but finitely many of the conjugacy classes $\{[\mathbf{L}_i]; i = 1, 2, 3, \dots\}$ admit an exchange move. We will then show that this implies the stronger assertion.

We may assume that $n \geq 4$. For, clearly $n \neq 1$, because the unknot is the only link of braid index 1. Also, $n \geq 2$, because B_2 is infinite cyclic, and since we are considering *oriented* links, each type $(2, p)$ torus link has a unique conjugacy class of 2-braid representatives, whereas the unknot has exactly 2. The case $n = 3$ is non-trivial; however it is proved in [B-M,III] that a link of braid index 3 (respectively 2, 1) has 1 or 2 (respectively 2, 3) conjugacy classes of 3-braid representatives. So $n \geq 4$.

Let $L = L_1$. Assume that L is the boundary of a not necessarily connected surface F which is oriented so that the positive normal bundle to F has the orientation induced by that on L . Assume that F has been chosen to have maximum Euler characteristic χ among all such oriented spanning surfaces. A link L will in general have a multiplicity of such spanning surfaces; however having selected one we will stick with it throughout this paper. Thus each link L_i in our sequence will be assumed to be the boundary of F_i , where if $i \neq k$ there is a homeomorphism h_{ik} of S^3 such that $h_{ik}(F_k) = F_i$. In general h_{ik} will not fix the braid axis A , because by hypothesis $L_i = \partial F_i$ is in a different conjugacy class from $L_k = \partial F_k$.

The complement of the braid axis A in R^3 is an open cylinder, which is fibered by half-planes H_θ through A . Let H denote a choice of this fibration. The proof begins with a study of the foliation which is induced on F_i by its intersections with the fibers H_θ of H . This foliation was first studied by D. Bennequin in [Be], in connection with his studies of contact structures on R^3 . It was proved by Bennequin that F_i may be assumed to be a *Markov surface*. The first two properties of these surfaces are achieved by general position techniques:

(Mi) A intersects F_i transversally in a finite set of points p_1, \dots, p_k which we will refer to as "vertices." There is a neighborhood on F_i of each vertex which is foliated radially.

(Mii) All but finitely many fibers H_θ of H meet F_i transversally, and those which do not (the *singular fibers*) are each tangent to F_i at exactly one point in the interior of both F_i and H_θ . Moreover, the tangencies are assumed to be either local maxima or minima or saddle points.

A third property is obtained by Bennequin [Be] by small modifications in the F_i 's. A proof is also given in Lemma 2 of [B-M,I]:

(Miii) There are no simple closed curves in the foliation of F_i . If H_θ is a non-singular fiber, then each component of $H_\theta \cap F_i$ is an arc.

Following [B-M,I], pairs (F, H) and (F', H') which satisfy (Mi), (Mii) and (Miii) will be said to be *equivalent* if there is an isotopy h_t of R^3 , $t \in [0, 1]$, which takes $(\partial F, \partial H)$ to $(\partial F', \partial H')$ in such a way that each $h_t(F, H)$ satisfies (Mi), (Mii) and (Miii). Let $[F, H]$ be the equivalence class of (F, H) . Let $|A \cap F|$ denote the number of points of intersection of A with F . Let $|H \cdot F|$ be the number of tangencies between F and fibers of H . The *complexity* $C([F, H])$ is defined to be the pair $(|A \cap F|, |H \cdot F|)$. It is well-defined on equivalence classes.

We continue to investigate F_i . The next property, established in Lemma 1 of [B-M,I], follows from the fact that $L_i = \partial F_i$ is a closed braid, so that all its intersections with fibers of H are coherently oriented:

(Miv) An arc of intersection of F_i with a non-singular fiber of H never has both of its endpoints on L_i .

It follows from (Miv) that the arcs in $F_i \cap H_\theta$ for non-singular H_θ are restricted to two types:

- a-arcs:** one endpoint is on A and the other is on L_i ,
- b-arcs:** both endpoints are on A ,

and that the singularities are restricted to three types:

- aa** (if the arcs which come together are both type **a**),
- bb** (if the arcs which come together are both type **b**),
- ab** (if one is type **a** and the other type **b**).

A **b**-arc β in $H_\theta \cap F_i$ is *essential* if both sides of H_θ split along β are pierced by L . If β is inessential, then β and a subarc α of A will co-bound a disc D on H_θ which is not pierced by L_i . The axis A may then be pushed across D to eliminate two points of $A \cap F_i$, and so to reduce the number of points in $A \cap F_i$. After the reduction, one may recover (Mi)–(Miv) without increasing the complexity (see [B-M,IV] for details), so from now on we may assume:

(Mv) Every **b**-arc in the foliation is essential.

Finally, a component of F_i is *trivially foliated* if F_i is a disc which is pierced once by A , and if F_i is radially foliated by its arcs of intersection with fibers of H . Since the boundary of a trivially foliated component is necessarily a 1-braid representative of the unknot, a case which is not of interest in this paper, we assume from now on that:

(Mvi) No component of F_i is trivially foliated.

We begin to study the combinatorics of the foliation of F_i . To simplify the notation we will assume (until near the end of the proof) that F_i is connected. Later we will show how to modify things if F_i is disconnected. As in [B-M,I] we may choose a finite collection \mathcal{E}_i of **a**-arcs and **b**-arcs such that F_i split along \mathcal{E}_i is a union of foliated 2-cells \mathcal{T}_i , which we call *tiles*. See Figure 3. The arcs in \mathcal{E}_i are chosen so that there is exactly one singularity of the foliation in the interior of each tile. The left column in Figure 3 shows the three types of tiles which occur. The tile *edges* are the arcs in \mathcal{E}_i , shown as dotted lines. The singular leaves as solid lines, and subarcs of **L** are thick solid lines. The singularities are indicated by black dots. The tile *vertices* \mathcal{V}_i are the points where **A** pierces F_i . They are labeled p_1, p_2, \dots . We have also labeled the **L**-endpoints of the singular leaves with symbols $\lambda_1, \lambda_2, \dots$. The tiles are denoted as being *types* **aa**, **ab** and **bb**, according as the singularity is type **aa**, **ab** or **bb**. There are four **a**-arcs in the boundary of an **aa**-tile and four **b**-arcs in the boundary of a **bb**-tile, and two of each type in the boundary of an **ab**-tile.

For future use, we record at this time two other features of the combinatorics which we will use in this paper. The foliation of F_i was assumed to be radial about each vertex, so as we push forward in the positive direction through the fibration, the arcs of intersection of F_i with fibers of **H** will flow radially about the vertices. This flow will be anticlockwise or clockwise, according as the oriented axis **A** intersects F_i from the positive or from the negative side. Call a vertex *positive* in the former case and *negative* in the latter. Notice that the braid index is the number of positive vertices minus the number of negative vertices.

The *type* of a vertex is the cyclically ordered array of **a**'s and **b**'s which records the tile edges meeting at that vertex. We shall not need it here as an oriented symbol, but we might as well orient it by the flow about the vertex. Later we will need the fact (see Figure 3) that the flow is always positive about a vertex which contains an **a** in its type symbol. On the other hand, a vertex of type **bb...b** could be either positive or negative. The *valence* of a vertex is the number of symbols in its type symbol.


In the right column of Figure 3 we have shown how the singular leaves would look if they are viewed on the singular fiber H_θ of **H**. We have used the same cyclic order for the four endpoints of the singular leaves in the left and right columns; this means that the view

| type | Foliated tile, viewed on F_2 | Singular leaves, viewed on H_0 |
|------|--------------------------------|----------------------------------|
| aa | | |
| ab | | |
| bb | | |

FIGURE 3

on F_i must be interpreted as being on the positive or negative side of F_i , according as the oriented normal to F at the singularity points in the direction of increasing or decreasing θ . Figure 4 (see next page) gives an example to indicate to the reader how distinct tile types might fit together, in the case where F_i is a disc. (This example can be realized by the unknot, represented as a 2-braid L .)

Recall that \mathcal{V}_i , \mathcal{E}_i and \mathcal{T}_i are the sets of tile vertices, edges and tiles in the tiling of F_i . Notice that our method of counting is rather special. We do not count subarcs of L_i which are in the boundary of a tile as an edge. Thus each tile has exactly four "edges." Similarly, we do not include in \mathcal{V}_i the points where a tile edge meets L_i . Let




Example: a tiling of a disc with no vertices of valence 1

FIGURE 4

V_i, E_i and T_i be the cardinality of $\mathcal{V}_i, \mathcal{E}_i$ and \mathcal{T}_i respectively. The next two lemmas relate to results established by the authors in other papers of this series.

LEMMA 3. *If the tiling of F_i has a vertex of valence 1, then L_i admits an exchange move.*

Proof. It is proved in Lemma 5 of [B-M,V] that if there is a vertex of valence 1, then L_i has a trivial loop. But then, as is shown in Figure 5, we may reconfigure the closed braid so that L_i is seen to admit an exchange move.



closed braid which has a trivial loop

the same closed braid, re-configured so it admits exchange move

FIGURE 5

Since our initial goal was to show that all but finitely many L_i admit an exchange, we will have achieved our goal if there is a vertex of valence 1 in the tiling of F_i . Therefore we may assume from now on that the tiling of F_i has no vertices of valence 1. Let $V_i(\alpha, \beta)$ denote the number of vertices in the tiling of F_i which have α **a**-arcs and β **b**-arcs in their type symbols.

LEMMA 4. *If any one of $V_i(1, 1)$ or $V_i(0, 2)$ or $V_i(0, 3)$ is non-zero, then L_i admits a complexity-reducing exchange move.*

Proof. It was proved in Lemma 5 of [B-M,IV] that if there is a vertex of type **bb** in the tiling of F_i , then $L_i = \partial F_i$ admits a complexity-reducing exchange. The corresponding statement for vertices of type **ab** is proved in Lemma 4 of [B-M,V], when the array of signs of the singularities in the tiles which meet at the vertex are $(+-)$. The case of vertices of type **bbb** is reduced to the case of vertices of type **bb** by Lemma 8 of [B-M,V]. Type **ab**, with sign sequence $(++)$ or $(--)$, is reduced to the case where there is a vertex of valence 1 by Lemma 9 of [B-M,V]. (Remark: this sign sequence is independent of the sign of a vertex, as we defined it earlier.) Thus, in every case L_i admits a complexity-reducing exchange. \square

We now proceed to analyze the consequences of our hypothesis that we have an infinite sequence $\{L_i; i \in N\}$ of closed n -braid representatives of \mathcal{L} , with L_i in a different conjugacy class from L_k if $i \neq k$. We look to the tiling of F_i for information about the existence of exchange moves.

LEMMA 5. *The number V_{i+} of positive vertices in the tiling of F_i and also the number V_{i-} of negative vertices goes to infinity as $i \rightarrow \infty$, with $V_{i+} - V_{i-}$ remaining constant.*

Proof. The surface F_i admits a non-trivial tiling, relative to our particular choice of fibration H of $S^3 - A$. We now use the tiling to make an Euler characteristic count. Since there are no vertices of valence 1, it follows that if one shrinks ∂F_i to a point, the image of each tile under the collapsing map will be a 2-cell with 4 edges and 4 vertices. Thus:

$$(1) \qquad 2T_i = E_i.$$

The collapsing map increases the number of tile vertices by 1, but it also raises the Euler characteristic of F_i by 1. The two effects cancel,

so we also have:

$$(2) \quad V_i - T_i = \chi.$$

Recall that the complexity of $[F_i, H]$ is the pair (V_i, T_i) , or equivalently (in view of equation (2)) the pair $(V_i, V_i - \chi)$. It was proved in Theorem 2 of [B-M,I] that there are at most finitely many equivalence classes $[F_i, H]$ for each fixed value of the complexity. This means that there are at most finitely many distinct conjugacy classes $[L_i]$ for each fixed value of the complexity. Since χ is fixed, and since by hypothesis infinitely many distinct conjugacy classes $[L_i]$ occur, the only possibility is that V_i goes to infinity as $i \rightarrow \infty$.

Now recall that $V_i = V_{i+} + V_{i-}$ and $V_{i+} - V_{i-} = n$, the braid index. Since n is by hypothesis independent of i , the assertion follows. \square

We continue our proof of Theorem 1. The symbol $V_i(\alpha, v - \alpha)$ denotes the number of vertices in the tiling of F_i which have valence v and have α **a**-arcs as edges. Thus, we can express V_i as the following sum of the $V_i(\alpha, v - \alpha)$'s:

$$(3) \quad V_i = \sum_{v=2}^{\infty} \sum_{\alpha=0}^v V_i(\alpha, v - \alpha).$$

Let $E_i(a)$ and $E_i(b)$ denote the number of type **a** and **b** edges, respectively, in the tiling, so that $E_i = E_i(a) + E_i(b)$. Since each type **a** edge is incident at one vertex, whereas each type **b** edge is incident at two vertices, we have related sums $E_i(a)$ and $E_i(b)$:

$$(4) \quad E_i(a) = \sum_{v=2}^{\infty} \sum_{\alpha=0}^v \alpha V_i(\alpha, v - \alpha),$$

$$(5) \quad 2E_i(b) = \sum_{v=2}^{\infty} \sum_{\alpha=0}^v (v - \alpha) V_i(\alpha, v - \alpha).$$

Using equation (1) to rewrite equation (2) in the form $4V_i - 2E_i(a) - 2E_i(b) = 4\chi$, we may then combine it with equations (3), (4) and (5) to obtain:

$$(6) \quad 4\chi = \sum_{v=2}^{\infty} \sum_{\alpha=0}^v (4 - v - \alpha) V_i(\alpha, v - \alpha).$$

Note that when $v \geq 4$ the coefficient $(4 - v - \alpha)$ will be non-positive. We may thus alter equation (6) so that all terms on both sides of the

equation, except possibly 4χ , are non-negative. This gives:

$$\begin{aligned}
 &V_i(1, 1) + 2V_i(0, 2) + V_i(0, 3) \\
 &= 4\chi + V_i(2, 1) + 2V_i(3, 0) \\
 (7) \quad &+ \sum_{v=4}^{\infty} \sum_{\alpha=0}^v (v + \alpha - 4)V_i(\alpha, v - \alpha).
 \end{aligned}$$

Recall that the *type* of a vertex p in the tiling of \mathbf{F}_i was defined to be the cyclic array of **a**'s and **b**'s which records the types of the tile edges which meet at p . Thus if $V_i(1, 1) \neq 0$ (respectively $V_i(0, 2) \neq 0$, $V_i(0, 3) \neq 0$) there is a vertex of type **ab** (respectively **bb**, **bbb**) in the tiling of \mathbf{F}_i .

We now consider the possible ways in which $V_i \rightarrow \infty$. Equation (7) will record any change in the differing values of the $V_i(\alpha, v - \alpha)$'s, except for the cases $(\alpha, v - \alpha) = (1, 2), (2, 0)$ and $(0, 4)$, since in those cases the coefficient $(v + \alpha - 4)$ which occurred in equation (6) was zero, so that the corresponding terms do not appear in equation (7). For all other values of $(\alpha, v - \alpha)$ we see that if $V_i(\alpha, v - \alpha)$ increases without bound the left-hand side of equation (7) would be forced to be non-zero. Thus there are two possibilities:

possibility 1:

At least one of $V_i(1, 1), V_i(0, 2)$ or $V_i(0, 3) \neq 0$.

possibility 2:

$V_i(1, 1) = V_i(0, 2) = V_i(0, 3) = 0$ for all i , also at least one of the following holds:

$$\begin{aligned}
 &V_i(1, 2) \rightarrow \infty \quad \text{as } i \rightarrow \infty, \\
 &V_i(2, 0) \rightarrow \infty \quad \text{as } i \rightarrow \infty, \\
 &V_i(0, 4) \rightarrow \infty \quad \text{as } i \rightarrow \infty,
 \end{aligned}$$

and also (passing to a subsequence if necessary) $V_i(\alpha, \beta)$ is independent of i for all other (α, β) .

If possibility 1 occurs, then by Lemma 4 we conclude that \mathbf{L}_i admits an exchange move. Thus we are reduced to possibility 2.

LEMMA 6. *In the situation of possibility 2, the cases $V_i(2, 0) \rightarrow \infty$ and $V_i(1, 2) \rightarrow \infty$ do not occur.*

Proof of Lemma 6. We first show that $V_i(2, 0) \rightarrow \infty$ is impossible, i.e. that the number of vertices of type **aa** cannot grow without bound as $i \rightarrow \infty$. Such a vertex can only occur when two **aa** tiles are joined




FIGURE 6

together along a pair of common edges. We need to see how this situation would look in 3-space, and to do so we ask how two **aa** tiles appear imbedded in 3-space. The left picture in Figure 6 shows two **aa** tiles glued together along two common **a**-edges, to give a vertex of type **aa**. We focus first on the tile with vertices p_0 and p_1 . Its singular leaves lie on some fiber H_θ of H , with one of them having its endpoints at points p_0 and p_1 on the axis A , and the other having its endpoints at λ and λ' on $L_i \cap H_\theta$. The right picture in Figure 6 shows these leaves in 3-space (to be imagined as lying on a single fiber). Since the embedded tile is transverse to the fibers everywhere else, the final picture (which is determined up to reversal of the half-twist in the band) must be as in the right picture in Figure 6. The second tile will have a similar imbedding (ambiguous up to the choice of the order of the three vertices on A and the senses of the half-twists), but in all cases we see that if $V_i(2, 0)$ were to grow without bound the braid index n would too, contrary to hypothesis.

Suppose next that $V_i(1, 2) \rightarrow \infty$ as $i \rightarrow \infty$, i.e. the number of vertices of type **abb** grows without bound. Let p be a vertex of type




FIGURE 7

abb. Then p is positive, and the **a** edge which is incident at p must be on an **ab** tile. Thus there are necessarily two **ab** tiles glued together along a common **a**-edge which meet p . We can think of them as a pair of attached houses, as in the left picture of Figure 7. Since the unique vertex which is at the peak of the roof in an **ab** tile is a negative vertex, we cannot glue an **ab** tile into the trough. So the only way to fill in the trough is with a **bb** tile, as illustrated. So, if the number of vertices of type **abb** goes to infinity as $i \rightarrow \infty$, we will necessarily find tiles in our sequence with arbitrarily long rows of attached houses glued together in a row as in the right picture in Figure 7, with a **bb** tile in the trough between each pair of adjacent roofs.

We now investigate the contributions to V_{i+} from our row of houses. The sum of the signs in each **bb** tile is zero, but the sum of the signs in each **ab** tile is $+1$. If our sequence contained k_i tiles of type **ab**, we would also have k_i tiles of type **bb**, and so there would be a net partial contribution of k_i to V_{i+} . Since $(V_{i+} - V_{i-})$ must be independent of i , this means that we will need to have k_i extra negative vertices somewhere else. However, that is impossible, because the sum of the signs of the vertices in our 3 tile types is $+2, +1$ and 0 . Thus, if $V_i(1, 2) \rightarrow \infty$ as $i \rightarrow \infty$, the braid index will be forced to increase without bound, contrary to hypothesis. \square

Possibility 2 has been reduced to the situation where $V_i(0, 4) \rightarrow \infty$ as $i \rightarrow \infty$. Passing to a subsequence if necessary, we may assume that $V_i(1, 2)$ and $V_i(2, 0)$ are independent of i .

LEMMA 7. *Allowing $V_i(0, 4)$ to go to ∞ as $i \rightarrow \infty$ only gives finitely many distinct conjugacy classes of closed n -braids representing our link \mathcal{L} .*




FIGURE 8

Proof of Lemma 7. Assume that $V_i(0, 4) \rightarrow \infty$ as $i \rightarrow \infty$. By Lemma 5 the number of positive vertices of type **bbbb** and also the number of negative vertices of type **bbbb** must go to infinity as $i \rightarrow \infty$, with $V_{i+} - V_{i-}$ remaining constant. From Figure 3 we see that the only tile type which contains a positive vertex at which **b**-arcs are incident is a type **bb** tile, so the only way to obtain a positive vertex of type **bbbb** is as in Figure 8. So this picture occurs infinitely often.

Now notice that the sum of the signs of the vertices in a **bb** tile is zero. It then follows that the *negative* vertices of type **bbbb** must also lie at the intersection of four **bb** tiles, because if there were infinitely many such negative vertices which included an **ab** tile, the difference $V_{i+} - V_{i-}$ would not remain constant. Thus there must be interior regions of F_i which are tiled entirely by **bb** tiles. Moreover (passing to a subsequence if necessary) we may assume that the tiling remains fixed outside one such region, whereas the number of **bb** tiles in the region increases without bound as i is increased. Since the Euler characteristic is independent of i , the region in question must be an annulus. The tiling on the boundary of our annulus must be fixed, so the only way things can change is if the number of tiles between the two boundary components grows without bound as $i \rightarrow \infty$. See Figure 9 for an example of one way this could occur. The difference $V_{i+} - V_{i-}$ will thus remain independent of tube length, as will the Euler characteristic of F_i .

Recall our assumption that $V_i(\alpha, \beta)$ is independent of i if $(\alpha, \beta) \neq (0, 4)$. Even more, by passing to a subsequence if necessary, we may assume that the tiling of F_i is fixed in the component of our annulus.




FIGURE 9

An additional property then follows immediately from equation (4):

$$(8) \quad E_i(a) \text{ is independent of } i.$$

Let $T_i(aa)$ and $T_i(ab)$ denote the number of tiles of type **aa** and **ab** in the tiling of F_i . Notice that a tile of type **aa** (respectively **ab**) contains 4 (respectively 2) edges of type **a**. Since each edge is an edge of two tiles, we then have:

$$(9) \quad E_i(a) = 2T_i(aa) + T_i(ab).$$

Since $E_i(a)$ is independent of i , by (8), we conclude that $2T_i(aa) + T_i(ab)$ is too. Thus

$$(10) \quad T_i(aa) \text{ is independent of } i,$$

$$(11) \quad T_i(ab) \text{ is independent of } i.$$

It will be convenient now to assume that \mathcal{L} is a knot. Later we will modify the proof to the case where \mathcal{L} is a link. Our idea is to show that a finite set of combinatorial data in the tiling of F_i determines L_i as an embedded simple closed curve, relative to \mathbf{A} and fibers of \mathbf{H} . It will turn out that the embedding of L_i is determined entirely by data in the **aa** and **ab** tiles, and is independent of the **bb** tiles (which lie in the interior of F_i). Since, by (10) and (11), the number of **aa** and **ab** tiles is independent of i , we will thus be able to conclude that even though the tiling of F_i is changing, the changes do not affect the boundary, so there cannot be infinitely many conjugacy classes of closed braids.

With that plan in mind, let

τ = the number of singularities of types **aa** and **ab**, i.e. $|\mathbf{aa}| + |\mathbf{ab}|$.

ω = the number of points of $\mathbf{A} \cap \mathbf{F}_i$ which are vertices of these τ tiles.

ρ = the number of endpoints of singular leaves which are on \mathbf{L}_i .

Since each singular leaf on an **aa** (respectively **ab**) tile meets \mathbf{L}_i twice (respectively once) the integer $\rho = 2|\mathbf{aa}| + |\mathbf{ab}|$.

Let s_1, \dots, s_τ be the singular points on the τ tiles of type **aa** and **ab**, ordered to correspond to their cyclic order in the fibers of \mathbf{H} . That is, each s_k is in $\mathbf{H}_{\theta_k} \cap \mathbf{F}_i$, where $0 \leq \theta_1 < \theta_2 < \dots < \theta_\tau \leq 2\pi$. Let p_1, \dots, p_ω be the vertices of these τ tiles, ordered to correspond to their cyclic order on \mathbf{A} . Finally, let $\lambda_1, \dots, \lambda_\rho$ be the points where the \mathbf{L} -endpoints of the singular leaves in the tiling intersect the singular fibers of \mathbf{H} , ordered to correspond to their natural cyclic order on \mathbf{L}_i .

We now associate signs to the s_k 's and p_j 's. The sign ξ_k of s_k is + or - according as the sense of increasing θ agrees or disagrees with that of the outward-drawn normal to \mathbf{F}_i at s_k . The sign δ_j of p_j is + or -, according as the orientation of \mathbf{A} agrees or disagrees with that of the outward-drawn normal to \mathbf{F}_i at p_j .

Finally, we associate to the tiling of \mathbf{F}_i a combinatorial symbol. Let \mathbf{H}_{θ_k} be a singular fiber containing the singularity s_k on an **aa** tile, as depicted in the right column of Figure 3. The singular leaves through s_k have four endpoints, and these are alternately in the sets $\{p_1, \dots, p_\omega\}$ and $\{\lambda_1, \dots, \lambda_\rho\}$, so we can associate to \mathbf{H}_{θ_k} a cyclically ordered 4-tuple $\mathbf{4}_k = \xi_k(p_{1_k}, \lambda_{2_k}, p_{3_k}, \lambda_{4_k})$. The sign ξ_k is the sign of the singularity at s_k . The order of the 4 points is determined by their cyclic order on \mathbf{H}_{θ_k} . In the case of an **ab** tile (again see Figure 3) three of the four endpoints of the singular leaves are in $\{p_1, \dots, p_\omega\}$ and one is in $\{\lambda_1, \dots, \lambda_\rho\}$. There is a signed and cyclically ordered 4-tuple $\mathbf{4}_k = \xi_k(p_{1_k}, p_{2_k}, p_{3_k}, \lambda_{4_k})$ associated to the tile. Our *combinatorial symbol* is the array:

$$((\tau, \omega, \rho), (s_1, \dots, s_\tau), (\delta_1, \dots, \delta_\omega), \{\lambda_1, \dots, \lambda_\rho\}, (\mathbf{4}_1, \dots, \mathbf{4}_\tau)).$$

We now claim:

* *The combinatorial symbol determines the conjugacy class of \mathbf{L}_i as a closed braid, relative to the fibers of \mathbf{H} .*

Parenthetical remark. The reader who is familiar with [B-M,I] will recognize similarities between the proof we are about to give and a related proof in [B-M,I]; however there is a difference. In [B-M,I] we

were interested in constructing F_i as an embedded surface, whereas our focus in this proof is entirely on ∂F_i . We will see that the embedding of ∂F_i is completely determined by combinatorial data in the **aa** and **ab** tiles.

We return to the matter at hand. To prove $*$, we assume that we are given the combinatorial symbol, and proceed to construct L_i as a closed braid. We assume that the braid axis A is the z axis in R^3 , and that the fibers of H are the half-planes through A . The combinatorial symbol gives us the integer ω , and the first step is to choose ω points on A and label them p_1, \dots, p_ω , in order. Up to an isotopy of 3-space which preserves A it will not matter where we put them. The combinatorial symbol also tells us the integer τ , so we choose τ fibers, selecting a point in the interior of each and labeling them s_1, \dots, s_τ , in the cyclic order in which the fibers occur. Passing to the k th fiber, we may then construct the singular leaves in that fiber, as in the right column in Figure 3, with the help of the symbol 4_k . One of the singular leaves has both of its endpoints on A , at p_{1_k} and p_{3_k} (in the notation of Figure 3). The combinatorial symbol 4_k will tell us which points on A to use for p_{1_k} and p_{3_k} . The other singular leaf crosses this one, and its two endpoints will be at λ_{2_k} and λ_{4_k} (if the tile is type **aa**) or at p_{3_k} and λ_{4_k} (if it is type **ab**). The points on L can be anywhere in the interior of H_θ split along $p_{1_k}p_{3_k}$. Moving them will simply modify the embedding of L by an isotopy of S^3 which fixes A and each fiber of H setwise.

We have just described how to embed the singular leaves of the tiles which are adjacent to L . The next task is to extend this embedding to a neighborhood of the singular leaves. The first thing to notice is that we know how to embed a neighborhood of p_1, \dots, p_ω on F_i . For, if we choose a neighborhood which is small enough it is radially foliated by its intersection with H , so it must be a disc transverse to A . The sign δ_j of the point p_j tells us the side of the surface which is pierced by A . Next, notice that we can embed a little neighborhood of the singular leaves, for such a neighborhood is transverse to fibers everywhere except at the singular point, moreover the sign ξ_k of the singularity tells us which side of F faces in the direction of increasing θ at the singular point. The last thing to do is to fill in regions on F which lie between adjacent points λ_i and λ_{i+1} on L . See Figure 10 (on next page). These regions are foliated by radial arcs which emanate from the vertex, so they are everywhere transverse to the fibers of H . The leaves of the foliation are level sets for the polar




FIGURE 10

angle function, so they determine the embedding uniquely, up to an isotopy of S^3 which fixes each fiber setwise. When we are done we will have embedded a tiled neighborhood of the part of F_i which is closest to L_i , and in particular we will have embedded L_i in S^3 .

Recall that we have passed to a subsequence in which the changes which occur in the tiling of F_i as i is increased are restricted to the interior of F_i and to vertices of type **bbbb** which are at the intersection of four type **bb** tiles. From this it follows that the integers ρ , ω and τ are independent of i . Since at most finitely many distinct combinatorial symbols are possible for fixed values of ρ , ω and τ , and since the combinatorial symbol determines the embedding, we conclude that there are at most finitely many embeddings which are possible for $L_1 = \partial F_1$. Thus only a finite number of conjugacy classes can occur among the links L_i in our sequence. Since this violates the hypotheses of Theorem 1, we conclude that this case does not occur.

The only thing we need to change if L_i has more than one component is the instructions for filling in the pie-shaped regions. Modify the combinatorial data by changing the labeling of the L -endpoints of the singular leaves to

$$((\lambda_{1,1}, \dots, \lambda_{1,\rho_1}), (\lambda_{2,1}, \dots, \lambda_{2,\rho_2}), \dots, (\lambda_{t,1}, \dots, \lambda_{t,\rho_t})),$$

where the second subscript indicates the component. We may then proceed as before, one boundary component at a time, to fill in the regions next to the link.

If F is not connected, we proceed as before up to Lemma 5. We then replace Lemma 5 by the assertion that some component of F_i must have a tiling F_{i1} in which the number of vertices grows without




FIGURE 11

bound. In general, there will be, say, t components. So, instead of a sequence of tiled surfaces $\{F_i; i = 1, 2, 3, \dots\}$ we will have tiled surfaces:

$$\{F_{i1} \cup F_{i2} \cup \dots \cup F_{it}; i = 1, 2, 3, \dots\},$$

where particular surfaces have tilings in which the number of vertices grows without bound. Some of these subsurfaces may have more than one boundary component, it will not matter. Lemma 5 applies to each subsurface. Lemmas 6 and 7 do too.

Thus we have proved: If our link \mathcal{L} has infinitely many conjugacy classes of n -braid representatives, then all but finitely many of them admit exchange moves. This is a weak version of Theorem 1.

To prove the stronger version, we must ask how the infinitely many conjugacy classes which admit exchange moves are related to one another. By our proof, each time that there is an exchange move, there is a vertex of type **a** or of type **bb** or **ab** with sign $+-$, or a vertex of type **bbb**. We first prove that we may assume there are no vertices of type **a**. Suppose, on the contrary, that there are infinitely many F_i 's which contain at least one vertex of type **a**. A vertex of type **a** always occurs on an "end tile," as in Figure 11. One of the singular leaves in that tile is a separating leaf β . If we cut F_i along β , and then modify the cut edge slightly so that the modified surface F'_i is transverse to the foliation near the cut, the boundary of the modified surface will be a new closed braid L'_i which is, in effect, obtained from L_i by deleting the trivial loop. Notice that the changes we just made do not alter the tiling in the complement of the end tile. So, after we delete our trivial loop we will obtain a new tiled surface F'_i with all the properties of the old except: there is one less **aa** tile and

the braid index has been reduced by 1. This process may be repeated on F'_i if the tiling of F'_i contains a vertex of type **a**. However, it must end after $k \leq n$ such repetitions because each deletion reduces the braid index. Let $\{F''_i; i = 1, 2, 3, \dots\}$ be the infinite sequence of surfaces obtained after all type **a** vertices have been eliminated. Divide these into equivalence classes so that the braid index n'' is constant in each equivalence class. At least one of these equivalence classes must contain infinitely many members, because the original sequence did. Moreover, the surfaces in the new infinite sequence must have infinitely many distinct tilings, because we had infinitely many distinct tilings in our original sequence and all we did was to cut off finitely many end tiles from each F_i . But then by our earlier work, we conclude that the tiling of infinitely many F_i 's in the original sequence contains a vertex of type **bb** or **ab** with sign $+-$ or **bbb**.

Passing to a subsequence, we may assume that every F_i contains a vertex of type **bb** or **ab** with sign $+-$ or **bbb**. By Lemma 4, each closed braid $L_i = \partial F_i$ admits a complexity-reducing exchange move. When we change an essential **b**-arc to an inessential one we reduce the number of vertices in the tiling, and so reduce the complexity. For details see [B-M,IV and V]. The process of making exchange moves must therefore end.

Recall that when we began our work, we chose a spanning surface F for our link which had maximal Euler characteristic. The choice of F was not unique. The surfaces F_i which have been the object of our investigation here were then determined by finding homeomorphisms $h_i: (S^3, L) \rightarrow (S^3, L_i)$, and setting $F_i = h_i(F)$. Later we introduced several types of modifications. Some of those modifications involved the passage to a subsequence. Others involved an isotopic deformation of F_i . For example, when we changed F_i to a Markov surface, or when we cut off the end tiles, or when we modified L_i by an exchange move, we were changing F_i by isotopy. This means that each surface F'_i in our final sequence is still homeomorphic to our fixed surface F . If the set of complexities $(C(F'_i, H); i = 1, 2, 3, \dots)$ is bounded, then by the main theorem in [B-M,I] there can be only finitely many distinct conjugacy classes among the links $L'_i = \partial F'_i$. If, on the other hand, the set of complexities is unbounded, there will be a contradiction to Lemmas 4, 5, 6, 7 of this paper. The proof of Theorem 1 is complete. \square

Proof of the Corollary. Notice that the exchange move as defined in Figure 1 does not change the exponent sum of an n -braid. Since, by

the proof of Theorem 1, it is possible after repeated applications of exchange moves to produce one of the finitely many conjugacy classes of Theorem 2 of [B-M,I], we can only have finitely many possible exponent sums. \square

Proof of Theorem 2. The proof is the same as the proof of Theorem 1. Just notice that nowhere is it used that all the L_i 's represent the same link type. The only fact we needed was that all the L_i 's bound incompressible Seifert surfaces of fixed genus.

REFERENCES

- [Be] D. Bennequin, *Entrelacements et equations de Pfaff*, Asterisque, **107-108** (1983), 87–161.
- [B-M] J. Birman and W. Menasco, *A Calculus on Links in S^3* , to appear in the Proceedings of the International Conference on Knots, Osaka, Japan, August 14–18, 1990.
- [B-M,I] —, *Studying links via closed braids I: A finiteness theorem*, Pacific J. Math., **154** (1992), 17–36.
- [B-M,II] —, *Studying links via closed braids II: On a theorem of Bennequin*, Topology Appl., **40** (1991), 71–82.
- [B-M,III] —, *Studying links via closed braids III: Classifying links which are closed 3-braids*, Pacific J. Math., to appear.
- [B-M,IV] —, *Studying links via closed braids IV: Split and composite links*, Invent. Math., **102**, Fasc. 1 (1990), 115–139.
- [B-M,V] —, *Studying links via closed braids V: The unlink*, Trans. Amer. Math. Soc., **329** (1992), 585–606.
- [J] V. R. F. Jones, *Hecke algebra representations of braid groups and link polynomials*, Annals of Math., **126** (1987), 335–388.
- [VB] James Van Buskirk, *Prime positive knots with inconjugate minimal string braid representatives*, preprint, University of Oregon.

Received November 17, 1990 and in revised form July 15, 1991. Partially supported by NSF grants DMS-88-05672 and DMS-90-02673.

COLUMBIA UNIVERSITY
 NEW YORK, NY 10027
E-mail address: jb@math.columbia.edu

AND

SUNY AT BUFFALO
 BUFFALO, NY 14214
E-mail address: mthtwwm@ubvmsd.cc.buffalo.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982) F. WOLF (1904–1989)

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@henri.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Elaine Barth, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 50 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$190.00 a year (10 issues). Special rate: \$95.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Copyright © 1992 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 156 No. 2 December 1992

| | |
|--|-----|
| Surfaces in the 3-dimensional Lorentz-Minkowski space satisfying $\Delta x = Ax + B$ | 201 |
| LUIS ALÍAS, ANGEL FERRANDEZ and PASCUAL LUCAS | |
| Lie algebras of type D_4 over number fields | 209 |
| BRUCE ALLISON | |
| Subsemigroups of completely simple semigroups | 251 |
| ANNE ANTONIPPILLAI and FRANCIS PASTIJN | |
| Studying links via closed braids. VI. A nonfiniteness theorem | 265 |
| JOAN BIRMAN and WILLIAM W. MENASCO | |
| Minimal orbits at infinity in homogeneous spaces of nonpositive curvature | 287 |
| MARÍA J. DRUETTA | |
| Generalized horseshoe maps and inverse limits | 297 |
| SARAH ELIZABETH HOLTE | |
| Determinantal criteria for transversality of morphisms | 307 |
| DAN LAKSOV and ROBERT SPEISER | |
| Four dodecahedral spaces | 329 |
| PETER LORIMER | |
| Semifree actions on spheres | 337 |
| MONICA NICOLAU | |
| Conformal deformations preserving the Gauss map | 359 |
| ENALDO VERGASTA | |
| Hecke eigenforms and representation numbers of arbitrary rank lattices | 371 |
| LYNNE WALLING | |