

*Pacific
Journal of
Mathematics*

INTEGRAL SPINOR NORMS IN DYADIC LOCAL FIELDS. I

FEI XU

INTEGRAL SPINOR NORMS IN DYADIC LOCAL FIELDS I

FEI XU

The spinor norms of integral rotations on some quadratic forms over an arbitrary dyadic local field are determined. As an application, the results obtained in Bon Durant's paper are improved.

1. Introduction. Because of the absence of a local-global principle in the equivalence of integral quadratic forms over a global field, Eichler and Kneser developed the theory of spinor genera, which strongly depend on spinor norms of local integral rotations. For nondyadic cases the spinor norms are well understood (see [6]), but they become difficult to be determined when 2 is no longer a unit. Hsia [5], the author [10] computed the spinor norms of local integral rotations on modular quadratic forms completely. Earnest and Hsia [3] determined the spinor norms of integral rotations on arbitrary quadratic forms over a dyadic local field in which 2 is prime, and Bon Durant [1] recently considered the spinor norms of integral rotations over a local field which is a quadratic ramified extension field of Q_2 . In the present article we first determine the spinor norms of all binary quadratic forms over an arbitrary dyadic local field, and then use these results to obtain the spinor norms of some quadratic forms over an arbitrary dyadic local field which Jordan components are one dimension. As an application, we consider the spinor norms of quadratic forms over the dyadic local fields which ramification index of 2 is 2 and improve the sufficient condition for the class number of an indefinite quadratic form over the ring of integers of a number field to be a divisor of the class number of the field.

Notation and terminology used here is that of [8]. In particular, F denotes a dyadic local field, ϑ the ring of integers in F , \mathfrak{p} the maximal ideal of ϑ , U the group of units in ϑ , $e = \text{ord } 2$ the ramification index of 2 in F , π a fixed prime element in F , $D(\ , \)$ the quadratic defect function, Δ a fixed unit of quadratic defect 4ϑ , V a regular quadratic space over F associated symmetric bilinear form $B(x, y)$, L a lattice on V , $O^+(V)$ the group of rotations on V , $O^+(L)$ the corresponding subgroup of units of L , and $\theta(\ , \)$ the

spinor norm function. We use the symbol $\langle a, b, c, \dots \rangle$ for lattices, and $[a, b, c, \dots]$ for spaces.

2. Binary cases. In this section we determine the spinor norms of local integral rotations on binary lattices completely. All the results are expressed in convenient closed forms. Some cases have already been treated in [4], but we may repeat them for completeness.

Let L be a binary lattice. If L is modular then $\theta(O^+(L))$ has been determined in [5] and [10]. So we only consider the nonmodular cases. Since the spinor norm is not affected by scaling, we can put $L = \vartheta x \perp \vartheta y$ with $Q(x) = 1$ and $Q(y) = \varepsilon\pi^r$ where $r \geq 1$ and $\varepsilon \in U$.

PROPOSITION 2.1. *If $r > 4e$, then $\theta(O^+(L)) = \dot{F}^2 \cup \varepsilon\pi^r \dot{F}^2$.*

Proof. Take a symmetry τ_z in $O(L)$ where $z = ax + by$ with $a, b \in \vartheta$ and one of them is in U . By [4, Prop. 3.2], we get $\text{ord } a \leq e$; or $\text{ord } a \geq r - e$, $\text{ord } b = 0$, so $Q(x)$ is in \dot{F}^2 or $\varepsilon\pi^r \dot{F}^2$ respectively by the Local Square Theorem, and $\theta(O^+(L)) = \dot{F}^2 \cup \varepsilon\pi^r \dot{F}^2$ by [7, 1.1].

PROPOSITION 2.2. *Suppose $2e < r \leq 4e$.*

(i) *When r is odd, $\theta(O^+(L)) = ((1 + p^{r-2e})\dot{F}^2 \cup \varepsilon\pi(1 + p^{r-2e})\dot{F}^2) \cap Q(\dot{F}L)$.*

(ii) *When r is even, $D(\varepsilon) = p^d$ with $1 \leq d \leq 2e - r/2$, $\theta(O^+(L)) = ((1 + p^{r-2e+d})\dot{F}^2 \cup \varepsilon(1 + p^{r-2e+d})\dot{F}^2) \cap Q(\dot{F}L)$.*

(iii) *When r is even, $D(\varepsilon) = p^d$ with $d > 2e - r/2$, or 0; $\theta(O^+(L)) = (1 + p^{r/2})\dot{F}^2 \cup \varepsilon(1 + p^{r/2})\dot{F}^2$.*

Proof. Take a symmetry τ_z in $O(L)$ where $z = ax + by$ with $a, b \in \vartheta$ and one of them is in U . By [4, Prop. 3.2], we obtain $\text{ord } a = 0$, $\text{ord } b \geq 0$; or $0 < \text{ord } a \leq e$, $\text{ord } b = 0$; or $\text{ord } a \geq r - e$, $\text{ord } b = 0$.

Case (i). If $\text{ord } a \leq e$, then

$$\text{ord}(a^{-2}b^2\varepsilon\pi^r) \geq r - 2\text{ord } a \geq r - 2e.$$

So

$$Q(z) = a^2(1 + a^{-2}b^2\varepsilon\pi^r)$$

is in $(1 + p^{r-2e})\dot{F}^2 \cap Q(\dot{F}L)$.

If $\text{ord } a \geq r - e$, $\text{ord } b = 0$; then

$$\text{ord}(\varepsilon^{-1}\pi^{-r}(b^{-1}a)^2) = 2 \text{ord } a - r \geq r - 2e.$$

So

$$Q(z) = \varepsilon\pi^r b^2(1 + \varepsilon^{-1}\pi^{-r}(b^{-1}a)^2)$$

is in $\varepsilon\pi(1 + p^{r-2e})\dot{F}^2 \cap Q(\dot{F}L)$. Therefore

$$\theta(O^+(L)) \subseteq ((1 + p^{r-2e})\dot{F}^2 \cup \varepsilon\pi(1 + p^{r-2e})\dot{F}^2) \cap Q(\dot{F}L)$$

by [7, 1.1].

Conversely, take an element h in $(1 + p^{r-2e})\dot{F}^2 \cap Q(\dot{F}L)$, so there exists z in FL such that $Q(z) = h$. Without loss of generality, we assume $z = ax + by$ where $a, b \in \vartheta$ and one of them is a unit.

If $e < \text{ord } a < r/2$, then $\text{ord } b = 0$. Put

$$k = \text{ord}(\varepsilon\pi^r(ba^{-1})^2) = r - 2 \text{ord } a < r - 2e \leq 2e.$$

Note that k is a positive odd integer. Since h is in $(1 + p^{r-2e})\dot{F}^2$, we obtain

$$(1 + c\pi^{r-2e})f^2 = h = a^2 + \varepsilon\pi^r b^2 = a^2(1 + (a^{-1}b)^2\varepsilon\pi^r)$$

where $f \in \dot{F}^2$, $c \in \vartheta$. Let $\eta = af^{-1}$, so η is in U by the above equation, and

$$\eta^2(1 + \varepsilon\pi^r(ba^{-1})^2) = (1 + c\pi^{r-2e}).$$

But

$$D(\eta^2(1 + \varepsilon\pi^r(ba^{-1})^2)) = p^k \supset p^{r-2e} \supseteq D(1 + c\pi^{r-2e})$$

by [8, 63:5], a contradiction.

If $r - e > \text{ord } a > r/2$, then $\text{ord } b = 0$ again,

$$\text{ord } h = \text{ord}(a^2 + \varepsilon\pi^r b^2) = r$$

is odd. But h is in $(1 + p^{r-2e})\dot{F}^2$, so that $\text{ord } h$ is even. This is a contradiction. Now there remains $\text{ord } a = 0$; or $0 < \text{ord } a \leq e$, $\text{ord } b = 0$; or $\text{ord } a \geq r - e$, $\text{ord } b = 0$, so τ_z is in $O(L)$ by [4, Prop. 3.2]. Therefore

$$\theta(O^+(L)) = ((1 + p^{r-2e}) \cup \varepsilon\pi(1 + p^{r-2e})\dot{F}^2) \cap Q(\dot{F}L).$$

Case (ii). Since $D(\varepsilon) = p^d$, we can assume $\varepsilon = 1 + \sigma\pi^d$ for convenience, where $\sigma \in U$ and d is odd. Note $1 \leq d \leq 2e - r/2 < 2e$.

If $\text{ord } a = 0$, then

$$\text{ord}(\varepsilon\pi^r(a^{-1}b)^2) = r + 2 \text{ord } b \geq r > r - 2e + d.$$

So

$$Q(z) = a^2(1 + \varepsilon\pi^r(a^{-1}b)^2)$$

is in $(1 + p^{r-2e})\dot{F}^2 \cap Q(\dot{F}L)$. If $0 < \text{ord } a \leq e < r/2$, $\text{ord } b = 0$ then

$$\text{ord}(\sigma\pi^{d+r}(a^{-1}b)^2) = r + d - 2 \text{ord } a \geq r + d - 2e$$

and

$$\text{ord}(-2\pi^{r/2}(a^{-1}b)) = e + r/2 - \text{ord } a \geq r/2 \geq r + d - 2e.$$

So

$$\begin{aligned} Q(z) &= a^2(1 + \varepsilon\pi^r(a^{-1}b)^2) = a^2(1 + (1 + \sigma\pi^d)\pi^r(a^{-1}b)^2) \\ &= a^2(1 + \pi^r(a^{-1}b)^2 + \sigma\pi^{r+d}(a^{-1}b)^2) \\ &= a^2((1 + \pi^{r/2})^2 - 2\pi^{r/2}(a^{-1}b) + \sigma\pi^{d+r}(a^{-1}b)^2) \end{aligned}$$

is in $(1 + p^{r-2e+d})\dot{F}^2 \cap Q(\dot{F}L)$.

If $\text{ord } a \geq r - e$, $\text{ord } b = 0$; then

$$\text{ord}(-2(\varepsilon^{-1}b^{-1})a\pi^{-r/2}) = e + \text{ord } a - r/2 \geq r - 2e + d$$

and

$$\text{ord}(\sigma(\varepsilon^{-1}b^{-1})^2a^2\pi^{-r+d}) = 2 \text{ord } a - r + d \geq r - 2e + d.$$

So

$$\begin{aligned} Q(z) &= \varepsilon\pi^r b^2(1 + \varepsilon(\varepsilon^{-1}b^{-1})^2a^2\pi^{-r}) \\ &= \varepsilon\pi^r b^2(1 + (1 + \sigma\pi^d)(\varepsilon^{-1}b^{-1})^2a^2\pi^{-r}) \\ &= \varepsilon\pi^r b^2(1 + (\varepsilon^{-1}b^{-1})^2a^2\pi^{-r} + \sigma(\varepsilon^{-1}b^{-1})^2a^2\pi^{-r+d}) \\ &= \varepsilon\pi^r b^2((1 + \varepsilon^{-1}b^{-1}a\pi^{-r/2})^2 - 2\varepsilon^{-1}b^{-1}a\pi^{-r/2} \\ &\quad + \sigma(\varepsilon^{-1}b^{-1})^2a^2\pi^{-r+d}) \end{aligned}$$

is in $\varepsilon(1 + p^{r-2e+d})\dot{F}^2 \cap Q(\dot{F}L)$. Therefore

$$\theta(O^+(L)) \subseteq ((1 + p^{r-2e+d})\dot{F}^2 \cup \varepsilon(1 + p^{r-2e+d})\dot{F}^2) \cap Q(\dot{F}L).$$

Conversely, take an element h in $(1 + p^{r-2e+d})\dot{F}^2 \cap Q(\dot{F}L)$, so there exists z in FL such that $Q(z) = h$. Without loss of generality, we assume $z = ax + by$ where $a, b \in \vartheta$ and one of them is a unit.

If $e < \text{ord } a < r/2$, then $\text{ord } b = 0$. Put

$$\begin{aligned} k &= \text{ord}(\sigma(a^{-1}b)^2\pi^{r+d}) = r + d - 2 \text{ord } a, \\ &= (e + r/2 - \text{ord } a) + (r/2 + d - e - \text{ord } a) \\ &< \min\{(e + r/2 - \text{ord } a) + (r/2 + d - 2e), r + d - 2e\} \\ &\leq \min\{(e + r/2 - \text{ord } a), r + d - 2e\} \\ &= \min\{\text{ord}(-2a^{-1}b\pi^{r/2}), r + d - 2e\} \end{aligned}$$

and k is a positive odd integer. Since h is in $(1 + p^{r-2e+d})\dot{F}^2$, we obtain

$$f^2(1 + c\pi^{r-2e+d}) = h = (a^2 + \varepsilon\pi^r b^2) = a^2(1 + (a^{-1}b)^2\varepsilon\pi^r)$$

where $f \in \dot{F}$, $c \in \vartheta$. Let $\eta = af^{-1}$, so η is in U by the above equation, and

$$\begin{aligned} 1 + c\pi^{r-2e+d} &= \eta^2(1 + \varepsilon\pi^r(a^{-1}b)^2) \\ &= \eta^2(1 + (1 + \sigma\pi^d)\pi^r(a^{-1}b)^2) \\ &= \eta^2(1 + \pi^r(a^{-1}b)^2 + \sigma\pi^{r+d}(a^{-1}b)^2) \\ &= \eta^2((1 + (a^{-1}b)\pi^{r/2})^2 - 2(a^{-1}b)\pi^{r/2} + \sigma\pi^{r+d}(a^{-1}b)^2). \end{aligned}$$

But

$$\begin{aligned} D(\eta^2((1 + a^{-1}b\pi^{r/2})^2 - 2(a^{-1}b)\pi^{r/2} + \sigma\pi^{r+d}(a^{-1}b)^2)) \\ = p^k \supset p^{r+d-2e} \supseteq D(1 + c\pi^{r-2e+d}) \end{aligned}$$

by [8, 63:5], a contradiction.

If $\text{ord } a = r/2$, then $\text{ord } b = 0$. Let $\xi = a^{-1}\pi^{r/2}b$, so ξ is in U . Suppose

$$\begin{aligned} h &= a^2(1 + \varepsilon\pi^r(a^{-1}b)^2) = a^2(1 + \varepsilon\xi^2) = a^2(1 + (1 + \sigma\pi^d)\xi^2) \\ &= a^2(1 + \xi^2 + \sigma\xi^2\pi^d) = a^2((1 + \xi)^2 - 2\xi + \sigma\xi^2\pi^d) \end{aligned}$$

is in $(1 + p^{r-2e+d})\dot{F}^2$. Note $d \leq 2e - r/2 < e$, and d is odd, so $2 \text{ord}(1 + \xi) = \text{ord}(1 + \xi)^2 < d < e$.

Writing $1 + \xi = \delta\pi^s$ with $\delta \in U$, $s \geq 0$, we obtain

$$\eta^2(1 - \delta^{-2}\xi^2 2\pi^{-2s} + \sigma\delta^{-2}\xi^2\pi^{d-2s}) = (1 + c\pi^{r-2e+d})$$

where $\eta \in U$, $c \in \vartheta$. But

$$\begin{aligned} D(\eta^2(1 - \delta^{-2}\xi^2 2\pi^{-2s} + \sigma\delta^{-2}\xi^2\pi^{d-2s})) \\ = p^{d-2s} \supseteq p^d \supset p^{r-2e+d} \supseteq D(1 + c\pi^{r-2e+d}) \end{aligned}$$

by [8, 63:5], a contradiction

If $r/2 < \text{ord } a < r - e$, then $\text{ord } b = 0$. We have

$$\text{ord}(2b^{-1}a\pi^{-r/2}) = e + \text{ord } a - r/2 > e > 2e - r/2 \geq d \geq 1,$$

and

$$\begin{aligned} h &= Q(z) = b^2\pi^r(\varepsilon + (b^{-1}a\pi^{-r/2})^2) = b^2\pi^r(1 + \sigma\pi^d + (b^{-1}a\pi^{-r/2})^2) \\ &= b^2\pi^r((1 + b^{-1}a\pi^{-r/2})^2 - 2b^{-1}a\pi^{-r/2} + \sigma\pi^d) \end{aligned}$$

is in $(1 + p^{r-2e+d})\dot{F}^2$ by hypothesis. So

$$\eta^2((1 + b^{-1}a\pi^{-r/2})^2 - 2b^{-1}a\pi^{-r/2} + \sigma\pi^d) = (1 + c\pi^{r-2e+d})$$

where $\eta \in U$ and $c \in \mathfrak{O}$. But

$$\begin{aligned} D(\eta^2((1 + b^{-1}a(\pi^{-r/2})^2 - 2b^{-1}a\pi^{-r/2} + \sigma\pi^d)) \\ = p^d \supset p^{r-2e+d} \supseteq D(1 + c\pi^{r-2e+d}) \end{aligned}$$

by [8, 63:5], a contradiction.

Now there remains $\text{ord } a = 0$ or $0 < \text{ord } a \leq e$, $\text{ord } b = 0$; or $\text{ord } a \geq r - e$, $\text{ord } b = 0$, so τ_z is in $O(L)$ by [4, Prop. 3.2]. Therefore

$$\theta(O^+(L)) = ((1 + p^{r-2e+d})\dot{F}^2 \cup \varepsilon(1 + p^{r-2e+d})\dot{F}^2) \cap Q(\dot{F}L).$$

Case (iii). Write $\varepsilon = 1 + \sigma\pi^d$ with $d > 2e - r/2$, $\sigma \in U$.

If $\text{ord } a = 0$, then

$$\text{ord}(\varepsilon\pi^r(a^{-1}b)^2) = r + 2\text{ord } b \geq r \geq r/2,$$

so

$$Q(z) = a^2(1 + \varepsilon\pi^r(a^{-1}b)^2)$$

is in $(1 + p^{r/2})\dot{F}^2$.

If $0 < \text{ord } a \leq e < r/2$, $\text{ord } b = 0$; then

$$\begin{aligned} r/2 &\leq \text{ord}(-2(a^{-1}b)\pi^{r/2}) = e - \text{ord } a + r/2 \\ &= (r + d - 2\text{ord } a) + (\text{ord } a - r/2 - d + e) \\ &\leq (r + d - 2\text{ord } a) + (2e - r/2 - d) \\ &< r + d - 2\text{ord } a = \text{ord}(\sigma\pi^{r+d}(a^{-1}b)^2). \end{aligned}$$

So

$$\begin{aligned} Q(z) &= a^2(1 + \varepsilon\pi^r(a^{-1}b)^2) = a^2(1 + (1 + \sigma\pi^d)\pi^r(a^{-1}b)^2) \\ &= a^2(1 + (a^{-1}b)^2\pi^r + \sigma(a^{-1}b)^2\pi^{r+d}) \\ &= a^2(1 + a^{-1}b\pi^{r/2})^2 - 2a^{-1}b\pi^{r/2} + \sigma(a^{-1}b)^2\pi^{r+d} \end{aligned}$$

is in $(1 + p^{r/2})\dot{F}^2$.

If $\text{ord } a \geq r - e > r/2$, $\text{ord } b = 0$; then

$$\begin{aligned} r/2 &\leq \text{ord}(-2\varepsilon^{-1}b^{-1}a\pi^{-r/2}) = e + \text{ord } a - r/2 \\ &= (2\text{ord } a + d - r) + (r/2 - \text{ord } a + e - d) \\ &\leq (2\text{ord } a + d - r) + (2e - r/2 - d) \\ &< (2\text{ord } a + d - r) = \text{ord}(\sigma(\varepsilon^{-1}b^{-1}a)^2\pi^{d-r}). \end{aligned}$$

So

$$\begin{aligned} Q(z) &= \varepsilon\pi^r b^2(1 + \varepsilon(\varepsilon^{-1}b^{-1}a)^2\pi^{-r}) \\ &= \varepsilon\pi^r b^2(1 + (1 + \sigma\pi^d)(\varepsilon^{-1}b^{-1}a)^2\pi^{-r}) \\ &= \varepsilon\pi^r b^2(1 + (\varepsilon^{-1}b^{-1}a)^2\pi^{-r} + \sigma(\varepsilon^{-1}b^{-1}a)^2\pi^{d-r}) \\ &= \varepsilon\pi^r b^2((1 + \varepsilon^{-1}b^{-1}a\pi^{-r/2})^2 - 2\varepsilon^{-1}b^{-1}a\pi^{-r/2} \\ &\quad + \sigma(\varepsilon^{-1}b^{-1}a)^2\pi^{d-r}) \end{aligned}$$

is in $\varepsilon(1 + p^{r/2})\dot{F}^2$.

Therefore

$$\theta(O^+(L)) \subseteq (1 + p^{r/2})\dot{F}^2 \cup \varepsilon(1 + p^{r/2})\dot{F}^2$$

by [7, 1.1].

Conversely, take an element h in $(1 + p^{r/2})\dot{F}^2$, by the Local Square Theorem, we assume $h = 1 + c\pi^{r/2}$ with $0 \leq \text{ord } c < e$. Write $c = \lambda\pi^k$ with $\lambda \in U$. By Hensel's lemma, there exists η in U such that

$$(\pi^{r-2e+2k} - \sigma c^{-1}\pi^{d+r/2-2e+2k})\eta^2(2c^{-1}\pi^{k-e} + 2\pi^{r/2-e+k})\eta + 1 = 0.$$

Put $z = \pi^{e-k}x + \eta y$, so τ_z is in $O(L)$ by [4, Prop. 3.2], and

$$\begin{aligned} Q(z) &= \pi^{2e-2k} + \varepsilon\pi^r \eta^2 = \pi^{2e-2k} + (1 + \sigma\pi^d)\pi^r \eta^2 \\ &= \pi^{2e-2k}(1 + \pi^{r+2k-2e}\eta^2 + \sigma\pi^{d+r+2k-2e}\eta^2) \\ &= \pi^{2e-2k}((1 + \pi^{r/2+k-e}\eta)^2 - 2\pi^{r/2+k-e}\eta + \sigma\pi^{d+r+2k-2e}\eta^2) \\ &= \pi^{2e-2k}((1 + \pi^{r/2+k-e}\eta)^2 \\ &\quad + (-2c^{-1}\pi^{k-e}\eta + \sigma c^{-1}\pi^{d+r/2+2k-2e}\eta^2)c\pi^{r/2}) \\ &= \pi^{2e-2k}((1 + \pi^{r/2+k-e}\eta)^2 + (1 + \pi^{r/2-e+k}\eta)^2 c\pi^{r/2}) \\ &= \pi^{2e-2k}(1 + \pi^{r/2+k-e}\eta)^2(1 + c\pi^{r/2}) = \pi^{2e-2k}(1 + \pi^{r/2+k-e}\eta)^2 h. \end{aligned}$$

Therefore

$$\theta(O^+(L)) = (1 + p^{r/2})\dot{F}^2 \cup \varepsilon(1 + p^{r/2})\dot{F}^2.$$

PROPOSITION 2.3. *Suppose $0 < r \leq 2e$.*

(i) *When r is odd, or r is even and $D(-\varepsilon) = p^d$ with $1 \leq d \leq e - r/2$; $\theta(O^+(L)) = Q(\dot{F}L)$.*

(ii) *When r is even and $D(-\varepsilon) = p^d$ with $(3e - r/2)/2 \geq d > e - r/2$; $\theta(O^+(L)) = ((1 + p^{d-e+r/2})\dot{F}^2) \cap Q(\dot{F}L)$.*

(iii) *When r is even and $D(-\varepsilon) = p^d$ with $d > (3e - r/2)/2$, or 0; $\theta(O^+(L)) = (1 + p^{e-[e/2-r/4]})\dot{F}^2$.*

Proof. Take a maximal anisotropic vector $z = ax + by$ in L , where $a, b \in \vartheta$ and one of them is a unit, we know τ_z is in $O(L)$ if and only if $2B(z, L) \subseteq Q(z)\vartheta$. This is equivalent to

$$\text{ord}(a^2 + \varepsilon\pi^r b^2) \leq \min\{\text{ord } a, r + \text{ord } b\} + e.$$

In fact, this inequality is the same as $\text{ord } a \neq r/2$; or $\text{ord } a = r/2$, $\text{ord } b = 0$, $0 \leq \text{ord}((a\pi^{-r/2}b^{-1})^2 + \varepsilon) \leq e - r/2$.

Case (i). When r is odd, we always have $\text{ord } a \neq r/2$; when r is even and $\text{ord } a = r/2$, we have $0 \leq \text{ord}(\varepsilon + (\pi^{-r/2}b^{-1}a)^2) \leq d \leq e - r/2$ since $D(-\varepsilon) = p^d$ with $1 \leq d \leq e - r/2$. Note FL is anisotropic in this case, so $\theta(O^+(L)) = Q(\dot{F}L)$.

Case (ii). Write $-\varepsilon = 1 + \sigma\pi^d$ with $\sigma \in U$ and $2e > (3e - r/2)/2 \geq d > e - r/2$.

If $\text{ord } a = 0$, then

$$\begin{aligned} \text{ord}(\sigma(\pi^{r/2}ba^{-1})^2\pi^d) &= d + r + 2 \text{ord } b \\ &= (e + r/2 + \text{ord } b) + \text{ord } b + (d + r/2 - e) \\ &> e + r/2 + \text{ord } b = \text{ord}(-2\pi^{r/2}(ba^{-1})(1 + a^{-1}b\pi^{r/2})) \\ &\geq e + r/2 > d - e + r/2. \end{aligned}$$

So

$$\begin{aligned} Q(z) &= a^2(1 + \varepsilon\pi^r(a^{-1}b)^2) = a^2(1 - (1 + \sigma\pi^d)(a^{-1}b)^2\pi^r) \\ &= a^2((1 + a^{-1}b\pi^{r/2})^2 \\ &\quad - 2(a^{-1}b\pi^{r/2})(1 + a^{-1}b\pi^{r/2}) - \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \end{aligned}$$

is in $(1 + p^{d-e+r/2})\dot{F}^2 \cap Q(\dot{F}L)$.

If $0 < \text{ord } a < r/2$, $\text{ord } b = 0$; then

$$\begin{aligned} \text{ord}(-2a^{-1}b\pi^{r/2}(1 + a^{-1}b\pi^{r/2})) &= e + r/2 - \text{ord } a \\ &> e = (e + e/2 - r/4) + r/4 - e/2 \geq d + (r/2 - e)/2 \\ &\geq d - e + r/2 \end{aligned}$$

and

$$\text{ord}(-\sigma(a^{-1}b\pi^{r/2})^2\pi^d) = d + r - 2 \text{ord } a \geq d \geq d - e + r/2.$$

So

$$\begin{aligned} Q(z) &= a^2((1 + a^{-1}b\pi^{r/2})^2 \\ &\quad - 2(a^{-1}b\pi^{r/2})(1 + a^{-1}b\pi^{r/2}) - \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \end{aligned}$$

is in $(1 + p^{d-e+r/2})\dot{F}^2 \cap Q(\dot{F}L)$.

If $\text{ord } a > r/2$, $\text{ord } b = 0$; then

$$\begin{aligned} & \text{ord}(-2\varepsilon^{-1}b^{-1}a\pi^{-r/2}(1 + \varepsilon^{-1}b^{-1}a\pi^{-r/2})) \\ & = e + \text{ord } a - r/2 > e \geq d - e + r/2 \end{aligned}$$

and

$$\text{ord}(-\sigma(\varepsilon^{-1}b^{-1}a\pi^{-r/2})^2\pi^d) = -r + 2 \text{ord } a + d > d \geq d - e + r/2.$$

So

$$\begin{aligned} Q(z) &= \varepsilon\pi^r b^2(1 + \varepsilon(\varepsilon^{-1}b^{-1}a\pi^{-r/2})^2) \\ &= \varepsilon\pi^r b^2(1 - (1 + \sigma\pi^d)(\varepsilon^{-1}b^{-1}a\pi^{-r/2})^2) \\ &= \varepsilon\pi^r b^2((1 + \varepsilon^{-1}b^{-1}a\pi^{-r/2})^2 \\ &\quad - 2\varepsilon^{-1}b^{-1}a\pi^{-r/2}(1 + \varepsilon^{-1}b^{-1}a\pi^{-r/2}) \\ &\quad - \sigma(\varepsilon^{-1}b^{-1}a\pi^{-r/2})^2\pi^d) \end{aligned}$$

is in $\varepsilon(1 + p^{d-e+r/2})\dot{F}^2 \cap Q(\dot{F}L)$.

If $\text{ord } a = r/2$, $\text{ord } b = 0$, and $\text{ord}(\varepsilon + (\pi^{-r/2}ab^{-1})^2) \leq e - r/2$, note

$$\text{ord}(2a^{-1}b\pi^{r/2}(1 + a^{-1}b\pi^{r/2})) = e + \text{ord}(1 + a^{-1}b\pi^{r/2}) \geq e > e - r/2$$

and

$$\text{ord}(\sigma(a^{-1}b\pi^{r/2})^2\pi^d) = d > e - r/2.$$

Then

$$\begin{aligned} & \text{ord}(1 + a^{-1}b\pi^{r/2})^2 \\ &= \text{ord}((1 + a^{-1}b\pi^{r/2})^2 \\ &\quad - 2a^{-1}b\pi^{r/2}(1 + a^{-1}b\pi^{r/2})^2 - \sigma(a^{-1}b\pi^{r/2})^2\pi^d \\ &\quad + 2a^{-1}b\pi^{r/2}(1 + a^{-1}b\pi^{r/2}) + \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ &= \text{ord}(1 - (a^{-1}b\pi^{r/2})^2 - \sigma(a^{-1}b\pi^{r/2})^2\pi^d \\ &\quad + 2a^{-1}b\pi^{r/2}(1 + a^{-1}b\pi^{r/2}) + \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ &= \text{ord}(1 + (a^{-1}b\pi^{r/2})^2(-1 - \sigma\pi^d) \\ &\quad + 2a^{-1}b\pi^{r/2}(1 + a^{-1}b\pi^{r/2}) + \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ &= \text{ord}((1 + \varepsilon(a^{-1}b\pi^{r/2})^2) \\ &\quad + 2a^{-1}b\pi^{r/2}(1 + a^{-1}b\pi^{r/2}) + \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ &= \text{ord}(1 + \varepsilon(a^{-1}b\pi^{r/2})^2) \leq e - r/2. \end{aligned}$$

So

$$\begin{aligned} & \text{ord}(\sigma\pi^d(a^{-1}b\pi^{r/2})^2(1+a^{-1}b\pi^{r/2})^{-2}) \\ & = d - 2 \text{ord}(1+a^{-1}b\pi^{r/2}) \geq d - e + r/2 \end{aligned}$$

and

$$\begin{aligned} & \text{ord}((-2a^{-1}b\pi^{r/2})(1+a^{-1}b\pi^{r/2})^{-1}) = e - \text{ord}(1+a^{-1}b\pi^{r/2}) \\ & \geq e - (e - r/2)/2 \geq d - e + r/2. \end{aligned}$$

Thus

$$\begin{aligned} Q(z) & = a^2(1+(a^{-1}b\pi^{r/2})\varepsilon) = a^2(1-(1+\sigma\pi^d)(a^{-1}b\pi^{r/2})^2) \\ & = a^2((1+a^{-1}b\pi^{r/2})^2 \\ & \quad - 2(1+a^{-1}b\pi^{r/2})a^{-1}b\pi^{r/2} - \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ & = a^2(1+a^{-1}b\pi^{r/2})^2(1-2a^{-1}b\pi^{r/2}(1+a^{-1}b\pi^{r/2})^{-1} \\ & \quad - \sigma\pi^d(a^{-1}b\pi^{r/2})^2(1+a^{-1}b\pi^{r/2})^{-2}) \end{aligned}$$

is in $(1+p^{d-e+r/2})\dot{F}^2 \cap Q(\dot{F}L)$. Therefore

$$\theta(O^+(L)) \subseteq ((1+p^{d-e+r/2})\dot{F}^2 \cap \varepsilon(1+p^{d-e+r/2})\dot{F}^2) \cap Q(\dot{F}L).$$

Conversely, take an element h in $(1+p^{d-e+r/2})\dot{F}^2 \cap Q(\dot{F}L)$, so there exists z in $\dot{F}L$ such that $h = Q(z)$. We can assume $z = ax + by$ with $a, b \in \vartheta$ and one of them is in U . We claim that τ_z is in $O(L)$. In fact, if $\text{ord } a = r/2$, $\text{ord } b = 0$, and $\text{ord}((\pi^{-r/2}ab^{-1})^2 + \varepsilon) > e - r/2$; note

$$\begin{aligned} h = Q(z) & = a^2(1 + \varepsilon(a^{-1}b\pi^{r/2})^2) = a^2(1 + (-1 - \sigma\pi^d)(a^{-1}b\pi^{r/2})^2) \\ & = a^2((1+a^{-1}b\pi^{r/2})^2 - 2a^{-1}b\pi^{r/2}(1+a^{-1}b\pi^{r/2}) - \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \end{aligned}$$

is in $(1+p^{d-e+r/2})\dot{F}^2$, and d is odd, $d < 2e$; then

$$\begin{aligned} & \text{ord}(1+a^{-1}b\pi^{r/2})^2 \\ & = \text{ord}((1+a^{-1}b\pi^{r/2})^2 \\ & \quad - 2a^{-1}b\pi^{r/2}(1+a^{-1}b\pi^{r/2}) - \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ & = \text{ord}(1 + \varepsilon(a^{-1}b\pi^{r/2})^2) > e - r/2. \end{aligned}$$

Write $1+a^{-1}b\pi^{r/2} = \lambda\pi^s$ with $\lambda \in U$, $s \geq 0$. So $d > 2s > e - r/2$, and

$$\eta^2(1 - (\lambda^{-1}a^{-1}b\pi^{r/2})(2\pi^{-s}) - \sigma(\lambda^{-1}a^{-1}b\pi^{r/2})^2\pi^{d-2s}) = 1 + c\pi^{d-e+r/2}$$

where $\eta \in U$, $c \in \vartheta$. Since $d \leq e + (e - r/2)/2 < e + s$, that is $0 < d - 2s < e - s$, and $d - 2s$ is odd, we have

$$\begin{aligned} & D(\eta^2(1 - (\lambda^{-1}a^{-1}b\pi^{r/2})(2\pi^{-s}) - \sigma(\lambda^{-1}a^{-1}b\pi^{r/2})^2\pi^{d-2s})) = p^{d-2s} \\ & \supset p^{d-e+r/2} \supseteq D(1 + c\pi^{d-e+r/2}). \end{aligned}$$

This leads to a contradiction. So τ_z is in $O(L)$, and

$$\theta(O^+(L)) = ((1 + p^{d-e+r/2})\dot{F}^2 \cup \varepsilon(1 + p^{d-e+r/2})\dot{F}^2) \cap Q(\dot{F}L).$$

Case (iii). Write $-\varepsilon = 1 + \sigma\pi^d$ with $\sigma \in U$ and $d > (3e - r/2)/2$.

If $\text{ord } a = 0$, then

$$\begin{aligned} \text{ord}(-2(1 + a^{-1}b\pi^{r/2})a^{-1}b\pi^{r/2}) &= e + \text{ord } b + r/2 \\ &\geq e + r/2 > e - [e/2 - r/4] \end{aligned}$$

and

$$\begin{aligned} \text{ord}(-\sigma(a^{-1}b\pi^{r/2})^2\pi^d) &= 2 \text{ord } b + r + d \\ &\geq r + d > d > e + [e/2 - r/4] \geq e - [e/2 - r/4]. \end{aligned}$$

So

$$\begin{aligned} Q(z) &= a^2(1 - (a^{-1}b\pi^{r/2})^2(1 + \sigma\pi^d)) \\ &= a^2((1 + a^{-1}b\pi^{r/2})^2 \\ &\quad - 2(1 + a^{-1}b\pi^{r/2})a^{-1}b\pi^{r/2} - \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \end{aligned}$$

is in $(1 + p^{e-[e/2-r/4]})\dot{F}^2$.

If $\text{ord } a > r/2$, $\text{ord } b = 0$; then

$$\begin{aligned} \text{ord}(-2(\varepsilon^{-1}b^{-1}\pi^{-r/2}a)(1 + \varepsilon^{-1}b^{-1}\pi^{-r/2}a)) \\ = e - r/2 + \text{ord } a > e \geq e - [e/2 - r/4] \end{aligned}$$

and

$$\begin{aligned} \text{ord}(-\sigma(\varepsilon^{-1}b^{-1}\pi^{-r/2}a)^2\pi^d) &= d + 2 \text{ord } a - r \\ &> d > e + [e/2 - r/4] \geq e - [e/2 - r/4]. \end{aligned}$$

So

$$\begin{aligned} Q(z) &= \varepsilon\pi^r b^2(1 + \varepsilon(\varepsilon^{-1}b^{-1}\pi^{-r/2}a)^2) \\ &= \varepsilon\pi^r b^2(1 - (1 + \sigma\pi^d)(\varepsilon^{-1}b^{-1}\pi^{-r/2}a)^2) \\ &= \varepsilon\pi^r b^2((1 + \varepsilon^{-1}b^{-1}\pi^{-r/2}a)^2 \\ &\quad - 2(1 + \varepsilon^{-1}b^{-1}\pi^{-r/2}a)(\varepsilon^{-1}b^{-1}\pi^{-r/2}a) \\ &\quad - \sigma(\varepsilon^{-1}b^{-1}\pi^{-r/2}a)^2\pi^d) \end{aligned}$$

is in $\varepsilon(1 + p^{e-[e/2-r/4]})\dot{F}^2$.

If $\text{ord } a < r/2$, $\text{ord } b = 0$; then

$$\begin{aligned} \text{ord}(-2(1 + a^{-1}b\pi^{r/2})a^{-1}b\pi^{r/2}) \\ = e - \text{ord } a + r/2 > e \geq e - [e/2 - r/4] \end{aligned}$$

and

$$\begin{aligned} \text{ord}(-\sigma(a^{-1}b\pi^{r/2})^2\pi^d) &= r + d - 2 \text{ord } a \\ &> d > e + [e/2 - r/4] \geq e - [e/2 - r/4]. \end{aligned}$$

So

$$Q(z) = a^2((1+a^{-1}b\pi^{r/2})^2 - 2(1+a^{-1}b\pi^{r/2})a^{-1}b\pi^{r/2} - \sigma(a^{-1}b\pi^{r/2})^2\pi^d)$$

is in $(1 + p^{e-[e/2-r/4]})\dot{F}^2$. If $\text{ord } a = r/2$, $\text{ord } b = 0$, and

$$\text{ord}(\varepsilon + (b^{-1}a\pi^{-r/2})^2) \leq e - r/2,$$

note

$$\text{ord}(\sigma(a^{-1}b\pi^{r/2})^2\pi^d) = d > e + (e/2 - r/4) \geq e > e - r/2$$

and

$$\text{ord}(2(1+a^{-1}b\pi^{r/2})(a^{-1}b\pi^{r/2})) = e + \text{ord}(1+a^{-1}b\pi^{r/2}) \geq e > e - r/2$$

Then

$$\begin{aligned} &\text{ord}(1+a^{-1}b\pi^{r/2})^2 \\ &= \text{ord}((1+a^{-1}b\pi^{r/2})^2 - 2(1+a^{-1}b\pi^{r/2})(a^{-1}b\pi^{r/2}) \\ &\quad - \sigma(a^{-1}b\pi^{r/2})^2\pi^d \\ &\quad + 2(1+a^{-1}b\pi^{r/2})(a^{-1}b\pi^{r/2}) + \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ &= \text{ord}((1+\varepsilon(a^{-1}b\pi^{r/2})^2) + 2(1+a^{-1}b\pi^{r/2})(a^{-1}b\pi^{r/2}) \\ &\quad + \sigma(a^{-1}b\pi^{r/2})^2\pi^d) \\ &= \text{ord}(1+\varepsilon(a^{-1}b\pi^{r/2})^2) \leq e - r/2. \end{aligned}$$

So

$$\begin{aligned} &\text{ord}(-2(1+a^{-1}b\pi^{r/2})^{-1}(a^{-1}b\pi^{r/2})) \\ &= e - \text{ord}(1+a^{-1}b\pi^{r/2}) \geq e - [e/2 - r/4] \end{aligned}$$

and

$$\begin{aligned} &\text{ord}(-\sigma(a^{-1}b\pi^{r/2})^2\pi^d(1+a^{-1}b\pi^{r/2})^{-2}) \\ &= d - 2 \text{ord}(1+a^{-1}b\pi^{r/2}) \geq d - e + r/2 \geq e - [e/2 - r/4]. \end{aligned}$$

Consider

$$Q(z) = a^2(1+a^{-1}b\pi^{r/2})^2(1-2a^{-1}b\pi^{r/2}(1+a^{-1}b\pi^{r/2})^{-1} - \sigma(a^{-1}b\pi^{r/2})^2\pi^d(1+a^{-1}b\pi^{r/2})^{-2})$$

is in $(1 + p^{e-[e/2-r/4]})\dot{F}^2$. Therefore

$$\theta(O^+(L)) \subseteq (1 + p^{e-[e/2-r/4]})\dot{F}^2 \cup \varepsilon(1 + p^{e-[e/2-r/4]})\dot{F}^2.$$

Conversely, take an element $h = 1 + c\pi^{e-[e/2-r/4]}$ in $(1+p^{e-[e/2-r/4]})\dot{F}^2$, and write $c = \lambda\pi^k$ with $\lambda \in U$, $k \geq 0$. If $0 \leq k \leq [e/2 - r/4]$, there exists η in U such that

$$\sigma\pi^{d-e-[e/2-r/4]+k}\eta^2 + 2\pi^{-e}\eta + \lambda = 0$$

by Hensel's lemma. Put $z = (-1 + \eta^{-1}\pi^{[e/2-r/4]-k})\pi^{r/2}x + y$ when $-1 + \eta^{-1}\pi^{[e/2-r/4]-k}$ is in U , note

$$\text{ord}(-\sigma\pi^d) = d > e + [e/2 - r/4] > 2[e/2 - r/4] \geq 2[e/2 - r/4] - 2k$$

and

$$\text{ord}(-2\eta^{-1}\pi^{[e/2-r/4]-k}) = e + [e/2 - r/4] - k > 2[e/2 - r/4] - 2k$$

So

$$\begin{aligned} & \text{ord}((-1 + \eta^{-1}\pi^{[e/2-r/4]-k})^2 + \varepsilon) \\ &= \text{ord}(1 + \eta^{-2}\pi^{2[e/2-r/4]-2k} - 2\eta^{-1}\pi^{[e/2-r/4]-k} - 1 - \sigma\pi^d) \\ &= \text{ord}(\eta^{-2}\pi^{2[e/2-r/4]-2k} - 2\eta^{-1}\pi^{[e/2-r/4]-k} - \sigma\pi^d) \\ &= \text{ord}(\eta^{-2}\pi^{2[e/2-r/4]-2k}) \\ &= 2[e/2 - r/4] - 2k \leq 2[e/2 - r/4] \leq e - r/2. \end{aligned}$$

Thus, τ_z is in $O(L)$, and

$$\begin{aligned} Q(z) &= \pi^r((-1 + \eta^{-1}\pi^{[e/2-r/4]-k})^2 + \varepsilon) \\ &= \pi^r(\eta^{-2}\pi^{2[e/2-r/4]-2k} - 2\eta^{-1}\pi^{[e/2-r/4]-k} - \sigma\pi^d) \\ &= \pi^{r+2[e/2-r/4]-2k}\eta^{-2}(1 - 2\eta\pi^{k-[e/2-r/4]} - \sigma\eta^2\pi^{d+2k-[e/2-r/4]}) \\ &= \eta^{-2}\pi^{r+2[e/2-r/4]-2k} \\ &\quad \times (1 + \pi^{k-[e/2-r/4]+e}(-2\pi^{-e}\eta - \sigma\eta^2\pi^{d+k-[e/2-r/4]-e})) \\ &= \eta^{-2}\pi^{r+2[e/2-r/4]-2k}(1 + \lambda\pi^{k-[e/2-r/4]+e}) \\ &= \eta^{-2}\pi^{r+2[e/2-r/4]-2k}(1 + c\pi^{e-[e/2-r/4]}) \\ &= \eta^{-2}\pi^{r+2[e/2-r/4]-2k}h. \end{aligned}$$

If $k > [e/2 - r/4]$, there exists η in U such that

$$\begin{aligned} & \lambda\eta^2 + (2\pi^{-e} + 2\lambda\pi^{k-[e/2-r/4]})\eta + \lambda\pi^{2k-2[e/2-r/4]} \\ & \quad + 2\pi^{-e+k-[e/2-r/4]} + \sigma\pi^{d-e+k-[e/2-r/4]} = 0 \end{aligned}$$

by Hensel’s lemma. Put $z = \eta\pi^{r/2+[e/2-r/4]-k}x + y$, note $r/2 + [e/2 - r/4] - k < r/2$, so τ_z is in $O(L)$, and

$$\begin{aligned} Q(z) &= \eta^2\pi^{r+2[e/2-r/4]-2k} + \varepsilon\pi^r \\ &= \pi^{r+2[e/2-r/4]-2k}(\eta^2 - \pi^{2k-2[e/2-r/4]} - \sigma\pi^{d-2[e/2-r/4]+2k}) \\ &= \pi^{r+2[e/2-r/4]-2k}((\eta + \pi^{k-[e/2-r/4]})^2 + \pi^{e+k-[e/2-r/4]} \\ &\quad \times (-2\eta\pi^{-e}2\pi^{-e+k-[e/2-r/4]} - \sigma\pi^{d-e+k-[e/2-r/4]})) \\ &= \pi^{r+2[e/2-r/4]-2k}((\eta + \pi^{k-[e/2-r/4]})^2 + \pi^{e+k-[e/2-r/4]} \\ &\quad \times (\lambda\eta^2 + 2\lambda\pi^{k-[e/2-r/4]}\eta + \lambda\pi^{2k-2[e/2-r/4]})) \\ &= \pi^{r+2[e/2-r/4]-2k}(\eta + \pi^{k-[e/2-r/4]})^2(1 + \lambda\pi^k\pi^{e-[e/2-r/4]}) \\ &= \pi^{r+2[e/2-r/4]-2k}(\eta + \pi^{k-[e/2-r/4]})^2h. \end{aligned}$$

Therefore

$$\theta(O^+(L)) = (1 + p^{e-[e/2-r/4]})\dot{F}^2 \cup \varepsilon(1 + p^{e-[e/2-r/4]})\dot{F}^2$$

REMARK 1. Suppose two units ε, η with $D(\varepsilon) = p^s$ and $D(\eta) = p^t$, where s, t are odd integers, and $s + t > 2e$; then the Hilbert symbol $(\varepsilon, \eta)_p = 1$. This fact can be proved by an argument similar to that of the case (iii) of Proposition 2.2 or [5, Lemma 4]. On the other hand, consider a unit ε with $D(\varepsilon) = p^s$ where s is an odd integer. For any positive odd integer t , if $s = t \leq 2e$, then there exists a unit η with $D(\eta) = p^t$ and the Hilbert symbol $(\varepsilon, \eta)_p = -1$. This result can be obtained from [5, Lemma 3].

REMARK 2. Suppose E is a finite extension of F and \bar{L} is the lifting of L to E , then we can check $N_{E/F}(\theta(O^+(\bar{L}))) \subseteq \theta(O^+(L))$ by the results obtained in [5], [10] and this section. These can give an alternative proof about spinor genus extension of binary lattices (see [2], [4]).

3. Main results. In this section let L be a lattice with $\dim FL \geq 3$ and Jordan splitting $L = \vartheta x_1 \perp \vartheta x_2 \perp \dots \perp \vartheta x_n$ where $Q(x_1) = 1$, $Q(x_i) = \varepsilon_i\pi^{r_i}$, $r_i \in \mathbb{Z}$, $\varepsilon_i \in U$, $i = 2, 3, \dots, n$; and $0 < r_i < r_{i+1}$, $i = 2, 3, \dots, n - 1$. We assume $r_1 = 0$, $\varepsilon_1 = 1$. First we generalize [3, Th. 2.2] to an arbitrary dyadic local field.

THEOREM 3.1. *Suppose there is at least one k with $1 \leq k \leq n-1$ for which $r_{k+1} - r_k \leq 2e + 1$ and $r_{k+1} - r_k$ is odd. Then if $0 < r_s - r_t \leq 4e$ and $r_s - r_t$ is even for any $s, t = 1, \dots, n$, we have $\theta(O^+(L)) = \dot{F}$.*

Proof. Put $L_{k+1,k} = \vartheta x_k \perp \vartheta x_{k+1}$. When $r_{k+1} - r_k = 2e + 1$, then

$$\begin{aligned} \theta(O^+(L)) &= ((1+p)\dot{F}^2 \cup \varepsilon\pi(1+p)\dot{F}^2) \cap Q([1, \varepsilon_k \dot{\varepsilon}_{k+1}\pi]) \\ &= (U\dot{F}^2 \cup \pi U\dot{F}^2) \cap Q([1, \varepsilon_k \dot{\varepsilon}_{k+1}\pi]) = Q([1, \varepsilon_k \dot{\varepsilon}_{k+1}\pi]) \end{aligned}$$

by case (i) of Proposition 2.2. When $r_{k+1} - r_k < 2e$, we also obtain $\theta(O^+(L_{k+1,k})) = Q([1, \varepsilon_k \dot{\varepsilon}_{k+1}\pi])$ by case (i) of Proposition 2.3. Note $Q([1, \varepsilon_k \dot{\varepsilon}_{k+1}\pi])$ is a subgroup of \dot{F} with index 2 which does not contain Δ .

Put $L_{s,t} = \vartheta x_s \perp \vartheta x_t$. By cases (ii), (iii) of Proposition 2.2 and Proposition 2.3, we check Δ in $\theta(O^+(L_{s,t}))$. Hence $\theta(O^+(L)) = \dot{F}$.

Let $S(L)$ be the group generated by symmetries in $O(L)$ (see [7]). We need the following lemma.

LEMMA 3.2. *If $r_{i+1} - r_i \geq e$, for all $i = 1, 2, \dots, n - 1$, with at most one exception, then $O(L) = S(L)$.*

Proof. First we assume $r_2 \geq e$, otherwise we scale the dual lattice of L by $\varepsilon_n \pi^{r_n}$ and obtain the lattice L' which satisfies this assumption, and we know $O(L) = O(L')$.

Take σ in $O(L)$. Suppose $\sigma x_1 = \sum_{i=1}^n a_i x_i$, $a_i \in \vartheta$. So

$$1 = Q(\sigma x_1) = Q\left(\sum_{i=1}^n a_i x_i\right) = \sum_{i=1}^n \varepsilon_i \pi^{r_i} a_i^2.$$

If $\text{ord}(a_1 - 1) \leq e$, then

$$\text{ord } Q(\sigma x_1 - x_1) = \text{ord} \left((a_1 - 1)^2 + \sum_{i=2}^n a_i^2 \varepsilon_i \pi^{r_i} \right) = \text{ord } 2(a_1 - 1)$$

and

$$2B(\sigma x_1 - x_1, L)\vartheta = 2(a_1 - 1)\vartheta = Q(\sigma x_1 - x_1)\vartheta.$$

Thus, $\tau_{\sigma x_1 - x_1}$ is in $O(L)$.

If $\text{ord}(a_1 - 1) > e$, then $\text{ord}(a_1 + 1) = \text{ord}(2 + (a_1 - 1)) = e$, and

$$\text{ord } Q(\sigma x_1 + x_1) = \text{ord} \left((a_1 + 1)^2 + \sum_{i=2}^n a_i^2 \varepsilon_i \pi^{r_i} \right) = \text{ord } 2(a_1 + 1)$$

and

$$2B(\sigma x_1 + x_1, L)\vartheta = 2(a_1 + 1)\vartheta = Q(\sigma x_1 + x_1)\vartheta.$$

Thus, $\tau_{\sigma x_1 + x_1}$ is in $O(L)$.

Since $\tau_{\sigma x_1 - x_i} \sigma$ and $\tau_{x_1} \tau_{\sigma x_1 + x_i} \sigma$ can be regarded as elements in $O(K)$ where $K = \vartheta x_2 \perp \cdots \perp \vartheta x_n$, by induction on n , we obtain $O(L) = S(L)$.

Now we deduce the Kneser-type result about spinor norms.

THEOREM 3.3. *If $r_{i+1} - r_i > 4e$, for all $i = 1, 2, \dots, n - 1$; then*

$$\theta(O^+(L)) = \bigcup_{k \text{ even}} \varepsilon_{i_1} \cdots \varepsilon_{i_k} \pi^{r_{i_1} + \cdots + r_{i_k}} \dot{F}^2.$$

Proof. Take a symmetry τ_z in $O(L)$, where z is a maximal anisotropic vector in L . Then $2B(z, L)\vartheta \subseteq Q(z)\vartheta$. Write $z = \sum_{i=1}^n a_i x_i$ with $a_i \in \vartheta$, so

$$\text{ord } Q(z) = \text{ord} \left(\sum_{i=1}^n a_i^2 \varepsilon_i \pi^{r_i} \right) \leq \min_{1 \leq i \leq n} \{r_i + \text{ord } a_i\} + e.$$

Choose k so that $\text{ord}(a_k^2 \varepsilon_k \pi^{r_k}) = \min_{1 \leq i \leq n} \{\text{ord}(a_i^2 \varepsilon_i \pi^{r_i})\}$. Then

$$2 \text{ord } a_k + r_k \leq \text{ord } Q(z) \leq \min_{1 \leq i \leq n} \{r_i + \text{ord } a_i\} + e \leq r_j + \text{ord } a_j + e$$

for $j = 1, 2, \dots, n$. When we take $j = k$, $\text{ord } a_k \leq e$ is obtained.

When $j < k$, we have

$$\begin{aligned} \text{ord}((\varepsilon_j \varepsilon_k^{-1})(a_j a_k^{-1})^2 \pi^{r_j - r_k}) &= 2 \text{ord } a_j - 2 \text{ord } a_k + r_j - r_k \\ &= 2(\text{ord } a_j - 2 \text{ord } a_k - r_k + r_j + e) + 2 \text{ord } a_k + (r_k - r_j) - 2e \\ &\geq (r_k - r_j) - 2e > 2e. \end{aligned}$$

When $j > k$, consider

$$\begin{aligned} \text{ord}((\varepsilon_j \varepsilon_k^{-1})(a_j a_k^{-1})^2 \pi^{r_j - r_k}) &= 2 \text{ord } a_j - 2 \text{ord } a_k + r_j - r_k \\ &\geq r_j - r_k - 2 \text{ord } a_k \geq r_j - r_k - 2e > 2e. \end{aligned}$$

So

$$Q(z) = \varepsilon_k \pi^{r_k} a_k^2 \left(1 + \sum_{i=1}^{k-1} (\varepsilon_k^{-1} \varepsilon_i)(a_k^{-1} a_i)^2 \pi^{r_i - r_k} + \sum_{i=k+1}^n (\varepsilon_k^{-1} \varepsilon_i)(a_k^{-1} a_i)^2 \pi^{r_i - r_k} \right)$$

is in $\varepsilon_k \pi^{r_k} \dot{F}^2$ by the Local Square Theorem. By Lemma 3.2, we obtain $\theta(O^+(L)) = \bigcup_{k \text{ even}} \varepsilon_{i_1} \cdots \varepsilon_{i_k} \pi^{r_{i_1} + \cdots + r_{i_k}} \dot{F}^2$.

4. Application. In this section we will show that the restriction in [1], that F be a ramified quadratic extension of Q_2 , is unnecessary. In fact we have the following theorem.

THEOREM 4.1. *Suppose ϑ is the ring of integers of dyadic local field F with $e = \text{ord } 2 = 2$. Let L be a regular ϑ -lattice with $sL \subseteq \vartheta$ and $\text{rank } L \geq 3$. If $\text{ord}(dL) \leq 3$ then $\theta(O^+(L)) \supseteq U\dot{F}^2$.*

Proof. Let $L = L_1 \perp \dots \perp L_t$ be a Jordan splitting of L . We can assume $t \geq 2$ and $\dim FL_i \leq 2$, $i = 1, \dots, t$, by [8, 93:20], and $2sL_i \subset nL_i$ by [5, Lemma 1], $i = 1, \dots, t$. Since $\text{ord}(dL) \leq 3$, $t \leq 3$.

(1) $L = L_1 \perp L_2$ with $\text{rank } L_1 = \text{rank } L_2 = 2$, so L_1 is unimodular and $sL_2 = p$. By [8, 93:17] we may write $L_i = \vartheta x_i + \vartheta y_i$ with $Q(x_i) = a_i$, $Q(y_i) = -\delta_i a_i^{-1}$, $B(x_i, y_i) = \pi^{i-1}$, and $a_i \vartheta = nL_i$, $\delta_i \vartheta = D(-dL_i)$, $i = 1, 2$.

If $nL_1 = psL_1 = p$, we only need to consider $D(-dL_1) = p^3$ by [5, Prop. C], and $\theta(O^+(L_1))$ contains Δ in this case by [5, Prop. B].

Suppose $nL_2 = psL_2 = p^2$. Put $K = \vartheta x_1 \perp \vartheta x_2$. Consider any maximal vector of K , say z , which is then also a maximal anisotropic vector of L . We can check that $2B(z, L)\vartheta \subseteq Q(z)\vartheta$, so τ_z is in $O(L)$, and $\theta(O^+(L)) \supseteq Q([1, \dot{a}_1 a_2])\dot{F}^2$ which is a subgroup of \dot{F} with index 2 and does not contain Δ . Therefore $\theta(O^+(L)) = \dot{F}$.

Suppose $nL_2 = sL_2 = p$. Put $K = \vartheta y_1 \perp \vartheta x_2$, by taking the same argument as above, we obtain $\theta(O^+(L)) = \dot{F}$ again.

If $nL_1 = sL_1 = \vartheta$, then $\theta(O^+(L_1))$ contains Δ by the results obtained in [5] and [10].

Suppose $nL_2 = psL_2 = p^2$. By [5, Prop. C], we only need to treat the case of $D(-dL_2) = p^5$. Put $K = \vartheta x_1 \perp \vartheta y_2$. Note any maximal vector of K , say z , which is also a maximal anisotropic vector of L , satisfies $2B(z, L)\vartheta \subseteq Q(z)\vartheta$. So τ_z is in $O(L)$, $\theta(O^+(L)) \supseteq Q([1, \dot{a}_1 b_2])$ which does not contain Δ . Therefore $\theta(O^+(L)) = \dot{F}$.

Suppose $nL_2 = sL_2 = p$. Put $K = \vartheta x_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$.

(2) $L = L_1 \perp L_2$ with $\text{rank } L_1 = 2$, $\text{rank } L_2 = 1$, so L_1 is unimodular.

Write $L_1 = \vartheta x_1 + \vartheta y_1$ with $Q(x_1) = a_1$, $Q(y_1) = -\delta_1 a_1^{-1}$, $B(x_1, y_1) = 1$, and $nL_1 = a_1 \vartheta$, $D(-dL_1) = \delta_1 \vartheta$ by [8, 93:17]. Let $L_2 = \vartheta x_2$ with $Q(x_2) = a_2 \in p$. If $nL_1 = psL_1 = p$, by [5, Prop. C], we only need to consider $D(-dL_1) = p^3$, and $\theta(O^+(L_1))$ contains Δ in this case by [5, Prop. B].

When $Q(x_2)\vartheta = a_2\vartheta = p$, put $K = \vartheta y_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$. When $Q(x_2)\vartheta = p^2$, put $K = \vartheta x_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$. When $Q(x_2)\vartheta = p^3$, put $K = \vartheta y_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$.

If $nL_1 = sL_1 = \vartheta$, [5, Prop. C], we only need to consider $\delta_1\vartheta = p$ or p^3 , and we know that $\theta(O^+(L_1))$ contains Δ in these two cases by [5, Prop. B,E]. When $Q(x_2)\vartheta = p$, put $K = \vartheta x_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$. When $Q(x_2)\vartheta = p^3$, put $K = \vartheta x_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$. When $Q(x_2)\vartheta = p^2$, and $\delta_1\vartheta = p$, put $K = \vartheta y_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$. When $Q(x_2)\vartheta = a_2\vartheta = p^2$, and $\delta_1\vartheta = p^3$; we know $(1+p^2)\dot{F}^2 \subseteq \theta(O^+(L_1))$ by [5, Prop. E]. Write $a_2 = \sigma\pi^2$ with $\sigma \in U$, $-a_1^{-1}\delta_1 = \eta\pi^3$ with $\eta \in U$. Take an element $1 + \lambda\pi$ in $1 + p$ with $\lambda \in U$.

Suppose $\lambda\sigma\eta^{-1} = \xi^2 + \varepsilon\pi^d$ with $\xi, \varepsilon \in U$ and $d \geq 4$; then there exists t in ϑ such that $a_1t^2 + 2\pi^{-2}\xi t - \eta\varepsilon\pi^{d-1} = 0$ by Hensel's lemma. Put $z = \pi^2tx_1 + \xi y_1 + x_2$, which is a maximal vector of L , and

$$\begin{aligned} Q(z) &= \pi^4t^2a_1 + 2\pi^2\xi t + \xi^2\eta\pi^3 + \sigma\pi^2 \\ &= \pi^4\eta\varepsilon\pi^{d-1} + \xi^2\eta\pi^3 + \sigma\pi^2 = \eta\pi^3(\xi^2 + \varepsilon\pi^d) + \sigma\pi^2 \\ &= \lambda\sigma\pi^3 + \sigma\pi^2 = \sigma\pi^2(1 + \lambda\pi). \end{aligned}$$

So $2B(z, L)\vartheta \subseteq Q(z)\vartheta$, and τ_z is in $O(L)$. Then $(1 + \lambda\pi)$ is in $\theta(O^+(L))$. Suppose $\lambda\sigma\eta^{-1} = \xi^2 + \varepsilon\pi^d$ with $\xi, \varepsilon \in U$ and $1 \leq d \leq 3$, and d is odd. Put $\sigma^{-1}\eta\varepsilon = \omega^2 + \delta\pi^h$ with $\omega, \delta \in U$. Then there exists t in ϑ such that $\sigma^{-1}a_1t^2 + (2\pi^{-2})\sigma^{-1}\xi t - \delta\pi^{d+h-1} + (2\pi^{-2})\omega\pi^{(d+1)/2} = 0$ by Hensel's lemma. Put $z = \pi^2tx_1 + \xi y_1 + (1 + \omega\pi^{(d+1)/2})x_2$ which is a maximal vector of L , and

$$\begin{aligned} Q(z) &= \tau^4t^2a_1 + 2\pi^2t\xi + \xi^2\pi^3\eta + (1 + \omega\pi^{(d+1)/2})^2\sigma\pi^2 \\ &= \sigma\pi^2(\pi^2(\sigma^{-1}a_1t^2 + 2\pi^{-2}\sigma^{-1}\xi t) \\ &\quad + \sigma^{-1}\xi^2\pi\eta + 1 + 2\omega\pi^{(d+1)/2} + \omega^2\pi^{d+1}) \\ &= \sigma\pi^2(\delta\pi^{d+h+1} - 2\omega\pi^{(d+1)/2} + \sigma^{-1}\xi^2\pi\eta + 1 \\ &\quad + 2\omega\pi^{(d+1)/2} + \omega^2\pi^{d+1}) \\ &= \sigma\pi^2(1 + \sigma^{-1}\xi^2\eta\pi + \pi^{d+1}(\omega^2 + \delta\pi^h)) \\ &= \sigma\pi^2(1 + \sigma^{-1}\xi^2\eta\pi + \sigma^{-1}\eta\varepsilon\pi^{d+1}) \\ &= \sigma\pi^2(1 + \sigma^{-1}\eta\pi(\xi^2 + \varepsilon\pi^d)) \\ &= \sigma\pi^2(1 + \sigma^{-1}\eta\pi\lambda\sigma\eta^{-1}) = \sigma\pi^2(1 + \lambda\pi). \end{aligned}$$

So $2B(z, L) \subseteq Q(z)\vartheta$, and τ_z is in $O(L)$, then $(1 + \lambda\pi)$ is in $\theta(O^+(L))$. Therefore, $\theta(O^+(L)) \supseteq (1 + p)\dot{F}^2 = U\dot{F}^2$.

(3) $L = L_1 \perp L_2$ with $\text{rank } L_1 = 1, \text{rank } L_2 = 2$. We scale the dual lattice of L by π and reduce to case (2).

(4) $L = L_1 \perp L_2 \perp L_3$ with $\text{rank } L_1 = \text{rank } L_2 = \text{rank } L_3 = 1$. Let $L_i = \vartheta x_i, i = 1, 2, 3$; so $Q(x_i)\vartheta = p^{i-1}, 1 \leq i \leq 3$. Therefore $\theta(O^+(L)) = \dot{F}$ by Theorem 3.1.

(5) $L = L_1 \perp L_2 \perp L_3$ with $\text{rank } L_1 = 2, \text{rank } L_2 = \text{rank } L_3 = 1$. So L_1 is unimodular, and $L_i = \vartheta x_i$ with $Q(x_i)\vartheta = p^{i-1}, i = 2, 3$. Since Δ is in $\theta(O^+(L_1))$ by the results obtained in [5] and [10], put $K = \vartheta x_2 \perp \vartheta x_3$, obtain $\theta(O^+(L)) = \dot{F}$.

LEMMA 4.2. *Suppose ϑ is the ring of integers of a dyadic local field with $e = \text{ord } 2 = 2$. Let L be a regular ϑ -lattice with $sL \subseteq \vartheta$ and $\text{rank } L \geq 4$. If $\text{ord}(dL) \leq 7$, then $\theta(O^+(L)) \supseteq U\dot{F}^2$.*

Proof. Using the above theorem, considering components and dual lattices when necessary, there remain two cases to be treated.

(1) $L = L_1 \perp L_2$ where L_1 is binary unimodular, L_2 is binary p^2 -modular. Write $L_1 = \vartheta x_1 \perp \vartheta y_1$ with $Q(x_1) = a_1, B(x_1, y_1) = 1, Q(y_1) = -\delta_1 a^{-1}$, and $nL_1 = a_1\vartheta, D(-dL_1) = \delta_1\vartheta$; and $L_2 = \vartheta x_2 + \vartheta y_2$ with $Q(x_2) = a_2, B(x_2, y_2) = \pi^2, Q(y_2) = -\delta_2 a_2^{-1}$, and $nL_2 = a_2\vartheta. D(-dL_2) = \delta_2\vartheta$ by [8, 93:17].

We know that L_i has an orthogonal base if and only if $nL_i = sL_i (i = 1, 2)$. Using the above theorem, considering components and dual lattices when necessary, we assume $nL_1 = p(sL_1) = p$, and $nL_2 = p(sL_2) = p^3$. By [5, Prop. C], we only need to consider $D(-dL_1) = p^3$, and Δ is in $\theta(O^+(L_1))$ by [5, Prop. B] in this case. Put $K = \vartheta y_1 \perp \vartheta x_2$. Then $\theta(O^+(L)) = \dot{F}$.

(2) $L = L_1 \perp L_2$ where L_1 is binary unimodular and L_2 is binary p^3 -modular. We know L_i has an orthogonal base if and only if $nL_i = sL_i (i = 1, 2)$. Using the above theorem, considering components and dual lattices when necessary, we assume $nL_1 = p(sL_1) = p$ and $nL_2 = psL_2 = p^4$. Write $L_1 = \vartheta x_1 + \vartheta y_1$ with $Q(x_1)\vartheta = nL_1 = p$, and $L_2 = \vartheta x_2 + \vartheta y_2$ with $Q(x_2)\vartheta = nL_2 = p^4$, note Δ is in $\theta(O^+(L_1))$ by [5] and [10], put $K = \vartheta x_1 \perp \vartheta x_2$, then $\theta(O^+(L)) = \dot{F}$.

By using Theorem 4.1, Lemma 4.2 and Theorem 3.1, we can also prove [1, Lemma 3.4] and [1, Theorem 3.5] when ϑ is the ring of integers of a dyadic local field with $e = \text{ord } 2 = 2$. Finally we consider an example.

EXAMPLE 4.3. The bound $\text{ord}(dL) \leq 7$ given in Lemma 4.2 cannot be unconditionally improved. Consider the lattice $L = L_1 \perp L_2$ over

the ring of integers of a ramified quadratic extension of Q_2 , where $L_1 = \vartheta x_1 + \vartheta y_1$ with $Q(x_1) = 1$, $Q(y_1) = \pi^3$, $B(x_1, y_1) = 1$, and $L_2 = \vartheta x_2 + \vartheta y_2$ with $Q(x_2) = \pi^4$, $Q(y_2) = \pi^7$, $B(x_2, y_2) = \pi^4$. We claim $\theta(O^+(L)) = (1 + p^2)\dot{F}^2 \subset U\dot{F}^2$.

First we prove that $O(L)$ is generated by symmetries of L . Note L_i has an orthogonal base ($i = 1, 2$), and write $L_1 = \vartheta z \perp \vartheta w$. Put $K = \vartheta w \perp L_2$. Then $O(K)$ is generated by symmetries of K by the proof of Lemma 3.2. Let $\sigma \in O(L)$, and $\sigma z = az + bw + u$ with $u \in L_2$, so $Q(z) = a^2Q(z) + b^2Q(w) + Q(u)$ and $Q(\sigma z - z) = 2(1 - a)Q(z)$, $Q(\sigma z + z) = 2(1 + a)Q(z)$.

If $\text{ord}(1 - a) = 0$, then $2B(\sigma z - z, L) = Q(\sigma z - z)\vartheta$, and $\tau_{\sigma z - z} \in O(L)$. If $\text{ord}(1 - a) = 1$, then $b^2Q(w) = (1 - a)(1 + a)Q(z) - Q(u) \in p$, and $\text{ord } b \geq 1$. So $2B(\sigma z - z, L) = Q(\sigma z - z)\vartheta$, and $\tau_{\sigma z - z} \in O(L)$.

If $\text{ord}(1 - a) = 2$, then $\text{ord}(1 + a) = \text{ord}(2 - (1 - a)) > 2$, and $b^2Q(w) = (1 + a)(1 - a)Q(z) - Q(u) \in p^4$, so $\text{ord } b \geq 2$, and $2B(\sigma z - z, L) = Q(\sigma z - z)\vartheta$. Thus $\tau_{\sigma z - z} \in O(L)$.

If $\text{ord}(1 - a) > 2$, then $\text{ord}(1 + a) = \text{ord}(2 - (1 - a)) = 2$, and $b^2Q(w) = (1 + a)(1 - a)Q(z) - Q(u) \in p^4$, so $\text{ord } b \geq 2$, and $2B(\sigma z + z, L) \subseteq Q(\sigma z + z)\vartheta$. Thus $\tau_{\sigma z + z} \in O(L)$.

Note that $\tau_{\sigma z - z}\sigma$ and $\tau_z\tau_{\sigma z + z}\sigma$ can be regarded as elements in $O(K)$ which is generated by symmetries of K . Therefore $O(L)$ is generated by symmetries of L .

Now we calculate $\theta(O^+(L))$. Let $\tau_t \in O(L)$ where t is a maximal anisotropic vector of L , write $t = cx_1 + dy_1 + fx_2 + gy_2$ with $c, d, f, g \in \vartheta$.

When $\text{ord } c \leq 1$, then $2c^{-1}d + c^{-2}d^2\pi^3$ is always in p^2 whenever c and d are units or not. So

$$Q(t) = c^2(1 + (2c^{-1}d + c^{-2}d^2\pi^3) + c^{-2}\pi^4(f^2 + 2fg + g^2\pi^3))$$

is in $(1 + p^2)\dot{F}^2$.

When $\text{ord}(d) = 0$, and $\text{ord}(c) > 1$; then $\text{ord } Q(t) = 3$, and $2B(t, L) = p^2 \supset Q(t)\vartheta$, contradicting the assumption that τ_t is in $O(L)$.

When $\text{ord}(f) = 0$, $\text{ord}(d) > 0$, and $\text{ord}(c) > 1$; then $\text{ord}(c) \geq 2$, and the inclusion $2B(t, L) \subseteq Q(t)\vartheta$ forces $\text{ord}(d) \geq 2$.

Suppose $\text{ord}(c) = 2$. Then $\text{ord}(c^2 + f^2\pi^4) \geq 5$. So

$$\text{ord } Q(t) = \text{ord}((c^2 + f^2\pi^4) + 2cd + d^2\pi^3 + 2\pi^4fg + g^2\pi^7) \geq 5$$

and $2B(t, L) = p^4 \supset Q(t)\vartheta$, a contradiction.

Therefore $\text{ord}(c) \geq 3$, and

$$Q(t) = f^2\pi^4(1 + f^{-2}c^2\pi^{-4} + 2f^{-2}cd\pi^{-4} + d^2f^{-2}\pi^{-1} + 2f^{-1}g + f^{-2}g^2\pi^3)$$

is in $(1 + p^2)\dot{F}^2$.

When $\text{ord}(g) = 0$, $\text{ord}(f) > 0$, $\text{ord}(d) > 0$, and $\text{ord}(c) > 1$; then $\text{ord}(c) \geq 2$, and $\text{ord}(d) \geq 2$.

Suppose $\text{ord}(c) = 2$. Then

$$Q(t) = c^2(1 + 2c^{-1}d + (c^{-1}d)^2\pi^3 + c^{-2}\pi^4(f^2 + 2fg + g^2\pi^3))$$

is in $(1 + p^2)\dot{F}^2$.

Suppose $\text{ord}(c) \geq 3$. Then $\text{ord}(c) \geq 4$ and $\text{ord}(d) \geq 4$.

If $\text{ord}(f) = 1$, then $2f^{-1}g + (f^{-1}g)^2\pi^3 \in p^2$, and

$$Q(t) = f^2\pi^4(1 + (f^{-1}c\pi^{-2})^2 + 2cdf^{-2}\pi^{-4} + d^2f^{-2}\pi^{-1} + (2f^{-1}g + (f^{-1}g)^2\pi^3))$$

is in $(1 + p^2)\dot{F}^2$.

If $\text{ord}(f) \geq 2$, then $\text{ord } Q(t) = 7$, and $2B(t, L) = p^6 \supset Q(t)\vartheta$. But τ_t is in $O(L)$ and t is a maximal vector, a contradiction. In any case, we have $\theta(O^+(L)) \subseteq (1 + p^2)\dot{F}^2$. On the other hand, $\theta(O^+(L)) \supseteq \theta(O^+(L_1)) \supseteq (1 + p^2)\dot{F}^2$ by [5, Prop. E]. Thus, we conclude $\theta(O^+(L)) = (1 + p^2)\dot{F}^2$.

Acknowledgment. I wish to thank Professor Zhu Fu-zu for introducing me to the theory of quadratic forms and giving me much help in this field.

REFERENCES

[1] W. R. Bon Durant, *Spinor norms of rotations of local integral quadratic forms*, J. Number Theory, **33** (1989), 83–94.
 [2] A. G. Earnest and D. R. Estes, *Glass groups in the genus and spinor genus of binary quadratic lattices*, Proc. London Math. Soc., **40** (1980), 40–52.
 [3] A. G. Earnest and J. S. Hsia, *Spinor norms of local integral rotations*, II, Pacific J. Math., **61**, No. 1 (1975), 71–86; errata **115** (1984), 493–494.
 [4] —, *Spinor genera under field extensions II: 2 unramified in the bottom field*, Amer. J. Math., **100** (1978), 523–538.
 [5] J. S. Hsia, *Spinor norms of local integral rotations*, I, Pacific J. Math., **57** (1975), 199–206.
 [6] M. Kneser, *Klassenzahlen indefiniter quadratischer Formen in drei oder mehr Veränderlichen.*, Arch. Math., **7** (1956), 323–332.
 [7] O. T. O’Meara and Barth Pollak, *Generation of local integral orthogonal groups*, II, Math. Z., **93** (1966), 171–188.
 [8] O. T. O’Meara, *Introduction to Quadratic Forms*, Springer, New York, 1963.

- [9] C. Riehm, *On the integral representations of quadratic forms over local fields*, Amer. J. Math., **86** (1964), 25–62.
- [10] Fei Xu, *A remark on spinor norms of local integral rotations, I*, Pacific J. Math., **136**, No. 1 (1989), 81–84.

Received April 13, 1990 and in revised form October 16, 1991.

NANKAI INSTITUTE OF MATHEMATICS
WEIJIN ROAD 94, TIANJIN 300071
PEOPLE'S REPUBLIC OF CHINA

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982) F. WOLF (1904–1989)

EDITORS

V. S. VARADARAJAN
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

C. C. MOORE
University of California
Berkeley, CA 94720

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@henri.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

PACIFIC JOURNAL OF MATHEMATICS

Volume 157 No. 1 January 1993

Permutation enumeration symmetric functions, and unimodality	1
FRANCESCO BRENTI	
On the analytic reflection of a minimal surface	29
JAIGYOUNG CHOE	
Contractive zero-divisors in Bergman spaces	37
PETER LARKIN DUREN, DMITRY KHAVINSON, HAROLD SEYMOUR SHAPIRO and CARL SUNDBERG	
On the ideal structure of positive, eventually compact linear operators on Banach lattices	57
RUEY-JEN JANG and HAROLD DEAN VICTORY, JR.	
A note on the set of periods for Klein bottle maps	87
JAUME LLIBRE	
Asymptotic expansion at a corner for the capillary problem: the singular case	95
ERICH MIERSEMANN	
A state model for the multivariable Alexander polynomial	109
JUN MURAKAMI	
Free Banach-Lie algebras, couniversal Banach-Lie groups, and more	137
VLADIMIR G. PESTOV	
Four manifold topology and groups of polynomial growth	145
RICHARD ANDREW STONG	
A remark on Leray's inequality	151
AKIRA TAKESHITA	
A_∞ and the Green function	159
JANG-MEI GLORIA WU	
Integral spinor norms in dyadic local fields. I	179
FEI XU	