

*Pacific  
Journal of  
Mathematics*

**DETERMINANT IDENTITIES**

GEORGE W. EYRE ANDREWS AND WILLIAM H. BURGE

## DETERMINANT IDENTITIES

GEORGE E. ANDREWS AND WILLIAM H. BURGE

**A number of determinants are evaluated in closed form including**

$$\det \left( \left( \begin{matrix} i+j+x \\ 2i-j \end{matrix} \right) + \left( \begin{matrix} i+j+y \\ 2i-j \end{matrix} \right) \right)_{0 \leq i, j \leq n-1}.$$

**1. Introduction.** In one of their series of papers on plane partitions and related questions, Mills, Robbins and Rumsey [9; p. 53] prove the following determinant formula.

$$(1.1) \quad m_n(x) = \det \left( \left( \begin{matrix} i+j+x \\ 2i-j \end{matrix} \right) \right)_{0 \leq i, j \leq n-1} = \frac{1}{2^n} \prod_{k=0}^{n-1} \Delta_{2k}(2x),$$

where  $\Delta_0(u) = 2$  and for  $j > 0$

$$(1.2) \quad \Delta_{2j}(u) = \frac{(u+2j+2)_j (\frac{1}{2}u+2j+\frac{3}{2})_{j-1}}{(j)_j (\frac{1}{2}u+j+\frac{3}{2})_{j-1}}$$

with

$$(1.3) \quad (A)_j = A \cdot (A+1) \cdots (A+j-1).$$

Our object here is primarily to prove the following generalization of (1.1).

**THEOREM 1.** *Let*

$$(1.4) \quad M_n(x, y) = \det \left( \left( \begin{matrix} i+j+x \\ 2i-j \end{matrix} \right) + \left( \begin{matrix} i+j+y \\ 2i-j \end{matrix} \right) \right)_{0 \leq i, j \leq n-1};$$

$$(1.5) \quad N_n(x, y) = \det \left( \frac{2}{x+1-y} \left\{ \begin{matrix} i+j+x+1 \\ 2i-j+1 \end{matrix} \right. \right. \\ \left. \left. - \begin{matrix} i+j+y \\ 2i-j+1 \end{matrix} \right\} \right)_{0 \leq i, j \leq n-1}.$$

*Then*

$$(1.6) \quad M_n(x, y) = N_n(x, y) = \prod_{k=0}^{n-1} \Delta_{2k}(x+y).$$

Sections 2 and 3 will be devoted to the proof of this result. In §§4 and 5 we shall show how our work leads to two alternative proofs of the T.S.S.C.P.P. conjecture [2], and we shall mention a related application of Ishikawa [6].

**2. Bailey's balanced  ${}_4F_3$  summation.** In this section, we consider summation formulas for hypergeometric series [4; p. 8], [10, p. 41]:

$$(2.1) \quad {}_{n+1}F_n \left[ \begin{matrix} a_0, a_1, \dots, a_n; t \\ b_1, \dots, b_n \end{matrix} \right] = \sum_{j=0}^{\infty} \frac{(a_0)_j (a_1)_j \cdots (a_n)_j t^n}{j! (b_1)_j \cdots (b_n)_j}.$$

The formula of Bailey [3; p. 512, (c)] [10; p. 245, (III.20)] alluded to in the title of this section is

$$(2.2) \quad {}_4F_3 \left[ \begin{matrix} \frac{a}{2}, \frac{a+1}{2}, b+n, -n; 1 \\ \frac{b}{2}, \frac{b+1}{2}, a+1 \end{matrix} \right] = \frac{(b-a)_n}{(b)_n},$$

and a closely related companion is

$$(2.3) \quad {}_4F_3 \left[ \begin{matrix} \frac{a}{2}, \frac{a+1}{2}, b+n, -n; 1 \\ \frac{b+1}{2}, \frac{b+2}{2}, a \end{matrix} \right] = \frac{(b-a+1)_n}{(b+1)_{n-1} (b+2n)}.$$

Also useful for our work is a transformation due to F.J.W. Whipple [14; p. 537, eq. (10.1)]. If one of  $z$  and  $n$  is a nonnegative integer, then

$$(2.4) \quad {}_4F_3 \left[ \begin{matrix} a, b, -z, -n; 1 \\ u, v, w \end{matrix} \right] \\ = \frac{\Gamma(v+z+n)\Gamma(w+z+n)\Gamma(v)\Gamma(w)}{\Gamma(v+z)\Gamma(v+n)\Gamma(w+n)\Gamma(w+z)} \\ \times {}_4F_3 \left[ \begin{matrix} u-a, u-b, -z, -n; 1 \\ 1-v-z-n, 1-w-z-n, u \end{matrix} \right].$$

We note in passing that (2.4) specialized to  $a = \alpha/2$ ,  $b = (\alpha+1)/2$ ,  $z = -\beta - \nu$ ,  $n = \nu$ ,  $v = (\beta+1)/2$ ,  $w = (\beta+2)/2$ ,  $u = \alpha$  yields

$$(2.5) \quad {}_4F_3 \left[ \begin{matrix} \frac{\alpha}{2}, \frac{\alpha+1}{2}, \beta + \nu, -\nu; 1 \\ \frac{\beta+1}{2}, \frac{\beta+2}{2}, \alpha \end{matrix} \right] \\ = \frac{\beta}{\beta+2\nu} {}_4F_3 \left[ \begin{matrix} \frac{\alpha-1}{2}, \frac{\alpha}{2}, \beta + \nu, -\nu; 1 \\ \frac{\beta+1}{2}, \frac{\beta}{2}, 1 + (\alpha-1) \end{matrix} \right] \\ = \frac{\beta}{\beta+2\nu} \frac{(\beta-\alpha+1)_\nu}{(\beta)_\nu} \quad (\text{by (2.2)}).$$

Thus (2.3) is an immediate consequence of (2.2) and (2.4).

**LEMMA 1.** *Let  $\alpha$  be a positive integer, then*

$$(2.6) \quad {}_4F_3 \left[ \begin{matrix} -\frac{\alpha}{2}, -\frac{\alpha-1}{2}, b+z, -z; 1 \\ \frac{b}{2}, \frac{b+1}{2}, 1-\alpha \end{matrix} \right] = \frac{(b+z)_\alpha}{(b)_\alpha} + \frac{(-z)_\alpha}{(b)_\alpha}.$$

*Proof.* We begin by considering (2.2) when  $a = -\alpha$  a negative integer. The index of summation  $j$  runs from 0 to  $n$  (when  $j > n$  all terms = 0). Furthermore, the terms with  $\alpha/2 < j < \alpha$  are all identically zero. For  $n \geq j \geq \alpha$  there are cancelling zeros in numerator and denominator.

Thus by (2.2)

$$\begin{aligned} \frac{(b+\alpha)_n}{(b)_n} &= \sum_{0 \leq 2j \leq \alpha} \frac{\left(-\frac{\alpha}{2}\right)_j \left(\frac{1-\alpha}{2}\right)_j (b+n)_j (-n)_j}{j! \left(\frac{b}{2}\right)_j \left(\frac{b+1}{2}\right)_j (1-\alpha)_j} \\ &\quad + \sum_{j=\alpha}^n \frac{(-\alpha)(-\alpha+1)\cdots(-1)(1)(2)\cdots(-\alpha+2j-1)(b+n)_j (-n)_j}{j!(b)_{2j}(1-\alpha)(2-\alpha)\cdots(-1)\cdot(1)\cdot(2)\cdots(j-\alpha)} \\ &= S_1 + S_2. \end{aligned}$$

Now

$$\begin{aligned} S_2 &= \sum_{j=0}^{n-\alpha} \frac{(-1)^\alpha \alpha! (\alpha+2j-1)! (b+n)_{j+\alpha} (-n)_{j+\alpha}}{(j+\alpha)! (b)_{2j+2\alpha} (-1)^{\alpha-1} (\alpha-1)! j!} \\ &= -\frac{(b+n)_\alpha (-n)_\alpha}{(b)_{2\alpha}} \sum_{j=0}^{n-\alpha} \frac{(\alpha)_{2j} (b+n+\alpha)_j (-n+\alpha)_j}{j! (b+2\alpha)_{2j} (\alpha+1)_j} \\ &= -\frac{(b+n)_\alpha (-n)_\alpha}{(b)_{2\alpha}} {}_4F_3 \left( \begin{matrix} \frac{\alpha}{2}, \frac{\alpha+1}{2}, b+2\alpha+(n-\alpha), -(n-\alpha); 1 \\ \frac{b+2\alpha}{2}, \frac{b+2\alpha+1}{2}, 1+\alpha \end{matrix} \right) \\ &= -\frac{(b+n)_\alpha (-n)_\alpha}{(b)_{2\alpha}} \frac{(b+\alpha)_{n-\alpha}}{(b+2\alpha)_{n-\alpha}} \quad (\text{by (2.2)}) \\ &= -\frac{(-n)_\alpha}{(b)_\alpha}. \end{aligned}$$

Hence

$$\begin{aligned} (2.7) \quad &\sum_{0 \leq 2j \leq \alpha} \frac{\left(-\frac{\alpha}{2}\right)_j \left(\frac{1-\alpha}{2}\right)_j (b+n)_j (-n)_j}{j! \left(\frac{b}{2}\right)_j \left(\frac{b+1}{2}\right)_j (1-\alpha)_j} \\ &= \frac{(b+\alpha)_n}{(b)_n} + \frac{(-n)_\alpha}{(b)_\alpha} = \frac{(b+n)_\alpha}{(b)_\alpha} + \frac{(-n)_\alpha}{(b)_\alpha}, \end{aligned}$$

which is precisely (2.6) when  $z$  is any positive integer,  $n$ . However, both sides of (2.6) are polynomials in  $z$  of degree at most  $\alpha$ , and since they agree for all positive integral  $z$  we see that (2.6) holds for all real  $z$ .  $\square$

LEMMA 2. *Let  $\alpha$  be a nonnegative integer, then*

$$(2.8) \quad {}_4F_3 \left[ \begin{matrix} -\frac{\alpha}{2}, \frac{-\alpha+1}{2}, b+z, -z; 1 \\ \frac{b+1}{2}, \frac{b+2}{2}, -\alpha \end{matrix} \right] \\ = \frac{(b+z)_{\alpha+1}}{(b+2z)(b+1)_{\alpha}} - \frac{(-z)_{\alpha+1}}{(b+1)_{\alpha}(b+2z)}.$$

*Proof.* In parallel with Lemma 1, we begin by considering (2.3) with  $a = -\alpha$  a negative integer. If  $j$  is the index of summation in the  ${}_4F_3$ , then the nonzero terms of the sum occur for  $0 \leq j \leq \alpha/2$  and  $\alpha < j \leq n$ . If we call the two resulting sums  $T_1$  and  $T_2$ , then

$$(2.9) \quad \frac{(b+\alpha+1)_n}{(b+1)_{n-1}(b+2n)} = T_1 + T_2.$$

Now

$$\begin{aligned} T_2 &= \sum_{j=\alpha+1}^n \frac{(-\alpha)_{\alpha}(1)_{2j-\alpha-1}(b+n)_j(-n)_j}{j!(b+1)_{2j}(-\alpha)_{\alpha}(1)_{j-\alpha-1}} \\ &= \sum_{j=0}^{n-\alpha-1} \frac{(1)_{2j+\alpha+1}(b+n)_{j+\alpha+1}(-n)_{j+\alpha+1}}{(j+\alpha+1)!(b+1)_{2j+2\alpha+2}j!} \\ &= \frac{(\alpha+1)!(b+n)_{\alpha+1}(-n)_{\alpha+1}}{(\alpha+1)!(b+1)_{2\alpha+2}} \\ &\quad \times {}_4F_3 \left[ \begin{matrix} \frac{\alpha}{2}+1, \frac{\alpha}{2}+\frac{3}{2}, b+n+\alpha+1, -n+\alpha+1; 1 \\ \frac{b+2\alpha+3}{2}, \frac{b+2\alpha+4}{2}, \alpha+2 \end{matrix} \right] \\ &= \frac{(b+1)_{\alpha+n}(-n)_{\alpha+1}}{(b+1)_{n-1}(b+1)_{2\alpha+2}} \frac{(b+\alpha+1)_{n-\alpha-1}}{(b+2\alpha+3)_{n-\alpha-2}(b+2n)} \quad (\text{by (2.3)}) \\ &= \frac{(-n)_{\alpha+1}}{(b+1)_{\alpha}(b+2n)}. \end{aligned}$$

Hence

$$(2.10) \quad {}_4F_3 \left[ \begin{matrix} -\frac{\alpha}{2}, \frac{-\alpha+1}{2}, b+n, -n; 1 \\ \frac{b+1}{2}, \frac{b+2}{2}, -\alpha \end{matrix} \right] \\ = T_1 = \frac{(b+\alpha+1)_n}{(b+1)_{n-1}(b+2n)} - \frac{(-n)_{\alpha+1}}{(b+1)_{\alpha}(b+2n)} \\ = \frac{(b+n)_{\alpha+1}}{(b+1)_{\alpha}(b+2n)} - \frac{(-n)_{\alpha+1}}{(b+1)_{\alpha}(b+2n)},$$

which is (2.8) when  $z$  is any positive integer. Since (2.8) is an identity of rational functions in  $z$ , the result in full generality follows immediately.  $\square$

**3. The main theorem.** Our proof of Theorem 1 relies on the following binomial coefficient summations.

**LEMMA 3.** For integers  $i, j \geq 0$

$$(3.1) \quad 2 \sum_{k=0}^i \frac{(y-x)}{(y-x+i-k)} \binom{y-x+i-k}{2i-2k} \binom{k+j+x+y}{2k-j} \\ = \binom{i+j+2x}{2i-j} + \binom{i+j+2y}{2i-j}.$$

$$(3.2) \quad \sum_{k=0}^i \frac{(2x-2y+1)}{(y-x+i-k)} \binom{y-x+i-k}{2i-2k+1} \binom{k+j+x+y}{2k-j} \\ = \binom{i+j+2x+1}{2i-j+1} - \binom{i+j+2y}{2i-j+1}.$$

*Proof.* We note that the only nonzero terms on the left-hand side of (3.1) occur for  $i \geq k \geq j/2$ . Consequently, if  $j > 2i$  then both sides of (3.1) are zero. Hence we may assume  $2i \geq j$ .

Therefore

$$(3.3) \quad 2 \sum_{k=0}^i \frac{(y-x)}{(y-x+i-k)} \binom{y-x+i-k}{2i-2k} \binom{k+j+x+y}{2k-j} \\ = 2 \sum_{k=0}^i \frac{(y-x)}{(y-x+k)} \binom{y-x+k}{2k} \binom{i+j-k+x+y}{2i-j-2k} \\ = 2 \binom{i+j+x+y}{2i-j} \sum_{k=0}^j \frac{(y-x)_k (-y+x)_k (-2i+j)_{2k}}{(2k)! (-i-j-x-y)_k (2j-i+x+y+1)_k} \\ = 2 \binom{i+j+x+y}{2i-j} {}_4F_3 \left[ \begin{matrix} y-x, -y+x, -i+\frac{j}{2}, -i+\frac{i+1}{2}; 1 \end{matrix} \right] \\ = 2 \binom{i+j+x+y}{2i-j} \\ \times \frac{\Gamma(i+j+x+y+1/2)\Gamma(2i-j)\Gamma(2j-i+x+y+1)\Gamma(1/2)}{\Gamma(3j/2+x+y+1)\Gamma(1/2+i-j/2)\Gamma(3j/2+x+y+1/2)\Gamma(i-j/2)} \\ \times {}_4F_3 \left[ \begin{matrix} -i-j-2y, -i-j-2x, -i+\frac{j}{2}, -i+\frac{i+1}{2}; 1 \end{matrix} \right] \quad (\text{by (2.4)}) \\ = \binom{i+j+x+y}{2i-j} \frac{\Gamma(i+j+x+y+1/2)\Gamma(2j-i+x+y+1)2^{2i+2j+2x+2y}}{\Gamma(1/2)\Gamma(3j+2x+2y+1)} \\ \times \left( \frac{(-i-j-2x)_{2i-j}}{(-2i-2j-2x-2y)_{2i-j}} + \frac{(-i-j-2y)_{2i-j}}{(-2i-2j-2x-2y)_{2i-j}} \right) \\ (\text{by Lemma 1 and Gauss's duplication formula [5; p. 5, eq. (15)]}) \\ = \binom{i+j+2x}{2i-j} + \binom{i+j+2y}{2i-j}$$

as desired for (3.1).

For (3.2) again we obtain zero equals zero unless  $2i \geq j$ . Hence

$$\begin{aligned}
 (3.4) \quad & \sum_{k=0}^i \frac{(2x-2y+1)}{(y-x+i-k)} \binom{y-x+i-k}{2i-2k+1} \binom{k+j+x+y}{2k-j} \\
 &= \sum_{k=0}^i \frac{(2x-2y+1)}{(y-x+k)} \binom{y-x+k}{2k+1} \binom{i-k+j+x+y}{2i-j-2k} \\
 &= (2x-2y+1) \binom{i+j+x+y}{2i-j} \\
 &\quad \times {}_4F_3 \left[ \begin{matrix} y-x, 1+x-y, -i+\frac{j}{2}, -i+\frac{j}{2}+\frac{1}{2}; 1 \\ \frac{3}{2}, -i-j-x-y, x+y+2j-i+1 \end{matrix} \right] \\
 &= (2x-2y+1) \binom{i+j+x+y}{2i-j} \\
 &\quad \times \frac{\Gamma(2i-j+1)\Gamma(i+j+x+y+1/2)\Gamma(3/2)\Gamma(2j-i+x+y+1)}{\Gamma(i-\frac{j}{2}+\frac{3}{2})\Gamma(i-\frac{j}{2}+1)\Gamma(\frac{3j}{2}+x+y+1)\Gamma(\frac{3j}{2}+x+y+\frac{1}{2})} \\
 &\quad \times {}_4F_3 \left[ \begin{matrix} -i-j-2y, -i-j-2x-1, -i+\frac{j}{2}, -i+\frac{j}{2}+\frac{1}{2}; 1 \\ -i-j-x-y, -2i+j, -i-j-x-y+\frac{j}{2} \end{matrix} \right] \quad (\text{by (2.4)}) \\
 &= (2x-2y+1) \binom{i+j+x+y}{2i-j} \\
 &\quad \times \frac{\Gamma(i+j+x+y+\frac{1}{2})2^{2i+2j+2x+2y}\Gamma(2j-i+x+y+1)}{\Gamma(\frac{1}{2})\Gamma(3j+2x+2y+1)(2i-j+1)} \\
 &\quad \times \left( \frac{(-i-j-2x-1)_{2i-j+1}}{(2y-2x-1)(-2i-2j-2x-2y)_{2i-j}} \right. \\
 &\quad \quad \left. - \frac{(-i-j-2y)_{2i-j+1}}{(2y-2x-1)(-2i-2j-2x-2y)_{2i-j}} \right) \\
 &= \binom{i+j+2x+1}{2i-j+1} - \binom{i+j+2y}{2i-j+1}. \quad \square
 \end{aligned}$$

*Proof of Theorem 1.* We define five matrices

$$(3.5) \quad \mathbf{M}_n(x) = \left( \binom{i+j+x}{2i-j} \right)_{0 \leq i, j \leq n-1},$$

$$(3.6) \quad \mu_n(x, y) = \left( \binom{i+j+x}{2i-j} + \binom{i+j+y}{2i-j} \right)_{0 \leq i, j \leq n-1},$$

$$(3.7) \quad \nu_n(x, y) = \left( \frac{2}{x+1-y} \left( \binom{i+j+x+1}{2i-j+1} - \binom{i+j+y}{2i-j+1} \right) \right)_{0 \leq i, j \leq n-1},$$

$$(3.8) \quad \tau_n(x, y) = \left( \frac{2(y-x)}{(y-x+i-j)} \binom{y-x+i-j}{2i-2j} \right)_{0 \leq i, j \leq n-1},$$

$$(3.9) \quad \sigma_n(x, y) = \left( \frac{2}{(y-x+i-j)} \binom{y-x+i-j}{2i-2j+1} \right)_{0 \leq i, j \leq n-1}.$$

Clearly

$$(3.10) \quad m_n(x) = \det(\mathbf{M}_n(x)),$$

$$(3.11) \quad M_n(x, y) = \det(\mu_n(x, y)),$$

$$(3.12) \quad N_n(x, y) = \det(\nu_n(x, y)),$$

$$(3.13) \quad \det(\tau_n(x, y)) = 2^n,$$

and

$$(3.14) \quad \det(\sigma_n(x, y)) = 2^n.$$

Equations (3.10)–(3.12) are restatements of (1.1), (1.4) and (1.5), while (3.13) and (3.14) are obvious since each matrix in question is lower triangular.

Now

$$(3.15) \quad \begin{aligned} & \tau_n(x, y) \mathbf{M}_n(x+y) \\ &= \left( \sum_{k=0}^i \frac{2(y-x)}{(y-x+i-k)} \binom{y-x+i-k}{2i-2k} \binom{k+j+x+y}{2k-j} \right)_{0 \leq i, j \leq n-1} \\ &= \left( \binom{i+j+2x}{2i-j} + \binom{i+j+2y}{2i-j} \right)_{0 \leq i, j \leq n-1} \quad (\text{by Lemma 3, eq. (3.1)}) \\ &= \mu_n(2x, 2y). \end{aligned}$$

Hence by (3.15) and (1.1)

$$\begin{aligned} M_n(x, y) &= \det(\mu_n(x, y)) \\ &= \det \left( \tau_n \left( \frac{x}{2}, \frac{y}{2} \right) \mathbf{M}_n \left( \frac{x+y}{2} \right) \right) \\ &= 2^n \cdot \det \left( \mathbf{M}_n \left( \frac{x+y}{2} \right) \right) \\ &= \prod_{k=0}^{n-1} \Delta_{2k}(x+y), \end{aligned}$$

which proves the first part of (1.6).



Similarly

$$\begin{aligned}
 (3.16) \quad & \sigma_n(x, y) \mathbf{M}_n(x+y) \\
 &= \left( \sum_{k=0}^i \frac{2}{(y-x+i-k)} \binom{y-x+i-k}{2i-2k+1} \binom{k+j+x+y}{2k-j} \right)_{0 \leq i, j \leq n-1} \\
 &= \left( \frac{2}{(2x-2y+1)} \left( \binom{i+j+2x+1}{2i-j+1} - \binom{i+j+2y}{2i-j+1} \right) \right)_{0 \leq i, j \leq n-1} \\
 &\quad \text{(by Lemma 3, eq. (3.2))} \\
 &= \nu_n(2x, 2y).
 \end{aligned}$$

Therefore by (3.16) and (1.1)

$$\begin{aligned}
 N_n(x, y) &= \det(\nu_n(x, y)) \\
 &= \det \left( \sigma_n \left( \frac{x}{2}, \frac{y}{2} \right) \mathbf{M}_n \left( \frac{x+y}{2} \right) \right) \\
 &= 2^n \cdot \det \left( \mathbf{M}_n \left( \frac{x+y}{2} \right) \right) \\
 &= \prod_{k=0}^{n-1} \Delta_{2k}(x+y). \quad \square
 \end{aligned}$$

**4. Applications to the T.S.S.C.P.P. conjecture.** We shall consider in this section a few instances of the results we have obtained. We begin with a rather odd determinant for generalized harmonic numbers.

**COROLLARY 1.** *Let  $H_n(x) = \sum_{j=0}^n \frac{1}{x+j}$ , then*

$$\begin{aligned}
 (4.1) \quad & \det \left( \binom{i+j+x}{2i-j+1} (H_{i+j}(x) - H_{2j-i-1}(x)) \right)_{0 \leq i, j \leq n-1} \\
 &= \frac{1}{2^n} \prod_{k=0}^{n-1} \Delta_{2k}(2x-1).
 \end{aligned}$$

*Proof.* From (1.5) and (1.6) with  $x$  replaced by  $x-1$  we find

$$\begin{aligned}
 (4.2) \quad & \det \left( \frac{1}{x-y} \left( \binom{i+j+x}{2i-j+1} - \binom{i+j+y}{2i-j+1} \right) \right)_{0 \leq i, j \leq n-1} \\
 &= 2^{-n} \prod_{k=0}^{n-1} \Delta_{2k}(x+y-1).
 \end{aligned}$$

Now let  $y \rightarrow x$ , and we obtain

$$(4.3) \quad \det \left( \frac{d}{dx} \binom{i+j+x}{2i-j+1} \right)_{0 \leq i, j \leq n-1} = 2^{-n} \prod_{k=0}^{n-1} \Delta_{2k}(2x-1).$$

Identity (4.1) is merely (4.3) after the differentiation has been completed.  $\square$

In [8], Mills, Robbins and Rumsey define “a totally symmetric plane partition of size  $n$  (to be) a plane partition whose three-dimensional Ferrers graph is contained in the box

$$X_n = [1, n] \times [1, n] \times [1, n]$$

and which is mapped to itself under all permutations of the coordinate axes. The complement of the Ferrers graph of such a plane partition (that is, the set of lattice points in the box  $X_n$  that do not belong to the Ferrers graph) is again totally symmetric when viewed from the vantage point of the vertex  $(n + 1, n + 1, n + 1)$ . A totally symmetric plane partition is self complementary if it is congruent (in the geometrical sense) to its complement. This cannot occur unless  $n = 2m$  is even”.

If we define  $A_n$  by the recurrence (4.6), then Mills et al. [8] conjecture that  $A_n$  is the total number of TSSCPP’s in  $X_{2n}$ .

In [12], J. Stembridge essentially proved that the TSSCPP conjecture reduces to the following result (the details of Stembridge’s result and its equivalence to the following are provided in [2; Sec. 2]). It should be noted that our proof of Corollary 2 in the final analysis relies on (1.1) and thus is quite different from the proof of the T.S.S.C.P.P. Conjecture given in [2].

**COROLLARY 2.**

$$(4.4) \quad \det(a_{ij})_{0 \leq i, j \leq n-1} = A_n^2,$$

where

$$(4.5) \quad a_{ij} = \begin{cases} 1 & \text{if } i = j = 0, \\ 0 & \text{if } i = j > 0, \\ \sum_{s=2i-j+1}^{2j-i} \binom{i+j}{s} & \text{if } i < j, \\ -a_{ji} & \text{if } i > j, \end{cases}$$

and

$$(4.6) \quad A_n = \prod_{i=0}^{n-1} \frac{(3i+1)!}{(n+i)!} = \frac{(3n-2)!(n-1)!}{(2n-1)!(2n-2)!} A_{n-1}.$$

*Proof.* We define several new matrices:

$$(4.7) \quad w(n) = \left( \binom{i+j+1}{2j-i} + \binom{i+j}{2j-i-1} \right)_{0 \leq i, j \leq n-1},$$

$$(4.8) \quad u(n) = (\delta_{ij} - 2\delta_{i, j+1})_{0 \leq i, j \leq n-1},$$

$$(4.9) \quad v(n) = \left( \binom{i+j+2}{2j-i+1} \frac{(3i+1)(3j+1)(3j-3i)}{(i+j)(i+j+1)(i+j+2)} \right)_{0 \leq i, j \leq n-1}$$

(where we define the  $(0, 0)$ -th entry of  $v(n)$  to be 1),

$$(4.10) \quad \begin{aligned} u_1(n) &= u(n) + (2\delta_{(i-1)^2+j, 0})_{0 \leq i, j \leq n-1} \\ &= \begin{pmatrix} 1 & 0 & 0 & 0 & \cdots \\ 0 & 1 & 0 & 0 & \cdots \\ 0 & -2 & 1 & 0 & \cdots \\ 0 & 0 & -2 & 1 & \cdots \end{pmatrix}_{n \times n}, \end{aligned}$$

$$(4.11) \quad st(n) = (a_{ij})_{0 \leq i, j \leq n-1}.$$

The matrices  $u$  and  $u_1$  are introduced to perform certain simple row and column operations. In particular, elementary algebra reveals that

$$(4.12) \quad u(n)w(n) = v(n)$$

and

$$(4.13) \quad u_1(n)st(n)(u_1(n))^T = v(n).$$

Finally, if we expand  $M_{n+1}(-2, -1)$  along the top row we find

$$(4.14) \quad \begin{aligned} M_{n+1}(-2, -1) &= 2 \det \left( \binom{i+j}{2i-j+1} + \binom{i+j+1}{2i-j+1} \right)_{0 \leq i, j \leq n-1} \\ &= 2 \det \left( \binom{i+j}{2j-i-1} + \binom{i+j+1}{2j-i} \right)_{0 \leq i, j \leq n-1} \\ &\quad \left( \text{since } \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} A \\ A-B \end{pmatrix} \right) \\ &= 2 \det(w(n)). \end{aligned}$$

Consequently, since the determinants of  $u$  and  $u_1$  are each 1, it follows from (1.6), (4.12), (4.13) and (4.14) that

$$\begin{aligned}
 (4.15) \quad \det(a_{ij})_{0 \leq i, j \leq n-1} &= \det(st(n)) = \det(v(n)) \\
 &= \det(w(n)) = \frac{1}{2} M_{n+1}(-2, -1) \\
 &= \prod_{k=1}^n \Delta_{2k}(-3) = A_n^2
 \end{aligned}$$

because setting  $u = -3$  in (1.2) we find

$$(4.16) \quad \Delta_{2k}(-3) = \left( \frac{(3k-2)!(k-1)!}{(2k-2)!(2k-1)!} \right)^2. \quad \square$$

**5. A related identity and another proof of the T.S.S.C.P.P. conjecture.** We have not found any related identities for two variables that are genuinely different from (1.6). However, there is one with one variable that merits mention. To this end we need the  $\Delta_j(n)$  with odd subscript [1; p. 196]:

$$(5.1) \quad \Delta_{2j-1}(\mu) = \frac{(\mu+2j)_{j-1}(\frac{1}{2}\mu+2j+\frac{1}{2})_j}{(j)_j(\frac{1}{2}\mu+j+\frac{1}{2})_{j-1}}.$$

The next formula is implicit in the work of Mills-Robbins-Rumsey [9]. However, they do not state it so we record it here: Let

$$(5.2) \quad P_n(\mu) = \left( \binom{i+j+\mu}{2i-j} + 2 \binom{i+j+\mu+2}{2i-j+1} \right)_{0 \leq i, j \leq n-1}.$$

In the notation of [9; p. 50], the determinant of  $P_n(\mu)$  is  $R_n(1, \mu)$ . This is easily seen by setting  $x = 1$  in their definition of  $R_n(x, \mu)$  [9; p. 50] and applying the Chu-Vandermonde summation. Furthermore, from their Theorems 5 and 7, it is easy to see that

$$\begin{aligned}
 (5.3) \quad \det(P_n(\mu)) &= \frac{\det(\delta_{ij} + \binom{i+j+2\mu}{i})_{0 \leq i, j \leq 2n-1}}{\det(\binom{i+j+\mu}{2i-j})_{0 \leq i, j \leq n-1}} \\
 &= \frac{2^n \prod_{k=0}^{2n-1} \Delta_k(2\mu)}{\prod_{k=0}^{n-1} \Delta_{2k}(2\mu)} = 2^n \prod_{k=1}^n \Delta_{2k-1}(2\mu).
 \end{aligned}$$

Our final result gives the determinant for

$$(5.4) \quad W_n(x) = \left( \binom{i+j+x+1}{2i-j} + \binom{i+j+x}{2i-j-1} \right)_{1 \leq i, j \leq n}.$$

## THEOREM 2.

$$(5.5) \quad \det(W_n(x)) = \prod_{k=1}^n \Delta_{2k-1}(2x+3).$$

*Proof.* We require an auxiliary matrix

$$(5.6) \quad \mathcal{S}_n(x) = \left( \frac{(-1/2)_{i-j}(-1)^{i-j}}{2^{2i-2j-1}(i-j)!} \right)_{0 \leq i, j \leq n-1}.$$

Consequently,

$$(5.7) \quad \mathcal{S}_n(x) \cdot W_n(x) = P_n(x+3/2).$$

This is easily seen if we introduce

$$(5.8) \quad f(i, j, X) = \sum_{k=0}^i \frac{(-1/2)_{i-k}(-1)^{i-k}}{2^{2i-2k-1}(i-k)!} \binom{k+j+X}{2k-j},$$

and note that as an immediate corollary of the Pfaff-Saalschultz summation [4; p. 9]:

$$(5.9) \quad f(i, j, X) = \binom{X+i+j-1/2}{2i-j} + \binom{X+i+j+1/2}{2i-j}.$$

Thus

$$\begin{aligned} & \mathcal{S}_n(x)W_n(x) \\ &= \left( \sum_{k \geq 0} \frac{(-1/2)_{i-k}(-1)^{i-k}}{2^{2i-2k-1}(i-k)!} \left\{ \binom{k+j+x+3}{2k-j+1} + \binom{k+j+x+2}{2k-j} \right\} \right)_{0 \leq i, j \leq n-1} \\ &= (f(i, j-1, x+4) + f(i, j, x+2))_{0 \leq i, j \leq n-1} \\ &= \left( \binom{x+i+j+5/2}{2i-j+1} + \binom{x+i+j+7/2}{2i-j+1} + \binom{x+i+j+3/2}{2i-j} + \binom{x+i+j+5/2}{2i-j} \right)_{0 \leq i, j \leq n-1} \\ &= \left( 2 \binom{x+i+j+7/2}{2i-j+1} + \binom{x+i+j+3/2}{2i-j} \right)_{0 \leq i, j \leq n-1} \\ &= P_n(x+3/2), \end{aligned}$$

as asserted in (5.7). Consequently since  $\det(\mathcal{S}_n(x)) = 2^n$ , we see from (5.7) and (5.3) that

$$\det(W_n(x)) = \prod_{k=1}^n \Delta_{2k-1}(2x+3)$$

as asserted in Theorem 2. □

We note that Corollary 2 is also derivable from Theorem 2. This is because from (4.7), (5.4) and (5.5)

$$\det(w(n)) = \det(W_{n-1}(0)) = \prod_{k=1}^{n-1} \Delta_{2k-1}(3) = A_n^2,$$

since

$$\Delta_{2k-1}(3) = \frac{(3+2k)_{k-1}(2+2k)_k}{(k)_k(2+k)_{k-1}} = \left( \frac{(3k+1)!k!}{(2k)!(2k+1)!} \right)^2.$$

We also remark that other special values for  $W_n(x)$  can be derived from Theorem 2 besides  $\det(W_n(0)) = A_{n+1}^2$ . Namely

$$\begin{aligned} \det(W_n(-1)) &= H_{2n+1}, \\ \det(W_n(-2)) &= H_{2n}, \\ \det(W_n(-3)) &= A_{n-1}A_n, \end{aligned}$$

where the sequence  $H_n$  is defined by  $H_0 = 1$ ,

$$\frac{H_{2n+1}}{H_{2n}} = \binom{3n}{n} / \binom{2n}{n}, \quad \frac{H_{2n}}{H_{2n-1}} = \frac{4}{3} \frac{\binom{3n}{n}}{\binom{2n}{n}}.$$

The sequences  $A_n$  and  $H_n$  occur in a number of unsolved problems of Mills-Robbins-Rumsey (cf. [11] for a survey of the problems).

**6. Conclusion.** The problem of enumerating symmetry classes of plane partitions is considered extensively in [12]. Indeed the identity (1.1) which we rely on heavily throughout our work was used by Mills et al. to treat plane partitions in a different symmetry class [9]. G. Kuperberg [7] has recently prepared an appealing survey of this topic.

Also recently M. Ishikawa [6] has found a nice plane partition theoretic interpretation of  $\det(v(n)) = A_n^2$  (see (4.9) and (4.15)).

**Acknowledgment.** Every stage of this work and each theorem and lemma was found empirically using the symbolic algebra package AXIOM. While our proofs do not rely on the computer, each of our discoveries would have been impossible without the flexibility and power of AXIOM.

## REFERENCES

- [1] G. E. Andrews, *Plane partitions III: the weak Macdonald conjecture*, Invent. Math., **53** (1979), 193–225.
- [2] ———, *Plane partitions V: the T.S.S.C.P.P. conjecture*, J. Combin. Theory Ser. A, (to appear).
- [3] W. N. Bailey, *Some identities involving generalized hypergeometric series*, Proc. London Math. Soc. Ser. 2, **29** (1929), 503–516.
- [4] ———, *Generalized Hypergeometric Series* (Reprinted: Hafner, New York, 1964), Cambridge University Press, London and New York, 1935.
- [5] A. Erdelyi et al., *Higher Transcendental Functions*, vol. 1, McGraw-Hill, New York, 1953.
- [6] M. Ishikawa, *A remark on totally symmetric self-complementary plane partition*, (to appear).
- [7] G. Kuperberg, *Symmetries of plane partitions and the permanent-determinant method*, (to appear).
- [8] W. H. Mills, D. P. Robbins and H. Rumsey, *Self-complementary, totally symmetric plane partitions*, J. Combin. Theory Ser. A, **42** (1986), 277–292.
- [9] ———, *Enumeration of a symmetry class of plane partitions*, Discrete Math., **67** (1987), 43–55.
- [10] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, Cambridge, 1966.
- [11] R. P. Stanley, *A baker's dozen of conjectures concerning plane partitions*, from Combinatoire énumérative (G. Labelle and P. Leroux, eds.), Lecture Notes in Math. No. 1234, Springer, Berlin/New York, 1986, pp. 285–293.
- [12] ———, *Symmetries of plane partitions*, J. Combin. Theory Ser. A, **3** (1986), 103–113.
- [13] J. Stembridge, *Nonintersecting paths, Pfaffians, and plane partitions*, Advances in Math., **83** (1990), 96–131.
- [14] F. J. W. Whipple, *Well-poised series and other generalized hypergeometric series*, Proc. London Math. Soc. Ser. 2, **25** (1926), 525–544.

Received May 13, 1991 and in revised form March 16, 1992. Partially supported by NFS Grant DMS 8702695-03 and the IBM Thomas J. Watson Research Center.

PENNSYLVANIA STATE UNIVERSITY  
UNIVERSITY PARK, PA 16802

AND

IBM THOMAS J. WATSON RESEARCH CENTER  
YORKTOWN HEIGHTS, NY 10598

# PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982)      F. WOLF (1904–1989)

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555  
vsv@math.ucla.edu

F. MICHAEL CHRIST  
University of California  
Los Angeles, CA 90024-1555  
christ@math.ucla.edu

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112  
clemens@math.utah.edu

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093  
tenright@ucsd.edu

NICHOLAS ERCOLANI  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

R. FINN  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720  
vfr@math.berkeley.edu

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106  
mgscharl@henri.ucsb.edu

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
UNIVERSITY OF MONTANA  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON



# PACIFIC JOURNAL OF MATHEMATICS

Volume 158    No. 1    March 1993

---

Determinant identities	1
GEORGE W. EYRE ANDREWS and WILLIAM H. BURGE	
A spectral theory for solvable Lie algebras of operators	15
E. BOASSO and ANGEL RAFAEL LAROTONDA	
Simple group actions on hyperbolic Riemann surfaces of least area	23
S. ALLEN BROUGHTON	
Duality for finite bipartite graphs (with an application to $\text{II}_1$ factors)	49
MARIE CHODA	
Szegő maps and highest weight representations	67
MARK GREGORY DAVIDSON and RON STANKE	
Optimal approximation class for multivariate Bernstein operators	93
ZEEV DITZIAN and XINLONG ZHOU	
Witt rings under odd degree extensions	121
ROBERT FITZGERALD	
Congruence properties of functions related to the partition function	145
ANTHONY D. FORBES	
Bilinear operators on $L^\infty(G)$ of locally compact groups	157
COLIN C. GRAHAM and ANTHONY TO-MING LAU	
Nonuniqueness of the metric in Lorentzian manifolds	177
GEOFFREY K. MARTIN and GERARD THOMPSON	
Index theory and Toeplitz algebras on one-parameter subgroups of Lie groups	189
EFTON PARK	