

*Pacific  
Journal of  
Mathematics*

**SOLUTIONS OF THE STATIONARY AND NONSTATIONARY  
NAVIER-STOKES EQUATIONS IN EXTERIOR DOMAINS**

ZHI MIN CHEN

# SOLUTIONS OF THE STATIONARY AND NONSTATIONARY NAVIER-STOKES EQUATIONS IN EXTERIOR DOMAINS

ZHI-MIN CHEN

**It is shown that a nonstationary exterior Navier-Stokes flow tends to a small stationary flow in  $L^2$  like  $t^{-3/4}$  as  $t \rightarrow \infty$ .**

**0. Introduction.** In this paper we are concerned with the stationary Navier-Stokes equations

$$(0.1) \quad \begin{aligned} (w \cdot D)w - \Delta w + D\bar{p} &= f, \quad D \cdot w = 0 \quad \text{in } G, \\ w &= 0 \quad \text{on } \partial G \quad (D = \text{grad}), \end{aligned}$$

and the nonstationary Navier-Stokes equations

$$\begin{aligned} v_t + (v \cdot D)v - \Delta v + D\bar{\bar{p}} &= f \quad \text{in } G \times (0, \infty), \\ D \cdot v &= 0 \quad \text{in } G \times (0, \infty), \\ v &= 0 \quad \text{on } \partial G \times (0, \infty), \\ v|_{t=0} &= a + w \quad \text{in } G \quad (v_t = \partial v / \partial t). \end{aligned}$$

Here and in what follows  $G$  denotes a smooth exterior domain of  $R^3$ ,  $f = f(x)$  is a prescribed vector field, and  $\bar{p}$  (resp.  $\bar{\bar{p}}$ ) represents unknown stationary (resp. nonstationary) scalar pressure which can be determined by the stationary solution  $w$  via (0.1) (resp. nonstationary solution  $v$  via (0.2)).

As is well known, it was shown by Finn [8, 9] that (0.1) admits a small solution

$$(0.3) \quad \begin{aligned} w \in L^\infty(G; R^3), \quad Dw \in L^3(G; R^9), \\ C_0 = \sup_{x \in G} |x| |w(x)| < \infty. \end{aligned}$$

If  $C_0 < 1/2$  the Finn's solution  $w$  may be formed as a limit of a nonstationary solution  $v$  as  $t \rightarrow \infty$  in local or global  $L^2$ -norms (cf. Heywood [15, 14], Galdi and Rionero [11], Miyakawa and Sohr [23]) and in other norms (cf. Heywood [16], Masuda [20]). Moreover it has recently proved (cf. Borchers and Miyakawa [4]) that every weak solution of (0.2) tends the Finn's solution in  $L^2(G; R^3)$

like  $t^{-(3/p-3/2)/2}$  with  $6/5 < p < 2$ , provided  $C_0 < 1/2$  and  $a \in L^2(G; R^3) \cap L^p(G; R^3)$ .

In this paper we are only interested in the case  $w \in L^3(G; R^3)$ ,  $Dw \in L^{3/2}(G; R^9)$ , or  $Dw \in L^r(G; R^9) \cap L^p(G; R^9)$  with  $1 < r < 3/2 < p < 2$ . Under certain smallness assumptions on  $w$  we show now that every weak solution of (0.2) tends to the stationary solution  $w$  in  $L^2(G; R^3)$  like the sharp decay rate  $t^{-3/4}$ .

**1. Notation and main result.** In this paper we use the following spaces.

$L^p$  = the Lebesgue spaces  $L^p(G; R^3)$ , with  $\|\cdot\|_p$  the associated norm,

$C_\sigma^\infty$  = the set of compactly supported solenoidal in  $C^\infty(G; R^3)$ ,

$W^{k,p}$  = the Sobolev space  $W^{k,p}(G; R^3)$ ,

$J^p$  = the completion of  $C_\sigma^\infty$  in  $L^p$ ,

$W_\sigma^{1,p}$  = the completion of  $C_\sigma^\infty$  in  $W^{1,p}$ ,

$\widehat{W}_\sigma^{1,p}$  = the completion of  $C_\sigma^\infty$  under the norm  $\|D \cdot\|_p$ ,

$\widehat{W}_\sigma^{2,p}$  = the space  $\{u \in \widehat{W}_\sigma^{1,3p/(3-p)}; D^2u \in L^p(G; R^{27})\}$

for  $1 < p < 3$ ,

$W^{-1,2}$  = the dual of  $W_\sigma^{1,2}$ ,

$\widehat{W}^{-1,p}$  = the dual of  $\widehat{W}_\sigma^{1,p/(p-1)}$ , with  $\|\cdot\|_{-1,p}$  the associated norm.

Moreover for  $1 < r < \infty$  and  $n \geq 1$ , we denote by  $r'$  the real  $r/(r-1)$ , by  $(\cdot, \cdot)$  the inner product in  $L^2(G; R^n)$ , by  $P$  the bounded projection from  $L^r$  onto  $J^r$  (cf. [22]), by  $A$  the Stokes operators  $-P\Delta$  with the domain  $W_\sigma^{1,r} \cap W^{2,r}$ , by  $\bar{A}$  the Laplacian  $-\Delta$  with the domain  $W^{2,r}(R^3; R^3)$ , and by  $C$  a positive constant which may vary from line to line, but is always independent of the quantities  $t, T, u, v, w, f, u_k$ , and  $a$ .

Now we make preparations for stating our main result. The existence of the stationary solutions  $w$  is guaranteed by the following.

**LEMMA 1.1.** *Let  $1 < r \leq 3/2 < p < 2$ , and  $f \in C_\sigma^\infty$ . Then there is a small  $h > 0$  such that (0.1) admits a unique solution within the class*

$$\{w \in \widehat{W}_\sigma^{1,r} \cap \widehat{W}_\sigma^{1,p}; \|Dw\|_{3/2} \leq h\},$$

*provided that  $\|f\|_{-1,3/2} \leq h^2$ . Moreover*

$$\|Dw\|_r + \|Dw\|_p \leq C(\|f\|_{-1,r} + \|f\|_{-1,p}).$$

From (0.1) and (0.2) we see that  $u = v - w$  and  $\hat{p} = \bar{p} - \bar{\bar{p}}$  solve the problem

$$\begin{aligned}
 (1.1) \quad & u_t + (u \cdot D)u - \Delta u + (u \cdot D)w + (w \cdot D)u + D\hat{p} = 0, \\
 & D \cdot u = 0 \quad \text{in } G \times (0, \infty), \\
 & u = 0 \quad \text{on } \partial G \times (0, \infty), \\
 & u|_{t=0} = a \quad \text{in } G.
 \end{aligned}$$

Weak solutions are given in the following sense.

DEFINITION 1.1. Let  $a \in J^2$ , and  $w \in \widehat{W}_\sigma^{1,3/2}$  solve (0.1). A weakly continuous function  $u: [0, \infty) \rightarrow J^2$  is said to be a weak solution of (1.1) if  $u(0) = a$ ,  $u \in L^\infty(0, \infty; J^2) \cap L^2(0, \infty; \widehat{W}_\sigma^{1,2})$ ,

$$(1.2) \quad \|u(t)\|_2^2 + \int_s^t \|Du(z)\|_2^2 dz \leq \|u(s)\|_2^2,$$

$$\begin{aligned}
 (1.3) \quad & (u(t), g(t)) + \int_s^t ((Du, Dg) + ((u \cdot D)w, g) \\
 & \quad + ((w \cdot D)u, g) - (u, g_z)) dz \\
 & = (u(s), g(s)) - \int_s^t ((u \cdot D)u, g) dz
 \end{aligned}$$

for all  $t > s \geq 0$  and all  $g \in C([0, \infty); W_\sigma^{1,2}) \cap C^1([0, \infty); J^2)$ , where  $g_z = \partial g / \partial z$ .

The existence of weak solutions to (1.1) is guaranteed by the following.

LEMMA 1.2. Let  $a \in J^2$ , and  $w \in \widehat{W}_\sigma^{1,3/2}$  such that  $\|Dw\|_{3/2} < 1/8$ . Then (1.1) admits a weak solution.

We are now in a position to state our main result.

THEOREM 1.1. Let  $1 < r < 3/2 < p < 2$ ,  $a \in J^2 \cap L^1$ , and let  $w \in W_\sigma^{1,r} \cap W_\sigma^{1,p}$  such that  $w$  solves (0.1) and  $\|Dw\|_r + \|Dw\|_p$  is sufficiently small. Then every weak solution of (1.1) possesses the sharp decay property

$$\|u(t)\|_2 = O(t^{-3/4}).$$

Section 2 is concerned with the proof of Lemmas 1.1 and 1.2. In [23], it has been obtained an existence result on weak solutions of (1.1) with  $w$  the Finn's solution such that  $C_0 < 1/2$ . However,

the argument of [23] heavily depends on the property (0.3). In §3, with the use of the approach developed from [7], we shall show sharp decay estimates of solutions to the linearized equations of (1.1). If  $w$  only satisfies (0.3) and  $C_0 < 1/2$ , such estimates seem unavailable. Theorem 1.1 will be proved in §4 by making use of the estimates carried out in §3 and studying the time average  $t^{-1} \int_0^t \|u(s)\|_2 ds$ . A similar technique has been used in [23, 4]. However, we have not used the spectral decomposition of the Stokes operator  $A$  in  $L^2$  as usually used in earlier work concerning the  $L^2$  decay problem. Moreover our proof seems much simpler.

It should be noted that the  $L^2$  decay problem of (1.1) with  $w = 0$  stems from Leray [19], and has affirmatively been solved (cf. [24, 3, 2] and the references therein). If  $1 < p < 2$  and  $u$  is a weak solution of (1.1) with  $w = 0$ , it has been proved that  $\|u(t)\|_2 = O(t^{-(3/p-3/2)/2})$  provided  $u(0) \in J^2 \cap L^p$  (cf. [2]), and  $\|u(t)\|_2 = O(t^{-3/4})$  provided  $u(0) \in J^2 \cap L^1$  and  $\|e^{-tA}a\|_2 \leq Ct^{-3/4}\|a\|_1$  (cf. [3]).

**2. Proof of Lemmas 1.1, 1.2.** To begin with we give the estimate (cf. [2, Theorem 3.6] or [12, 18] for a similar consideration)

$$(2.1) \quad \|Du\|_p \leq C \sup\{|(Du, Dv)|; v \in C_\sigma^\infty, \|Dv\|_{p'} = 1\} \\ \text{for } 1 < p < n, u \in \widehat{W}_\sigma^{1,p},$$

and the Sobolev inequality (cf. [13])

$$(2.2) \quad \|u\|_{3p/(3-p)} \leq 2p(3-p)^{-1}3^{-1/2}\|Du\|_p \\ \text{for } 1 < p < n, u \in \widehat{W}_\sigma^{1,p}.$$

*Proof of Lemma 1.1.* Let  $r$  and  $p$  be given in Lemma 1.1. We rewrite (0.1) in the abstract form  $Aw + P(w \cdot D)w = f$ ,  $w \in \widehat{W}_\sigma^{1,r} \cap \widehat{W}_\sigma^{1,p}$ . Since the proof of [5, (3.1)] implies that  $A$  can be extended as a bounded and invertible operator from  $\widehat{W}_\sigma^{2,q}$  onto  $J^q$  with  $1 < q < 3/2$ , we can set

$$H: \widehat{W}_\sigma^{1,r} \cap \widehat{W}_\sigma^{1,p} \rightarrow \widehat{W}_\sigma^{2,3p/(6-p)} \quad \text{such that } Hw = A^{-1}(f - P(w \cdot D)w).$$

Let  $w \in \widehat{W}_\sigma^{1,r} \cap \widehat{W}_\sigma^{1,p}$ ,  $r < s < p$ , and  $v \in C_\sigma^\infty$  with  $\|Dv\|_{s'} = 1$ . Integrating by parts and using the divergence condition  $D \cdot w = 0$ , we have

$$\begin{aligned} (DHw, Dv) &= (f, v) - ((w \cdot D)w, v) \\ &= (f, v) + ((w \cdot D)v, w) \\ &\leq (f, v) + \|w\|_3 \|w\|_{3s/(3-s)} \|Dv\|_{s'}, \end{aligned}$$

that is, by (2.1)–(2.2),

$$\|DHw\|_s \leq C(\|f\|_{-1,s} + \|Dw\|_s \|Dw\|_{3/2}).$$

Similarly, for  $w, w^* \in W_\sigma^{1,r} \cap W_\sigma^{1,p}$  we have

$$\|DHw - DHw^*\|_s \leq C(\|Dw\|_{3/2} + \|Dw^*\|_{3/2})\|Dw - Dw^*\|_s.$$

Consequently, the desired assertion follows immediately from the contraction mapping principle. The proof is complete.

In [23], Miyakawa and Sohr proved that (1.1) admits a weak solution in case  $w$  is the Finn’s solution and  $C_0 < 1/2$ . However, as for our case, the argument of [23] does not work somewhere. Now we give our proof in a slightly different way. Similar to [23], we also study approximate solutions of (1.1) by applying a technique developed from [6].

*Proof of Lemma 1.2.* Let  $k > 1$ . We set  $J_k = k(k + A)^{-1}$  and  $I_k = k(k + \bar{A})^{-1}E$ , where  $E$  denotes the extension operator such that  $Eu = u$  in  $G$  and  $Eu = 0$  outside  $G$ . With the use of the notation above, we have

$$(2.3) \quad \|J_k u\|_p \leq C(k)\|u\|_r, \quad \|I_k u\|_p \leq C(k)\|u\|_r$$

for  $1 < r < p \leq \infty, u \in J^r,$

$$(2.4) \quad \|I_k u\|_r \leq \|u\|_r, \quad \|J_k u\|_r \leq C\|u\|_r \quad \text{for } 1 < r < \infty, u \in J^r,$$

where  $C$  is independent of  $k$ . (2.3) is a consequence of the Sobolev embedding theorem and  $L^r$ -estimates. The first inequality in (2.4) follows from the proof of [1, Lemma 10.1], and the second one from [2, Theorem 1.2].

Now we proceed to the evolution equation

$$(2.5) \quad (d/dt)u_k + Au_k = F_k(u_k), \quad u_k(0) = J_k a \quad \text{in } J^2,$$

where  $F_k(u) = F_k(u, u)$  with

$$F_k(u, v) = -P(J_k u \cdot D)v - P(J_k w \cdot D)u - P(I_k u \cdot D)I_k w.$$

For  $u, v \in W_\sigma^{1,2}$ , we have

$$(2.6) \quad \|F_k(u, v)\|_2 + \|P(J_k v \cdot D)u\|_2$$

$$\leq \|J_k u\|_\infty \|Dv\|_2 + \|J_k w\|_\infty \|Du\|_2$$

$$+ \|I_k u\|_6 \|DI_k w\|_3 + \|J_k v\|_\infty \|Du\|_2$$

$$\leq C(k)(\|u\|_6 \|Dv\|_2 + \|w\|_3 \|Du\|_2$$

$$+ \|u\|_6 \|I_k DEw\|_3 + \|v\|_6 \|Du\|_2), \quad \text{by (2.3),}$$

$$\leq C(k)\|Du\|_2(\|Dv\|_2 + \|Dw\|_{3/2}), \quad \text{by (2.2).}$$

On the other hand, given  $k$  and  $T > 0$ , we suppose that  $u_k$  solve (2.5) over  $[0, T)$ , and  $u_k \in L^2(0, T; W_\sigma^{1,2} \cap W^{2,2}) \cap W^{1,2}(0, T; J^2)$ . Then multiplying (2.5) by  $2u_k$  and  $2Au_k$ , respectively, we have

$$\begin{aligned} (d/dt)\|u_k\|_2^2 + 2\|Du_k\|_2^2 &= 2(F_k(u_k), u_k), \\ (d/dt)\|Du_k\|_2^2 + 2\|Au_k\|_2^2 &= 2(F_k(u_k), Au_k). \end{aligned}$$

The estimation of the right-hand side terms of the preceding identities can be achieved as follows.

$$\begin{aligned} 2(F_k(u_k), u_k) &= 2((I_k u_k \cdot D)u_k, I_k w), \\ &\quad \text{since } D \cdot J_k u_k = D \cdot J_k w = D \cdot I_k u = 0, \\ &\leq 2\|I_k u_k\|_6 \|Du_k\|_2 \|I_k w\|_3 \\ &\leq (12/3^{-1/2})\|w\|_3 \|Du_k\|_2^2, \quad \text{by (2.4) and (2.2),} \\ &\leq 8\|Dw\|_{3/2} \|Du_k\|_2^2, \quad \text{by (2.2),} \\ &\leq \|Du_k\|_2^2, \quad \text{by setting } \|Dw\|_{3/2} < 1/8, \end{aligned}$$

$$\begin{aligned} 2(F_k(u_k), Au_k) &\leq 2\|Au_k\|_2 (\|J_k u_k\|_\infty \|Du_k\|_2 + \|J_k w\|_\infty \|Du_k\|_2 \\ &\quad + \|I_k u\|_\infty \|I_k DEw\|_2) \\ &\leq C(k)\|Au_k\|_2 \|Du_k\|_2 (\|u_k\|_2 + \|Dw\|_{3/2} + \|DEw\|_{3/2}), \\ &\quad \text{by (2.3) and (2.2),} \\ &\leq C(k)\|Au_k\|_2 \|Du_k\|_2 (\|u_k\|_2 + \|Dw\|_{3/2}) \\ &\leq 2\|Au_k\|_2^2 + C(k)\|Du_k\|_2^2 (\|u_k\|_2^2 + \|Dw\|_{3/2}^2). \end{aligned}$$

Consequently, we have

$$(2.7) \quad \|u_k(t)\|_2^2 + \int_s^t \|Du_k(z)\|_2^2 dz \leq \|u_k(s)\|_2^2, \quad 0 \leq s < t < T,$$

$$\begin{aligned} (2.8) \quad \|Du_k(t)\|_2^2 &\leq \|DJ_k a\|_2^2 + C(k) \int_0^t \|Du_k(s)\|_2^2 (\|u_k(s)\|_2^2 + \|Dw\|_{3/2}^2) ds \\ &\leq \|DJ_k a\|_2^2 + C(k)\|J_k a\|_2^2 (\|J_k a\|_2^2 + \|Dw\|_{3/2}^2), \quad \text{by (2.7)} \end{aligned}$$

Thus, following the same way as in the proof of [23, Proposition 3.4] by making use of (2.6)–(2.8), we conclude that (2.5) admits a unique global solution  $u_k$  satisfying (2.6), and  $u_k \in L^2(0, T; W_\sigma^{1,2} \cap W^{2,2}) \cap W^{1,2}(0, T; J^2)$  for all  $T > 0$ .

To obtain a weak solution of (1.1), we need to study compactness of the sequence  $u_k$ . Let  $v \in W_\sigma^{1,2}$ . Applying (2.2) and (2.4) repeatedly, we have, from (2.5),

$$\begin{aligned} & ((d/dt)u_k, v) \\ & \leq \|Du_k\|_2 \|Dv\|_2 + \|J_k u_k\|_3 \|Du_k\|_2 \|v\|_6 + \|J_k w\|_3 \|Du_k\|_2 \|v\|_6 \\ & \quad + \|I_k u_k\|_6 \|DI_k w\|_{3/2} \|v\|_6 \\ & \leq \|Du_k\|_2 \|Dv\|_2 + C \|v\|_6 (\|u_k\|_3 \|Du_k\|_2 + \|w\|_3 \|Du_k\|_2 \\ & \quad \quad \quad + \|u_k\|_6 \|DEw\|_{3/2}) \\ & \leq C \|Dv\|_2 (\|Du_k\|_2 + \|u_k\|_2^{1/2} \|Du_k\|_2^{3/2} + \|Du_k\|_2 \|Dw\|_{3/2}) \\ & \leq C \|Dv\|_2 (1 + \|a\|_2^{1/2} + \|Dw\|_{3/2}) (\|Du_k\|_2 + \|Du_k\|_2^{3/2}), \\ & \quad \quad \quad \text{by (2.7) and (2.4),} \end{aligned}$$

with  $C$  independent of  $k$ . This together with (2.7) implies that the sequence  $u_k$  is bounded in

$$L^\infty(0, \infty; J^2) \cap L^2(0, \infty; \widehat{W}_\sigma^{1,2}) \cap W^{1,4/3}(0, T; W^{-1,2})$$

for all  $0 < T < \infty$ . From [26, Theorem 2.1 in Chapter III] it follows readily that there are a function  $u$  and a subsequence of  $u_k$ , denoted again  $u_k$ , satisfying

$$\begin{aligned} u_k & \xrightarrow{w^*} u \text{ in } L^\infty(0, \infty; J^2), \\ u_k & \xrightarrow{w} u \text{ in } L^2(0, \infty; \widehat{W}_\sigma^{1,2}), \\ u_k & \rightarrow u \text{ strongly in } L^2_{\text{loc}}(G \times (0, \infty)). \end{aligned}$$

As in [21], we can check that the limit  $u$  is a weak solution of (1.1). The proof is complete.

**3. Decay estimates.** In this section, we let  $t > 0$ ,  $1 < r < 3/2 < p < 2$ , and  $w$  be a solution of (0.1) such that  $w \in \widehat{W}_\sigma^{1,r} \cap \widehat{W}_\sigma^{1,p}$ , and set

$$\begin{aligned} Lu & = Au + P(u \cdot D)w + P(w \cdot D)u, \\ B^*u & = -p(w \cdot D)u + P \sum_{i=1}^n u^i Dw^i, \\ L^*u & = Au + B^*u. \end{aligned}$$

Thus, we see that

$$(Lu, v) = (u, L^*v) \quad \text{for } u, v \in W_\sigma^{1,2} \cap W^{2,2},$$



and the linearized equation of (1.1) can be stated in the form

$$(d/dt)v + Lv = 0, \quad v(0) = u.$$

Denote by  $e^{-tL}u$  the solution of the preceding equation. It is the purpose of this section to prove the following.

**PROPOSITION 3.1.** *Suppose that  $\|Dw\|_r + \|Dw\|_p$  is sufficiently small. Then there holds*

$$(3.1) \quad \|e^{-tL}Pu\|_2 \leq Ct^{-3/4}\|u\|_1$$

for  $u \in L^1 \cap L^{6/5}$ .

The preceding proposition is based on the following decay estimates.

$$(3.2) \quad \|e^{-tA}u\|_\infty \leq Ct^{-1/4}\|u\|_6 \quad \text{for } u \in J^6,$$

$$(3.3) \quad \|e^{-tA}u\|_s \leq Ct^{-(3/q-3/s)/2}\|u\|_q \quad \text{for } 1 < q \leq s < \infty, u \in J^q,$$

$$(3.4) \quad \|De^{-tA}u\|_s \leq Ct^{-(1+3/q-3/s)/2}\|u\|_q \quad \text{for } 1 < q \leq s \leq 3, u \in J^q.$$

The estimates (3.3) and (3.4) were recently obtained by Iwashita (cf. [17, Theorems 1.2, 1.3]). (3.2) will be proved in the Appendix by using the argument of [17].

With the use of (3.2)–(3.4), we can now prove the following.

**LEMMA 3.1.** *Let  $u \in C_\sigma^\infty$ . Then there hold*

$$(3.5) \quad \|e^{-tA}u\|_\infty \leq Ct^{-3/4}\|u\|_2,$$

$$(3.6) \quad \|e^{-tA}B^*u\|_\infty + \|De^{-tA}B^*u\|_3 \leq Ct^{-3/2p}(t+1)^{-(3/r-3/p)/2}(\|u\|_\infty + \|Du\|_3)(\|Dw\|_r + \|Dw\|_p).$$

*Proof.* From (3.2), (3.3), (2.2) and the semigroup property of  $e^{-tA}$  we get (3.5) and

$$\begin{aligned} \|e^{-tA}B^*u\|_\infty &\leq Ct^{-3/2b}\|B^*u\|_b \\ &\leq Ct^{-3/2b}\|Dw\|_b(\|u\|_\infty + \|Du\|_3) \end{aligned}$$

for  $b = r, p$ . Moreover (3.4) and (2.2) yield

$$\|De^{-tA}B^*u\|_3 \leq Ct^{-3/2b}\|Dw\|_b(\|u\|_\infty + \|Du\|_3) \quad \text{for } b = r, p.$$

Collecting terms, we get readily (3.6) and complete the proof.

*Proof of Proposition 3.1.* Setting  $v(t) = e^{-tL^*}u$  with  $u \in C_\sigma^\infty$ , we have obviously that  $v \in C([0, \infty); L^\infty \cap W_\sigma^{1,3})$  and

$$v(t) = e^{-tA}u + \int_0^t e^{-(t-s)A}B^*v(s) ds.$$

This gives, by (3.4)–(3.6),

$$\begin{aligned} & \|v(t)\|_\infty + \|Dv(t)\|_3 \\ & \leq Ct^{-3/4}\|u\|_2 + C \int_0^t (t-s)^{-3/2p}(t-s+1)^{-(3/r-3/p)/2} \\ & \quad \times (\|v\|_\infty + \|Dv\|_3) ds (\|Dw\|_r + \|Dw\|_p). \end{aligned}$$

Setting  $\|v\|_t = \sup_{0 < s < t} s^{3/4}(\|v(s)\|_\infty + \|Dv(s)\|_3)$ , we have

$$\begin{aligned} & \|v(t)\|_\infty + \|Dv(t)\|_3 \\ & \leq Ct^{-3/4}\|u\|_2 + C(\|Dw\|_r + \|Dw\|_p)\|v\|_t \\ & \quad \times \int_0^t (t-s)^{-3/2p}(t-s+1)^{-(3/r-3/p)/2} s^{-3/4} ds \\ & \leq Ct^{-3/4}\|u\|_2 + Ct^{-3/4}(\|Dw\|_r + \|Dw\|_p)\|v\|_t \\ & \quad \times \int_0^t s^{-3/2p}(s+1)^{-(3/r-3/p)/2} ds \\ & \quad + Ct^{1/4-3/2p}(t+1)^{-(3/r-3/p)/2}(\|Dw\|_r + \|Dw\|_p)\|v\|_t \\ & \leq Ct^{-3/4}(\|u\|_2 + (\|Dw\|_r + \|Dw\|_p)\|v\|_t), \end{aligned}$$

where we have used the condition  $r < 3/2 < p$ . Hence, if we presuppose that

$$(3.7) \quad C(\|Dw\|_r + \|Dw\|_p) < 1/2$$

with the constant  $C$  given in the last term above, we obtain

$$(3.8) \quad \|e^{-tL^*}u\|_\infty \leq Ct^{-3/4}\|u\|_2.$$

Now we take  $u \in L^1 \cap L^{6/5}$  and  $v \in L^2$ . By (3.8) we have

$$(e^{-tL}Pu, v) = (u, e^{-tL^*}Pv) \leq \|u\|_1 \|e^{-tL^*}Pv\|_\infty \leq Ct^{-3/4}\|u\|_1 \|v\|_2$$

and therefore the validity of (3.1). The proof is complete.

**4. Proof of Theorem 1.1.** In this section we always suppose that the stationary solution  $w \in \widehat{W}_\sigma^{1,r} \cap \widehat{W}_\sigma^{1,p}$  with  $1 < r < 3/2 < p < 2$  such that (3.7) holds. Let  $u$  be a weak solution of (1.1). Then (1.2) implies

$$(4.1) \quad \|u(t)\|_2 \leq t^{-1} \int_0^t \|u(s)\|_2 ds.$$

On the other hand, taking  $v \in C_\sigma^\infty$  and applying (1.3) with  $g(z) = e^{-(t-z)L^*}v$ , we have

$$\begin{aligned} & (u(t), v) + \int_0^t (Lu(s), e^{-(t-s)L^*}v) ds - \int_0^t (u(s), L^*e^{-(t-s)L^*}v) ds \\ & = (a, e^{-tL^*}v) - \int_0^t ((u \cdot D)u, e^{-(t-s)L^*}v) ds, \end{aligned}$$

that is,

$$\begin{aligned} (u(t), v) &= (e^{-tL}a, v) - \int_0^t (e^{-(t-s)L}P(u \cdot D)u(s), v) ds \\ &\leq \|e^{-tL}a\|_2 \|v\|_2 + \int_0^t \|e^{-(t-s)L}P(u \cdot D)u(s)\|_2 ds \|v\|_2 \\ &\leq C \|v\|_2 \left( t^{-3/4} \|a\|_1 + \int_0^t (t-s)^{-3/4} \|u(s)\|_2 \|Du(s)\|_2 ds \right), \end{aligned}$$

where we have used (3.1). We then get

$$\|u(s)\|_2 \leq Cs^{-3/4} \|a\|_1 + C \int_0^s (s-z)^{-3/4} \|u(z)\|_2 \|Du(z)\|_2 dz.$$

Integrating the above inequality from 0 to  $t$ , we have

$$\begin{aligned} \int_0^t \|u(s)\|_2 ds &\leq Ct^{1/4} \|a\|_1 + C \int_0^t dz \int_z^t (s-z)^{-3/4} \|u(z)\|_2 \|Du(z)\|_2 ds \\ &\leq Ct^{1/4} \|a\|_1 + Ct^{1/4} \int_0^t \|u(s)\|_2 \|Du(s)\|_2 ds \\ &\leq Ct^{1/4} \|a\|_1 + Ct^{1/4} \|a\|_2 \left( \int_0^t \|u(s)\|_2^2 ds \right)^{1/2}, \quad \text{by (1.2)}. \end{aligned}$$

Combining this with (4.1), we have

$$\|u(t)\|_2 \leq Ct^{-3/4} \|a\|_1 + Ct^{-3/4} \|a\|_2 \left( \int_0^t \|u(s)\|_2^2 ds \right)^{1/2},$$

that is,

$$(4.2) \quad \|u(t)\|_2 \leq C_1 t^{-3/4} \left( 1 + \left( \int_0^t \|u(s)\|_2^2 ds \right)^{1/2} \right),$$

where and in what follows  $C_1 = C_1(\|a\|_1, \|a\|_2)$  may vary from line to line.

Now we apply (4.2) and (1.2) to complete our proof via a boot strap iteration argument.

Note that

$$(4.3) \quad \|u(t)\|_2 \leq C_1, \quad \text{by (1.2),}$$

and

$$(4.4) \quad \|u(t)\|_2 \leq C_1 t^{-3/4} (1 + t^{1/2}), \quad \text{by (4.2) and (4.3).}$$

Combining (4.4) with (4.3), we have

$$(4.5) \quad \|u(t)\|_2 \leq C_1 t^{-1/4}.$$

Moreover, taking (4.2) and (4.5) into account, we have

$$\|u(t)\|_2 \leq C_1 t^{-3/4} (1 + t^{1/4}).$$

This together with (4.3) implies

$$(4.6) \quad \|u(t)\|_2 \leq C_1 (t + 1)^{-1/2}.$$

Similarly, (4.2) and (4.6) yield

$$\|u(t)\|_2 \leq C_1 t^{-3/4} (1 + \ln(t + 1)),$$

and so, by (4.3),

$$(4.7) \quad \|u(t)\|_2 \leq C_1 (t + 1)^{-2/3}.$$

Finally, by (4.2) and (4.7), we arrive at the desired estimate

$$\|u(t)\|_2 \leq C_1 t^{-3/4}$$

and complete the proof.

**REMARK 4.1.** It should be noted that the validity of the assumption of Lemma 1.2 follows from the inequality  $\|Dw\|_{3/2} \leq \|Dw\|_r + \|Dw\|_p$  and (2.7).

**Appendix: Proof of (3.2).** Let  $Q$  be a domain of  $R^3$ . By  $\|\cdot\|_{k,p,Q}$  and  $\|\cdot\|_{p,Q}$  we denote respectively the norms of the Sobolev space  $W^{k,p}(Q; R^3)$  and the Lebesgue space  $L^p(Q; R^3)$ . Of course,  $\|\cdot\|_{k,p} = \|\cdot\|_{k,p,G}$  and  $\|\cdot\|_p = \|\cdot\|_{p,G}$ .  $\bar{P}$  is the bounded projection from  $L^p(R^3; R^3)$  onto  $J^p(R^3; R^3)$ , where  $J^p(R^3; R^3)$  denotes the completion of the set of compactly supported solenoidal in  $C^\infty(R^3; R^3)$ . Let  $h$  be a constant such that  $|x| < h - 1$  for  $x \in \partial G$ , and let  $g \in C^\infty(R^3; R)$  be a fixed function such that  $g = 1$  for  $|x| > h$  and  $g = 0$  for  $|x| < h - 1$ . Moreover we set  $G_h = \{x \in G; |x| < h\}$ .

In arriving at (3.2), we need the following lemmas.

**LEMMA A.1.** *Let  $1 < p \leq q < \infty$ ,  $t > 0$ ,  $v \in L^p(R^3; R^3) \cap L^q(R^3; R^3)$ ,  $n > 1$ , and  $u \in J^6$ . Then we have*

$$(A.1) \quad \|e^{-t\bar{A}}v\|_{\infty, R^3} \leq C t^{-3/2q} (t + 1)^{-(3/p-3/q)/2} (\|v\|_{p, R^3} + \|v\|_{q, R^3}),$$

$$(A.3) \quad \|e^{-tA}u\|_{2n, 6} \leq C (t^{-n} + 1) \|u\|_6.$$

(A.1) is deduced immediately by an elementary calculation. (A.2) is a consequence of  $L^p$ -estimates (cf. [25]) and the Sobolev embedding theorem. One can also refer to [17] for details.

LEMMA A.2 ([17, Lemmas 5.3, 5.4] and (A.2)). Let  $t > 0$ ,  $v \in J^6$ , and  $P^*$  be a certain pressure such that  $p^* = Ae^{-(t+1)A}v + \Delta e^{-(t+1)A}v$ . Then

$$\|e^{-(t+1)A}v\|_{2,6,G_h} + \|Ae^{-(t+1)A}v\|_{2,6,G_h} + \|p^*(t)\|_{3,6,G_h} \leq Ct^{-1/4}\|v\|_6.$$

LEMMA A.3 ([17, (5.18)] and (A.2)). Let  $v \in J^6$ , and  $t > 0$ . Then there is a function  $v^*$  such that

$$\begin{aligned} D \cdot v^* &= D \cdot (ge^{-(t+1)A}v), \\ \text{supp } v^*(t) &\subset \{x \in R^3; h-1 < |x| < h\}, \\ \|v^*(t)\|_{2,6} + \|(\partial/\partial t)v^*(t)\|_6 &\leq C(t+1)^{-1/4}\|v\|_6. \end{aligned}$$

LEMMA A.4. Let  $t > 0$ ,  $v$  and  $v^*$  be given in Lemma A.3. Then we have

$$\|ge^{-(t+1)A}v - v^*(t)\|_\infty \leq C(t+1)^{-1/4}\|v\|_6.$$

*Proof.* Set  $u(t) = ge^{-(t+1)A}v - v^*(t)$ ,  $u_0 = u(0)$ , and

$$\begin{aligned} F(t) &= p^*(t)Dg - 2(Dg \cdot D)e^{-(t+1)A}v - (\Delta g)e^{-(t+1)A}v \\ &\quad + \Delta v^*(t) - (\partial/\partial t)v^*(t), \end{aligned}$$

where  $p^*$  is given in Lemma A.2. By Lemmas A.2, A.3 we have that the support of  $F(t)$  is contained in  $\{x \in R^3; h-1 < |x| < h\}$ , and

$$\begin{aligned} \text{(A.3)} \quad &(t+1)^{1/4}\|F(t)\|_6 + \|u_0\|_{1,6} \leq C\|v\|_6, \\ &u_t - \Delta u + D(gp^*) = F, \quad D \cdot u = 0 \text{ in } R^3 \times (0, \infty). \end{aligned}$$

We thus rewrite  $u$  in the integral form

$$\text{(A.4)} \quad u(t) = e^{-t\bar{A}}u_0 + \int_0^t e^{-(t-s)\bar{A}}\bar{P}F(s) ds.$$

From (A.1), (A.3), and Sobolev's embedding theorem it follows that

$$\|e^{-t\bar{A}}u_0\|_{\infty, R^3} \leq C(t+1)^{-1/4}(\|u_0\|_\infty + \|u_0\|_6) \leq Ct^{-1/4}\|v\|_6,$$

and

$$\begin{aligned} & \left\| \int_0^t e^{-(t-s)\bar{A}} \bar{P} F(s) ds \right\|_{\infty, R^3} \\ & \leq C \int_0^t (t-s)^{-1/2} (t-s+1)^{-3/4} (\|F(s)\|_{3, G_h} + \|F(s)\|_{6/5, G_h}) ds \\ & \leq C \int_0^t (t-s)^{-1/2} (t-s+1)^{-3/4} \|F(s)\|_6 ds \\ & \leq C \|v\|_6 \int_0^t (t-s)^{-1/2} (t-s+1)^{-3/4} (s+1)^{-1/4} ds \\ & \leq C(t+1)^{-1/4} \|v\|_6. \end{aligned}$$

Taking (A.4) into account, we have the desired estimate and complete the proof.

*Proof of (3.2).* Let  $v \in J^6$ . By Lemmas A.1, A.2, A.3, Sobolev inequality, and Gagliardo-Nirenberg inequality (cf. [10]), we have

$$\begin{aligned} \|e^{-(t+1)A} v\|_{\infty} & \leq \|ge^{-(t+1)A} v\|_{\infty} + \|e^{-(t+1)A} v\|_{\infty, G_h} \\ & \leq \|ge^{-(t+1)A} v - v^*(t)\|_{\infty} + C\|v^*(t)\|_{1,6} \\ & \quad + C\|e^{-(t+1)A} v\|_{1,6, G_h} \\ & \leq C(t+1)^{-1/4} \|v\|_6 \quad \text{for } t > 0, \\ \|e^{-tA} v\|_{\infty} & \leq C\|e^{-tA} v\|_6^{3/4} \|e^{-tA} v\|_{2,6}^{1/4} \\ & \leq C(t^{-1} + 1)^{1/4} \|v\|_6 \leq Ct^{-1/4} \|v\|_6 \end{aligned}$$

for  $1 > t > 0$ . The proof is complete.

The author would like to thank T. Miyakawa for sending [2, 3, 4]. He would also like to thank the referee for his valuable suggestions.

REFERENCES

[1] H. Amann, *Dual semigroups and second order linear elliptic boundary value problems*, Israel J. Math., **45** (1983), 225–254.  
 [2] W. Borchers and T. Miyakawa, *Algebraic  $L^2$  decay for Navier-Stokes flows in exterior domains*, Acta Math., **165** (1990), 189–227.  
 [3] ———, *Algebraic  $L^2$  decay for Navier-Stokes flows in exterior domains II*, Hiroshima Math. J., to appear.  
 [4] ———,  *$L^2$  decay for Navier-Stokes flows in unbounded domains, with application to exterior stationary flows*, Arch. Rational Mech. Anal., **118** (1992), 273–295.  
 [5] W. Borchers and H. Sohr, *On the semigroup of the Stokes operator for exterior domains in  $L^q$ -spaces*, Math. Z., **196** (1987), 415–425.

- [6] L. Caffarelli, R. Kohn and L. Nirenberg, *Partial regularity of suitable weak solutions of the Navier-Stokes equations*, Comm. Pure Appl. Math., **35** (1982), 771–831.
- [7] Z.-M. Chen,  *$L^n$  solutions of the stationary and nonstationary Navier-Stokes equations in  $R^n$* , Pacific J. Math., to appear.
- [8] R. Finn, *On the exterior stationary problem for the Navier-Stokes equations, and associated perturbation problems*, Arch. Rational Mech. Anal., **19** (1965), 363–406.
- [9] —, *Mathematical questions relating to viscous fluid flow in an exterior domain*, Rocky Mountain J. Math., **3** (1973), 107–140.
- [10] A. Friedman, *Partial Differential Equations*, Academic Press, New York, 1969.
- [11] G. P. Galdi and S. Rionero, *Weighted Energy Methods in Fluid Dynamics and Elasticity*, Lecture Notes in Math., **1134**, Springer-Verlag, Berlin-Heidelberg-New York, 1985.
- [12] G. Galdi and C. G. Simader, *Existence, uniqueness and  $L^q$ -estimates for the Stokes problem in an exterior domain*, Arch. Rational Mech. Anal., **112** (1990), 291–318.
- [13] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer-Verlag, Berlin-Heidelberg-New York, 1983.
- [14] J. G. Heywood, *On stationary solutions of the Navier-Stokes equations as limits of nonstationary solutions*, Arch. Rational Mech. Anal., **37** (1970), 48–60.
- [15] —, *The exterior nonstationary problem for the Navier-Stokes equations*, Acta Math., **129** (1972), 11–34.
- [16] —, *The Navier-Stokes equations: on the existence, regularity and decay of solutions*, Indiana Univ. Math. J., **29** (1980), 639–681.
- [17] H. Iwashita,  *$L_q-L_r$  estimates for solutions of the nonstationary Stokes equations in an exterior domain and the Navier-Stokes initial value problem in  $L_q$  spaces*, Math. Ann., **285** (1989), 265–288.
- [18] H. Kozono and H. Sohr, *New a priori estimates for the Stokes equations in exterior domains*, Indiana Univ. Math. J., **40** (1991), 1–27.
- [19] J. Leray, *Sur le mouvement d'un liquide visqueux emplissant l'espace*, Acta Math., **63** (1934), 193–248.
- [20] K. Masuda, *On the stability of incompressible viscous fluid motions past objects*, J. Math. Soc. Japan, **27** (1975), 294–327.
- [21] —, *Weak solutions of the Navier-Stokes equations in an exterior domain*, Tohoku Math. J., **36** (1984), 623–646.
- [22] T. Miyakawa, *On nonstationary solutions of the Navier-Stokes equations in an exterior domain*, Hiroshima Math. J., **12** (1982), 115–140.
- [23] T. Miyakawa and H. Sohr, *On energy inequality, smoothness and large time behavior in  $L^2$  for weak solutions of the Navier-Stokes equations in exterior domains*, Math. Z., **199** (1988), 455–478.
- [24] M. E. Schonbek,  *$L^2$  decay for weak solutions of the Navier-Stokes equations*, Arch. Rational Mech. Anal., **88** (1985), 209–222.
- [25] V. A. Solonnikov, *Estimates for solutions of nonstationary Navier-Stokes equations*, J. Soviet Math., **8** (1977), 467–529.
- [26] R. Teman, *Navier-Stokes Equations*, North-Holland, Amsterdam, 1984.

Received October 7, 1991 and in revised form March 13, 1992.

# PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982)      F. WOLF (1904–1989)

## EDITORS

V. S. VARADARAJAN  
(Managing Editor)  
University of California  
Los Angeles, CA 90024-1555  
vsv@math.ucla.edu

F. MICHAEL CHRIST  
University of California  
Los Angeles, CA 90024-1555  
christ@math.ucla.edu

HERBERT CLEMENS  
University of Utah  
Salt Lake City, UT 84112  
clemens@math.utah.edu

THOMAS ENRIGHT  
University of California, San Diego  
La Jolla, CA 92093  
tenright@ucsd.edu

NICHOLAS ERCOLANI  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

R. FINN  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

VAUGHAN F. R. JONES  
University of California  
Berkeley, CA 94720  
vfr@math.berkeley.edu

STEVEN KERCKHOFF  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

MARTIN SCHARLEMANN  
University of California  
Santa Barbara, CA 93106  
mgscharl@henri.ucsb.edu

HAROLD STARK  
University of California, San Diego  
La Jolla, CA 92093

## SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
UNIVERSITY OF MONTANA  
UNIVERSITY OF NEVADA, RENO  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
STANFORD UNIVERSITY  
UNIVERSITY OF HAWAII  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON

---

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Julie Speckart, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the Pacific Journal of Mathematics. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 75 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$200.00 a year (10 issues). Special rate: \$100.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at P.O. Box 969, Carmel Valley, CA 93924 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Carmel Valley, California 93924, and additional mailing offices. Postmaster: send address changes to Pacific Journal of Mathematics, P.O. Box 969, Carmel Valley, CA 93924.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

This publication was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$ ,  
the American Mathematical Society's  $\mathcal{T}\mathcal{E}\mathcal{X}$  macro system.  
Copyright © 1993 by Pacific Journal of Mathematics



# PACIFIC JOURNAL OF MATHEMATICS

Volume 159    No. 2    June 1993

---

- L<sup>p</sup>*-integrability of the second order derivatives of Green potentials in convex domains 201  
VILHELM ADOLFSSON
- Solutions of the stationary and nonstationary Navier-Stokes equations in exterior domains 227  
ZHI MIN CHEN
- Mesures de Patterson-Sullivan sur le bord d'un espace hyperbolique au sens de Gromov 241  
MICHEL COORNAERT
- Differential-difference operators and monodromy representations of Hecke algebras 271  
CHARLES F. DUNKL
- Between the unitary and similarity orbits of normal operators 299  
PAUL GUINAND and LAURENT WALSH MARCOUX
- Skeins and handlebodies 337  
W. B. RAYMOND LICKORISH
- The Plancherel formula for homogeneous spaces with polynomial spectrum 351  
RONALD LESLIE LIPSMAN
- On the uniform approximation problem for the square of the Cauchy-Riemann operator 379  
JOAN MANUEL VERDERA MELENCHÓN