

*Pacific
Journal of
Mathematics*

LACUNARY STATISTICAL CONVERGENCE

JOHN ALBERT FRIDY AND CIHAN ORHAN

Volume 160 No. 1

September 1993

LACUNARY STATISTICAL CONVERGENCE

J. A. FRIDY AND C. ORHAN

The sequence x is statistically convergent to L provided that for each $\varepsilon > 0$,

$$\lim_n n^{-1} \{ \text{the number of } k \leq n : |x_k - L| \geq \varepsilon \} = 0.$$

In this paper we study a related concept of convergence in which the set $\{k : k \leq n\}$ is replaced by $\{k : k_{r-1} < k \leq k_r\}$, for some lacunary sequence $\{k_r\}$. The resulting summability method is compared to statistical convergence and other summability methods, and questions of uniqueness of the limit value are considered.

1. Introduction. A complex number sequence x is said to be *statistically convergent* to the number L if for every $\varepsilon > 0$,

$$(1) \quad \lim_n \frac{1}{n} |\{k \leq n : |x_k - L| \geq \varepsilon\}| = 0,$$

where the vertical bars indicate the number of elements in the enclosed set. In this case we write $S\text{-}\lim x = L$ or $x_k \rightarrow L(S)$. We shall also use S to denote the set of all statistically convergent sequences. The idea of statistical convergence was introduced by Fast [4] and studied by several authors [2], [3], [5], [6], [11]. There is a natural relationship [2] between statistical convergence and strong Cesàro summability:

$$|\sigma_1| := \left\{ x : \text{for some } L, \lim_n \left(\frac{1}{n} \sum_{k=1}^n |x_k - L| \right) = 0 \right\}.$$

By a *lacunary sequence* we mean an increasing integer sequence $\theta = \{k_r\}$ such that $k_0 = 0$ and $h_r := k_r - k_{r-1} \rightarrow \infty$ as $r \rightarrow \infty$. Throughout this paper the intervals determined by θ will be denoted by $I_r := (k_{r-1}, k_r]$, and the ratio k_r/k_{r-1} will be abbreviated by q_r . There is a strong connection [7] between $|\sigma_1|$ and the sequence space N_θ , which is defined by

$$N_\theta := \left\{ x : \text{for some } L, \lim_r \left(\frac{1}{h_r} \sum_{k \in I_r} |x_k - L| \right) = 0 \right\}.$$

The purpose of this paper is to introduce and study a concept of convergence that is related to statistical convergence (1) in the same way that N_θ is related to $|\sigma_1|$.

DEFINITION. Let θ be a lacunary sequence; the number sequence x is S_θ -convergent to L provided that for every $\varepsilon > 0$,

$$(2) \quad \lim_r \frac{1}{h_r} |\{k \in I_r : |x_k - L| \geq \varepsilon\}| = 0.$$

In this case we write $S_\theta\text{-}\lim x = L$ or $x_k \rightarrow L(S_\theta)$, and we define

$$S_\theta := \{x : \text{for some } L, S_\theta\text{-}\lim x = L\}.$$

The limits in (1) and (2) can be expressed using matrix transformations of the characteristic function χ_K of the set

$$K = K(x, L, \varepsilon) := \{k \in \mathbb{N} : |x_k - L| \geq \varepsilon\}.$$

The limit in (1) is $\lim_n (C_1 \chi_K)_n = 0$, where C_1 is the Cesàro mean; the limit in (2) is $\lim_n (C_\theta \chi_K)_n = 0$, where C_θ is the matrix given by

$$C_\theta[n, k] := \begin{cases} \frac{1}{h_r}, & \text{if } k \in I_r, \\ 0, & \text{if } k \notin I_r. \end{cases}$$

In this form S_θ -convergence is seen to be a part of ‘‘A-density convergence’’ as defined in [8] and [3].

In the next section we establish inclusion relations between S_θ and N_θ and also between S_θ and S . In §3 we show that the S_θ -limit of a given sequence x is not necessarily unique for different θ 's, but different S_θ -limits cannot occur if $x \in S$. In the final section we get a relationship between S_θ -convergence and strong almost convergence, a concept introduced by Maddox [10] and (independently) by Freedman et al. [7].

2. Inclusion theorems. In this section we first give some inclusion relations between N_θ - and S_θ -convergence and show that they are equivalent for bounded sequences. We also study the inclusions $S \subseteq S_\theta$ and $S_\theta \subseteq S$ under certain restrictions on $\theta = \{k_r\}$.

THEOREM 1. *Let $\theta = \{k_r\}$ be a lacunary sequence; then*

- (i) (a) $x_k \rightarrow L(N_\theta)$ implies $x_k \rightarrow L(S_\theta)$, and
- (b) N_θ is a proper subset of S_θ ;
- (ii) $x \in l_\infty$ and $x_k \rightarrow L(S_\theta)$ imply $x_k \rightarrow L(N_\theta)$;
- (iii) $S_\theta \cap l_\infty = N_\theta \cap l_\infty$,

where l_∞ denotes the set of bounded sequences.

Before proving this theorem we remark that this result is included by Theorem 8 in [3], where Connor bases the proof on the concept of ideals in l_∞ ; we give a direct proof.

Proof. (a) If $\varepsilon > 0$ and $x_k \rightarrow L(N_\theta)$ we can write

$$\sum_{k \in I_r} |x_k - L| \geq \sum_{\substack{k \in I_r \\ |x_k - L| \geq \varepsilon}} |x_k - L| \geq \varepsilon |\{k \in I_r : |x_k - L| \geq \varepsilon\}|,$$

which yields the result.

(b) In order to establish that the inclusion $N_\theta \subseteq S_\theta$ in (i) is proper, let θ be given and define x_k to be $1, 2, \dots, [\sqrt{h_r}]$ at the first $[\sqrt{h_r}]$ integers in I_r , and $x_k = 0$ otherwise. Note that x is not bounded. We have, for every $\varepsilon > 0$,

$$\frac{1}{h_r} |\{k \in I_r : |x_k - 0| \geq \varepsilon\}| = \frac{[\sqrt{h_r}]}{h_r} \rightarrow 0 \quad \text{as } r \rightarrow \infty,$$

i.e., $x_k \rightarrow 0(S_\theta)$. On the other hand,

$$\frac{1}{h_r} \sum_{k \in I_r} |x_k - 0| = \frac{1}{h_r} \frac{[\sqrt{h_r}]([\sqrt{h_r}] + 1)}{2} \rightarrow \frac{1}{2} \neq 0;$$

hence $x_k \not\rightarrow 0(N_\theta)$.

(ii) Suppose that $x_k \rightarrow L(S_\theta)$ and $x \in l_\infty$, say $|x_k - L| \leq M$ for all k . Given $\varepsilon > 0$, we get

$$\begin{aligned} \frac{1}{h_r} \sum_{k \in I_r} |x_k - L| &= \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |x_k - L| \geq \varepsilon}} |x_k - L| + \frac{1}{h_r} \sum_{\substack{k \in I_r \\ |x_k - L| < \varepsilon}} |x_k - L| \\ &\leq \frac{M}{h_r} |\{k \in I_r : |x_k - L| \geq \varepsilon\}| + \varepsilon, \end{aligned}$$

from which the result follows.

We remark that the example given in (i) shows that the boundedness condition cannot be omitted from the hypothesis of Theorem 1 (ii).

(iii) This is an immediate consequence of (i) and (ii).

Since any N_θ -summable sequence is C_θ -summable, we conclude from Theorem 1 (ii) that any bounded S_θ -summable sequence is also C_θ -summable.

LEMMA 2. *For any lacunary sequence θ , $S\text{-lim } x = L$ implies $S_\theta\text{-lim } x = L$ if and only if $\liminf_r q_r > 1$. If $\liminf_r q_r = 1$, then there exists a bounded S_θ -summable sequence that is not S -summable (to any limit).*

Proof. Suppose first that $\liminf_r q_r > 1$; then there exists a $\delta > 0$ such that $q_r \geq 1 + \delta$ for sufficiently large r , which implies that

$$\frac{h_r}{k_r} \geq \frac{\delta}{1 + \delta}.$$

If $x_k \rightarrow L(S)$, then for every $\varepsilon > 0$ and for sufficiently large r , we have

$$\begin{aligned} \frac{1}{k_r} |\{k \leq k_r : |x_k - L| \geq \varepsilon\}| &\geq \frac{1}{k_r} |\{k \in I_r : |x_k - L| \geq \varepsilon\}| \\ &\geq \frac{\delta}{1 + \delta} \cdot \frac{1}{h_r} |\{k \in I_r : |x_k - L| \geq \varepsilon\}|; \end{aligned}$$

this proves the sufficiency.

Conversely, suppose that $\liminf_r q_r = 1$. Proceeding as in [7; p. 510] we can select a subsequence $\{k_{r(j)}\}$ of the lacunary sequence θ such that

$$\frac{k_{r(j)}}{k_{r(j)-1}} < 1 + \frac{1}{j} \quad \text{and} \quad \frac{k_{r(j)-1}}{k_{r(j-1)}} > j, \quad \text{where } r(j) \geq r(j-1) + 2.$$

Now define a bounded sequence x by $x_i = 1$ if $i \in I_{r(j)}$ for some $j = 1, 2, \dots$ and $x_i = 0$ otherwise. It is shown in [7; p. 510] that $x \notin N_\theta$ but $x \in |\sigma_1|$. The above Theorem 1 (ii) implies that $x \notin S_\theta$, but it follows from Theorem 2.1 of [2] that $x \in S$. Hence $S \not\subseteq S_\theta$, and the proof is complete.

LEMMA 3. *For any lacunary sequence θ , S - $\lim x = L$ implies S_θ - $\lim x = L$ if and only if $\limsup_r q_r < \infty$. If $\limsup_r q_r = \infty$, then there exists a bounded S -summable sequence that is not S_θ -summable (to any limit).*

Proof. If $\limsup_r q_r < \infty$, then there is an $H > 0$ such that $q_r < H$ for all r . Suppose that $x_k \rightarrow L(S_\theta)$, and let $N_r := |\{k \in I_r : |x_k - L| \geq \varepsilon\}|$. By (2), given $\varepsilon > 0$, there is an $r_0 \in \mathbb{N}$ such that

$$(3) \quad \frac{N_r}{h_r} < \varepsilon \quad \text{for all } r > r_0.$$

Now let $M := \max\{N_r : 1 \leq r \leq r_0\}$ and let n be any integer satisfying

$k_{r-1} < n \leq k_r$; then we can write

$$\begin{aligned}
 \frac{1}{n} |\{k \leq n : |x_k - L| \geq \varepsilon\}| &\leq \frac{1}{k_{r-1}} |\{k \leq k_r : |x_k - L| \geq \varepsilon\}| \\
 &= \frac{1}{k_{r-1}} \{N_1 + N_2 + \cdots + N_{r_0} + N_{r_0+1} + \cdots + N_r\} \\
 &\leq \frac{M}{k_{r-1}} \cdot r_0 + \frac{1}{k_{r-1}} \left\{ h_{r_0+1} \frac{N_{r_0+1}}{h_{r_0+1}} + \cdots + h_r \frac{N_r}{h_r} \right\} \\
 &\leq \frac{r_0 \cdot M}{k_{r-1}} + \frac{1}{k_{r-1}} \left(\sup_{r > r_0} \frac{N_r}{h_r} \right) \{h_{r_0+1} + \cdots + h_r\} \\
 &\leq \frac{r_0 \cdot M}{k_{r-1}} + \varepsilon \cdot \frac{k_r - k_{r_0}}{k_{r-1}}, \quad \text{by (3),} \\
 &\leq \frac{r_0 \cdot M}{k_{r-1}} + \varepsilon \cdot q_r \leq \frac{r_0 \cdot M}{k_{r-1}} + \varepsilon H,
 \end{aligned}$$

and the sufficiency follows immediately.

Conversely, suppose that $\limsup_r q_r = \infty$. Following the idea in [7; p. 511] we can select a subsequence $\{k_{r(j)}\}$ of the lacunary sequence $\theta = \{k_r\}$ such that $q_{r(j)} > j$, and define a bounded sequence by $x_i = 1$ if $k_{r(j)-1} < i \leq 2k_{r(j)-1}$ for some $j = 1, 2, \dots$, and $x_i = 0$ otherwise. It is shown in [7; p. 5.11] that $x \in N_\theta$ but $x \notin |\sigma_1|$. By Theorem 1 (i) we conclude that $x \in S_\theta$, but Theorem 2.1 of [2] implies that $x \notin S$. Hence, $S_\theta \not\subseteq S$.

Combining Lemma 2 and Lemma 3 we get

THEOREM 4. *Let θ be a lacunary sequence; then $S = S_\theta$ if and only if*

$$1 < \liminf_r q_r \leq \limsup_r q_r < \infty;$$

then S -lim $x = L$ implies S_θ -lim $x = L$.

For an example of a lacunary sequence satisfying the conditions of Theorem 4, we can take $k_r = 2^r$ for $r > 0$, whence $S_{\{2^r\}} = S$. We remark that the examples given in Lemmas 2 and 3 illustrate the difference between S -convergence and S_θ -convergence.

We conclude this section with the following observation. Buck [1, Theorem 3.2] proved that if a real sequence is C_1 -summable to its finite limit inferior, then the sequence “converges to that point for almost all n ” (i.e., it is statistically convergent to its limit inferior [2]). Note that this result remains true if we replace limit inferior by

limit superior. For each subset K of \mathbb{N} , define

$$D(K) := \lim_r (C_\theta \chi_K)_r = \lim_r \frac{|K \cap I_r|}{h_r};$$

then D is a density [8; p. 296], and it is not hard to get a result for S_θ -convergence that is analogous to Buck's. To be precise, the following result is such an analogue.

PROPOSITION 5. *If the real number sequence x is C_θ -summable to either its finite limit inferior or finite limit superior, then x is S_θ -convergent to that value.*

3. Uniqueness of S_θ -limit and lacunary refinements. It is easy to see that, for any fixed θ , the S_θ -limit is unique. It is possible, however, for a sequence—even a bounded one—to have different S_θ -limits for different θ 's. This can be seen by applying Theorem 1 (i) to the sequence x given in [7, proof of Theorem 2.1] for which $N_{\theta_1}\text{-lim } x = 0$ and $N_{\theta_2}\text{-lim } x = 1$. The next theorem shows that this situation cannot occur if $x \in S$; in other words, every S_θ method is consistent with the S -method.

THEOREM 6. *If $x \in S \cap S_\theta$, then $S_\theta\text{-lim } x = S\text{-lim } x$.*

Proof. Suppose $S\text{-lim } x = L$ and $S_\theta\text{-lim } x = L'$, and $L \neq L'$. For $\varepsilon < \frac{1}{2}|L - L'|$ we get

$$\lim_n \frac{1}{n} |\{k \leq n : |x_k - L'| \geq \varepsilon\}| = 1.$$

Consider the k_m th term of the statistical limit expression $n^{-1}|\{k \leq n : |x_k - L'| \geq \varepsilon\}|$:

$$(4) \quad \frac{1}{k_m} \left| \left\{ k \in \bigcup_{r=1}^m I_r : |x_k - L'| \geq \varepsilon \right\} \right| \\ = \frac{1}{k_m} \sum_{r=1}^m |\{k \in I_r : |x_k - L'| \geq \varepsilon\}| = \frac{1}{\sum_{r=1}^m h_r} \sum_{r=1}^m h_r t_r,$$

where $t_r = h_r^{-1}|\{k \in I_r : |x_k - L'| \geq \varepsilon\}| \rightarrow 0$ because $x_k \rightarrow L'(S_\theta)$. Since θ is a lacunary sequence, (4) is a regular weighted mean transform of t , and therefore it, too, tends to zero as $m \rightarrow \infty$. Also, since this is a subsequence of $\{n^{-1}|\{k \leq n : |x_k - L'| \geq \varepsilon\}|\}_{n=1}^\infty$, we infer that

$$\frac{1}{n} |\{k \leq n : |x_k - L'| \geq \varepsilon\}| \rightarrow 1,$$

and this contradiction shows that we cannot have $L \neq L'$.

We now consider the inclusion of $S_{\theta'}$ by S_{θ} , where θ' is a lacunary refinement of θ . Recall [7] that the lacunary sequence $\theta' = \{k'_r\}$ is called a *lacunary refinement* of the lacunary sequence $\theta = \{k_r\}$ if $\{k_r\} \subseteq \{k'_r\}$.

THEOREM 7. *If θ' is a lacunary refinement of θ and $x_k \rightarrow L(S_{\theta'})$, then $x_k \rightarrow L(S_{\theta})$.*

Proof. Suppose each I_r of θ contains the points $\{k'_{r,i}\}_{i=1}^{\nu(r)}$ of θ' so that

$$k_{r-1} < k'_{r,1} < k'_{r,2} < \dots < k'_{r,\nu(r)} = k_r, \quad \text{where } I'_{r,i} = (k'_{r,i-1}, k'_{r,i}].$$

Note that for all r , $\nu(r) \geq 1$ because $\{k_r\} \subseteq \{k'_r\}$. Let $\{I_j^*\}_{j=1}^{\infty}$ be the sequence of abutting intervals $\{I'_{r,i}\}$ ordered by increasing right end points. Since $x_k \rightarrow L(S_{\theta'})$, we get, for each $\varepsilon > 0$,

$$(5) \quad \lim_j \sum_{I_j^* \subset I_r} \frac{1}{h_r^*} |\{k \in I_j^* : |x_k - L| \geq \varepsilon\}| = 0.$$

As before we write, $h_r = k_r - k_{r-1}$, $h'_{r,i} = k'_{r,i} - k'_{r,i-1}$, and $h'_{r,1} = k'_{r,1} - k_{r-1}$. For each $\varepsilon > 0$ we have

$$(6) \quad \begin{aligned} & \frac{1}{h_r} |\{k \in I_r : |x_k - L| \geq \varepsilon\}| \\ &= \frac{1}{h_r} \sum_{I_j^* \subset I_r} h_j^* \frac{1}{h_j^*} |\{k \in I_j^* : |x_k - L| \geq \varepsilon\}| \\ &= \frac{1}{h_r} \sum_{I_j^* \subset I_r} h_j^* (C_{\theta'} \chi_K)_j, \end{aligned}$$

where χ_K is the characteristic function of the set $K := \{k \in \mathbb{N} : |x_k - L| \geq \varepsilon\}$. By (5), $C_{\theta'} \chi_K$ is a null sequence, and (6) is a regular weighted mean transform of $C_{\theta'} \chi_K$. Hence, the transform (6) also tends to zero as $r \rightarrow \infty$.

We conclude this section by observing that Theorem 7 establishes inclusion between two lacunary methods *only* when one sequence is a lacunary refinement of the other. The example cited at the beginning of this section shows that S_{θ} can be inconsistent with $S_{\theta'}$. A general description of inclusion between two arbitrary lacunary methods is left as an open problem.

4. Strong almost convergence and S_θ -convergence. The idea of almost convergence was introduced by Lorentz [9]: the sequence x is said to be *almost convergent* to L if

$$\lim_n \frac{1}{n} \sum_{i=m+1}^{m+n} (x_i - L) = 0, \quad \text{uniformly in } m.$$

Maddox [10] and (independently) Freedman et al. [7] introduced the notion of strong almost convergence: the sequence x is said to be *strongly almost convergent* to L if

$$\lim_n \frac{1}{n} \sum_{i=m+1}^{m+n} |x_i - L| = 0, \quad \text{uniformly in } m.$$

Let c , AC and $[AC]$, respectively, denote the sets of all convergent, almost convergent, and strongly almost convergent sequences. It is known [10] that

$$(7) \quad c \subsetneq [AC] \subsetneq AC \subsetneq l_\infty.$$

THEOREM 8. *If \mathcal{L} denotes the set of all lacunary sequences, then*

$$[AC] = l_\infty \cap \left(\bigcap_{\theta \in \mathcal{L}} S_\theta \right).$$

Proof. By [7, Theorem 3.1], the relations (7) and Theorem 1 (iii), we have

$$\begin{aligned} l_\infty \supset [AC] &= \bigcap_{\theta \in \mathcal{L}} N_\theta = l_\infty \cap \left(\bigcap_{\theta \in \mathcal{L}} N_\theta \right) \bigcap_{\theta \in \mathcal{L}} (l_\infty \cap N_\theta) \\ &= \bigcap_{\theta \in \mathcal{L}} (l_\infty \cap S_\theta) = l_\infty \cap \left(\bigcap_{\theta \in \mathcal{L}} S_\theta \right). \end{aligned}$$

Finally we remark that in contrast to [7, Theorem 3.1] where it was proved that $[AC] = \bigcap N_\theta$, the factor l_∞ cannot be omitted from Theorem 8. For, $\bigcap S_\theta \not\subset l_\infty$ and $\bigcap N_\theta = [AC]$ is a proper subset of $\bigcap S_\theta$. To see this consider the sequence x defined by $x_k = m$, if $k = m^2$ for $m = 1, 2, \dots$, and $x_k = 0$ otherwise. Observe that x is not bounded, so it is not strongly almost convergent. On the other hand, for any lacunary sequence θ , we have

$$\frac{1}{h_r} |\{k \in I_r : x_k \neq 0\}| \leq \frac{\sqrt{h_r}}{h_r} \rightarrow 0, \quad \text{as } r \rightarrow \infty;$$

hence, $x_k \rightarrow O(S_\theta)$.

The authors wish to thank the referee for several very helpful suggestions that have improved the exposition of these results.

REFERENCES

- [1] R. C. Buck, *Generalized asymptotic density*, Amer. J. Math., **75** (1953), 335–346.
- [2] J. Connor, *The statistical and strong p -Cesàro convergence of sequences*, Analysis, **8** (1988), 47–63.
- [3] ———, *On strong matrix summability with respect to a modulus and statistical convergence*, Canad. Math. Bull., **32** (1989), 194–198.
- [4] H. Fast, *Sur la convergence statistique*, Colloq. Math., **2** (1951), 241–244.
- [5] J. A. Fridy, *On statistical convergence*, Analysis, **5** (1985), 301–313.
- [6] J. A. Fridy and H. I. Miller, *A matrix characterization of statistical convergence*, Analysis, **11** (1991), 55–66.
- [7] A. R. Freedman, J. J. Sember, and M. Raphael, *Some Cesàro type summability spaces*, Proc. London Math. Soc., **37** (1978), 508–520.
- [8] A. R. Freedman and J. J. Sember, *Densities and summability*, Pacific J. Math., **95** (1981), 293–305.
- [9] G. G. Lorentz, *A contribution to the theory of divergent sequences*, Acta Math., **80** (1948), 167–190.
- [10] I. J. Maddox, *A new type of convergence*, Math. Proc. Cambridge Phil. Soc., **83** (1978), 61–64.
- [11] I. J. Schoenberg, *The integrability of certain functions and related summability methods*, Amer. Math. Monthly, **66** (1959), 361–375.

Received November 11, 1990 and in revised form March 16, 1992. The second author's research was supported by the Scientific and Technical Research Council of Turkey.

KENT STATE UNIVERSITY
KENT, OH 44242
U.S.A.

AND

ANKARA UNIVERSITY
ANKARA, 06100
TURKEY

CONTENTS

G. D. Anderson, M. K. Vamanamurthy, and M. Vuorinen , Inequalities for quasi-conformal mappings in space	1
T. Bhattacharya , A nonexistence result for the n -Laplacian	19
J. A. Cima, K. Stroethoff, and K. Yale , Bourgain algebras on the unit disk	27
J. A. Fridy and C. Orhan , Lacunary statistical convergence	43
D. Grenier , On the shape of fundamental domains in $GL(n, \mathbf{R})/O(n)$	53
B. Jiang and J. Guo , Fixed points of surface diffeomorphisms	67
P. Lejarraga , The moduli of rational Weierstrass fibrations over \mathbf{P}^1 : singularities	91
G. J. Martin , On discrete isometry groups of negative curvature	109
T. Nakashima , Adjoint linear systems on a surface of general type in positive characteristic	129
B. Ralph , A homotopy transfer for finite group actions	133
Y. Rong , Maps between Seifert fibered spaces of infinite π_1	143
J.-Y. Shi , Some numeric results on root systems	155
E. Spanier , Singular homology and cohomology with local coefficients and duality for manifolds	165

PACIFIC JOURNAL OF MATHEMATICS

Volume 160 No. 1 September 1993

Inequalities for quasiconformal mappings in space	1
GLEN DOUGLAS ANDERSON, MAVINA KRISHNA VAMANAMURTHY and MATTI VUORINEN	
A nonexistence result for the n -Laplacian	19
TILAK BHATTACHARYA	
Bourgain algebras on the unit disk	27
JOSEPH A. CIMA, KAREL M. STROETHOFF and KEITH YALE	
Lacunary statistical convergence	43
JOHN ALBERT FRIDY and CIHAN ORHAN	
On the shape of fundamental domains in $GL(n, \mathbf{R})/O(n)$	53
DOUGLAS MARTIN GRENIER	
Fixed points of surface diffeomorphisms	67
BOJU JIANG and JIANHAN GUO	
The moduli of rational Weierstrass fibrations over \mathbf{P}^1 : singularities	91
PABLO LEJARRAGA	
On discrete isometry groups of negative curvature	109
GAVEN MARTIN	
Adjoint linear systems on a surface of general type in positive characteristic	129
TOHRU NAKASHIMA	
A homotopy transfer for finite group actions	133
WILLIAM J. RALPH	
Maps between Seifert fibered spaces of infinite π_1	143
YONGWU RONG	
Some numeric results on root systems	155
J. Y. SHI	
Singular homology and cohomology with local coefficients and duality for manifolds	165
EDWIN SPANIER	