

Pacific Journal of Mathematics

PRODUCTIVE POLYNOMIALS

RICHARD ARENS

Volume 162 No. 2

February 1994

PRODUCTIVE POLYNOMIALS

RICHARD F. ARENS

The problem addressed is: When is a class B of polynomials in n non-commuting indeterminates closed under substitution into a given polynomial q ?

1. Introduction. Let \mathbb{F} be a field and let $\mathbb{F}\langle x_1, \dots, x_n \rangle$ be the linear algebra of polynomials in the non-commuting indeterminates x_1, \dots, x_n . Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let A be an associative¹ algebra over \mathbb{F} . q defines a mapping \hat{q} of $A \times \dots \times A = A^n$ into A whose value $\hat{q}(a_1, \dots, a_n)$ at (a_1, \dots, a_n) is the result of replacing each x_i in q by the corresponding a_i , and then carrying out the algebraic operations proper to A . A linear subspace B of the algebra A will be called q -closed if whenever $\mathbf{a} = (a_1, \dots, a_n) \in A^n$ then $\hat{q}(\mathbf{a}) \in B$. Let $q((B))$ be the smallest q -closed linear subspace containing B . We study mainly the case that A is $\mathbb{F}\langle x_1, \dots, x_n \rangle$ itself, and B is the linear subspace generated by x_1, \dots, x_n and the unit 1. The q -closed set generated by x_1, \dots, x_n and 1 will be denoted in this case simply by $((q))$.

We will usually use just P to stand for $\mathbb{F}\langle x_1, \dots, x_n \rangle$. $q \in P$ will be called *productive* if $((q)) = P$; and otherwise, *non-productive*.

Two questions interest us:

1.1. When is a given $q \in P$ productive,
and

1.2. If it is not, how to find elements p which are not in $((q))$?

A clear-cut answer to 1.1 is given by 3.9. An answer to 1.2 is given in §4, illustrated by an example 8.5. We regard q as an n -ary operation and prepare a suitable ideal theory.

2. Theorems establishing productivity. Consider $q = x_1x_2$. Then a linear subspace B is q -closed if it contains the product of any pair of members: B is a subalgebra.² Thus, if B is the linear subspace generated by x_1, \dots, x_n and 1, then $((q))$ is the algebra generated by x_1, \dots, x_n and 1. This being P , x_1x_2 is productive.

¹In this paper, all algebras are supposed to be associative, and so we omit the term.

²Note that B is x_1x_2 -closed if and only if it is x_ix_j -closed, where i, j are any two distinct indices.

We will usually write x, y for x_1, x_2 . The main example of a productive element is xy .³ As observed, $((xy))$ is the subalgebra generated by x_1, \dots, x_n and the unit 1, and this is patently P .

For finding sufficient conditions for elements to be productive, we are aided by the following concept. Let $E = (e(1), \dots, e(n))$ be n non-negative integers. Let $p \in P$. Then p is *homogeneous of type E* if it is homogeneous of degree $e(i)$ in x_i , for each i .

2.1. DEFINITION. Given $q \in P$, its *homogeneous constituent q_E* shall be the sum of its monomial terms homogeneous of type E .

Obviously, q is the sum of its homogeneous constituents.

2.2. LEMMA. *If the field \mathbb{F} is infinite, then $q_E \in ((q))$ for each homogeneity type E .*

Proof. Let $\lambda = (\lambda_1, \dots, \lambda_n)$, and define $q\lambda$ as $\hat{q}(\lambda_1 x_1, \dots, \lambda_n x_n)$. We have $q\lambda = \sum q_E \lambda = \sum \lambda^E q_E$. By making enough different choices for λ , we get a system of linear equations for the various q_E . These can then be expressed in terms of the left-hand sides. The left-hand sides belongs to $((q))$ no matter how the λ are selected. Therefore, so do the q_E .

2.3. THEOREM. *Let $q \in P$. If $p \in ((q))$ then $((q))$ is p -closed and $((p))$ is contained in $((q))$.*

This can be deduced from the following representation of $q((C_1))$ as the union of sets C_m where C_1 is a subset of P . We will define C_m for $m = 2, 3, \dots$. Suppose now that C_m has been defined. Then C_{m+1} shall consist of the elements of C_m , their linear combinations, and the values of $q(\mathbf{u})$ where \mathbf{u} varies over the n -member subsets of C_m . Let C^∞ be the union of the sets C_m . Using induction, we see that each C_m lies in $q((C_1))$, so C^∞ is contained in $q((C_1))$. Since C^∞ is obviously q -closed, it must include $q((C_1))$. Thus

$$(2.31) \quad q((C_1)) = C^\infty.$$

Such a sequence of sets $\{C_m\}$ may be called a q -system. There is a q -system Q with $C_1 = B$, the span of 1 and x_1, \dots, x_n . Now suppose p lies in the set C_m and let r_1, \dots, r_n lie in C_k . We want to show that $\hat{p}(r_1, \dots, r_n)$ lies in $((q))$.

In the system Q replace each x_i by r_i , giving Q' . This is a q -system $\{C'_m\}$ and it has $\hat{p}(r_1, \dots, r_n)$ in the m th set. C'_1 is

³See also 2.4.

contained in C_k . Consequently C'_m is contained in C_{k+m-1} , so $\hat{p}(r_1, \dots, r_n)$ lies in $((q))$. This shows that $((q))$ is p -closed, whence the p -closure of B , namely $((p))$ is contained in $((q))$.

Thus 2.3 is proved. We mention an immediate generalization.

COROLLARY. *Let $q \in P$. Let C be a subset of P . Let $p \in q((C))$. Then $p((C))$ is contained in $q((C))$.*

2.4. COROLLARY. *Let $q \in P$. Then $P = ((q))$ if and only if $xy \in ((q))$.*

This follows from 2.3 with $p = xy$.

2.5. COROLLARY. *Let $q \in P$. If any q_E is productive⁴ then q is productive.*

Proof. $((q_E))$ is contained in $((q))$, so if the former contains xy , so does the latter.

The next proposition can be as easily proved as was 2.4.

2.6. PROPOSITION. *Let $q \in P$. Select an index i . Define $q_i(x_1, \dots, x_n)$ as the coefficient of t in $q(x_1, \dots, x_i + t1, \dots, x_n)$ where t ranges over \mathbb{F} and 1 is the unit of $\mathbb{F}\langle x_1, \dots, x_n \rangle$. Then q_i belongs to $((q))$.*

This can be used to show that $q = x^2y + xyx + xy^2$ is productive. First we get $x^2y + xyx \in ((q))$ by 2.2. Then 2.6 tells us that $2xy + yx + xy \in ((q))$, whence $3xy + yx \in ((q))$, and by substitution of x for y and y for x , $3yx + xy \in ((q))$. Linear combination gives us xy .

COROLLARY. *If f and g have positive degree, then $f(x)g(y)$ is productive.*

Proof. By differentiating an appropriate number⁵ of times and applying 2.6, we can obtain $xy \in ((f(x)g(y)))$. By 2.3, $((f(x)g(y)))$ includes $((xy))$, $= \mathbb{F}\langle x_1, \dots, x_n \rangle$.

The connection with Jordan algebras may be noted. (See [BK].)

⁴However, if all the q_E are non-productive, the sum k may still be productive. See 4.5.

⁵Using 2.3 after each differentiation.

REMARK. If q is a polynomial in one variable and has degree at least 2, then $((q)) = ((x^2))$, i.e., it is the special Jordan algebra $\mathbb{F}\langle x_1, \dots, x_n \rangle^+$.

Proof. Let q be a polynomial in x . Replace x by $1+x$ and expand. One sees from 2.2 that $x^2 \in ((q))$. On the other hand, replace x by x^2 and see that $x^4, x^8, x^{16}, \dots \in ((x^2))$. Keep this up until the exponent exceeds the degree of q . Replacing x by $1+x$ and using 2.2, one sees that all the powers in q are in $((x^2))$. Forming linear combinations, one obtains $q \in ((x^2))$. Thus, by 2.3 applied twice, $((q)) = ((x^2))$.

3. Characterizing productivity. For $k = 1, 2, \dots$ let P_k stand for the set of homogeneous polynomials of degree k . For $p \in P$ let $p_{,k}$ be the $P_0 + \dots + P_k$ component of p . We will use $\text{Quad}(p)$ to denote the P_2 component of p .

We are going to be dealing with n -tuples (p_1, \dots, p_n) of elements of P . We will abbreviate such an n -tuple by \mathbf{p} . We will write $\mathbf{p}_{,k}$ for $(p_{1,k}, \dots, p_{n,k})$.

We will also make use of the linear automorphism $p \rightarrow p^*$ of P defined inductively by $1^* = 1, (px_i)^* = x_i p^*$. An element u for which $u^* = u$ will be called symmetric, and one for which $u^* = -u$, skew.

We will let S_+ denote the linear subspace of P of those elements p for which $\text{Quad}(p)$ is symmetric, and S_- for those for which it is skew. In fact p is in S_+ or S_- if and only if $p_{,2}$ is symmetric or skew, respectively.

PROPOSITION. *Let $q \in P$. Then one of the three sentences 3.1, 3.2, or 3.3 must be true.*

3.1. *If f_1, \dots, f_n belong to $P_0 + P_1$, then $\text{Quad } q(\mathbf{f})$ is symmetric.*

3.2. *If f_1, \dots, f_n belong to $P_0 + P_1$, then $\text{Quad } q(\mathbf{f})$ is skew.*

3.3. *Both of the following hold:*

3.31. *There are f_1, \dots, f_n in $P_0 + P_1$, and $\text{Quad } q(\mathbf{f})$ is not symmetric.*

3.32. *There are f_1, \dots, f_n in $P_0 + P_1$, and $\text{Quad } q(\mathbf{f})$ is not skew.*

We pass on to consider the consequences of each of these sentences or conditions.

LEMMA. *Condition 3.1 holds \Leftrightarrow*

3.4. *$\text{Quad}(q(\lambda + \mathbf{x}))$ is symmetric for all $\lambda = (\lambda_1, \dots, \lambda_n)$.*

Proof. \Rightarrow : Choose $f_i = \lambda_i + x_i$. By 3.1, $\text{Quad}(q(\lambda + \mathbf{x}))$ is symmetric. \Leftarrow : Suppose $\text{Quad}(q(\lambda + \mathbf{x})) = \sum \alpha_{ij} x_i x_j$. This forces α_{ij} to be symmetric as a matrix, which makes $\sum \alpha_{ij} X_i X_j$ symmetric when the X_i are any linear forms. So $\text{Quad}(q(\lambda + \mathbf{x})) = \sum \alpha_{ij} X_i X_j$ is symmetric as 3.1 requires.

In the same way one can show

LEMMA. Condition 3.2 holds \Leftrightarrow

3.5. $\text{Quad}(q(\lambda + \mathbf{x}))$ is skew for all $\lambda = (\lambda_1, \dots, \lambda_n)$.

3.6. LEMMA. If 3.1 holds, then $p_1, \dots, p_n \in S_+$ implies that $q(p_1, \dots, p_n) \in S_+$.

Proof. $\text{Quad } q(\mathbf{p})$ is identical to $\text{Quad}(q(\mathbf{p}, 2))$. Now $\text{Quad}(q(\mathbf{p}, 2)) = \text{Quad}(q(\mathbf{p}, 1))$ plus a linear combination of terms $p_{i,2}$. Because $p_1, \dots, p_n \in S_+$, these extra terms are symmetric. By 3.1, $\text{Quad}(q(\mathbf{p}, 1))$ is symmetric, whence $\text{Quad}(q(\mathbf{p}))$ is symmetric, and $q(p_1, \dots, p_n) \in S_+$.

3.7. LEMMA. If 3.2 holds, then $p_1, \dots, p_n \in S_-$ implies that $q(p_1, \dots, p_n) \in S_-$.

A proof may be obtained by replacing “symmetric” by “skew”.

3.71. LEMMA. If 3.31 holds, then $x_1 x_2 - x_2 x_1$ belongs to $((q))$.

Proof. From 2.2, we see that $\text{Quad}(q(\mathbf{f}))$ belongs to $((q))$. Thus there is an element $p = \sum \alpha_{ij} x_i x_j$ in $((q))$ such that $p - p^* \neq 0$. This leads to an element $\sum \alpha_{ij} (x_i x_j - x_j x_i)$ in $((q))$ and not 0. By 2.2 again, some $x_i x_j - x_j x_i$ is in $((q))$. Permuting the x_k gives us 3.71.

3.72. LEMMA. If 3.32 holds, then $x_1 x_2 + x_2 x_1$ belongs to $((q))$.

A proof can be assembled from the preceding, except that we might arrive at the element $x_1 x_1 + x_1 x_1$. From here it is easy to get to $x_1 x_2 + x_2 x_1$ by polarization. We note an obvious consequence.

3.8. LEMMA. If 3.31 and 3.32 both hold then $x_1 x_2$ belongs to $((q))$.

3.9. THEOREM. *Let $q \in P$. If 3.4 holds then $((q))$ is included in S_+ and q is nonproductive. If 3.5 holds then $((q))$ is included in S_- and q is non-productive. If neither of these hold, then q is productive.*

Proof. Suppose 3.4 holds. Therefore, 3.1 holds, so by 3.6, S_+ is invariant under q . Now $1, x_1, \dots, x_n$ are symmetric and thus in S_+ . Therefore, $((q))$ is included in S_+ . Obviously S_+ is not all of P .

Supposing 3.5 holds, we proceed in an analogous fashion, and conclude that q is non-productive.

If neither 3.4 nor 3.5 holds then we obtain 3.31 and 3.32. We appeal to 3.9 and to 2.4, and conclude that q is productive.

4. Examples.

4.1. THEOREM. *Suppose either $\alpha_{ij} = \alpha_{ji}$ for all i, j or $\alpha_{ij} = -\alpha_{ji}$ for all i, j . Then, $\sum \alpha_{ij} x_i x_j$ is non-productive, and conversely.*

Proof. Suppose $\alpha_{ij} = \alpha_{ji}$ for all i, j . In $q = \sum \alpha_{ij} x_i x_j$ replace x_i by $\lambda_i + x_i$ where λ_i is a scalar. Obviously $\text{Quad}(\lambda + \mathbf{x})$ is q itself, which is surely symmetric when $\alpha_{ij} = \alpha_{ji}$. The case $\alpha_{ij} = -\alpha_{ji}$ is treated in a like manner. By 3.9, q is nonproductive.

If neither $\alpha_{ij} = \alpha_{ji}$ for all i, j or $\alpha_{ij} = -\alpha_{ji}$ for all i, j then neither 3.4 nor 3.5 holds. This completes the proof of 4.1

4.2. THEOREM. *Let $q = x^i y^j x^k$, where the exponents are all positive. Here x is x_1 and y is x_2 . Then q is non-productive precisely when $i = k$.*

Proof. Suppose 3.4 is true. Let the $\lambda_i = 1$. It is verifiable that $\text{Quad}((1+x)^i(1+y)^j(1+x)^k) = x^2(ik+i'+k') + xyij + yxjk + y^2j'$, where i' means $i(i-1)/2$, etc. So $j = k$, and conversely, if $j = k$ then $\text{Quad}(q(\lambda + \mathbf{x}))$ is symmetric. But we have not yet shown that $j = k$ implies $\text{Quad}(q(\lambda + \mathbf{x}))$ is symmetric even when some λ_i is 0. In this case we have to examine $\text{Quad}(x^i(\mu + y)^j x^k)$ which is either $\mu^j x^{i+k}$ or 0, in any case, symmetric.

This theorem implies that $x^i y^j x^i$ is non-productive. Observe that $x^i y^j x^i$ is symmetric. We indicate another way to see this.

4.3. THEOREM. *If q is symmetric, then every element of $((q))$ is symmetric and thus q is non-productive.*

Proof. If f_1, \dots, f_n are symmetric and q is symmetric, then $q(f_1, \dots, f_n)$ is symmetric. Now 1, and the x_i are symmetric, so every element of $((q))$ is symmetric.

The converse is not true. Examples abound, by virtue of the next theorem.

4.4. THEOREM. *Let $q = x^h y^i x^j y^k$. Then q is non-productive precisely when*

$$(4.41) \quad hk + hi + ji = ij.$$

Proof. This condition is necessary for 3.4 and 3.5, as can be seen by putting $\lambda = (1, \dots, 1)$. Then it can be shown sufficient for 3.4, in case $\lambda = (\lambda_1, \dots, \lambda_n)$ when no λ_i is 0, and finally when some are 0.

4.5. EXAMPLE (see 2.5). One can have q productive but every q_E non-productive.

Let $q_1 = xyx$. This is non-productive by 4.2. In the proof we showed that it generates $xy + yx$. Let $q_2 = xy - yx$. This is non-productive by 4.1. Let q be their sum. From q we can get q_1 and q_2 back, by 2.2. Hence we can get $xy + yx$ and $xy - yx$, whence by 2.4, q is productive.

5. Remarks about the rest of this paper. We want to present some ideas which will enable us to assert, for example, that xyx does not belong to $((xy^2))$. See 8.5 below. To establish such propositions we apparently have to bring up the concept of *seminorms*, which is familiar, and that of *ideals*, which may not be familiar in this context.

6. Seminorms. From now on, \mathbb{F} will be either the real field \mathbb{R} or the complex field \mathbb{C} .

DEFINITION. Suppose A is a linear algebra over \mathbb{F} . Suppose that a real-valued function S defined on A is a seminorm with respect to the linear space structure of A . A seminorm defines a topology in A , and thus also in $A \times \dots \times A$. It makes sense to ask whether this makes \hat{q} continuous at any selected point of $A \times \dots \times A$. If so, we will say that \hat{q} is *continuous in the topologies defined by S* , or that S renders \hat{q} continuous⁶ at that point.

⁶We will abbreviate this to “ q is continuous in the topology defined by S ”. If no particular point is mentioned, continuity at all points is implied.

6.1. LEMMA. *Suppose $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let A be an algebra over \mathbb{F} . Let S be a seminorm on A . Suppose q is continuous in the topology defined by S . If $p \in ((q))$, the q -closed subset of $\mathbb{F}\langle x_1, \dots, x_n \rangle$ generated by $\{1, x_1, \dots, x_n\}$ then p is continuous in the topology defined by S .*

Proof. Construct the q -system $\{C_i\}$ as in 2.3, with C_1 being the set $\{1, x_1, \dots, x_n\}$. The union of the C_i is $((q))$. Every element of C_1 is certainly continuous in the topology defined by S . Assume it is true for every element of C_m . Each element of C_{m+1} is either a linear combination of these, or the value of \hat{q} on n of them, and thus surely also continuous.

We will actually need a slight but immediate corollary of this.

6.2. COROLLARY. *Suppose $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let A be an algebra over \mathbb{F} . Let S be a seminorm on A . Suppose q has 0 constant term and is continuous at $(0, \dots, 0)$ in the topology defined by S . Suppose $p \in ((q))$, and suppose that p has constant term 0. Then p is continuous at $(0, \dots, 0)$ in the topology defined by S .*

This result will enable us to exhibit some polynomials p and q where $p \notin ((q))$. We just have to find a seminorm such that \hat{p} is not continuous. To do that we develop an appropriate ideal theory.

7. q -ideals.

DEFINITION. Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let A be an algebra. Then a linear subspace J of A will be called a q -ideal if whenever $\mathbf{a} = (a_1, \dots, a_n) \in A^n$ and $\mathbf{j} \in J^n$, then $\hat{q}(\mathbf{a} + \mathbf{j}) - \hat{q}(\mathbf{a}) \in J$.

One example is $q = x_1 x_2$. Here a q -ideal is just an ordinary two-sided ideal of the algebra A . Another example is $q = (x_1)^2$. In this case a q -ideal is an ideal of the Jordan algebra A^+ (see [BK]).

LEMMA. *Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let A be an algebra. Let J and K be q -ideals. Then $J \cap K$ and $J + K$ are q -ideals.*

Proof. The assertion about $J \cap K$ is elementary. As to the other, $\hat{q}(\mathbf{a} + \mathbf{j} + \mathbf{k}) - \hat{q}(\mathbf{a}) = [\hat{q}(\mathbf{a} + \mathbf{j} + \mathbf{k}) - \hat{q}(\mathbf{a} + \mathbf{j})] + [\hat{q}(\mathbf{a} + \mathbf{j}) - \hat{q}(\mathbf{a})]$. The first bracket belongs to K and the second to J . Thus the lemma is proved.

7.1. DEFINITION. The smallest q -ideal containing a given subset B of A is the q -ideal generated by B and will be denoted by $I_q(B)$.

The next lemma is helpful in discovering what elements belong to $I_q(B)$ in specific cases.

DEFINITION. Let q be an element of $\mathbb{F}\langle x_1, \dots, x_n \rangle$. Let J be a linear subspace of a linear algebra A . Let $(a_1, \dots, a_n) \in A^n$ and let $(j_1, \dots, j_n) \in J^n$. For each i , $j \leq 1 \leq n$, let

$$(7.2) \quad q_i(a_1, \dots, a_n, j_1, \dots, j_n) \\ = \frac{d}{dt} \hat{q}(a_1, \dots, a_i + t j_i, \dots, a_n) |_{t=0}.$$

LEMMA. A linear subspace J is a q -ideal if and only if

$$(7.3) \quad q_i(a_1, \dots, a_n, j_1, \dots, j_n) \in J$$

whenever $a_1, \dots, a_n \in A$ and $j_1, \dots, j_n \in J$.

Proof. It seems adequate to us to give a proof only for a special example, say $q = xyx$. Here we write x for x_1 and y for x_2 and $n = 2$. Also let $a_1 = a$, $a_2 = b$, $j_1 = j$, and $j_2 = k$. In this case of $q = xyx$, conditions 7.3 say

$$(7.4) \quad jba + abj \text{ and } aka \in J \text{ whenever } j, k \in J \text{ and } a, b \in A.$$

We have to show that $(a + j)(b + k)(a + j) - aba \in J$ whenever $j, k \in J$ and $a, b \in A$, if and only if 7.4 holds.

If $(a + sj)(b + tk)(a + sj) - aba$ lies in J for all s and t , we can deduce, by algebraic procedures, that the derivatives in 7.4, that is the derivatives 7.3, lie in J .

Conversely, does $(a + j)(b + k)(a + j) - aba$ lie in J if 7.4 holds? Actually, we undertake to prove that $(a + sj)(b + tk)(a + sj) - aba$ lies in J for all real s and t , if 7.4 holds. Let

$$f(s) = (a + sj)(b + tk)(a + sj) - aba.$$

Then $f(s) = \sum c_h s^h$, a sum of three terms. Conditions 7.4 imply that all derivatives of $f(s)$ are in J . Therefore, all the c_h are in J for $h > 0$. The constant of integration c_0 is also in J because $f(0) = atka \in J$. Hence $f(s) \in J$, as we promised to show.

PROPOSITION. Consider the element $q = xyx$ of $\mathbb{F}\langle x, y \rangle$. Let $I_{xyx}(x)$ be the q -ideal generated by the single element x of $\mathbb{F}\langle x, y \rangle$. Then a basis for the elements of this ideal of degree not greater than 3 is

$$(7.5) \quad \{x, x^2, xy + yx, x^3, x^2y, xyx, yx^2, xy^2 + y^2x\}.$$

Proof. By repeated use of 4.52 one can show that any xyx -ideal which contains x , must contain x^2 and $xy + yx$. Also one can show that an ideal which contains these must contain the entire list 7.5. The list is clearly linearly independent. Then, one can show that on the other hand, that the span of 7.5 together with all polynomials of degree at least four, is an xyx -ideal. This sketch should suffice for a proof.

PROPOSITION. *Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let $A = \mathbb{F}\langle x_1, \dots, x_m \rangle$. Let k be a positive integer. Let X^k be the set of polynomials whose constituent homogeneous summands are all of degree k or more. Then X^k is a q -ideal.*

Proof. Let a_1, \dots, a_m be members of A and let j_1, \dots, j_m be members of X^k . The expression $q(\dots, a_i + j_i, \dots) - q(\dots, a_i, \dots)$ is a sum of monomials in the a_i and j_i . After the obvious cancellation, every term must have at least one factor j_h in it, and so the sum must belong to X^k .

7.6. LEMMA. *Let $q \in P \equiv \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let m be an integer not necessarily related to n . Let $A = \mathbb{F}\langle x_1, \dots, x_m \rangle$. Let J be a q -ideal in A and let p be an element of A which does not lie in J . Then there is a q -ideal K to which p does not belong, which has finite codimension in A , and contains J .*

Proof. Let k be greater than the degree of p . Let $K = J + X^k$. This is a q -ideal, and it has finite codimension, because X^k does. If the element p were to be in K , say $p = j + z$, then the homogeneous constituents of z could not appear in p because the degree of p is too small. Thus they would find and cancel their negatives in j , and p would lie in J .

8. Establishing non-productivity. For convenience, we introduce another term.

DEFINITION. Suppose A is a linear algebra over \mathbb{F} . Suppose that S is a seminorm on A , and μ is a real number satisfying $S(\hat{q}(a_1, \dots, a_n)) \leq \mu$ whenever $S(a_i) \leq 1$ for all i . Then μ is a q -factor for S , and S will be said to have a q -factor.

8.1. LEMMA. *Let $q \in P \equiv \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let A be an algebra and let J be a q -ideal such that A/J is finite dimensional. Then there is a seminorm S on A that has a q -factor and has kernel J .*

In the following proof, we always intend summation over repeated indices. Also, any lower case j stands for a member of J .

Let A/J have dimension m . Let e_1, \dots, e_m be a basis for A/J . Now $\hat{q}(\sum \alpha_{1h}e_h, \dots, \sum \alpha_{nh}e_h)$ is well defined because J is a q -ideal, which implies that evaluating \hat{q} is well defined modulo J . Since q is a polynomial, $\hat{q}(\sum \alpha_{1h}e_h, \dots, \sum \alpha_{nh}e_h) = P_i e_i$, where the P_i are polynomials in the α_{ih} .

Choose a norm $\|\cdot\|$ for A/J . Define the seminorm S on A by $S(a) = \|a + J\|$. Then $S(\hat{q}(\sum \alpha_{1h}e_h, \dots, \sum \alpha_{nh}e_h)) = \|P_i e_i\|$.

The condition $\|a + J\| \leq 1$ defines a closed bounded set B in A/J . For a_1, \dots, a_n to have $S(a_i) \leq 1$ is the same as having the $a_i + J$ lie in B . This in turn, makes $\|P_i e_i\| \leq \mu$ for some real μ . We have shown that $S(a_i) \leq 1$ for $i = 1, \dots, n$ implies $S(\hat{q}(a_1, \dots, a_n)) \leq \mu$. Thus the seminorm has a q -factor.

The kernel of S is obviously J . This proves 8.1.

8.2. LEMMA. *Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let p be an element of $A = \mathbb{F}\langle x_1, \dots, x_m \rangle$ vanishing at $(0, \dots, 0)$ of $A \times \dots \times A = A^m$. Suppose that p does not lie in the q -ideal $I_q(x_1, \dots, x_m)$ generated by x_1, \dots, x_m in A . Then there exists a seminorm S on A that has q -factors, but does not render \hat{p} continuous at $(0, \dots, 0)$.*

Proof. By 7.6, there is a q -ideal K to which p does not belong, and such that A/K has finite dimension. By 8.1 there is a seminorm S having q -factors, and having K for its null space. So $S(p) > 0$ because p is not in K . However, $S(x_i) = 0$. Now suppose \hat{p} were continuous at $(0, \dots, 0)$. Its value there is of course the zero element of A . Hence for every $\varepsilon > 0$ there is a $\delta > 0$ such that for those a_1, \dots, a_m , with $S(a_i) < \delta$ for all i , one will have $S(\hat{p}(a_1, \dots, a_m)) < \varepsilon$. Take $\varepsilon = S(p)$ which is positive. Consider letting a_i have the value x_i . With that a_i , we surely have $S(a_i) < \delta$ for all i . So $S(\hat{p}(a_1, \dots, a_m)) < \varepsilon$. But now $S(\hat{p}(a_1, \dots, a_m)) = S(\hat{p}(x_1, \dots, x_m))$, and a little reflection on the definition of the function \hat{p} shows that $\hat{p}(x_1, \dots, x_m)$ is indeed p . So $S(p) < S(p)$, is obviously a contradiction. This completes our proof of 8.2.

8.3. THEOREM. *Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let p be an element of $A = \mathbb{F}\langle x_1, \dots, x_m, z_1, \dots, z_r \rangle$ that is homogeneous in each of z_1, \dots, z_r and homogeneous of positive degree in x_1 . Suppose p does not belong to the q -ideal generated by z_1, \dots, z_r . Then there is*

a seminorm S on A which has q -factors but renders \hat{p} discontinuous at $(0, \dots, 0)$.

Proof. As above, there is a q -ideal K including z_1, \dots, z_r , but not p , and such that A/K is finite dimensional. Consequently there is a seminorm S such that $S(z_i) = 0$, but $S(p) > 0$. Say $S(p) = \varepsilon$. Suppose \hat{p} were continuous at $(0, \dots, 0)$. Then there is a $\delta > 0$ such that if $S(a_i) < \delta$ for all i and $S(b_j) < \delta$ for all j then $S(\hat{p}(a_1, \dots, a_m, b_1, \dots, b_r)) < \varepsilon$. Let $a_i = \alpha_i x_i$ and $b_j = \beta_j z_j$. If the α_i are sufficiently small, then $S(a_i) < \delta$ for all i and $S(b_j) < \delta$ for all j , whereupon $S(\hat{p}(a_1, \dots, a_m, b_1, \dots, b_r)) < \varepsilon$. Now $S(\hat{p}(a_1, \dots, a_m, b_1, \dots, b_r))$ is a product of α 's and β 's times $S(\hat{p}(x_1, \dots, x_m, z_1, \dots, z_r))$. After the α 's have been chosen small enough, and β_2, \dots, β_n have been set equal to 1, the product of α 's and β_1 can be made arbitrarily large by varying β_1 . Thus $S(\hat{p}(x_1, \dots, x_m, z_1, \dots, z_r))$ is actually 0. But as mentioned at the end of the proof of 8.2, $S(\hat{p}(x_1, \dots, x_m, z_1, \dots, z_r))$ is $S(p)$. This contradiction proves 8.3.

8.4. THEOREM. *Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let p be an element of $\mathbb{F}\langle x_1, \dots, x_m, z_1, \dots, z_r \rangle$ that is homogeneous in each of z_1, \dots, z_r and homogeneous of positive degree in x_1 . Suppose p does not belong to the q -ideal generated by z_1, \dots, z_r . Then p does not belong to $((q))$.*

Proof. If $p \in ((q))$ then \hat{p} is continuous by 6.1. But this is contradicted by 8.3.

In order to apply this more smoothly to a situation met earlier, we make a mere alphabetic variation:

8.4'. THEOREM. *Let $q \in \mathbb{F}\langle x_1, \dots, x_n \rangle$. Let p be an element of $\mathbb{F}\langle y_1, \dots, y_m, x_1, \dots, x_r \rangle$ that is homogeneous in each of x_1, \dots, x_r and homogeneous of positive degree in y_1 . Suppose p does not belong to the q -ideal generated by x_1, \dots, x_r . Then p does not belong to $((q))$.*

8.5. THEOREM. *Consider the element $q = xyx$ of $\mathbb{F}\langle x, y \rangle$. Let p be an element of $\mathbb{F}\langle x, y \rangle$ that is homogeneous of positive degree in y , homogeneous in x , and homogeneous of the third degree in x, y . Suppose p is not a linear combination of the elements of the list 7.5. Then $p \notin ((xyx))$.*

Proof. By hypothesis, p does not belong to the ideal $I_{xyx}(x)$. We can appeal to 8.4' and conclude that $p \notin ((xyx))$.

REFERENCES

[BK] H. Braun and M. Koecher, *Jordan Algebren*, Springer-Verlag, 1966.

Received April 29, 1992.

UNIVERSITY OF CALIFORNIA
LOS ANGELES, CA 90024-1555

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906–1982) F. WOLF (1904–1989)

EDITORS

SUN-YUNG A. CHANG
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
chang@math.ucla.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

HERBERT CLEMENS
University of Utah
Salt Lake City, UT 84112
clemens@math.utah.edu

THOMAS ENRIGHT
University of California, San Diego
La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

HAROLD STARK
University of California, San Diego
La Jolla, CA 92093

V. S. VARADARAJAN
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Julie Honig, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 75 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$107.50 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to *Pacific Journal of Mathematics*, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The *Pacific Journal of Mathematics* at University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to *Pacific Journal of Mathematics*, P.O. Box 4163, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$,
the American Mathematical Society's $\mathcal{T}\mathcal{E}\mathcal{X}$ macro system.
Copyright © 1994 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 162 No. 2 February 1994

On the existence of convex classical solutions to multilayer fluid problems in arbitrary space dimensions	201
ANDREW FRENCH ACKER	
Extremal functions and the Chang-Marshall inequality	233
VALENTIN V. ANDREEV and ALEC LANE MATHESON	
Productive polynomials	247
RICHARD ARENS	
On factor representations of discrete rational nilpotent groups and the Plancherel formula	261
LAWRENCE JAY CORWIN and CAROLYN PFEFFER JOHNSTON	
Commutants of Toeplitz operators on the Bergman space	277
ZELJKO CUCKOVIC	
When L^1 of a vector measure is an AL-space	287
GUILLERMO P. CURBERA	
A convexity theorem for semisimple symmetric spaces	305
KARL-HERMANN NEEB	
Ideals of finite codimension in free algebras and the fc-localization	351
AMNON ROSENMAN and SHMUEL ROSSET	
Dec groups for arbitrarily high exponents	373
BHARATH AL SETHURAMAN	
Errata to: "The set of primes dividing the Lucas numbers has density $2/3$ "	393
JEFFREY C. LAGARIAS	