

*Pacific
Journal of
Mathematics*

**THE NONHOMOGENEOUS MINIMAL SURFACE EQUATION
INVOLVING A MEASURE**

WILLIAM KARL ZIEMER

THE NONHOMOGENEOUS MINIMAL SURFACE EQUATION INVOLVING A MEASURE

WILLIAM K. ZIEMER

We find existence of a minimum in BV for the variational problem associated with $\operatorname{div} A(Du) + \mu = 0$, where A is a mean curvature type operator and μ a nonnegative measure satisfying a suitable growth condition. We then show a local L^∞ estimate for the minimum. A similar local L^∞ estimate is shown for sub-solutions that are Sobolev rather than BV .

1. Introduction. In this paper we initiate an investigation of weak solutions of the

$$(1.1) \quad \operatorname{div} A(Du) + \mu = 0$$

in a bounded Lipschitz domain $\Omega \subset R^n$. Here A is a function for which the mean curvature operator is a prototype and μ is a nonnegative Radon measure supported in Ω that satisfies

$$(1.2) \quad \mu(B(r)) \leq Mr^{q(n-1)} \text{ for all } B(r) \subset \Omega,$$

where $M > 0$ and $1 < q \leq \frac{n}{n-1}$.

This paper has its origins in the work of [LS] where it was shown that if u is a weak solution of

$$\Delta u = \mu,$$

where μ is a measure that satisfies the growth condition

$$\mu(B(r)) \leq Mr^{n-2+\varepsilon}$$

for some $\varepsilon > 0$ and for all balls $B(r)$ of radius r , then u is Hölder continuous. In [RZ] this result was generalized to equations of the form

$$(1.3) \quad \operatorname{div} A(x, u, \nabla u) + B(x, u, \nabla u) + \mu = 0$$

where μ is a nonnegative Radon measure and A and B are Borel measurable functions satisfying structural conditions that allow, for example, the p -Laplacian. It is shown that if u is a Hölder continuous solution of 1.3, then μ satisfies

$$\mu(B(r)) \leq Mr^{n-p+\varepsilon}$$

for some $\varepsilon > 0$. Under further restrictions on the structural conditions, it was shown this growth condition on μ was sufficient for Hölder continuity of u .

Recently, Lieberman [L] improved the results in [RZ] by proving supremum inequalities for solutions of 1.3 without the restrictive structural conditions, thereby establishing necessary and sufficient conditions on the growth of μ for the Hölder continuity of solutions.

All of this analysis takes place in the framework surrounding the p -Laplacian, $p > 1$. It is our purpose to address the situation of $p = 1$. We first consider the question of existence of solutions of 1.5 in the case A is the mean curvature operator. We establish a variational solution by minimizing

$$(1.4) \quad \int_{\Omega} \sqrt{1 + |\nabla u|^2} dx + \int_{\Omega} u d\mu$$

in the class $u \in BV(\Omega)$ where u satisfies the Dirichlet condition $u^* = f$ on $\partial\Omega$, with f an integrable function on $\partial\Omega$. In order to ensure the existence of a minimum, it is necessary to assume that the constant M in 1.2 is chosen sufficiently small. This is analogous to the assumption made in [M], in which μ is taken as a bounded measurable function. We then show that the minimizer u is bounded. In this context, it is not possible to utilize the argument given in [L] to obtain an L^∞ bound since there is no variational equation associated with 1.4. Rather, we employ a technique used in [RZ] modeled on the method of DeGiorgi.

Next, we investigate an equation which contains the formal Euler-Lagrange equation of 1.4. Thus, we consider a weak solution $u \in W^{1,1}(\Omega) \cap L^\infty(\Omega)$ of the equation

$$(1.5) \quad \operatorname{div} A(Du) + \mu = 0$$

where we assume there exist non-negative constants a_1, a_2 such that

$$(1.6) \quad p \cdot A(p) \geq |p| - a_1$$

and

$$(1.7) \quad |A(p)| \leq a_2.$$

It is assumed that μ is a nonnegative Radon measure supported in the bounded domain Ω and satisfies 1.2. We show that if $u \in W^{1,1}(\Omega) \cap L^\infty(\Omega)$ is a weak solution of 1.5, then $|u|$ is bounded by the L^1 -norm of u with respect to the measure $d\nu = dx + d\mu$. Specifically, we show that u satisfies a supremum inequality, 6.4. The proof of this follows the proof in the corresponding result of [L]. The method of DeGiorgi will still work in this case, however the Moser iteration method used in [L] gives a slightly different result and is included for this reason. It is well known that weak solutions of 1.5 are not necessarily continuous, even under the assumption that μ is an absolutely continuous measure with bounded density (c.f. [M]). Therefore, it is not possible to obtain the weak Harnack inequality involving a lower bound for the solution.

The results of this paper are valid for equations with a more general structure. For the sake of simplicity, we employ this simple structure which fully illustrates the method. In a forthcoming paper, we will address the question of regularity of solutions of 1.4 in which almost everywhere continuity is established. The existence of an *a priori* L^∞ bound will be essential in this future investigation.

2. Preliminaries. Throughout, we assume that Ω is a bounded Lipschitz domain in R^n . The space $W^{1,1}(\Omega)$ is the space of $L^1(\Omega)$ functions whose distributional derivatives also lie in $L^1(\Omega)$.

The class of all functions in $L^1(\Omega)$ whose distributional partial derivatives are measures with finite total variation in Ω comprise the space $BV(\Omega)$. The notation

$$\int_{\Omega} |Du| \, dx$$

will be used to represent the total variation of the vector-valued measure, Du , the gradient of u . Specifically, the total variation of Du is

$$\sup \left\{ \int_{\Omega} u \operatorname{div} v \, dx : v = (v_1, \dots, v_n) \in C_0^\infty(\Omega; R^n), |v| \leq 1 \right\}.$$

We also make the notational definition

$$\begin{aligned} & \int_{\Omega} \sqrt{1 + |Du|^2} \, dx \\ &= \sup \left\{ \int_{\Omega} (f \operatorname{div} v + v_0) \, dx : v = (v_1, \dots, v_n) \in C_0^{\infty}(\Omega), \right. \\ & \quad \left. v_0 \in C_0^{\infty}(\Omega), |v|^2 + |v_0|^2 \leq 1 \right\}. \end{aligned}$$

The space $BV(\Omega)$ is equipped with the norm

$$\|u\|_{BV} = \int_{\Omega} |u| \, dx + \int_{\Omega} |Du| \, dx.$$

The trace of u on $\partial\Omega$ is denoted by u^* (c.f. [Z, Section 5.10]). We will make use of the following lemma on the convergence of the traces of BV functions.

LEMMA 2.1. *Let $\Omega \subset R^n$ a bounded Lipschitz domain, and let $\{u_k\}$, u in $BV(\Omega)$ with*

$$\begin{aligned} & \lim_{k \rightarrow \infty} \int_{\Omega} |u_k - u| \, dx = 0 \\ & \lim_{k \rightarrow \infty} \int_{\Omega} \sqrt{1 + |Du_k|^2} \, dx = \int_{\Omega} \sqrt{1 + |Du|^2} \, dx. \end{aligned}$$

Then

$$\lim_{k \rightarrow \infty} \int_{\partial\Omega} |u_k^* - u^*| \, dH^{n-1} = 0,$$

with H^{n-1} the $n - 1$ dimensional Hausdorff measure.

The proof follows directly from the proof in [G, Proposition 2.6; p.34].

We will also have need for the following compactness result for BV functions [Z, Corollary 5.3.4; p. 227].

THEOREM 2.2. *Let $\Omega \in R^n$ be a bounded Lipschitz domain. Then $BV(\Omega) \cap \{u : \|u\|_{BV(\Omega)} \leq 1\}$ is compact in $L^1(\Omega)$.*

It was shown in [MZ] that if μ satisfies the growth condition $\mu(B(r)) \leq Mr^{n-1}$ on all balls $B(r)$ (and therefore condition 1.2 in

particular), then μ can be identified with an element of the dual of $BV(\Omega)$. Furthermore, its norm

$$\tilde{M} = \|\mu\| = \sup \left\{ \int_{\Omega} u \, d\mu : \|u\|_{BV(\Omega)} \leq 1 \right\}$$

is comparable to M . Thus,

$$(2.1) \quad \begin{aligned} \left| \int_{\Omega} u \, d\mu \right| &\leq \int_{\Omega} |u| \, d\mu \\ &\leq \|\mu\| \|u\|_{BV(\Omega)} \\ &\leq \tilde{M} \|u\|_{BV(\Omega)} \end{aligned}$$

The following well known result, [M], will be used in the existence proof below.

$$(2.2) \quad \int_{\Omega} |u| \, dx \leq C \left(\int_{\Omega} |Du| \, dx + \int_{\partial\Omega} u^* \, dH^{n-1} \right)$$

with the constant $C = C(\Omega)$. This yields

$$(2.3) \quad \|u\|_{BV(\Omega)} \leq C \left(\int_{\Omega} |Du| \, dx + \int_{\partial\Omega} u^* \, dH^{n-1} \right)$$

Finally, we state the following Sobolev inequalities which are of critical importance in our development.

THEOREM 2.3. *Let Ω be a bounded Lipschitz domain and suppose μ is a measure supported in Ω satisfying condition 1.2. Then there exists a constant $C = C(\Omega, q, n)$ such that*

$$(2.4) \quad \left(\int_{\Omega} u^q \, d\mu \right)^{1/q} \leq CM^{1/q} \int_{\Omega} |Du| \, dx$$

whenever $u \in BV(\Omega)$ with compact support in Ω .

The proof may be found in [Z, Lemma 4.9.1; p. 209]. Also needed is the standard Sobolev inequality for $W^{1,1}$.

If $u \in W_0^{1,1}(\Omega)$ then there exists a constant $C = C(\Omega, q, n)$ such that

$$(2.5) \quad \left(\int_{\Omega} u^q \, dx \right)^{1/q} \leq C \|Du\|_1.$$

This is simply the above lemma in the special case that μ is Lebesgue measure.

3. Existence of a Minimum. With Ω a bounded Lipschitz domain and $f \in L^1(\partial\Omega)$, we define $I(u; \Omega)$ as follows,

$$I(u; \Omega) = \int_{\Omega} \sqrt{1 + |Du|^2} \, dx + \int_{\Omega} u \, d\mu + \int_{\partial\Omega} |u^* - f| \, dH^{n-1}.$$

We wish to minimize I over all $u \in BV(\Omega)$. That is, we wish to find a function $u \in BV(\Omega)$ such that

$$I(u; \text{supp } \varphi) \leq I(u + \varphi; \text{supp } \varphi), \quad \forall \varphi \in C_0^\infty(\Omega).$$

THEOREM 3.1. *Let Ω be a bounded Lipschitz domain. With I defined as above, there exists $u \in BV(\Omega)$ such that*

$$I(u; \Omega) = \min_{v \in BV(\Omega)} I(v; \Omega).$$

Proof. Following [G, Section 14.4], the first step is to consider a slightly different Dirichlet problem in the complement of Ω . For this purpose, let B be a ball that contains $\bar{\Omega}$, the closure of Ω . Use Theorem 2.16 of [G] to extend f to a $W^{1,1}$ function in $B - \bar{\Omega}$ that will still be denoted by f . Let

$$J(u; B) = \int_B \sqrt{1 + |Du|^2} \, dx + \int_B u \, d\mu.$$

Note that since $\text{supp } \mu \subset \Omega$, the second integral could have been taken over Ω . We wish to show that there exists $u \in BV(B)$, coinciding with f in $B - \bar{\Omega}$, that minimizes $J(u; B)$. We proceed by showing that J is bounded below if the constant M in 1.2 is sufficiently small.

$$\begin{aligned} J(u; B) &\geq \int_B |Du| \, dx + \int_{\Omega} u \, d\mu \\ \text{(by 2.1)} \quad &\geq \int_B |Du| \, dx - \tilde{M} \|u\|_{BV(\Omega)} \\ &\geq \int_B |Du| \, dx - \tilde{M} \left(C \int_{\partial\Omega} u^* \, dH^{n-1} \right. \\ \text{(by 2.3)} \quad &\quad \left. + (C + 1) \int_{\Omega} |Du| \, dx \right) \\ &\geq \frac{1}{2} \int_B |Du| \, dx - \tilde{M} C \int_{\partial\Omega} f \, dH^{n-1}. \end{aligned}$$

The last inequality is obtained when \tilde{M} is small enough to insure $1 - \tilde{M}(C + 1) \geq \frac{1}{2}$.

Let $J(u_k) \rightarrow \lambda$ a minimum of J . We wish to find $u \in BV(B)$ such that $J(u; B) = \lambda$. For sufficiently large k we obtain from the above inequality that

$$\lambda + 1 \geq \frac{1}{2} \int_B |Du_k| dx - MC \int_{\Omega} f dH^{n-1}.$$

Thus the terms $\int_B |Du_k| dx$ are uniformly bounded, which implies by 2.3 and Theorem 2.2 that there exists $u \in BV(B)$ with $u_k \rightarrow u$ in $L^1(B)$. The gradient is lower semi-continuous with respect to $L^1(B)$ convergence so that

$$\liminf_{k \rightarrow \infty} \int_B \sqrt{1 + |Du_k|^2} dx \geq \int_B \sqrt{1 + |Du|^2} dx.$$

From Theorem 2.3, the uniform bound on $\int_B |Du_k| dx$ also implies that the terms

$$\left(\int_{\Omega} u_k^q d\mu \right)^{1/q}$$

are uniformly bounded. Thus there exists a subsequence, denote it by $\{u_k\}$, that converges weakly in $L^q(\Omega; \mu)$ to some $w \in L^q(\Omega; \mu)$. The Banach–Saks Theorem implies that there exists a subsequence of $\{u_k\}$, again denote it by $\{u_k\}$, such that the sequence of Césaro sums, $\{v_k\}$, defined by

$$v_k = \frac{u_1 + \cdots + u_k}{k}$$

converges strongly to w in $L^q(\Omega; \mu)$. Moreover, the sequence v_k also converges strongly to u in $L^1(\Omega)$. This can be seen as follows: choose $\varepsilon > 0$ and let N denote an integer for which $\|u_j - u\|_{L^1(\Omega)} < \varepsilon$ for $j, k \geq N$. Then for $j \leq k$,

$$\begin{aligned} & \|v_k - u\| \\ &= \left\| \frac{(u_1 - u) + \cdots + (u_k - u)}{k} \right\| \\ &\leq \frac{\|u_1 - u\| + \cdots + \|u_{j-1} - u\|}{k} + \frac{\|u_j - u\| + \cdots + \|u_k - u\|}{k} \\ &\leq \frac{\|u_1 - u\| + \cdots + \|u_{j-1} - u\|}{k} + \frac{(k - j + 1)\varepsilon}{k}. \end{aligned}$$

Thus,

$$\limsup_{k \rightarrow \infty} \|v_k - u\| \leq \varepsilon,$$

which yields the desired result since ε is arbitrary. To show that $w = u$ almost everywhere in Ω note that the strong convergence of $\{v_k\}$ to w in $L^q(\Omega; \mu)$ implies the existence of a subsequence that converges pointwise to w μ -almost everywhere and therefore (Lebesgue) almost everywhere, since Lebesgue measure is absolutely continuous with respect to μ in Ω . But the strong convergence of $\{v_k\}$ to u in $L^1(\Omega)$ implies the almost everywhere pointwise convergence of a further subsequence to u in Ω . Hence, $u = w$ almost everywhere in Ω .

Since u_k converges weakly to u in $L^q(\Omega; \mu)$, the lower semicontinuity of the gradient with respect to $L^1(\Omega)$ convergence implies

$$(3.1) \quad \lambda = \liminf_{k \rightarrow \infty} J(u_k; B) \geq J(u; B).$$

Since u_k agrees almost everywhere with f in $B - \bar{\Omega}$, it follows that $u = f$ a.e. in $B - \bar{\Omega}$, thus showing that $J(u; B) \geq \lambda$. This completes the first step.

We now proceed with the second and final step of the proof. For each function $v \in BV(\Omega)$, define

$$v_f(x) = \begin{cases} v(x) & x \in \Omega \\ f(x) & x \in B - \Omega \end{cases}$$

Then $v_f \in BV(B)$ and by (2.15) of [G],

$$\begin{aligned} & \int_B \sqrt{1 + |Dv_f|^2} dx + \int_B v_f d\mu \\ &= \int_B \sqrt{1 + |Dv|^2} dx + \int_{B - \bar{\Omega}} \sqrt{1 + |Df|^2} dx \\ & \quad + \int_B v_f d\mu + \int_{\partial\Omega} |v_\Omega^* - f| dH^{n-1} \\ &= I(v; \Omega) + \int_{B - \bar{\Omega}} \sqrt{1 + |Df|^2} dx \end{aligned}$$

That is,

$$J(v_f; B) = I(v; \Omega) + \int_{B - \bar{\Omega}} \sqrt{1 + |Df|^2} dx.$$

Thus, a minimizer of $J(v; B)$ with $v = f$ on $B - \bar{\Omega}$ produces a minimizer of $I(v; \Omega)$. \square

4. An energy inequality. Now that we have obtained existence of a solution $u \in BV(\Omega)$ to 1.4, we will show that u is bounded. Before doing this we will obtain an energy estimate to be used in the DeGiorgi type argument of section 5.

Let B_R denote the ball of radius R in R^n . Let η be a cutoff function, $\eta = 1$ on B_r , $0 < r < r^* \leq R$, $\eta = 0$ on ∂B_{r^*} with $0 \leq \eta \leq 1$ on B_{r^*} and $|D\eta| \leq \frac{2}{r^* - r}$. Let $\varphi = -\eta(u - k)^+$, then $\text{supp } \varphi = A_k = \{u > k\} \cap B_{r^*}$ and

$$(4.1) \quad I(u; A_k) \leq I(u + \varphi; A_k)$$

Using

$$(4.2) \quad \int_{A_k} |Du| \, dx \leq \int_{A_k} \sqrt{1 + |Du|^2} \, dx \leq \int_{A_k} |Du| + 1 \, dx$$

and that on A_k

$$D(u + \varphi) = (1 - \eta)D(u - k)^+ - D\eta(u - k)^+,$$

we obtain from 4.1

$$\begin{aligned} \int_{A_k} |D(u - k)^+| \, dx &\leq \int_{A_k} (1 - \eta) |D(u - k)^+| \, dx \\ &\quad + \frac{2}{r^* - r} \int_{A_k} |(u - k)^+| \, dx \\ &\quad + \int_{A_k} \eta |(u - k)^+| \, d\mu + |A_k| \end{aligned}$$

where $|A_k|$ is the Lebesgue measure of A_k . This immediately implies

$$(4.3) \quad \begin{aligned} \int_{B_r} |D(u - k)^+| \, dx &\leq \int_{B_{r^*}} \eta |D(u - k)^+| \, dx \\ &\leq \frac{2}{r^* - r} \int_{B_{r^*}} |(u - k)^+| \, dx \\ &\quad + \int_{B_{r^*}} |(u - k)^+| \, d\mu + |A_k|. \end{aligned}$$

5. Supremum estimate for variational solutions.

THEOREM 5.1. *Let $\sigma \in (0, 1)$, Ω a bounded Lipschitz domain, and $B_R \subset \Omega$ with $R < 1$. Then for $u \in BV(\Omega)$ a minimum of I there exists a constant $C = C(\sigma, M)$ such that*

$$\sup_{B_{\sigma R}} u \leq C \left(R^{-n} \int_{B_R} u^+ dx + R^{-q(n-1)} \int_{B_R} u^+ d\mu \right)$$

where q is the constant from 1.2 and u^+ is the positive part of u .

Proof. Let k be a positive constant to be specified later. Set

$$k_i = k(1 - 2^{-i}), \quad r_i = \sigma R + 2^{-i}R(1 - \sigma),$$

$$\text{and } \tilde{r}_i = \frac{1}{2}(r_i + r_{i+1}).$$

For notational convenience, denote by B_i the ball of radius r_i , \tilde{B}_i the ball of radius \tilde{r}_i , and let

$$A_i = B_i \cap \{(u - k_{i+1})^+ > 0\}.$$

Note that $B_{i+1} \subset \tilde{B}_i \subset B_i$. Also, for all j we will use the notation

$$\int_{B_j} dx = R^{-n} \int_{B_j} dx \quad \text{and} \quad \int_{B_j} d\mu = R^{-q(n-1)} \int_{B_j} d\mu.$$

Let φ_i be the cutoff functions on \tilde{B}_i so that $\varphi_i \equiv 1$ on B_{i+1} and

$$(5.1) \quad |D\varphi_i| \leq \frac{2}{\tilde{r}_i - r_{i+1}} = \frac{2^{i+3}}{R(1 - \sigma)}.$$

Then 4.3 implies

$$(5.2) \quad \int_{B_{i+1}} |D(u - k_{i+1})^+| dx$$

$$\leq \frac{2^{i+3}}{R(1 - \sigma)} \int_{\tilde{B}_i} (u - k_{i+1})^+ dx$$

$$+ R^{-n+q(n-1)} \int_{\tilde{B}_i} (u - k_{i+1})^+ d\mu + R^{-n} |A_i|.$$

Now, by 2.4 and 5.1,

$$\begin{aligned}
 & \int_{B_{i+1}} (u - k_{i+1})^+ d\mu \\
 & \leq \int_{\tilde{B}_i} \varphi_i (u - k_{i+1})^+ d\mu \\
 & \leq \left(\int_{\tilde{B}_i} (\varphi_i (u - k_{i+1})^+)^q d\mu \right)^{1/q} (R^{-q(n-1)} \mu(A_i))^{1-1/q} \\
 & \leq CM^{1/q} R \int_{\tilde{B}_i} |D(\varphi_i (u - k_{i+1})^+)| dx (R^{-q(n-1)} \mu(A_i))^{1-1/q} \\
 & \leq CRM^{1/q} \left(\int_{\tilde{B}_i} |D(u - k_{i+1})^+| \varphi_i dx \right. \\
 & \quad \left. + \int_{\tilde{B}_i} (u - k_{i+1})^+ |D\varphi_i| dx \right) (R^{-q(n-1)} \mu(A_i))^{1-1/q} \\
 & \leq CRM^{1/q} \left(\int_{\tilde{B}_i} |D(u - k_{i+1})^+| dx \right. \\
 & \quad \left. + \frac{2^{i+3}}{R(1-\sigma)} \int_{\tilde{B}_i} (u - k_{i+1})^+ dx \right) (R^{-q(n-1)} \mu(A_i))^{1-1/q}.
 \end{aligned}$$

Applying 5.2 we have

$$\begin{aligned}
 & \int_{B_{i+1}} (u - k_{i+1})^+ d\mu \\
 & \leq CRM^{1/q} \left(\frac{2^{i+4}}{R(1-\sigma)} \int_{B_i} (u - k_{i+1})^+ dx \right. \\
 & \quad \left. + R^{-n+q(n-1)} \int_{B_i} (u - k_{i+1})^+ d\mu \right. \\
 & \quad \left. + R^{-n} |A_i| \right) (R^{-q(n-1)} \mu(A_i))^{1-1/q}.
 \end{aligned}$$

Thus we have the following iteration inequality,

(5.3)

$$\begin{aligned}
 & \int_{B_{i+1}} (u - k_{i+1})^+ d\mu \\
 & \leq CM^{1/q} \frac{2^{i+4}}{(1-\sigma)} \left(\int_{B_i} (u - k_i)^+ dx \right. \\
 & \quad \left. + \int_{B_i} (u - k_i)^+ d\mu + R^{-n} |A_i| \right) (R^{-q(n-1)} \mu(A_i))^{1-1/q}.
 \end{aligned}$$

To estimate the quantity $\mu(A_i)$ recall that $A_i = \{u > k_{i+1}\} \cap B_i$, and note that

$$\begin{aligned} k_{i+1} - k_i &= k(1 - 2^{-(i+1)}) - k(1 - 2^{-i}) \\ &= 2^{-i}k(1 - 2^{-1}) \\ &= 2^{-(i+1)}k. \end{aligned}$$

which implies

$$2^{-(i+1)}k < u - k_i \text{ on } A_i.$$

Thus

$$(5.4) \quad \begin{aligned} R^{-q(n-1)}\mu(A_i) &\leq 2^{i+1}k^{-1} \int_{B_i} (u - k_i)^+ d\mu \\ &\leq 2^{i+1}Y_i. \end{aligned}$$

where

$$Y_i = k^{-1} \int_{B_i} (u - k_i)^+ dx + k^{-1} \int_{B_i} (u - k_i)^+ d\mu.$$

We estimate $|A_i|$ in the same manner, obtaining

$$(5.5) \quad R^{-n} |A_i| \leq 2^{i+1}Y_i.$$

Using 5.4 and 5.5 in 5.3 we obtain

$$(5.6) \quad \begin{aligned} &k^{-1} \int_{B_{i+1}} (u - k_{i+1})^+ d\mu \\ &\leq CM^{1/q} \frac{2^{i+4}}{(1-\sigma)} \left(k^{-1} \int_{B_i} (u - k_i)^+ dx \right. \\ &\quad \left. + k^{-1} \int_{B_i} (u - k_i)^+ d\mu + k^{-1} 2^{i+1} Y_i \right) \left(2^{i+1} Y_i \right)^{1-1/q} \\ &\leq CM^{1/q} \frac{2^{i+4}}{(1-\sigma)} \left((1 + k^{-1} 2^{i+1}) Y_i \right) \left(2^{i+1} Y_i \right)^{1-1/q} \\ &\leq CM^{1/q} \frac{2^{i+4}}{(1-\sigma)} (k^{-1} + 2^{-i-1}) \left(2^{i+1} Y_i \right)^{1+\alpha}. \end{aligned}$$

where $\alpha = 1 - 1/q > 0$. Following the same analysis for dx instead of $d\mu$ we obtain

$$(5.7) \quad k^{-1} \int_{B_{i+1}} (u - k_{i+1})^+ dx \leq CM^{1/q} \frac{2^{i+4}}{(1-\sigma)} (k^{-1} + 2^{-i-1}) (2^{i+1}Y_i)^{1+\alpha}.$$

Combining 5.6 and 5.7, we have

$$(5.8) \quad Y_{i+1} \leq CM^{1/q} \frac{2^{i+4}}{(1-\sigma)} (k^{-1} + 2^{-i-1}) (2^{i+1}Y_i)^{1+\alpha} \leq CM^{1/q} \frac{2^{i+4}}{\kappa(1-\sigma)} (2^{i+1}Y_i)^{1+\alpha}$$

where $\kappa = \min(1, 1/(k^{-1} + 2^{-1}))$. The recursion lemma of [LU, lemma 4.7; p. 66] then implies that $Y_i \rightarrow 0$, and thus

$$\sup_{B_{\sigma R}} u \leq k,$$

provided that

$$Y_0 = k^{-1} \int_{B_R} u^+ dx + k^{-1} \int_{B_R} u^+ d\mu \leq \left(CM^{1/q} \frac{2^{5+\alpha}}{\kappa(1-\sigma)} \right)^{-1/\alpha} (2^{2+\alpha})^{-1/\alpha^2}.$$

This is true if

$$\kappa^{1/\alpha} k \geq \left(\frac{CM^{1/q} 2^{\alpha+6+2/\alpha}}{(1-\sigma)} \right)^{1/\alpha} \left(\int_{B_R} u^+ dx + \int_{B_R} u^+ d\mu \right).$$

Since $\kappa^{1/\alpha} \leq 1$, the result follows. \square

6. A supremum estimate for weak solutions. We will use a different version of the Sobolev inequalities 2.4 and 2.5.

COROLLARY 6.1. *Let B_R a ball of radius R in R^n . Suppose $u \in W_0^{1,1}(B_R)$ and μ is a measure satisfying 1.2, then there exists a constant $C = C(q, n)$ such that*

$$(6.1) \quad \left(R^{-q(n-1)} \int_{B_R} u^q d\mu \right)^{1/q} \leq M^{1/q} CR^{1-n} \int_{B_R} |Du| dx$$

and

$$(6.2) \quad \left(R^{-n} \int_{B_R} u^q dx \right)^{1/q} \leq C R^{1-n} \int_{B_R} |Du| dx.$$

Let u^+ denote the positive part of u .

THEOREM 6.2. *Let $B_R \subset R^n$ a ball of radius $R < 1$. Suppose that $u \in W^{1,1}(B_R) \cap L^\infty(B_R)$ satisfies the inequality*

$$(6.3) \quad \operatorname{div} A(Du) + \mu \geq 0 \quad \text{in } B_R$$

with A satisfying 1.6 and 1.7, and μ a Radon measure satisfying 1.2. Then for any $\varepsilon > 0$ there exists a constant $C = C(q, n, (a_1 + a_2)/\varepsilon)$ such that

$$(6.4) \quad \sup_{B_{R/2}} |u| \leq C \left(R^{-n} \int_{B_R} u^+ dx + R^{-q(n-1)} \int_{B_R} u^+ d\mu \right) + \varepsilon$$

Proof. Let $\varepsilon > 0$ and $R < 1$. Fix a cutoff function $\eta \in C_0^\infty(B_R)$ such that $\eta = 1$ in $B_{R/2}$, $\eta = 0$ on ∂B_R , and $0 \leq \eta \leq 1$ in B_R with $|D\eta| \leq 4/R$. Set $\zeta = \eta(1 - \frac{\varepsilon}{u})^+$ and $A_\varepsilon = \{\zeta > 0\} = \{u > \varepsilon\} \subset B_R$. Consider the weak formulation of 6.3 with test function $\zeta^{ks-t}u^s$, for constants k, s and t to be chosen later.

$$\begin{aligned} (ks - t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s D\zeta \cdot A(Du) dx \\ + s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} Du \cdot A(Du) dx \leq \int_{A_\varepsilon} \zeta^{ks-t} u^s d\mu. \end{aligned}$$

Use that $D\zeta = D\eta(1 - \frac{\varepsilon}{u}) + \eta\varepsilon u^{-2} Du$ and 1.6 to obtain

$$\begin{aligned} (ks - t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s (1 - \frac{\varepsilon}{u}) D\eta \cdot A(Du) dx \\ + (ks - t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \eta \varepsilon u^{-2} (|Du| - a_1) dx \\ + s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} (|Du| - a_1) dx \\ \leq \int_{A_\varepsilon} \zeta^{ks-t} u^s d\mu. \end{aligned}$$

Which implies that

$$\begin{aligned}
 s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} |Du| \, dx &\leq \int_{A_\varepsilon} \zeta^{ks-t} u^s \, d\mu \\
 &+ (ks-t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \left(1 - \frac{\varepsilon}{u}\right) D\eta \cdot A(Du) \, dx \\
 &+ (ks-t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \eta \varepsilon u^{-2} (a_1) \, dx \\
 &+ s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} (a_1) \, dx.
 \end{aligned}$$

Use 1.7 and that $\varepsilon/u < 1$ in A_ε to obtain

(6.5)

$$\begin{aligned}
 &s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} |Du| \, dx \\
 &\leq \int_{A_\varepsilon} \zeta^{ks-t} u^s \, d\mu + \frac{a_2^4 (ks-t)}{R} \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \, dx \\
 &\quad + (ks-t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s (a_1 u^{-1}) \, dx \\
 &\quad + s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} (a_1) \, dx \\
 &\leq \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \, d\mu + \frac{a_2^4 (ks-t)}{R} \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \, dx \\
 &\quad + \int_{A_\varepsilon} \zeta^{ks-t-1} u^s (a_1 \varepsilon^{-1} (ks-t+s)) \, dx \\
 &\leq \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \, d\mu \\
 &\quad + \frac{a_2^4 (ks-t) + a_1 (ks-t+s)}{\varepsilon R} \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \, dx.
 \end{aligned}$$

Set $w = \zeta^{ks-t} u^s$ and consider

$$\begin{aligned}
 \int_{A_\varepsilon} |Dw| \, dx &\leq s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} |Du| \, dx \\
 &\quad + (ks-t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s |D\zeta| \, dx
 \end{aligned}$$

$$\begin{aligned}
 &\leq s \int_{A_\varepsilon} \zeta^{ks-t} u^{s-1} |Du| \, dx \\
 &\quad + (ks - t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \left(\frac{1}{R} + u^{-1} |Du| \right) \, dx \\
 &\leq (s + ks - t) \int_{A_\varepsilon} \zeta^{ks-t-1} u^{s-1} |Du| \, dx \\
 &\quad + \frac{(ks - t)}{R} \int_{A_\varepsilon} \zeta^{ks-t-1} u^s \, dx.
 \end{aligned}$$

Then use 6.5 to obtain the energy type estimate

(6.6)

$$\begin{aligned}
 &\int_{A_\varepsilon} |Dw| \, dx \\
 &\leq \frac{s + ks - t}{s} \left(\int_{A_\varepsilon} \zeta^{ks-t-2} u^s \, d\mu \right. \\
 &\quad \left. + \frac{a_2 A(ks - t - 1) + a_1(ks - t - 1 + s)}{\varepsilon R} \int_{A_\varepsilon} \zeta^{ks-t-2} u^s \, dx \right) \\
 &\quad + \frac{(ks - t)}{R} \int_{A_\varepsilon} \zeta^{ks-t-2} u^s \, dx \\
 &\leq s(1 + k) \left(\int_{A_\varepsilon} \zeta^{ks-t-2} u^s \, d\mu + \left(4k \frac{a_1 + a_2}{\varepsilon} + 1 \right) \right. \\
 &\quad \left. \cdot \frac{1}{R} \int_{A_\varepsilon} \zeta^{ks-t-2} u^s \, dx \right), \text{ for } s \geq 1, t \geq 0, \text{ and } k \geq 1/5.
 \end{aligned}$$

Sobolev inequalities 6.1 and 6.2 imply

$$\begin{aligned}
 (6.7) \quad &\left(R^{-n} \int_{A_\varepsilon} w^q \, dx \right)^{1/q} + \left(M^{-1} R^{-q(n-1)} \int_{A_\varepsilon} w^q \, d\mu \right)^{1/q} \\
 &\leq CR^{-(n-1)} \int_{A_\varepsilon} |Dw| \, dx
 \end{aligned}$$

with $C = C(n, q)$. Define $v = \zeta^k u$ and set $t = \frac{2}{q-1}$, so that $tq = t+2$. Also, define a measure ν by

$$d\nu = \frac{dx}{R^n \zeta^{t+2}} + \frac{d\mu}{R^{q(n-1)} \zeta^{t+2}},$$

which is supported on $A_\varepsilon = \{u > \varepsilon\} \cap B_R$. We combine inequalities 6.6 and 6.7 to yield

$$(6.8) \quad \left(\int_{A_\varepsilon} v^{sq} d\nu \right)^{1/q} \leq C s \int_{A_\varepsilon} v^s d\nu.$$

where $C = C(q, n, (a_1 + a_2)/\varepsilon)$, since k will be chosen later to be $\frac{2}{q-1} + 2$ and $s \geq 1$ will be used.

We now iterate on the inequality 6.8. Take $s = 1$ in the first iteration,

$$\frac{1}{C} \left(\int_{A_\varepsilon} v^q d\nu \right)^{1/q} \leq \int_{A_\varepsilon} v d\nu.$$

Take $s = q$ in the second iteration,

$$\frac{1}{C} \left(\frac{1}{Cq} \left(\int_{A_\varepsilon} v^{q^2} d\nu \right)^{1/q} \right)^{1/q} \leq \int_{A_\varepsilon} v d\nu.$$

Proceeding with $s = q^{m-1}$ in the m^{th} iteration will yield

$$(6.9) \quad K_m \left(\frac{1}{C} \right)^{S_m} \left(\int_{A_\varepsilon} v^m d\nu \right)^{1/m} \leq \int_{A_\varepsilon} v d\nu.$$

with the constants K_m and S_m given by

$$K_m = \prod_{j=0}^{m-1} \left(\frac{1}{q^j} \right)^{\frac{1}{q^j}}, \quad S_m = \sum_{j=0}^{m-1} 1/q^j.$$

As $m \rightarrow \infty$ the constants $S_m \rightarrow \frac{q}{q-1}$ and $K_m \rightarrow K$, $0 < K < \infty$. Since $K_1 > K_2 > \dots > K$ we have, for all m , from 6.9

$$\begin{aligned} \left(\int_{A_\varepsilon} v^m d\nu \right)^{1/m} &\leq C^{S_m} \frac{1}{K} \int_{A_\varepsilon} v d\nu \\ &\leq \frac{C^{\frac{q}{q-1}}}{K} \int_{A_\varepsilon} v d\nu. \end{aligned}$$

This then implies (with C replacing $\frac{C^{\frac{q}{q-1}}}{K}$)

$$(6.10) \quad \sup_{A_\varepsilon} v \leq C \int_{A_\varepsilon} v d\nu.$$

On $B_{R/2}$ we have that $\zeta = (1 - \frac{\varepsilon}{u})^+$. Thus when $u \geq 2\varepsilon$, we have $\zeta \geq \frac{1}{2}$. Set $k = t + 2$, and 6.10 implies

$$\begin{aligned} \sup_{B_{R/2}} u &\leq 2^k \sup_{A_\varepsilon} u + 2\varepsilon \\ &\leq C \left(R^{-n} \int_{A_\varepsilon} u \, dx + R^{-q(n-1)} \int_{A_\varepsilon} u \, d\mu \right) + 2\varepsilon \end{aligned}$$

and the result follows, noting that $\int_{A_\varepsilon} u \, dx \leq \int_{B_R} u^+ \, dx$. \square

REFERENCES

- [G] E. Giusti, *Minimal Surfaces and Functions of Bounded Variation*, Birkhäuser (1984).
- [L] Gary Lieberman, *Sharp forms of estimates for subsolutions and supersolutions of quasilinear elliptic equations involving measures*, to appear.
- [LS] H. Levy and G. Stampacchia, *On the smoothness of superharmonics which solve the minimum problem*, J. Analyse Math., **23** (1970), 227-236.
- [LU] O.A. Ladyzhenskaia and N. Uraltseva, *Linear and Quasilinear Elliptic Equations*, Academic Press New York (1968).
- [M] Mario Miranda, *Dirichlet problem with L^1 data for the non-homogeneous minimal surface equation*, Indiana University Math. J., **24**, No. 3 (1974), 227-241.
- [MZ] N.G. Meyers and W. P. Ziemer, *Integral inequalities of Poincaré and Wirtinger type for BV functions*, Amer. J. of Math., **99** (1977), 1345-1360.
- [RZ] J.M. Rakotoson and W. P. Ziemer, *Local behavior of solutions of quasilinear elliptic equations with general structure*, Trans. Amer. Math. Soc., **319**, No. 2 (June 1990), 747-764.
- [Z] William P. Ziemer, *Weakly Differentiable Functions*, Springer-Verlag New York Inc. (1989).

Received August 13, 1992 and in revised form January 13, 1993.

CALIFORNIA STATE UNIVERSITY
 LONG BEACH, CA 90815
E-mail address: wziemer@csulb.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. BECKENBACH (1906-1982) F. WOLF (1904-1989)

EDITORS

SUN-YUNG A. CHANG
(Managing Editor)
University of California
Los Angeles, CA 90024-1555
pacific@math.ucla.edu

F. MICHAEL CHRIST
University of California
Los Angeles, CA 90024-1555
christ@math.ucla.edu

THOMAS ENRIGHT
University of California
San Diego, La Jolla, CA 92093
tenright@ucsd.edu

NICHOLAS ERCOLANI
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

R. FINN
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

VAUGHAN F. R. JONES
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

STEVEN KERCKHOFF
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

MARTIN SCHARLEMANN
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

GANG TIAN
Courant Institute
New York University
New York, NY 10012-1110
tiang@taotao.cims.nyu.edu

V. S. VARADARAJAN
University of California
Los Angeles, CA 90024-1555
vsv@math.ucla.edu

SUPPORTING INSTITUTIONS

UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF HAWAII
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced (not dittoed), double spaced with large margins. Please do not use built up fractions in the text of the manuscript. However, you may use them in the displayed equations. Underline Greek letters in red, German in green, and script in blue. The first paragraph must be capable of being used separately as a synopsis of the entire paper. In particular it should contain no bibliographic references. Please propose a heading for the odd numbered pages of less than 35 characters. Manuscripts, in triplicate, may be sent to any one of the editors. Please classify according to the 1991 *Mathematics Subject Classification* scheme which can be found in the December index volumes of *Mathematical Reviews*. Supply name and address of the author to whom proofs should be sent. All other communications should be addressed to the managing editor, or Julie Honig, University of California, Los Angeles, California 90024-1555.

There are page-charges associated with articles appearing in the *Pacific Journal of Mathematics*. These charges are expected to be paid by the author's University, Government Agency or Company. If the author or authors do not have access to such Institutional support these charges are waived. Single authors will receive 75 free reprints; joint authors will receive a total of 100 free reprints. Additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$108.00 a year to individual members of supporting institutions.

Subscriptions, orders for numbers issued in the last three calendar years, and changes of address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Old back numbers obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$,
the American Mathematical Society's $\mathcal{T}\mathcal{E}\mathcal{X}$ macro system.

Copyright ©1994 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 167 No. 1 January 1995

Local reproducing kernels on wedge-like domains with type 2 edges	1
AL BOGGESS and ALEXANDER NAGEL	
Discriminants of involutions on Henselian division algebras	49
MAURICE CHACRON, H. DHERTE, JEAN-PIERRE TIGNOL, ADRIAN R. WADSWORTH and V. I. YANCHEVSKIĀ	
Essential tori obtained by surgery on a knot	81
MARIO EUDAVE-MUÑOZ	
Non-compact totally peripheral 3-manifolds	119
LUKE HARRIS and PETER SCOTT	
Some representations of TAF algebras	129
JOHN LINDSAY ORR and JUSTIN PETERS	
A non-Haken hyperbolic 3-manifold covered by a surface bundle	163
ALAN W. REID	
The nonhomogeneous minimal surface equation involving a measure	183
WILLIAM KARL ZIEMER	