

Pacific Journal of Mathematics

LENGTH OF JULIA CURVES

DAVID H. HAMILTON

LENGTH OF JULIA CURVES

DAVID H. HAMILTON

Let the Fatou set of a meromorphic function f have two components with Denjoy-Wolff points at which f is not transcendental. Then the Julia set J is a “circle/line” or is nowhere rectifiable. In particular if f is rational and J is a Jordan curve then it follows it follows (in the second case) that $\dim(J) > 1$.

Let $F : \mathbb{C} \rightarrow \mathbb{C}$ be a meromorphic function. The Julia set $J = J(F)$ is the complement of the set of normality of the iterates F^n . Fatou (1920) proved that if F is rational and if J is a closed Jordan curve then either J is a “circle/line” or it has a dense set of points where there is no tangent. Under various special assumptions on the behaviour of F on J one can say more about the case where J is a closed Jordan curve not equal to a circle/line: Fatou claimed that if “ F is expanding on J ” then J has Hausdorff 1-measure $\lambda(J) = \infty$. (This was proved by Brolin (1965).) Also if “ F is expanding on J ” Sullivan (1983) proved that the Hausdorff dimension satisfies $\dim(J) > 1$. Sullivan does this by applying his construction of the conformally invariant measure to Brolin’s result. The main result of this paper is that these results hold for arbitrary rational functions.

THEOREM 1. *Let $F : \mathbb{C} \rightarrow \mathbb{C}$ be a rational function. Suppose that the Julia set J is a Jordan curve. Then J is a circle/line or $\dim(J) > 1$.*

REMARKS. (i) Actually Przytcki, Urbanski and Zdunik [14] had proved Theorem 1 for the “repellant case”. Also Zdunik [17] proves that if F is any polynomial with Julia set J a Jordan curve then $\dim(J) > 1$. We only prove $\lambda(J) = \infty$ and then make good use of Decker and Urbanski [5].

(ii) Note that we require J to be a closed Jordan curve. Our result is considerably more general than for Jordan curves. We only require that F has two forward invariant (disjoint simply connected)

domains Ω_1 and Ω_2 . Then we prove that either $\mathbb{C} - (\Omega_1 \cup \Omega_2)$ is a circle/line or $\lambda(\mathbb{C} - (\Omega_1 \cup \Omega_2)) = \infty$. The case that J is a Jordan arc has to be treated separately and follows from a general result we prove for “dendrites”, i.e. when both J and $\mathbb{C} - J$ are connected.

Our method depends primarily on the harmonic measure

$$\omega_j = \omega_j(z_j, E, \Omega_j),$$

i.e. the value of the function harmonic on Ω_j with boundary value 1 on E and 0 on $\partial\Omega_j - E$ taken at some point z_j inside Ω_j . The actual points z_j are unimportant so we henceforth refer only to the harmonic measures $\partial\omega_j$. Our basic result is:

THEOREM 2. *Let F be a rational function with two forward invariant (disjoint) domains Ω_1 and Ω_2 . Then either $\partial\Omega_1 = \partial\Omega_2$ is a circle/line or*

$$d\omega_1 \perp d\omega_2.$$

REMARKS. (i) This immediately implies (by classical results of Riesz, see [13]) that in the second case J is nowhere rectifiable. However we can improve this considerably by using results of Bishop, Carleson, Garnett and Jones [3]. They characterise the disjoint domains for which the harmonic measures are mutually singular. Let T be the set of points on J where there is a tangent. If the mutual boundary is a closed Jordan curve J singularity is equivalent to $\lambda(T) = 0$. In the general case one only has to replace tangents by “mutual sectorial accessibility”. In any case there is no “tangent” at almost all points, considerably strengthening the old result of Fatou.

We prove Theorem 2 directly as a consequence of our rigidity theory for inner functions f , see [8], [9] and [10]. Now f is inner if it is a bounded analytic function on the unit disk \mathbb{D} with radial boundary values $f(z) \in \partial\mathbb{D}$ for $z \in \partial\mathbb{D}$ (a.e.). Shub and Sullivan [15] dealt essentially with the special case where f is rational, i.e. a finite Blaschke product, and f fixes a point inside the disk. They prove that if there is a (mutually) absolutely continuous map $\phi : \partial\mathbb{D} \rightarrow \partial\mathbb{D}$ so that $g = \phi \circ f \circ \phi^{-1}$ is also Blaschke (on $\partial\mathbb{D}$) then ϕ is Mobius (bilinear or the complex conjugate of a bilinear

map). In the general case the real difficulty is dealing with functions with “Denjoy-Wolff” point on $\partial\mathbb{D}$, the generalization to inner functions being straight forward. The general result is that inner f have nontrivial conjugations by absolutely continuous ϕ to other inner functions if and only if f is not ergodic on $\partial\mathbb{D}$. By using the techniques of this method we shall obtain the stronger result on the singularity of harmonic measures.

Before proceeding consider some examples. If the Julia set J of a rational function f is a circle/line then f is essentially a Blaschke product. Examples where J is not a circle/line may be obtained by quasiconformal conjugation of a Blaschke product to a rational function F . Now if the fixed point is not on the Jordan curve J we have the expanding case. So this theory is new only for the case where we have a neutral attracting point, i.e. petals. Similar results are also seen in the usual family of suspects $z^2 + c$. For c close to 0 we have that J is a closed Jordan curve surrounding a strongly attracting point. However for some values like $c = 1/4$ although J remains a Jordan curve the bounded attracting point is also on J . For other values of c we obtain the so called “rabbit ears”. In this case f^2 is invariant on two bounded domains Ω_1 and Ω_2 as well as one unbounded domain Ω_3 . Now Theorem 2 implies that the relevant harmonic measures are mutually singular. This is hardly surprising for Ω_1 and Ω_2 as they only touch at one boundary point. However for Ω_1 and Ω_3 this is nontrivial. (Now we do not necessarily have that $\partial\Omega_j$ have infinite 1-measure, only that if they do have finite 1-measure then they meet on a set of zero measure.)

Also Theorems 1 and 2 will also be seen to hold for a large class of meromorphic functions F . However we need to restrict ourselves to meromorphic functions which are ergodic with respect to the harmonic measure on the invariant domains.

Now F is ergodic if there is no set $E \subset \partial\Omega_j$ with

$$0 < \omega_j(E) < 1 \text{ and } F^{-1}(E) = E.$$

A sufficient condition for this is that the “Denjoy-Wolff” point for F on a forward invariant domain Ω , i.e. the point w in the closure of Ω so that for all z in Ω $F^n(z) \rightarrow w$, should be a finite point (so that F is analytic at w). The function $F = \tan(z)$ is an example. However there are meromorphic functions, such as $F = 2z + \tan(z)$

(with “Denjoy-Wolff” point at ∞) which are not ergodic.

THEOREM 3. *Let F be a meromorphic function with two forward invariant (disjoint) domains Ω_1 and Ω_2 upon which F is ergodic with respect to the harmonic measures $d\omega_j$. Then either $\partial\Omega_1 = \partial\Omega_2$ is a circle/line or the harmonic measures are singular.*

REMARKS. (i) We also give an example of a meromorphic function F with Julia set J being a Jordan curve separating the plane into forward invariant unbounded domains with harmonic measures which are mutually absolutely continuous but J is not a circle/line. Indeed the harmonic measures are absolutely continuous with respect to $d\lambda$ on J . However we do not know if such J can be of locally finite length (although they are never smooth).

Finally we deal with the case where J is a Jordan arc or even a “dendrite”. The general case is for a simply connected forward invariant domain Ω . In many situations the points of $\partial\Omega$ are accessible from several different directions, i.e. to be more precise several prime ends correspond to the same boundary point. For example if J is a Jordan arc then (except at the end points) two prime ends correspond to each point of J . At the opposite extreme is the “rabbit” where a countable number of prime ends cut off some of the boundary points. Although we cannot speak of a homeomorphism of the circle generated by this correspondence we can obtain a bijection. Thus as above we shall prove:

THEOREM 4. *Let F be a rational function so that the Julia set J and its complement are connected. Then J is a circular arc/line interval or $\lambda(J) = \infty$.*

REMARKS. Actually one proves that the resulting bijection is trivial or singular. If J is actually a Jordan arc then Theorem 4 also holds for meromorphic functions.

The author would like to thank the referee and acknowledges conversations with M. Yacobson and P.W. Jones.

2. Conformal Pullback to the disk. Let F be meromorphic on \mathbb{C} with Julia set J . We consider a forward invariant simply connected component Ω of $\mathbb{C} - J$. Let h be the conformal mapping of the unit disk D onto Ω . We have by Fatou’s Theorem

$$\lim_{r \rightarrow 1} h(rz) \in \partial\Omega, z \in \partial\mathbb{D} \text{ (a.e.)}.$$

Consequently we have:

LEMMA 1. *The function*

$$f \equiv h^{-1} \circ F \circ h$$

is a well defined inner function of \mathbb{D} into \mathbb{D} .

Now f is a well defined analytic map of the unit disk into itself. So we have only to check that f is inner. If not there is a subset E of the circle with $\lambda(E) > 0$ and

$$\lim_{r \rightarrow 1} f(rz) \in \mathbb{D}, x \in E \text{ (a.e.)}.$$

Then the radial limits

$$G \equiv h(E_0), \lambda(E - E_0) = 0,$$

is a subset of $\partial\Omega$ but $F(G) \subset \Omega$! This of course is impossible.

Notice that f is bilinear if and only if F is conformal on Ω . Using the complete classification of dynamical behaviour this happens only if there is a bilinear mapping T of the disk onto itself so that $f = T \circ g \circ T^{-1}$ where

$$g(z) = e^{it}z,$$

where t/π is irrational. Thus f is ergodic, i.e. fixes no nontrivial set of positive length on $\partial\mathbb{D}$.

Now if F fixes a point w of Ω then by the principle of harmonic measure the measure $d\omega(w, z, \Omega)$ is F invariant. A simple argument shows that f is ergodic with respect to this finite measure.

Doering and Mané [6] considered the case for parabolic basins of rational functions and proved the corresponding f is ergodic on the unit circle even though there is no invariant finite measure. Their proof only depends on the following sufficient condition due to Aaronson [1]:

LEMMA A. *An inner function $f : \mathbb{D} \rightarrow \mathbb{D}$ is ergodic provided*

$$\sum_{n=1}^{\infty} (1 - |f^n(0)|) = \infty.$$

REMARK. Actually, see [7], this is equivalent to f is recurrent, i.e. for any subset A of positive measure and for all $z \in A$ (a.e.) $f^n(z) \in A$ for infinitely many positive integers n . (The relationship between recurrence and ergodicity of inner functions is not what one would expect, see Doering and Mané.)

Let Ω be a parabolic domain with attractive boundary point w (i.e. Denjoy-Wolff point). By the classical description of the petals we see that at w the domain Ω is asymptotic to a sector of angle $2\pi/k$, where we can assume (without loss of generality, see Beardon [2, p. 116-122])

$$F(z) = w + (z - w) - (z - w)^{k+1} + O((z - w)^{2k+1}).$$

Supposing that the conformal mapping h maps 1 to w then

$$h(z) = w + a_1(z - 1)^{k/2} + \dots$$

Thus by a simple computation for any $\alpha > 1/2$ for large enough n :

$$1 - |f^n(0)| > \frac{1}{n^\alpha}.$$

Thus as in [6] we obtain from Lemma A the first part of:

LEMMA 2. *Let Ω be a foreward invariant simply connected component of set of normality of a meromorphic function F with finite Denjoy-Wolff point. Then the corresponding inner function f is ergodic. Furthermore for any set $E \subset \partial\mathbb{D}$ with $f(E) \subset E$ (a.e.) we have $\lambda(E) = 0$ (a.e.) or $E = \partial\mathbb{D}$ (a.e.).*

To prove the last part of this lemma we define for fixed positive integer m

$$G = \{z \in \partial\mathbb{D} - E : f^m(z) \in E\}.$$

Now if $\lambda(E) > 0$ then $f^k(z) \in E$, which is disjoint from G , for $k > m - 1$ as f is recurrent $\lambda(G) = 0$, for all m . Therefore E is

backwards as well as forward invariant. Hence as f is ergodic we have the result required.

In order to prove our theorems we use conformal pullback to produce inner functions f_1 and f_2 . The unit circle has subsets E and G both of positive length and there is a (mutually) absolutely continuous bijection $\phi : E \rightarrow G$ so that on G

$$f_2 = \phi \circ f_1 \circ \phi^{-1}.$$

Our basic tool is the following result proved in [8],

THEOREM H1. *Suppose that f_1, f_2 are ergodic inner (rational) functions. Now suppose there is an absolutely continuous homeomorphism (bijection) $\phi : \partial\mathbb{D} \rightarrow \partial\mathbb{D}$ so that*

$$f_2 = \phi \circ f_1 \circ \phi^{-1}$$

on $\partial\mathbb{D}$ (a.e.). Then there is a Möbius transformation T of the disk onto itself:

$$\phi = T.$$


REMARK. Notice that our result is somewhat stronger for finite Blaschke products. This will be used in the proof of Theorem 4.

3. Boundary values of conformal mappings. In this section we collect some basic ideas on the boundary behaviour of conformal maps $h : \mathbb{D} \rightarrow \Omega$. A full account may be found in Pommerenke [13].

For a point $w \in \partial\mathbb{D}$ which is the endpoint of an open Jordan arc α contained inside Ω there is a point $z \in \partial\mathbb{D}$ so that z is the endpoint of $h^{-1}(\alpha)$ and h has radial limit w at z . We want to consider more complicated geometric behaviour. We say that w is accessible in a sector if there is an open triangle in Ω with w as one vertex. The set of points on $\partial\Omega$ which are sectorial limits is denoted by S . By a theorem of McMillan the harmonic measure $d\omega$ and length $d\lambda$ are mutually absolutely continuous on S . Also on S , except for a set of measure zero, every point has a “one sided tangent”. There may also be a nontrivial set R complementary to S , where up to a set of zero harmonic measure every point $w = h(z)$ is a twist point. This means that $\arg(h(rz) - w)$ is unbounded above and below as $r \rightarrow 1$.

Next we need to consider disjoint simply connected domains Ω_1 and Ω_2 with harmonic measures $d\omega_1$ and $d\omega_2$. We make use of a basic result of Bishop, Carleson, Garnett and Jones [3] who prove

THEOREM B. *The combined boundary $\partial\Omega_1 \cup \partial\Omega_2$ has a partition into disjoint sets $S_1 \cap S_2$, B_1 and B_2 where $\omega_1(B_2) = \omega_2(B_1) = 0$. The set $S_1 \cap S_2$ consists of all boundary points which are accessible in a sector from simultaneously both Ω_1 and Ω_2 . Furthermore almost all (harmonic measure or $d\lambda$ it does not matter) points of $S_1 \cap S_2$ are points where the two domains are mutually tangent.*



We are now ready to apply these concepts to ergodic inner functions. First we allow our domains Ω_1 and Ω_2 to be forward invariant simply connected domains of the region of normality of a meromorphic function F . Now, except for a countable number of points, F and its iterates are locally conformal. Thus the set $S_1 \cap S_2$ is forward and hence by Lemma 2 backward invariant. When we pullback to the inner functions f_j we obtain invariant sets $E_j = h_j^{-1}(S_1 \cap S_2)$. By

the previous section our inner functions are ergodic and have non-trivial invariant sets. Thus the sets $S_1 \cap S_2$ have harmonic measure 1 or 0 in both domains. Furthermore if the two harmonic measures are not singular then, by Theorem B, we have that $S_1 \cap S_2$ has harmonic measure 1 in both domains. From this we construct the mapping

$$\phi = h_2^{-1} \circ h_1$$

defined first on a subset of positive measure on the unit disk. Thus ϕ may be extended to an absolutely continuous bijection defined on the unit circle (a.e.). Finally by simple topology, as order is preserved, ϕ extends continuously to an absolutely continuous homeomorphism of the unit circle. Thus from the conjugation

$$f_j = h_j^{-1} \circ F \circ h_j$$

we have proved

LEMMA 3. *If $S_1 \cap S_2$ has positive measure then there is an absolutely continuous homeomorphism $\phi : \partial\mathbb{D} \rightarrow \partial\mathbb{D}$ so that $f_2 = \phi \circ f_1 \circ \phi^{-1}$ (a.e. on $\partial\mathbb{D}$).*

4. Proof of Theorems 1-3. From previous results we have first to prove Theorem 3. Assuming that harmonic measure is not singular then by applying Theorem H1 to Lemma 3 we obtain a Möbius map $T : \partial\mathbb{D} \rightarrow \partial\mathbb{D}$ so that

$$h_2 \circ T = h_1 \text{ (a.e. on } \partial\mathbb{D}\text{)}.$$

Up till now we had some freedom in our normalization of the Riemann maps. These may now be normalised so that we can assert that $T = z$ or $T = \bar{z}$. Consider the first case. Thus the conformal maps agree (a.e.) on the unit circle. Hence they have the same Laurent coefficients for the expansions in the powers of z . In other words h_2 is the analytic continuation of h_1 . So there is a bilinear mapping h so that $h_1(z) = h(z)$ and $h_2(z) = h(z)$ for $|z| > 1$. In particular the domains have mutual boundary which is a circle/line.

In the second case T must be antilinear. Consequently

$$h_2\left(\frac{1}{z}\right) = h_1(z) \text{ (a.e. on } \partial\mathbb{D}\text{)}.$$

By comparison of Laurent coefficients again we have

$$h_2\left(\frac{1}{z}\right) = h_1(z) \text{ for all } |z| \neq 1.$$

This implies that $\Omega_1 = \Omega_2$, contrary to our assumption. This completes the proof of Theorem 3. Theorem 2 immediately follows as by using a bilinear conjugation we can always assume any Denjoy-Wolff points of a rational function can be made finite. We complete the proof of Theorem 1 in the next section.

5. $\dim(J) > 1$. Decker and Urbanski [5] define a rational map F to be parabolic on J if there are no critical points on J but there exist at least one rationally indifferent periodic point. Now set $\delta = \dim(J)$. They generalise a result of Sullivan by proving that there is a unique δ -conformal probability measure $d\mu$ on J , i.e.

$$\mu(F(A)) = \int_A |F'|^\delta d\mu$$

for all A in J upon which F is 1 : 1. They prove:

LEMMA D. *If $\delta = 1$ then there is a positive finite constant c so that $d\mu = cd\lambda$.*

Let us now consider the case that J is a Jordan curve. There can be no critical points on J , as these would cause branching. If neither of the attractive points lie on J then F is expanding. So this is the Sullivan case. In the other case any attractive point on J must necessarily be rationally indifferent, in fact must be a petal of order 1 or 2. This is the parabolic case. Now by Lemma D and Theorem 2 we deduce that $\delta = 1$ only in the trivial case that J is a circle/line. This also completes the proof of Theorem 1.

This argument will not hold for Jordan arcs or dendrites which in general contain critical points.

6. Multiply connected case. It is natural to ask in what sense do these results generalise to nonsimply connected domains. That is we suppose we have a rational function F with two disjoint forward invariant components Ω_j on the set of normality, one of which at least is multiply connected.

First we fix one such domain which we call Ω . One sees that Lemma 1 and 2 generalise without any problem. For as $\partial\Omega$ has at

least 3 points there is a conformal mapping h of the unit disk \mathbb{D} onto the universal covering of Ω . Furthermore as $\partial\Omega$ in fact has positive logarithmic capacity a result of Nevanlinna for functions of bounded characteristic implies

$$\lim_{r \rightarrow 1} h(rz) \in \partial\Omega \quad (z \in \partial\mathbb{D} \text{ a.e.}).$$

Thus as before $f \equiv h^{-1} \circ F \circ h$ will be inner.

The simplest case of a nonsimply connected domain is when Ω_1 say is a Herman ring. This is the only other finitely connected case. Then F acting on Ω_1 is conformally conjugate to an irrational rotation f_1 of an annulus $D_1 = \{r < |z| < 1\}$:

$$f_1 \equiv h_1^{-1} \circ F \circ h_1$$

where $h_1 : D_1 \rightarrow \Omega_1$ is the conformal mapping which maps the boundary component $\partial\mathbb{D}$ to the outside of the boundary component $\partial_0\Omega_1$. Note that F leaves the boundary components of $\partial\Omega_1$ invariant so that the pullback to the unit disk f cannot be ergodic. Now without loss of generality the other invariant domain is in the unbounded component of $\mathbb{C} - \Omega_1$.

There are several possibilities:

- (i) Ω_2 is simply connected (and contains ∞).
- (ii) Ω_2 is doubly connected.
- (iii) Ω_2 is infinitely connected.

First we assume that Ω_2 is simply connected. Case (iii) shall be dealt with later. For case (i) we pullback Ω_2 to $D_2 = \{1 < |z|\}$ by conformal mapping $h_2 : D_2 \rightarrow \Omega_2$ taking $\partial\mathbb{D}$ to the boundary component $\partial\Omega_2$. So we obtain “inner functions” $f_j : D_j \rightarrow D_j$ as usual.

In this section we considerably weaken the hypothesis that the harmonic measures are nonsingular. All we require is that

$$\omega_j(\partial_0\Omega_1 \cap \partial\Omega_2) > 0, \text{ for } j = 1 \text{ and } 2.$$

Now as $\partial_0\Omega_1 \cap \partial\Omega_2$ is forward invariant under F by Lemma 2

$$\omega_2(\partial_0\Omega_1 \cap \partial\Omega_2) = 1.$$

Also as f_1 is an irrational rotation there are no nontrivial forward invariant sets. Thus

$$\omega_1(\partial_0\Omega_1 \cap \partial\Omega_2) = \omega_1(\partial_0\Omega_1).$$

Thus in particular $\partial_0\Omega_1 = \partial\Omega_2$ although this alone does not imply that there may be other components of the set of normality which lie between the two invariant domains. Under these circumstances it is easy to construct the bijection φ giving the conjugation from f_1 to f_2 on $\partial\mathbb{D}$. First observe that for any continua K in $\partial_0\Omega_1$ there is a unique subarc $I_j \subset \partial\mathbb{D}$ corresponding to K under h_j in the prime-end topology. Now by the above result on harmonic measure

$$\omega_1(K) > 0 \iff \omega_2(K) > 0.$$

Thus in mapping I_1 to I_2 there is no collapsing of arcs to points or vice-versa. Consequently, as by simple topology order is preserved, the bijection φ is defined, however we do not have that it is nonsingular merely that it is a homeomorphism.

Now as f_1 is just an irrational rotation we must have that its conjugate f_2 is 1:1 and is therefore a rotation too. By the classification of invariant domains in this case Ω_2 is also a Herman ring. As this was the second case we now assume it. So we deal with case (ii). In this case we pullback Ω_2 to $D_2 = \{1 < |z| < s\}$ by conformal mapping $h_2 : D_2 \rightarrow \Omega_2$ taking $\partial\mathbb{D}$ to the inner boundary component $\partial_i\Omega_2$. So we obtain "inner functions" $f_j : D_j \rightarrow D_j$ which are now irrational rotations:

$$f_j = e^{i\theta_j} z.$$

Now as above we have the dichotomy

$$\omega_j(\partial_0\Omega_1 \cap \partial\Omega_2) = 0, \text{ for } j = 1 \text{ or } 2,$$

or

$$\omega_j(\partial_0\Omega_1 \cap \partial\Omega_2) = 1, \text{ for } j = 1 \text{ or } 2.$$

Once again we concentrate on the second case and produce a homeomorphism φ of the circle onto itself so that

$$e^{i\theta_2}\varphi(z) = \varphi(e^{i\theta_1}z), \quad z \in \partial\mathbb{D},$$

where θ_j/π is irrational. Let φ have Fourier expansion

$$\sum_{-\infty}^{+\infty} a_k z^k, \quad |z| = 1.$$

Thus for each k :

$$e^{i\theta_2} a_k = e^{ik\theta_1} a_k.$$

Therefore $a_k = 0$ or $k\theta_1 - \theta_2 = 2\pi m_k$. But as θ_2/π is irrational there is at most one k for which $a_k \neq 0$. But as φ is a homeomorphism this implies that it is Möbius. Since we still have choice in the normalization of the conformal maps we may take $\varphi(z) = z$. Thus the conformal maps h_j satisfy

$$h_1(z) = h_2(z) \text{ on } \partial\mathbb{D} \text{ (a.e.)}.$$


Therefore by classical Laurent series h_2 is the analytic continuation of h_1 . This implies that the common boundary B of the Herman rings is the conformal image of $\partial\mathbb{D}$. Hence the ‘‘Herman Rings’’ are infact rings of a larger Herman Ring obtained by pasting across an analytic curve B . But then B cannot possibly be part of the set of nonnormality! Thus in all cases the harmonic measure of the common boundary must be zero.

Finally we have to deal with the case where $\Omega = \Omega_1$ is infinitely connected. Now F will have a (super) attractive point in Ω we see that the corresponding f may be assumed to fix 0. Then f will be ergodic and recurrent. We shall now prove that either the boundary components of $\mathbb{C} - \Omega$ each have harmonic measure zero or the boundary components of $\mathbb{C} - \Omega$ are all points (and there will be only one invariant domain). Consider any components K of $\mathbb{C} - \Omega$. Suppose that the harmonic measure $\omega(K) > 0$. Now let the pullback of ∂K under h be the set E . As f is recurrent there is a set $E_0 \subset E$ with $\lambda(E_0) > 0$ and a number n so that $f^n(E_0)$ is contained in E . But this means, as F maps components of $\mathbb{C} - \Omega$ to components of $\mathbb{C} - \Omega$, that $F^n(K) = K$. Without loss of generality $n = 1$. Thus E is forward invariant under f . But as f fixes 0 we see $\lambda(f^{-1}(E)) = \lambda(E)$ and therefore $f^{-1}(E) = E$ (a.e.). So as f is ergodic $\omega(K) = 1$. Thus the other components of $\mathbb{C} - \Omega$ are points. Now at least one of these components, Q say, is mapped by F onto

K otherwise f is normal on $\mathbb{C} - K$. But then Q must be a continuum too which is a contradiction. Therefore all components have zero harmonic measure. Thus we proved the multiply connected case is trivial for harmonic measures.

THEOREM 5. *If Ω_j are disjoint forward invariant domains at least one of which is multiply connected then $0 = \omega_1(\partial\Omega_1 \cap \partial\Omega_2) = \omega_2(\partial\Omega_1 \cap \partial\Omega_2)$.*

7. Dendrites.



It is known, see [2, p. 257], that if F is a polynomial then J is the interval $[-1, 1]$ if and only if F is one of the Tchebyshev polynomials T_n (or $-T_n$) where $T_n(\cos(t)) = \cos(nt)$. In general a rational function F has Julia set $[-1, 1]$ if and only if

$$F = h \circ f \circ h^{-1}, \text{ where } h = \frac{1}{2} \left(z + \frac{1}{z} \right),$$

where f is an ergodic and symmetric finite Blaschke product. Now, in general but not if F is a polynomial (by Hubbard's "Ecale" construction), the Teichmüller space of such rational functions is non-trivial. This means there is a quasiconformal mapping $\Phi : \mathbb{C} \rightarrow \mathbb{C}$ so that

$$F^* = \Phi \circ F \circ \Phi^{-1}$$

is a rational function not conjugate to F by any bilinear mapping. Consequently F^* has Julia set $J^* = \Phi(J)$ which in general is not an interval.

Actually we shall deal with more general objects than an open Jordan arc. Assuming that J and $\mathbb{C} - J$ are connected we have a single forward invariant simply connected domain Ω . In analogy to before we define S to be the set of points $\zeta \in S$ so that there exists $r > 0$ and directions t_j , $j = 1$ and 2 , so that there are disjoint sectors

$\Sigma_j = \{z : |z - \zeta| < r, \arg((z - \zeta)e^{-it_j}) < \theta_j\}$, of width $2\theta_j > \pi/3$, belonging to different components of $\Omega \cap \{z : |z - \zeta| < r\}$. Now there are at most two such sectors at each point. If J has (locally) finite length then almost all points of J belong to S . Notice that S is F invariant (a.e.). As the conformal pullback f is ergodic then either $\omega(S) = 0$ or $\omega(S) = 1$. In the latter case we define a bijection $\phi : \partial\mathbb{D} \rightarrow \partial\mathbb{D}$ (a.e.) as the pullback of the map swapping points in adjacent sectors - the identity, i.e. $\phi = h^{-1} \circ I \circ h$ which gives an involution (a.e.). By MacMillan's theorem ϕ is absolutely continuous. But as $f = h^{-1} \circ F \circ h$

$$\begin{aligned} \phi \circ f \circ \phi^{-1} &= h^{-1} \circ I \circ h \circ h^{-1} \circ F \circ h \circ h^{-1} \circ I \circ h \\ &= h^{-1} \circ I \circ F \circ I \circ h \\ &= h^{-1} \circ F \circ h = f, \end{aligned}$$

i.e. ϕ conjugates f to itself. Consequently Theorem H1 shows that ϕ is Möbius. After a trivial renormalisation, if necessary, we see that we may assume

$$\phi(z) = \bar{z}.$$

Therefore from above

$$f(\bar{z}) = \overline{f(z)} \text{ and } h(\bar{z}) = \overline{h(z)}.$$

This implies that J is symmetric with respect to the real axis. If J is not a subinterval of the real line there will be a subset of positive

harmonic measure where the bijection is not defined. This completes the proof of Theorem 4.

8. Counterexample. We use quasi-conformal deformation (see Beardon [2]) to prove

THEOREM 6. *There exists a meromorphic function F with Julia set J a Jordan curve through ∞ and with two forward invariant components Ω_1 and Ω_2 of $\mathbb{C} - J$ so that the harmonic measures satisfy $d\omega_1 \sim d\omega_2$, indeed $d\lambda \gg d\omega_j$.*

We begin with the functions

$$G_s(z) = sz + \tan(z), s > 0.$$

It is easy to see that for $s \geq 1$ these have Julia set J equal to the real line and the upper and lower half planes are invariant. Also by [6] for $s > 1$ the pullback to the disk, f , is not ergodic. Now in [8] we proved

THEOREM H2. *If $G_s : \mathbb{H} \rightarrow \mathbb{H}$ is inner but not ergodic there exists a quasiconformal mapping*

$$\Phi : \mathbb{C} \longrightarrow \mathbb{C}$$

preserving the upper half plane so that

- (i) $H_s = \Phi \circ G_s \circ \Phi^{-1}$ is meromorphic, inner on \mathbb{H} ,
 - (ii) $H_s \neq T \circ G_s \circ T^{-1}$ (a.e. on \mathbb{R}) for all Möbius T .
- Also the restriction of Φ to the real axis is absolutely continuous.*

Actually we do not use the last unadorned fact but the part of the construction that lies underneath it. Let Φ have complex dilatation

$$\mu(z) = \frac{\bar{\partial}\Phi}{\partial\Phi}.$$

By (i) the dilatation μ is G_s invariant, i.e.

$$\mu(G_s(z)) = \frac{G'_s(z)}{G'_s(z)}\mu(z)$$

and as the real line is mapped to the real line

$$\overline{\mu(z)} = \mu(\bar{z}).$$

The important property of μ is that it is supported by a set A which only has nontangential cluster points on a set E on the real axis with $\lambda(E) = 0$, proved in [8]. Recall that x is a nontangential cluster point of A if there is an angle θ , $0 < \theta < \pi/2$ with Stolz angle

$$\Sigma = \{\operatorname{Im}(z) > 0 : |z| < r, |\operatorname{Arg}(z - x)| < \theta\}.$$

So there is a sequence $z_n \in \Sigma \cap A$, $z_n \rightarrow x$.

We now define a second quasiconformal mapping

$$\psi : \mathbb{C} \longrightarrow \mathbb{C}$$

as a homeomorphic solution of the Beltrami equation

$$\frac{\bar{\partial}\psi}{\partial\psi} = \begin{cases} \mu(z), & \operatorname{Im}(z) > 0, \\ 0, & \text{otherwise,} \end{cases}$$

which fixes infinity, see [11].

As above ψ has a G_s invariant dilatation and therefore as in Bear-don

(iii) $F_s = \psi \circ G_s \circ \psi^{-1}$ is meromorphic.

We note

(iv) $F_s \neq T \circ G_s \circ T^{-1}$ for all Möbius T .

Otherwise as the restriction of ψ to the lower half plane is conformal we see that ψ will be bilinear on the lower half plane. But then by the usual Bers theory of universal Teichmüller space the dilatation μ is trivial, and Φ will be bilinear on the real line, contradicting (ii).

The image of the real line under ψ is a Jordan curve J through infinity. Furthermore as G_s has Julia set equal to the real line, F_s has Julia set equal to J . Thus there are two forward invariant components Ω_1 and Ω_2 of $\mathbb{C} - J$ with the harmonic measures $d\omega_1$, $d\omega_2$. By suitable choice of the conformal mappings

$$\begin{cases} h_1 : \{\operatorname{Im}(z) > 0\} \longrightarrow \Omega_1 \\ h_2 : \{\operatorname{Im}(z) < 0\} \longrightarrow \Omega_2 \end{cases}$$

we see that on \mathbb{R}

$$\Phi = h_2^{-1} \circ h_1.$$

Thus the harmonic measures are mutually absolutely continuous.

The second part of Theorem 6 is an application of the main theorem of [10] which with the above notation is:

THEOREM H3. *Suppose that the complex dilatation of ψ is supported by a set A which only has nontangential cluster points on a set E on the real axis with $\lambda(E) = 0$ then*

$$d\lambda_J \circ \psi \gg d\lambda_{\mathbb{R}}.$$

This concludes the proof of Theorem 6.

REFERENCES

- [1] Aaronson, J., *Ergodic theory for inner functions of the upper half-plane*, Ann. Inst. H. Poincaré, Sec. B, vol. XIV (1978), 233-253.
- [2] Beardon, A., *Iteration of rational functions*, Grad. Texts in Math., no. 132, Springer-Verlag 1991.
- [3] Bishop, C.J., Carleson, L., Garnett, J.B. and Jones, P.W. *Harmonic measures supported on curves*, Pacific J. Math., **138** (1989), 233-236.
- [4] Brolin, H., *Invariant sets under iteration of rational functions*, Ark. Mat., **6** (1965), 103-144.
- [5] Decker, M. and Urbanski, M., *Geometric measures for parabolic rational maps*, Ergod. Th. and Dynam. Sys., **12** (1992), 53-66.
- [6] Doering, C. and Mané, R., *The dynamics of inner functions*, Ensaissos Matematicos, **3** (1992), 1-79.
- [7] Eremenko, A.E. and Lyubich, M. Yu., *The dynamics of analytic transformations*, Leningrad Math. J., **1** (1990).
- [8] Hamilton, D.H., *Absolutely continuous conjugacies of Blaschke products*, to appear in Advances in Math.
- [9] ———, *Absolutely continuous conjugacies of Blaschke products III*, to appear in Journal d'Analyse.
- [10] ———, *Quasiconformal mappings absolutely continuous on the circle*, to appear.
- [11] Letho, O and Virtanen, K.I., *Quasiconformal mappings in the plane*, Springer-Verlag, 1973.
- [12] Nevanlinna, R., *Analytic functions*, Springer-Verlag 1953.
- [13] Pommerenke, C., *Boundary Behaviour of Conformal Maps*, Springer-Verlag Grundlehren der Math. Wissen., **299** (1991).
- [14] Przytycki, M., Urbanski, M., Zdunik, A., *Harmonic, Gibbs and Hausdorff measures on repellers for holomorphic maps*, Ann. Math., **130** (1989), 1-40.

- [15] Shub, M. and Sullivan, D., *Expanding endomorphisms of the circle revisited*, Ergod. Theo. and Dynam. Sys., **5** (1985), 285-289.
- [16] Sullivan, D., *Conformal dynamical systems*, 725-752, Lecture Notes in Math., vol. 1007, Springer-Verlag, 1983.
- [17] Zdunik, A., *Hausdorff measure of maximal measure for rational maps*, preprint, Inst. Math. Polish Acad. Sci. Warsaw (1987).

Received September 17, 1992 and in revised form May 24, 1994. This research was supported in part by the National Science foundation.

UNIVERSITY OF MARYLAND

COLLEGE PARK, MD 20742

E-mail address: dhh@math.umd.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

EDITORS

Sun-Yung Alice Chang (Managing Editor)
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

F. Michael Christ
University of California
Los Angeles, CA 90095-1555
christ@math.ucla.edu

Thomas Enright
University of California
San Diego, La Jolla, CA 92093
tenright@ucsd.edu

Nicholas Ercolani
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

Robert Finn
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

Vaughan F. R. Jones
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

Steven Kerckhoff
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

Martin Scharlemann
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

Gang Tian
Courant Institute
New York University
New York, NY 10012-1100
tiang@taotao.cims.nyu.edu

V. S. Varadarajan
University of California
Los Angeles, CA 90095-1555
vsv@math.ucla.edu

SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
STANFORD UNIVERSITY
UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
UNIVERSITY OF UTAH
UNIVERSITY OF WASHINGTON
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$108.00 a year to individual members of supporting institutions.

Subscriptions, orders for back issues published within the last three years, and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,
the American Mathematical Society's TEX macro system.
Copyright © 1995 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 169 No. 1 May 1995

Minimal sets of periods for torus maps via Nielsen numbers	1
LLUÍS ALSSEDÀ, STEWART BALDWIN, JAUME LLIBRE, RICHARD SWANSON and WIESLAW SZLENK	
Diagonalizing Hilbert cusp forms	33
TIMOTHY ATWILL	
A splitting criterion for rank 2 vector bundles on \mathbf{P}^n	51
EDOARDO BALLICO	
Controlling Tietze-Urysohn extensions	53
MARC FRANTZ	
Length of Julia curves	75
DAVID H. HAMILTON	
On the uniqueness of capillary surfaces over an infinite strip	95
JENN-FANG HWANG	
Volume estimates for log-concave densities with application to iterated convolutions	107
MARIUS JUNGE	
A reflection principle in complex space for a class of hypersurfaces and mappings	135
FRANCINE ANTOINETTE MEYLAN	
Jean Bourgain's analytic partition of unity via holomorphic martingales	161
PAUL F.X. MÜLLER	
Characters of Brauer's centralizer algebras	173
ARUN RAM	