

# *Pacific Journal of Mathematics*

**ON THE UNIQUENESS OF CAPILLARY SURFACES OVER AN  
INFINITE STRIP**

JENN-FANG HWANG

# ON THE UNIQUENESS OF CAPILLARY SURFACES OVER AN INFINITE STRIP

JENN-FANG HWANG

**In 1987, Tam proved that the solution of the capillary surface equation without gravity over an infinite strip must be a rigid rotation of a cylinder. Here we give a simple proof for Tam's Theorem and generalize his result.**

**1. Introduction.** Let  $\Omega$  be a domain in  $\mathbb{R}^n$ . Consider the equation of prescribed mean curvature

$$(1) \quad \operatorname{div} Tu = H \text{ in } \Omega$$

where  $Tu = \frac{Du}{\sqrt{1+|Du|^2}}$  and  $Du$  is the gradient of  $u$ .

Finn [4] proved that if  $H = n$ ,  $\Omega$  contains the unit ball  $B_1$  of  $\mathbb{R}^n$ , and (1) has a solution  $u$ , then  $\Omega$  has to be exactly  $B_1$  and  $u$  must be a lower hemisphere. We emphasize that no boundary condition is imposed.

In the case  $\Omega$  is unbounded, Finn [4] conjectured that the only solution of (1) with  $H = 2$  over an infinite strip of width 1 in  $\mathbb{R}^2$  is a cylinder. Wang [9] and Collin [2] independently showed that other different solutions can appear, so Finn's conjecture is not true.

In [7], [8] Tam considered the problem related to Finn's conjecture as follows:

$$(2) \quad \begin{cases} \operatorname{div} Tu = H & \text{in } \Omega, \\ Tu \cdot \nu = \cos \alpha & \text{on } \partial\Omega \end{cases}$$

where  $H$  and  $\alpha$  are constants,  $\Omega$  is the infinite strip  $(-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R}$ ,  $\nu$  is the unit outward normal of  $\partial\Omega$ . The boundary value problem (2) determines the height of a capillary surface without gravity. Note that if (2) has a solution, then  $H = 2 \cos \alpha$  ([7]).

Tam [7] proved that any solution of (2) is of the form  $\phi_\beta + \text{constant}$  for some  $|\beta| < 1$  if  $\frac{\pi}{2} > \alpha > 0$ , where

$$\phi_\beta = -\frac{1}{\sqrt{1-\beta^2}} \sqrt{\left(\frac{1}{2\cos\alpha}\right)^2 - x^2} + \frac{\beta}{\sqrt{1-\beta^2}} y,$$

that is, any solution of (2) must be a cylinder.

The statement above is still true for  $\alpha = 0$  and was proved by the author first, and then Tam modify his own method in [7] to give a unified proof for all cases  $\frac{\pi}{2} > \alpha \geq 0$ , (c.f. [8], where the author's name was spelled by Tam as "C. Wong").

In this paper, we do not give our original proof for the case  $\alpha = 0$ , but give a simple proof for the cases  $\frac{\pi}{2} > \alpha \geq 0$ . We shall prove that if  $\int_{\Gamma_{y_0}} Tu \cdot \nu_1 \, d\sigma$  is given where  $y_0$  is a constant,  $\Gamma_{y_0} = \left\{ (x, y_0) \mid -\frac{1}{2} \leq x \leq \frac{1}{2} \right\}$ ,  $\nu_1 = \langle 0, 1 \rangle$ . Then the solution of (2) is unique up to an additive constant. Therefore we not only obtain Tam's Theorem but also generalize it (§2).

The author is grateful to Prof. Finn's kind suggestion on this paper.

**2. Simple proof for Tam's Theorem.** To simplify the proof for Tam's Theorem, the well-known inequality

$$(Du - Dv) \cdot (Tu - Tv) \geq \frac{|Du - Dv|^2}{\max((1 + |Du|^2)^{\frac{3}{2}}, (1 + |Dv|^2)^{\frac{3}{2}})}$$

[6] will be sharpened, and actually we will use the following inequality (3) to deal with integral estimates:

LEMMA 1. *Let  $\Omega \subset \mathbb{R}^n$  and let  $u, v \in C^1(\Omega)$ . Then*

$$(3) \quad (Du - Dv) \cdot (Tu - Tv) \geq \frac{|Du - Dv|^2}{\sqrt{1 + (|Du| + |Du - Dv|)^2}} \left( 1 - \frac{|Du|}{\sqrt{1 + |Du|^2}} \right).$$

*Proof.* Let  $|Du| = \alpha$  and  $|Du - Dv| = b$ . After choosing a suitable coordinates, we may assume  $Du - Dv = (b, 0, \dots, 0)$ ,  $Du = (\alpha \cos \theta, \alpha_2, \dots, \alpha_n)$  where  $0 \leq \theta \leq \pi$ ,  $\alpha^2 \cos^2 \theta + \alpha_2^2 + \dots + \alpha_n^2 = \alpha^2$ .

Then  $Dv = (\alpha \cos \theta - b, \alpha_2, \dots, \alpha_n)$ . Hence

$$\begin{aligned}
 & \left( \frac{Du}{\sqrt{1+|Du|^2}} - \frac{Dv}{\sqrt{1+|Dv|^2}} \right) \cdot (Du - Dv) \\
 &= \left( \frac{\alpha \cos \theta}{\sqrt{1+\alpha^2}} - \frac{\alpha \cos \theta - b}{\sqrt{1+(\alpha \cos \theta - b)^2 + \alpha_2^2 + \dots + \alpha_n^2}} \right) b \\
 &= b \left( \alpha \cos \theta \left( \frac{1}{\sqrt{1+\alpha^2}} - \frac{1}{\gamma} \right) + \frac{b}{\gamma} \right) \\
 &= b \left( \alpha \cos \theta \left( \frac{-2\alpha b \cos \theta + b^2}{\sqrt{1+\alpha^2}\gamma(\sqrt{1+\alpha^2} + \gamma)} \right) + \frac{b}{\gamma} \right) \\
 &= \frac{b^2}{\gamma} \left( \frac{\alpha \cos \theta(-2\alpha \cos \theta + b)}{\sqrt{1+\alpha^2}(\sqrt{1+\alpha^2} + \gamma)} + 1 \right),
 \end{aligned}$$

where  $\gamma = \sqrt{1 + \alpha^2 - 2\alpha b \cos \theta + b^2}$ .

Since

$$\begin{aligned}
 \sqrt{1 + \alpha^2} + \sqrt{1 + \alpha^2 - 2\alpha b \cos \theta + b^2} &\geq |\alpha| + |\alpha \cos \theta - b| \\
 &\geq |-2\alpha \cos \theta + b|
 \end{aligned}$$

and since  $\sqrt{1 + \alpha^2 - 2\alpha b \cos \theta + b^2} \leq \sqrt{1 + (\alpha + b)^2}$ , we have

$$(Tu - Tv) \cdot (Du - Dv) \geq \frac{b^2}{\sqrt{1 + (\alpha + b)^2}} \left( 1 - \frac{\alpha}{\sqrt{1 + \alpha^2}} \right).$$

This completes the proof.  $\square$

Now we generalize Tam's Theorem.

**THEOREM 2.** *Let  $\Omega$  be the infinite strip  $(-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R}$  in  $\mathbb{R}^2$ . Let  $u, v$  be  $C^2(\Omega)$  functions satisfying  $Tu, Tv \in C^0(\bar{\Omega}) \cap C^1(\Omega)$ . Suppose that for every  $\varepsilon$  with  $0 < \varepsilon < \frac{1}{2}$ ,  $|Du|$  is uniformly bounded in  $[-\frac{1}{2} + \varepsilon, \frac{1}{2} - \varepsilon] \times \mathbb{R}$ . And if*

$$(4) \quad \begin{cases} \operatorname{div} Tv = \operatorname{div} Tu & \text{in } \Omega \\ Tv \cdot \nu = Tu \cdot \nu & \text{on } \partial\Omega \\ \int_{\Gamma_{y_0}} Tv \cdot \nu_1 \, d\sigma = \int_{\Gamma_{y_0}} Tu \cdot \nu_1 \, d\sigma & \text{for some constant } y_0, \end{cases}$$

where  $\nu$  is the unit outward normal of  $\partial\Omega$ ,  $\nu_1 = \langle 0, 1 \rangle$ ,  $\Gamma_{y_0} = \bar{\Omega} \cap \{(x, y) | y = y_0\}$ , then we have  $v \equiv u + \text{constant}$ .

**REMARK.** In (4), we assume neither  $\operatorname{div} Tu = \text{constant}$  nor  $Tu \cdot \nu = \text{constant}$ , hence we generalize Tam's Theorem.

As for more general domains  $\Omega$ , the results are stated in Theorem 4.

*Proof of Theorem 2.* For any two numbers  $y_1, y_2$  with  $y_1 < y_2$ , we set  $\Omega_{y_1, y_2} = \Omega \cap \{(x, y) \mid -y_1 < y < y_2\}$ . By divergence theorem, we have

$$\int_{\partial\Omega_{y_1, y_2}} (Tu \cdot \nu - Tv \cdot \nu) d\sigma = \int \int_{\Omega_{y_1, y_2}} \operatorname{div} Tu - \operatorname{div} Tv = 0.$$

Since  $(Tu - Tv) \cdot \nu = 0$  on  $\partial\Omega$ ,  $\nu = \nu_1$  on  $\Gamma_{y_2}$  and  $\nu = -\nu_1$  on  $\Gamma_{y_1}$ , we obtain

$$\int_{\Gamma_{y_2}} (Tu - Tv) \cdot \nu_1 d\sigma - \int_{\Gamma_{y_1}} (Tu - Tv) \cdot \nu_1 d\sigma = 0.$$

Hence

$$(5) \quad \int_{\Gamma_{y_1}} (Tu - Tv) \cdot \nu_1 d\sigma = \text{constant} = \int_{\Gamma_{y_0}} (Tu - Tv) \cdot \nu_1 d\sigma = 0$$

for every  $y_1 \in \mathbb{R}$ .

Similarly, applying divergence theorem again, we have

$$(6) \quad \begin{aligned} & \int_{\Gamma_{y_2}} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1 d\sigma \\ & - \int_{\Gamma_{y_1}} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1 d\sigma \\ & = \int \int_{\Omega_{y_1, y_2}} \frac{(Du - Dv)}{1 + (u - v)^2} \cdot (Tu - Tv) \\ & + \int \int_{\Omega_{y_1, y_2}} \tan^{-1}(u - v)(\operatorname{div} Tu - \operatorname{div} Tv) \\ & = \int \int_{\Omega_{y_1, y_2}} \frac{(Du - Dv)}{1 + (u - v)^2} \cdot (Tu - Tv) \geq 0, \quad (y_1 < y_2). \end{aligned}$$

For each  $y \in \mathbb{R}$ , let us write

$$(7) \quad f(y) = \int_{\Gamma_y} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1.$$

By (6),  $f(y)$  is increasing in  $y$ , hence  $\lim_{y \rightarrow \pm\infty} f(y)$  exist. We claim that  $\lim_{y \rightarrow +\infty} f(y) = 0$ . Otherwise, there exist two constants  $y'_0$  and  $C_1$  such that  $0 < C_1 < 1$  and

$$(8) \quad |f(y)| \geq C_1 \text{ for every } y \geq y'_0.$$

Define  $m(y) = \frac{1}{|\Gamma'_y|} \int_{\Gamma'_y} \tan^{-1}(u - v) d\sigma$  for every  $y \geq y'_0$  where

$$\Gamma'_y = \left[ -\frac{1}{2} + \frac{C_1}{8\pi}, \frac{1}{2} - \frac{C_1}{8\pi} \right] \times \{y\},$$

$|\Gamma'_y|$  is the length of  $\Gamma'_y$ . It follows that  $|m(y)| \leq \frac{\pi}{2}$  for every  $y \geq y'_0$ . From (5), we see that

$$(9) \quad \begin{aligned} & \int_{\Gamma_y} m(y)(Tu - Tv) \cdot \nu_1 \\ &= m(y) \int_{\Gamma_y} (Tu - Tv) \cdot \nu_1 = 0 \text{ for every } y \geq y'_0. \end{aligned}$$

By hypothesis,  $|Du|$  is uniformly bounded in

$$\left[ -\frac{1}{2} + \frac{C_1}{8\pi}, \frac{1}{2} - \frac{C_1}{8\pi} \right] \times \mathbb{R},$$

so there exists a positive constant  $C_2$  (independent of  $y$ ) such that

$$(10) \quad |Du| \leq C_2 \text{ in } \left[ -\frac{1}{2} + \frac{C_1}{8\pi}, \frac{1}{2} - \frac{C_1}{8\pi} \right] \times \mathbb{R}.$$

Hence for each  $y \geq y'_0$ ,

$$(11) \quad \begin{aligned} & \left| \int_{\Gamma'_y} (\tan^{-1}(u - v) - m(y))(Tu - Tv) \cdot \nu_1 \right| \\ & \geq \left| \int_{\Gamma_y} (\tan^{-1}(u - v) - m(y))(Tu - Tv) \cdot \nu_1 \right| \\ & \quad - \left| \int_{\Gamma_y \setminus \Gamma'_y} (\tan^{-1}(u - v) - m(y))(Tu - Tv) \cdot \nu_1 \right|. \end{aligned}$$

By direct computation, we have

$$\begin{aligned} & \left| \int_{\Gamma_y \setminus \Gamma'_y} (\tan^{-1}(u - v) - m(y))(Tu - Tv) \cdot \nu_1 \right| \\ & \leq 2 \cdot \frac{C_1}{8\pi} \left( \frac{\pi}{2} + \frac{\pi}{2} \right) \cdot 2 = \frac{C_1}{2}. \end{aligned}$$

Combining this result with (7)-(9) and (11), we have

$$(12) \quad \begin{aligned} & \left| \int_{\Gamma'_y} \tan^{-1}(u - v) - m(y)(Tu - Tv) \cdot \nu_1 \right| \\ & \geq \frac{C_1}{2} \text{ for every } y \geq y'_0. \end{aligned}$$

Applying Poincaré inequality to the left hand side of (12), we obtain

$$\begin{aligned}
\frac{C_1}{2} &\leq 2 \int_{\Gamma'_y} |\tan^{-1}(u - v) - m(y)| \\
&\leq 2|\Gamma'_y| \int_{\Gamma'_y} \frac{|Du - Dv|}{1 + (u - v)^2} \\
&\leq 2|\Gamma'_y| \left( \int_{\Gamma'_y \cap \{|Du - Dv| < 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{|Du - Dv|}{1 + (u - v)^2} \right. \\
&\quad \left. + \int_{\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{|Du - Dv|}{1 + (u - v)^2} \right) \\
&\leq \frac{C_1}{4} + 2|\Gamma'_y| \int_{\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{|Du - Dv|}{1 + (u - v)^2}.
\end{aligned}$$

Hence for every  $y \geq y'_0$ , we have

$$(13) \quad \frac{C_1}{8|\Gamma'_y|} \leq \int_{\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{|Du - Dv|}{1 + (u - v)^2}.$$

From (10), we have

$$\begin{aligned}
(14) \quad &\frac{|Du - Dv|^2}{\sqrt{1 + (|Du| + |Du - Dv|)^2}} \left( 1 - \frac{|Du|}{\sqrt{1 + |Du|^2}} \right) \\
&\geq \frac{|Du - Dv|^2}{\sqrt{1 + (C_2 + |Du - Dv|)^2}} \left( 1 - \frac{C_2}{\sqrt{1 + C_2^2}} \right) \\
&\geq |Du - Dv| \frac{8^{-1}|\Gamma'_y|^{-2}C_1}{\sqrt{1 + (C_2 + 8^{-1}|\Gamma'_y|^{-2}C_1)^2}} \left( 1 - \frac{C_2}{\sqrt{1 + C_2^2}} \right)
\end{aligned}$$

in  $\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}$ . Combining (13) with the inequalities (3) and (14), we obtain the estimate

$$\begin{aligned}
\frac{C_1}{8|\Gamma'_y|} &\leq 8|\Gamma'_y|^2 C_1^{-1} \sqrt{1 + (C_2 + 8^{-1}|\Gamma'_y|^{-2}C_1)^2} \left( 1 - \frac{C_2}{\sqrt{1 + C_2^2}} \right)^{-1} \\
&\quad \cdot \int_{\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{(Du - Dv) \cdot (Tu - Tv)}{1 + (u - v)^2}.
\end{aligned}$$

Since  $\Gamma'_y \subset \Gamma_y$ , it is easy to see that

$$(15) \quad C_3 \leq \int_{\Gamma_y} \frac{(Du - Dv) \cdot (Tu - Tv)}{1 + (u - v)^2}$$

for every  $y \geq y'_0$  where

$$C_3 = \frac{C_1^2}{64|\Gamma_y|^3 \sqrt{1 + (C_2 + 8^{-1}|\Gamma_y|^{-2}C_1)^2}} \left( 1 - \frac{C_2}{\sqrt{1 + C_2^2}} \right).$$

By (6), (7), (15) and Fubini's Theorem, for every  $y \geq y'_0$ , we have

$$\begin{aligned} (16) \quad & f(y) - f(y_0) \\ &= \int_{\Gamma_y} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1 \\ &\quad - \int_{\Gamma_{y'_0}} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1 \\ &= \int \int_{\Omega_{y'_0, y}} \frac{(Du - Dv)}{1 + (u - v)^2} \cdot (Tu - Tv) \\ &= \int_{y'_0}^y \left( \int_{\Gamma_t} \frac{(Du - Dv)}{1 + (u - v)^2} \cdot (Tu - Tv) \, d\sigma \right) dt \\ &\geq \int_{y'_0}^y C_3 |\Gamma_t| \, dt \\ &= C_3(y - y'_0). \end{aligned}$$

Note that  $|\int_{\Gamma_y} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1| \leq \pi \cdot |\Gamma_y| = \pi$ . Let  $y \rightarrow +\infty$  in (16), we get a contradiction. Thus we obtain  $\lim_{y \rightarrow +\infty} f(y) = 0$ . Similarly, we have  $\lim_{y \rightarrow -\infty} f(y) = 0$ . Using (6) again and letting  $y_2 \rightarrow +\infty, y_1 \rightarrow -\infty$ , we have  $\iint_{\Omega} \frac{(Du - Dv)}{1 + (u - v)^2} \cdot (Tu - Tv) = 0$ , hence  $(Du - Dv) \cdot (Tu - Tv) \equiv 0$  in  $\Omega$ , and  $Du \equiv Dv$  in  $\Omega$ . This completes the proof.  $\square$

Tam's Theorem is an immediate consequence of Theorem 2.

**COROLLARY 3.** *Every solution  $u$  of (2) is of the form  $\phi_\beta + \text{constant}$  for some  $|\beta| < 1$ , where  $\Omega$  is the infinite strip  $(-\frac{1}{2}, \frac{1}{2}) \times \mathbb{R}$ ,*

$$\phi_\beta = -\frac{1}{\sqrt{1 - \beta^2}} \sqrt{\left(\frac{1}{2 \cos \alpha}\right)^2 - x^2} + \frac{\beta}{\sqrt{1 - \beta^2}} y,$$



$\beta$  satisfies  $\int_{\Gamma_{y_0}} T\phi_\beta \cdot \nu_1 = \int_{\Gamma_{y_0}} Tu \cdot \nu_1$ , for some constant  $y_0$ .

Tam's Theorem for more general domains  $\Omega$  is stated in the following Theorem :

**THEOREM 4.** *Let  $g_1, g_2 \in C^1(\mathbb{R})$  satisfy  $g_2 < g_1$  in  $\mathbb{R}$ . Set  $\Omega = \{(x, y) | g_2(y) < x < g_1(y)\}$ . Let  $u, v \in C^2(\Omega)$  satisfy  $Tu, Tv \in C^0(\bar{\Omega}) \cap C^1(\Omega)$ . Suppose that for every  $\varepsilon > 0$ ,  $|Du|$  is uniformly bounded in  $\{(x, y) | g_2(y) + \varepsilon \leq x \text{ and } x \leq g_1(y) - \varepsilon\}$ . And if*

$$(4') \quad \begin{cases} \operatorname{div} Tv = \operatorname{div} Tu & \text{in } \Omega, \\ Tv \cdot \nu = Tu \cdot \nu & \text{on } \partial\Omega, \\ \int_{\Gamma_{y_0}} Tv \cdot \nu_1 d\sigma = \int_{\Gamma_{y_0}} Tu \cdot \nu_1 d\sigma & \text{for some constant } y_0, \\ |\Gamma_y| = O(|y|^{\frac{1}{3}-\alpha}) & \text{as } y \rightarrow \pm\infty \\ & \text{for some constant } \alpha \\ & \text{with } 0 < \alpha < \frac{1}{3}, \end{cases}$$

where  $\Gamma_y = [g_2(y), g_1(y)] \times \{y\}$ , then we have  $u \equiv v + \text{constant}$  in  $\Omega$ .

*Proof.* The proof is similar to that of Theorem 2. Here we only sketch the proof.

For each  $y \in \mathbb{R}$ , let us write

$$(7) \quad f(y) = \int_{\Gamma_y} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1.$$

By (6),  $f(y)$  is increasing in  $y$ , hence  $\lim_{y \rightarrow \pm\infty} f(y)$  exist. We claim that  $\lim_{y \rightarrow +\infty} f(y) = 0$ . Otherwise, there exist two positive constants  $y'_0$  and  $C_1$  such that

$$(8') \quad |f(y)| \geq C_1 \text{ for every } y \geq y'_0 > 0.$$

Hence  $C_1 \leq |\int_{\Gamma_y} \tan^{-1}(u - v)(Tu - Tv) \cdot \nu_1| \leq \pi |\Gamma_y|$ , and we have  $|\Gamma_y| \geq \frac{C_1}{\pi}$  for every  $y \geq y'_0$ . Define  $m(y) = \frac{1}{|\Gamma'_y|} \int_{\Gamma'_y} \tan^{-1}(u - v) d\sigma$  for every  $y \geq y'_0$ , where  $\Gamma'_y = [g_2(y) + \frac{C_1}{8\pi}, g_1(y) - \frac{C_1}{8\pi}] \times \{y\}$ .

By hypothesis,  $|Du|$  is uniformly bounded in  $\cup_{y \geq y'_0} \Gamma'_y$ , so there exists a positive constant  $C_2$  (independent of  $y$ ) such that

$$(10') \quad |Du| \leq C_2 \text{ in } \bigcup_{y \geq y'_0} \Gamma'_y.$$

Hence for each  $y \geq y'_0$ , we have

$$(12') \quad \left| \int_{\Gamma'_y} (\tan^{-1}(u - v) - m(y))(Tu - Tv) \cdot \nu_1 \right| \\ \geq |f(y)| - \frac{C_1}{2} \geq \frac{|f(y)|}{2}.$$

Applying Poincaré inequality to the left hand side of (12'), we obtain

$$\begin{aligned} \frac{|f(y)|}{2} &\leq 2 \int_{\Gamma'_y} |\tan^{-1}(u - v) - m(y)| \\ &\leq 2|\Gamma'_y| \int_{\Gamma'_y} \frac{|Du - Dv|}{1 + (u - v)^2} \\ &\leq \frac{C_1}{4} + 2|\Gamma'_y| \int_{\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{|Du - Dv|}{1 + (u - v)^2}. \end{aligned}$$

Hence for every  $y \geq y'_0$ , we have

$$(13') \quad \frac{|f(y)|}{8|\Gamma'_y|} \leq \int_{\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{|Du - Dv|}{1 + (u - v)^2}.$$

Combining (13') with the inequalities (3) and (14), we obtain

$$\begin{aligned} \frac{|f(y)|}{8|\Gamma'_y|} &\leq 8|\Gamma'_y|^2 C_1^{-1} \sqrt{1 + (C_2 + 8^{-1}|\Gamma'_y|^{-2}C_1)^2} \left(1 - \frac{C_2}{\sqrt{1 + C_2^2}}\right)^{-1} \\ &\quad \cdot \int_{\Gamma'_y \cap \{|Du - Dv| \geq 8^{-1}|\Gamma'_y|^{-2}C_1\}} \frac{(Du - Dv) \cdot (Tu - Tv)}{1 + (u - v)^2}. \end{aligned}$$

Since  $\Gamma'_y \subset \Gamma_y$ ,

$$\begin{aligned} |\Gamma'_y|^2 \sqrt{1 + (C_2 + 8^{-1}|\Gamma'_y|^{-2}C_1)^2} \\ \leq |\Gamma_y|^2 \sqrt{1 + (C_2 + 8^{-1}|\Gamma_y|^{-2}C_1)^2}, \end{aligned}$$

it is easy to see that

$$\begin{aligned} |f(y)| &\leq 64|\Gamma_y|^3 C_1^{-1} \sqrt{1 + (C_2 + 8^{-1}|\Gamma_y|^{-2}C_1)^2} \\ &\quad \cdot \left(1 - \frac{C_2}{\sqrt{1 + C_2^2}}\right)^{-1} \int_{\Gamma_y} \frac{(Du - Dv)(Tu - Tv)}{1 + (u - v)^2}, \end{aligned}$$

hence we have

$$(15') \quad C_4 |\Gamma_y|^{-3} |f(y)| \leq \int_{\Gamma_y} \frac{(Du - Dv) \cdot (Tu - Tv)}{1 + (u - v)^2}$$

where

$$C_4 = 64^{-1} C_1 \left( 1 + \left( C_2 + 8^{-1} \left( \frac{C_1}{\pi} \right)^{-2} C_1 \right)^2 \right)^{-\frac{1}{2}} \left( 1 - \frac{C_2}{\sqrt{1 + C_2^2}} \right).$$

By Fubini's Theorem, we have

$$\int_{\Gamma_y} \frac{(Du - Dv) \cdot (Tu - Tv)}{1 + (u - v)^2} = f'(y).$$

Hence

$$(17) \quad \frac{f'(y)}{|f(y)|} \geq C_4 |\Gamma_y|^{-3} \geq C_5 y^{-1+3\alpha} \text{ for } y \geq y'_0 > 0$$

where  $C_5$  is a positive constant. Integrate (17) from  $y'_0$  to  $y$ ,  $y > y'_0$ , we obtain

$$(18) \quad |\log |f(y)| - \log |f(y'_0)|| \geq \frac{C_5}{1 - 3\alpha} (y^{3\alpha} - y'^{3\alpha}_0).$$

Note that  $|f(y)| \leq \pi |\Gamma_y| = O(y^{\frac{1}{3}-\alpha})$ . Let  $y \rightarrow +\infty$  in (18), we get a contradiction. The remainder of the proof is similar to that of Theorem 2.  $\square$

**REMARK.** Let  $\Omega$  be a domain (bounded or unbounded) in  $\mathbb{R}^n$  and let  $k$  be a positive constant, where  $n \geq 2$  is an integer. Suppose that  $\alpha$  is defined on  $\partial\Omega$  with  $0 \leq \alpha \leq 2\pi$ . Then the boundary value problem

$$(19) \quad \begin{cases} \operatorname{div} Tu = ku & \text{in } \Omega \\ Tu \cdot \nu = \cos \alpha & \text{on } \partial\Omega \end{cases}$$

determines the height  $u$  of a capillary surface in a uniform gravitational field.

It was proved by Finn and the author [5] that if (19) has a solution then it is unique. We point out that neither growth condition of  $u$  nor condition on the form of  $\Omega$  at infinity is imposed.

## REFERENCES

- [1] J.T. Chen and W.-H. Huang, *Convexity of capillary surfaces in the outer spaces*, Invent. Math., **67** (1982), 253-259.
- [2] P. Collin, *Deux exemples de graphes courbure moyenne constante sur une bande de  $\mathbb{R}^2$* , C. R. Acad. Sci. Paris, **311** (1990), 539-542.
- [3] R. Finn, *New estimates for equations of minimal surface type*, Arch. Rational Mech. Anal., **14** (1963), 337-375.
- [4] ———, *Remarks relevant to minimal surfaces and to surfaces of prescribed mean curvature*, J. Analyse Math., **14** (1965), 139-160.
- [5] R. Finn and J.-F. Hwang, *On the comparison principle for capillary surfaces*, J. Fac. Sci. Univ. Tokyo Sect. IA, Math., **36** (1989), 131-134.
- [6] J.C.C. Nitsche, *Vorlesungen über Minimalflächen*, Springer-Verlag, Berlin-Heidelberg-New York, 1975.
- [7] L.F. Tam, *On the uniqueness of capillary surfaces without gravity over an infinite strip*, Indiana Univ. Math. J., **36** (1987), 79-89.
- [8] ———, *On the uniqueness of capillary surfaces*, Variational Methods for Free Surfaces Interfaces. (ed. P. Concus and R. Finn) Springer-Verlag, Berlin-Heidelberg-New York, 1987.
- [9] A.N. Wang, *Constant mean curvature surfaces on a strip*, Pacific J. Math., **145** (1990), 395-396.

Received October 29, 1992. Partially supported by grant NSC 82-0208-M-001-078.

INSTITUTE OF MATHEMATICS  
ACADEMIA SINICA  
TAIPEI 11529  
TAIWAN, R.O.C.



## PACIFIC JOURNAL OF MATHEMATICS

Founded by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

### EDITORS

Sun-Yung Alice Chang (Managing Editor)  
University of California  
Los Angeles, CA 90095-1555  
pacific@math.ucla.edu

F. Michael Christ  
University of California  
Los Angeles, CA 90095-1555  
christ@math.ucla.edu

Thomas Enright  
University of California  
San Diego, La Jolla, CA 92093  
tenright@ucsd.edu

Nicholas Ercolani  
University of Arizona  
Tucson, AZ 85721  
ercolani@math.arizona.edu

Robert Finn  
Stanford University  
Stanford, CA 94305  
finn@gauss.stanford.edu

Vaughan F. R. Jones  
University of California  
Berkeley, CA 94720  
vfr@math.berkeley.edu

Steven Kerckhoff  
Stanford University  
Stanford, CA 94305  
spk@gauss.stanford.edu

Martin Scharlemann  
University of California  
Santa Barbara, CA 93106  
mgscharl@math.ucsb.edu

Gang Tian  
Courant Institute  
New York University  
New York, NY 10012-1100  
tiang@taotao.cims.nyu.edu

V. S. Varadarajan  
University of California  
Los Angeles, CA 90095-1555  
vsv@math.ucla.edu

### SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
STANFORD UNIVERSITY  
UNIVERSITY OF ARIZONA  
UNIVERSITY OF BRITISH COLUMBIA  
UNIVERSITY OF CALIFORNIA  
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA  
UNIVERSITY OF NEVADA, RENO  
UNIVERSITY OF OREGON  
UNIVERSITY OF SOUTHERN CALIFORNIA  
UNIVERSITY OF UTAH  
UNIVERSITY OF WASHINGTON  
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

---

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$108.00 a year to individual members of supporting institutions.

Subscriptions, orders for back issues published within the last three years, and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

---

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,  
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,  
the American Mathematical Society's TEX macro system.  
Copyright © 1995 by Pacific Journal of Mathematics

# PACIFIC JOURNAL OF MATHEMATICS

Volume 169      No. 1      May 1995

---

Minimal sets of periods for torus maps via Nielsen numbers	1
LLUÍS ALSSEDÀ, STEWART BALDWIN, JAUME LLIBRE, RICHARD SWANSON and WIESLAW SZLENK	
Diagonalizing Hilbert cusp forms	33
TIMOTHY ATWILL	
A splitting criterion for rank 2 vector bundles on $\mathbf{P}^n$	51
EDOARDO BALLICO	
Controlling Tietze-Urysohn extensions	53
MARC FRANTZ	
Length of Julia curves	75
DAVID H. HAMILTON	
On the uniqueness of capillary surfaces over an infinite strip	95
JENN-FANG HWANG	
Volume estimates for log-concave densities with application to iterated convolutions	107
MARIUS JUNGE	
A reflection principle in complex space for a class of hypersurfaces and mappings	135
FRANCINE ANTOINETTE MEYLAN	
Jean Bourgain's analytic partition of unity via holomorphic martingales	161
PAUL F.X. MÜLLER	
Characters of Brauer's centralizer algebras	173
ARUN RAM	