

Pacific Journal of Mathematics

**THE CORESTRICTION OF VALUED DIVISION ALGEBRAS
OVER HENSELIAN FIELDS. II**

YOON SUNG HWANG

THE CORESTRICTION OF VALUED DIVISION ALGEBRAS OVER HENSELIAN FIELDS II

YOON SUNG HWANG

When L/F is a tame extension of Henselian fields (i.e. $\text{char}(\overline{F}) \nmid [L:F]$), we analyze the underlying division algebra cD of the corestriction $\text{cor}_{L/F}(D)$ of a tame division algebra D over L with respect to the unique valuations of cD and D extending the valuations on F and L . We show that the value group of cD lies in the value group of D and for the center of residue division algebra, $Z({}^c\overline{D}) \subseteq \mathcal{N}(Z(\overline{D})/\overline{F})^{1/k}$, where $\mathcal{N}(Z(\overline{D})/\overline{F})$ is the normal closure of $Z(\overline{D})$ over \overline{F} and k is an integer depending on which roots of unity lie in F and L .

Introduction.

This paper is a continuation of [H₂], where we analyzed the corestriction $\text{cor}_{L/F}(D)$ of a tame division algebra D over L when L/F is an inertial (unramified) extension of Henselian valued fields. We will follow terminology and notations in that paper. We will here concentrate on the cases when $\overline{L} = \overline{F}$, when L/F is a totally ramified of radical type (TRRT) extension (see below for definition) and when L/F is tame, where L/F is a finite separable extension of Henselian fields. We will consider only division algebras *finite-dimensional* over their centers.

Here is an overview of the paper: After a preliminary section, in section 2 we will analyze the underlying division algebra cD of the corestriction $\text{cor}_{L/F}(D)$ of inertially split division algebras D over L when $\overline{L} = \overline{F}$. In sections 3 and 4, we will consider the corestriction of tame division algebras when L/F is TRRT and when L/F is tame, respectively.

The following definition of a TRRT extension was given in [JW, Sec. 4]. For a finite extension L of a valued field (F, v) , we say that L is a *totally ramified* extension of F of *radical type* with respect to v (TRRT) if v extends to a valuation w on L such that L is totally ramified over F and there is a subgroup \mathcal{A} of L^*/F^* which maps via w isomorphically onto Γ_L/Γ_F .

Our basic results are summarized in the following table. Here Γ_D is the value group of the valuation on D and \overline{D} is the residue division ring of the valuation ring of D . Also $\mathcal{N}(Z(\overline{D})/\overline{F})$ denotes the normal closure of $Z(\overline{D})$ over \overline{F} , D^n is the underlying division algebra of the n -fold product

$D \otimes_L \cdots \otimes_L D$, and θ_D is the map of (1) below, so $\ker(\theta_D)$ is a subgroup of Γ_D/Γ_L .

$\bar{L} = \bar{F}$ (Th. 8)	D inertially split $\Gamma_{eD} = [L: F] \cdot \Gamma_D + \Gamma_F$ $Z(\overline{D^n}) \subseteq Z({}^c\bar{D}) \subseteq Z(\bar{D})$	D tame
L/F TRRT (Th. 15)	"	$[L: F]\Gamma_D \subseteq \Gamma_{eD} \subseteq \Gamma_D$ $Z({}^c\bar{D}) \subseteq Z(\bar{D})$
L/F tame (Th. 17, 18)	$\Gamma_{eD} \subseteq \Gamma_L: \Gamma_F \cdot \Gamma_D + \Gamma_F$ $Z({}^c\bar{D}) \subseteq \mathcal{N}(Z(\bar{D}/\bar{F}))$	$\Gamma_{eD} \subseteq \Gamma_D$ $Z({}^c\bar{D}) \subseteq \mathcal{N}(Z(\bar{D})/\bar{F})^{1/k}$ $k \mid \exp(\ker \theta_D)$

The integer k in the table above depends not only on Γ_D/Γ_L and $[\bar{L}: \bar{F}]$ but also on which roots of unity lie in F . One of the interesting results of this investigation is to see how heavily the corestriction depends on the roots of unity in F and L .

1. Preliminaries.

Let (D, v) be a valued division algebra, that is, a division ring D with valuation v . Associated to v , we have its value group $\Gamma_D = v(D^*)$; the valuation ring $V_D = \{d \in D^* \mid v(d) \geq 0\} \cup \{0\}$; the unique maximal left (and right) ideal M_D of V_D , $M_D = \{d \in D^* \mid v(d) > 0\} \cup \{0\}$; the group of v -units of D^* , $U_D = V_D - M_D = V_D^*$; the residue division ring $\bar{D} = V_D/M_D$. If F is the center $Z(D)$ of D , there is a well-defined epimorphism

$$(1) \quad \theta_D: \Gamma_D/\Gamma_F \rightarrow \text{Gal}(Z(\bar{D})/\bar{F}),$$

induced by $\alpha: D^* \rightarrow \text{Gal}(Z(\bar{D})/\bar{F})$ which is given by $d \mapsto \bar{c}_d$ where \bar{c}_d is the map induced by conjugation by d . (cf. [JW, 1.6]).

We recall two propositions which will be particularly useful for this paper.

Proposition 2 [M, Th. 1]. *Let D and E be division algebras over a field F with $[D: F] < \infty$. Suppose D has a valuation v and E has a valuation w with $v|_F = w|_F$. Suppose further*

- (i) D is defectless over F relative to v ;

(ii) $\overline{D} \otimes_{\overline{F}} \overline{E}$ is a division ring;

(iii) $\Gamma_D \cap \Gamma_E = \Gamma_F$.

Then $D \otimes_F E$ is a division ring with a unique valuation u such that $u|_D = v$, and $u|_E = w$. Furthermore, $\overline{D \otimes_F E} \cong \overline{D} \otimes_{\overline{F}} \overline{E}$ and $\Gamma_{D \otimes_F E} = \Gamma_D + \Gamma_E$.

Proposition 3 [JW, Lemma 6.2, Th. 6.3]. *If D is a tame division algebra over a Henselian field F , then there exist $S \in \mathcal{D}_{is}(F)$ and $T \in \mathcal{D}_{ttr}(F)$ such that $D \sim S \otimes_F T$ in $\text{Br}(F)$. (Such S and T are not unique.) Furthermore, if $D \sim S \otimes_F T$ is such a decomposition, $Z(\overline{D}) = \mathcal{F}(\theta_S((\Gamma_S \cap \Gamma_T) / \Gamma_F)) \subseteq Z(\overline{S})$, $\Gamma_D = \Gamma_S + \Gamma_T$ and $\ker(\theta_D) = \Gamma_T / \Gamma_F$.*

2. The case when $\overline{L} = \overline{F}$.

In this section, we assume that $(L, v) \supseteq (F, v)$ is a finite separable extension of Henselian fields with $\overline{L} = \overline{F}$. Recall that for $D \in \mathcal{D}(L)$, ${}^c D \in \mathcal{D}(F)$ denotes the underlying division algebra of $\text{cor}_{L/F}(D)$.

For any valued field (F, v) , let $\text{Br}(V_F)$ denote the Brauer group (of equivalence classes of Azumaya algebras) of the valuation ring V_F . There are canonical group homomorphisms $\alpha: \text{Br}(V_F) \rightarrow \text{Br}(F)$ given by $[A] \mapsto [A \otimes_{V_F} F]$, and $\beta: \text{Br}(V_F) \rightarrow \text{Br}(\overline{F})$ given by $[A] \mapsto [A / M_F A]$, where $[A]$ is the class of A , an Azumaya algebra over V_F . Then, by [JW, Prop. 2.5], α is injective.

Now assume that (F, v) is Henselian. Then define

$$I \text{Br}(F) = \{[D] \in \text{Br}(F) \mid D \in \mathcal{D}_i(F), \text{ i.e., } D \text{ is inertial over } F\}.$$

By [JW, Prop. 2.5 and Ex. 2.4 (ii)], $I \text{Br}(F) = \text{im}(\alpha)$, so $I \text{Br}(F)$ is a subgroup of $\text{Br}(F)$. Azumaya proved in [Az, Th. 31] that β is an isomorphism. The composite map $\beta \circ \alpha^{-1}: I \text{Br}(F) \rightarrow \text{Br}(\overline{F})$ is thus an isomorphism, and it maps $[D]$ to $[\overline{D}]$ for any $D \in \mathcal{D}_i(F)$.

Lemma 4. *If $D \in \mathcal{D}_i(L)$ then ${}^c D \in \mathcal{D}_i(F)$ and ${}^c \overline{D} \sim \overline{D}^{\otimes [L:F]}$ in $\text{Br}(\overline{F})$. (Recall that we assume $\overline{L} = \overline{F}$.)*

Proof. Consider the following commutative diagram.

$$\begin{array}{ccccc} I \text{Br}(F) & \xrightarrow{\alpha^{-1}} & \text{Br}(V_F) & \xrightarrow{\beta} & \text{Br}(\overline{F}) \\ \downarrow \otimes_F L & & \downarrow \otimes_{V_F} V_L & & \downarrow \otimes_{\overline{F}} \overline{L} \\ I \text{Br}(L) & \xrightarrow{\alpha^{-1}} & \text{Br}(V_L) & \xrightarrow{\beta} & \text{Br}(\overline{L}) \end{array}$$

Since $\overline{L} = \overline{F}$ by assumption, the restriction map $\text{res}_{\overline{L}/\overline{F}}: \text{Br}(\overline{F}) \rightarrow \text{Br}(\overline{L})$, given by $[\tilde{D}] \mapsto [\tilde{D} \otimes_{\overline{F}} \overline{L}]$ for any $\tilde{D} \in \mathcal{D}(\overline{F})$, is the identity map on $\text{Br}(\overline{F})$.

So the restriction map $\text{res}_{L/F}: I\text{Br}(F) \rightarrow I\text{Br}(L)$, given by $[D] \mapsto [D \otimes_F L]$ for any $D \in \mathcal{D}_i(F)$, is an isomorphism. So for any $D \in \mathcal{D}_i(L)$, there is a $D_0 \in \mathcal{D}_i(F)$ such that $[D] = \text{res}_{L/F}([D_0])$, i.e., $D \sim D_0 \otimes_F L$ in $\text{Br}(L)$. Then by the above commutative diagram, $[\overline{D}] = \beta \circ \alpha^{-1}([D]) = \beta \circ \alpha^{-1}(\text{res}_{L/F}([D_0])) = \text{res}_{\overline{L}/\overline{F}}(\beta \circ \alpha^{-1}([D_0])) = [\overline{D_0}]$ in $\text{Br}(\overline{L}) (= \text{Br}(\overline{F}))$. Also, by [Ti₂, Th. 2.5], ${}^c D \sim \text{cor}_{L/F}(D) \sim \text{cor}_{L/F}(D_0 \otimes_F L) \sim D_0^{\otimes [L:F]}$ in $\text{Br}(F)$. Since $[D_0] \in I\text{Br}(F)$ and $I\text{Br}(F)$ is a subgroup of $\text{Br}(F)$, $[{}^c D] = [D_0^{\otimes [L:F]}]$ is contained in $I\text{Br}(F)$ and $[{}^c \overline{D}] = [\overline{D_0^{\otimes [L:F]}}] = [\overline{D}^{\otimes [L:F]}]$ in $\text{Br}(\overline{F})$, as desired. \square

In Theorem 7, we will give relations between D and ${}^c D$ for $D \in \mathcal{D}_{is}(L)$ when $\overline{L} = \overline{F}$. To prove that theorem, we need the following information about the homological corestriction which is of interest in itself.

Let G be a group and A a left G -module. We write A^G for $\{a \in A \mid g(a) = a, \text{ all } g \in G\}$. Let H be a subgroup of G of index $n < \infty$, and N a normal subgroup of G . We have a set of representatives $\mathcal{R} = \{\rho_1, \dots, \rho_n\}$ of the left cosets of H in HN with $\rho_i \in N$. So, for $n \in N$ and any i there is a j with $n\rho_i = \rho_j h$, and $h \in H \cap N$. Thus, we have a map $\mathcal{N}: A^{H \cap N} \rightarrow A^N$ given by $\mathcal{N}(a) = \sum_{i=1}^n \rho_i(a)$. Observe that \mathcal{N} is independent of the choice of coset representatives used for \mathcal{R} . Then \mathcal{N} and the isomorphism from HN/N to $H/(H \cap N)$ induces the map $\mathcal{N}_{HN/H}^*: H^m(H/(H \cap N), A^{H \cap N}) \rightarrow H^m(G/N, A^N)$, $m \geq 0$ given by $(HN/N \xrightarrow{\cong} H/(H \cap N), A^{H \cap N} \xrightarrow{\mathcal{N}} A^N)$.

Theorem 5. *Let G, H, N, A and $\mathcal{N}_{HN/H}^*$ be as above. Suppose $f \in H^m(H/(H \cap N), A^{H \cap N})$, $m \geq 0$. Then*

$$\text{cor}_H^G \circ \left(\inf_{H/(H \cap N)}^H (f) \right) = \inf_{G/N}^G \circ \text{cor}_{HN/N}^{G/N} \circ \mathcal{N}_{HN/N}^* (f).$$

Proof. The theorem follows from the following formula for the special case when $G = HN$ since the corestriction is transitive and commutes with the inflation by [We, Prop. 2.4.5].

$$(6) \quad \text{cor}_H^G \left(\inf_{H/(H \cap N)}^H (f) \right) = \inf_{G/N}^G \left(\mathcal{N}_{G/H}^* (f) \right)$$

So, it suffices to prove (6) with assumption that $G = HN$.

For $m = 0$, this is clear. So we may assume $m \geq 1$. For any $\sigma \in G$, there are uniquely determined elements $\rho_\sigma \in \mathcal{R}$ and $h_\sigma \in H$ such that $\sigma = \rho_\sigma h_\sigma$. Also given $\rho_i \in \mathcal{R}$ and $\sigma \in G$, let $\rho_{\sigma_*(i)} \in \mathcal{R}$ and $\delta(\sigma, \rho_i) \in H$ be the elements such that $\sigma \rho_i = \rho_{\sigma_*(i)} \delta(\sigma, \rho_i)$. Since

$$\rho_{\sigma_*(i)} \delta(\sigma, \rho_i) = \sigma \rho_i = \rho_\sigma h_\sigma \rho_i = [\rho_\sigma h_\sigma \rho_i h_\sigma^{-1}] h_\sigma$$

and $\{\rho_{\sigma_*(i)}, \rho_\sigma(h_\sigma \rho_i h_\sigma^{-1})\} \subseteq N$ and $\{\delta(\sigma, \rho_i), h_\sigma\} \subseteq H$, we have $\delta(\sigma, \rho_i) \equiv h_\sigma \pmod{H \cap N}$. Also, as $\sigma = \rho_\sigma h_\sigma$ and $\rho_\sigma \in N$, $\sigma \equiv h_\sigma \pmod{N}$. For $h \in H$ (resp. $g \in G$), let \bar{h} (resp. \bar{g}) denote the left coset $h(H \cap N)$ (resp. gN) in $H/(H \cap N)$ (resp. G/N). So, for any $\sigma \in G$ and $\rho_i \in \mathcal{R}$,

$$(7) \quad \overline{\delta(\sigma, \rho_i)} = \overline{h_\sigma} \quad \text{and} \quad \tilde{\sigma} = \widetilde{h_\sigma}.$$

Let $f \in H^m(H/(H \cap N), A^{H \cap N})$ be represented by an inhomogeneous cocycle, say f again, in $Z^m(H/(H \cap N), A^{H \cap N})$. Then by [H₂, 1.3], for $\sigma_j \in G$, $1 \leq j \leq m$,

$$\begin{aligned} (1) \quad \text{cor}_H^G(\inf_{H/(H \cap N)}^H(f))(\sigma_1, \dots, \sigma_j, \dots, \sigma_m) \\ = \sum_{i=1}^n \rho_{(\sigma_1 \cdots \sigma_m)_*(i)} \left[\inf_{H/(H \cap N)}^H(f)(\delta(\sigma_1, \rho_{(\sigma_2 \cdots \sigma_m)_*(i)}), \dots, \right. \\ \left. \delta(\sigma_j, \rho_{(\sigma_{j+1} \cdots \sigma_m)_*(i)}), \dots, \delta(\sigma_m, \rho_i)) \right] \\ = \sum_{i=1}^n \rho_{(\sigma_1 \cdots \sigma_m)_*(i)} \left[f(\overline{h_{\sigma_1}}, \dots, \overline{h_{\sigma_j}}, \dots, \overline{h_{\sigma_m}}) \right] \quad \text{by (7)} \\ = \sum_{i=1}^n \rho_i \left[f(\overline{h_{\sigma_1}}, \dots, \overline{h_{\sigma_j}}, \dots, \overline{h_{\sigma_m}}) \right] \end{aligned}$$

as $(\sigma_1 \cdots \sigma_m)_* \in S_n$, the symmetric group, so we are just rearranging the order of summation. Then, as $\tilde{\sigma}_j \in G/N$ maps to $\overline{h_{\sigma_j}} \in H/(H \cap N)$ in $G/N \cong H/(H \cap N)$,

$$\begin{aligned} \text{cor}_H^G(\inf_{H/H \cap N}^H(f))(\sigma_1, \dots, \sigma_j, \dots, \sigma_m) \\ = \mathcal{N}_{G/H}^*(f)(\tilde{\sigma}_1, \dots, \tilde{\sigma}_j, \dots, \tilde{\sigma}_m) \\ = \inf_{G/N}^G(\mathcal{N}_{G/H}^*(f))(\sigma_1, \dots, \sigma_j, \dots, \sigma_m). \end{aligned}$$

□

Note that Th. 5 is valid for $f \in H_c^m(H/(H \cap N), A^{H \cap N})$, the m -th continuous cohomology group, if G is a profinite group, and H and N are also assumed closed in G , and A is a discrete G -module.

Recall we assume that $(L, v) \supseteq (F, v)$ is a separable extension of degree n of Henselian fields with $\bar{L} = \bar{F}$. Let L_{sep} (resp. F_{sep}) be the separable closure of L (resp. F). So $L_{sep} = F_{sep}$. Let $G = \text{Gal}(F_{sep}/F)$ and $H = \text{Gal}(L_{sep}/L)$. So H is a closed subgroup of G of index $n = [L:F]$. Let L_{nr} (resp. F_{nr}) be the maximal inertial extension of L (resp. F) in F_{sep} . Since F_{nr}/F is Galois and $L \cap F_{nr} = F$, L and F_{nr} are linearly disjoint over F and $L \otimes_F F_{nr}$ is the field $L \cdot F_{nr}$. Also by [JW, 1.9], $L \cdot F_{nr} = L_{nr}$.

Let $N = \text{Gal}(F_{\text{sep}}/F_{nr})$. Then since F_{nr}/F is Galois and $L \cap F_{nr} = F$, N is normal in G and $G = HN$. Also, $\text{Gal}(F_{nr}/F) \cong \text{Gal}(\overline{F}_{\text{sep}}/\overline{F}) \cong G/N$, and $\text{Gal}(L_{nr}/L) \cong \text{Gal}(\overline{L}_{\text{sep}}/\overline{L}) \cong H/(H \cap N)$ as $H \cap N = \text{Gal}(L_{\text{sep}}/L_{nr})$ with $L_{nr} = L \cdot F_{nr}$. But since $\overline{L} = \overline{F}$, $\text{Gal}(\overline{F}_{\text{sep}}/\overline{F}) = \text{Gal}(\overline{L}_{\text{sep}}/\overline{L})$. So by identifying $\text{Gal}(F_{nr}/F)$ and $\text{Gal}(L_{nr}/L)$ with $\text{Gal}(\overline{F}_{\text{sep}}/\overline{F}) = \text{Gal}(\overline{L}_{\text{sep}}/\overline{L})$ via canonical isomorphisms, we can identify G/N with $H/(H \cap N)$.

Via the crossed product construction, we have the isomorphisms $\text{Br}(L) \cong H_c^2(H, L_{\text{sep}}^*)$, $\text{Br}(F) \cong H_c^2(G, F_{\text{sep}}^*)$, $\text{Br}(L_{nr}/L) \cong H_c^2(H/H \cap N, L_{nr}^*)$, and $\text{Br}(F_{nr}/F) \cong H_c^2(G/N, F_{nr}^*)$.

Theorem 8. *Let $(L, v) \supseteq (F, v)$ be a separable extension of degree n of Henselian fields with $\overline{L} = \overline{F}$. Suppose $D \in \mathcal{D}_{is}(L)$, and θ_D is the map of (1). Then, ${}^cD \in \mathcal{D}_{is}(F)$, $\Gamma_{{}^cD} = n\Gamma_D + \Gamma_F$, and $Z({}^cD) = \mathcal{F}(\theta_D(\tilde{\Gamma}))$, where $\tilde{\Gamma} = \{\alpha + \Gamma_L \in \Gamma_D/\Gamma_L \mid n\alpha \in \Gamma_F\}$. So $Z(\overline{D^n}) \subseteq Z({}^cD) \subseteq Z(\overline{D})$, where D^n is the underlying division algebra of $D^{\otimes n}$, the n -fold product $D \otimes_L \cdots \otimes_L D$. (So if $D \in \mathcal{D}_t(L)$, then ${}^cD \in \mathcal{D}_t(F)$.)*

Proof. Since $L \otimes_F F_{nr}$ is the field L_{nr} , by [D₁, p. 56, Ex. 1] ${}^cD \otimes_F F_{nr} \sim \text{cor}_{L_{nr}/F_{nr}}(D \otimes_L L_{nr}) \sim \text{cor}_{L_{nr}/F_{nr}}(L_{nr}) \sim F_{nr}$ in $\text{Br}(F_{nr})$. So ${}^cD \in \mathcal{D}_{is}(F)$.

Since $[D] \in \text{Br}(L_{nr}/L) \subseteq \text{Br}(L)$, in $\text{Br}(L)$ $[D]$ is represented by $\inf_{H/(H \cap N)}^H(f)$ for some $f \in H_c^2(H/(H \cap N), L_{nr}^*)$. Since the algebraic corestriction corresponds to the homological corestriction, in $\text{Br}(F)$, $[{}^cD]$ is represented by $\text{cor}_H^G(\inf_{H/(H \cap N)}^H(f))$. But, by Th. 5 above

$$\text{cor}_H^G(\inf_{H/(H \cap N)}^H(f)) = \inf_{G/N}^G(\mathcal{N}_{G/H}^*(f)),$$

where $\mathcal{N}_{G/H}^*: H_c^2(H/(H \cap N), L_{nr}^*) \rightarrow H_c^2(G/N, F_{nr}^*)$ is induced by the norm map from L_{nr}^* to F_{nr}^* . Since $[{}^cD] \in \text{Br}(F_{nr}/F) \cong H_c^2(G/N, F_{nr}^*)$, $[{}^cD]$ is represented by $\mathcal{N}_{G/H}^*(f)$.

Let $H' = H/(H \cap N)$ and $G' = G/N$. Since $H' = \text{Gal}(L_{nr}/L)$ and $G' = \text{Gal}(F_{nr}/F)$, we have homomorphisms

$$\gamma: H_c^2(H', L_{nr}^*) \rightarrow \text{Hom}_c(H', \Delta/\Gamma_L)$$

and

$$\gamma: H_c^2(G', F_{nr}^*) \rightarrow \text{Hom}_c(G', \Delta/\Gamma_F),$$

which is helpful when we work with inertially split division algebras. (Δ is the divisible hull of Γ_F .)

Let $(\cdot n): H_c^2(H', \Gamma_L) \rightarrow H_c^2(G', \Gamma_F)$ be the map induced by multiplication map $\cdot n$ from Γ_L to Γ_F given by $\alpha \mapsto n\alpha$. (Note that $n\Gamma_L \subseteq \Gamma_F$ as $|\Gamma_L : \Gamma_F|$

divides n .) Let (v) be the maps from $H_c^2(H', L_{nr}^*)$ (resp. $H_c^2(G', F_{nr}^*)$) to $H_c^2(H', \Gamma_L)$ (resp. $H_c^2(G', \Gamma_F)$) induced by valuation v . Since v is Henselian, $v|_{F_{nr}}$ has a unique extension to F_{alg} , the algebraic closure of F . So by the argument in the proof of the theorem in [W₁], $v(N_{L_{nr}/F_{nr}}(a)) = nv(a)$ for any $a \in L_{nr}^*$ where $n = [L : F] = [L_{nr} : F_{nr}]$. So we have the following commutative diagram:

$$\begin{array}{ccccc}
 H_c^2(H', L_{nr}^*) & \xrightarrow{(v)} & & H_c^2(H', \Gamma_L) & \\
 \downarrow \mathcal{N}_{G/H}^* & \searrow N_{L_{nr}/F_{nr}}^* & & \swarrow (\cdot n) & \downarrow (\cdot n) \\
 & H_c^2(H', F_{nr}^*) & \xrightarrow{(v)} & H_c^2(H', \Gamma_F) & \\
 & \swarrow \cong & & \searrow \cong & \\
 H_c^2(G', F_{nr}^*) & \xrightarrow{(v)} & & H_c^2(G', \Gamma_F) &
 \end{array}$$

Since Δ is uniquely k -divisible for each integer $k \geq 1$, the connecting homomorphism $\delta : H_c^1(H', \Delta/\Gamma_L) \rightarrow H_c^2(H', \Gamma_L)$ is an isomorphism. So, from the diagram above we have the following commutative diagram:

$$\begin{array}{ccc}
 H_c^2(H', L_{nr}^*) & \xrightarrow{\gamma} & \text{Hom}_c(H', \Delta/\Gamma_L) \\
 \mathcal{N}_{G/H}^* \downarrow & & \downarrow (\cdot n) \\
 H_c^2(G', F_{nr}^*) & \xrightarrow{\gamma} & \text{Hom}_c(G', \Delta/\Gamma_F)
 \end{array} \quad (9)$$

We now identify $H' = \text{Gal}(L_{nr}/L)$ and $G' = \text{Gal}(F_{nr}/F)$ with

$$\text{Gal}(\overline{L}_{sep}/\overline{L}) \quad (= \text{Gal}(\overline{F}_{sep}/\overline{F})).$$

Also, we identify $H_c^2(H', L_{nr}^*)$ and $H_c^2(G', F_{nr}^*)$ with $\text{Br}(L_{nr}/L)$ and $\text{Br}(F_{nr}/F)$, respectively. Let $h_D = \gamma([D]) = \gamma(f)$, and $h_{cD} = \gamma([{}^cD]) = \gamma(\mathcal{N}_{G/H}^*(f))$. Then by [JW, Th. 5.6], the fixed field $\mathcal{F}(\ker(h_D))$ of $\ker(h_D)$ is $Z(\overline{D})$. Let $\tilde{h}_D : \text{Gal}(Z(\overline{D})/\overline{L}) \rightarrow \Gamma_D/\Gamma_L$ be the isomorphism induced by h_D (after identifying $H' / \ker(h_D)$ with $\text{Gal}(Z(\overline{D})/\overline{L})$). Then by [JW, Th. 5.6] again,

$$\begin{aligned}
 \Gamma_D/\Gamma_L &= \text{im}(h_D), \quad \ker(h_D) = \text{Gal}(\overline{L}_{sep}/Z(\overline{D})), \\
 \Gamma_{cD}/\Gamma_F &= \text{im}(h_{cD}), \quad \ker(h_{cD}) = \text{Gal}(\overline{F}_{sep}/Z({}^c\overline{D})),
 \end{aligned}$$

and $\tilde{h}_D = \theta_D^{-1}$.

Now by the commutative diagram (9), $h_{cD} = \gamma \circ \mathcal{N}_{G/H}^*(f) = (n \cdot) \circ \gamma(f) = (n \cdot)(h_D)$, so we have $\Gamma_{cD}/\Gamma_F = \text{im}(h_{cD}) = (n \cdot)(\text{im}(h_D)) = (n \cdot)(\Gamma_D/\Gamma_L) =$

$(n\Gamma_D + \Gamma_F)/\Gamma_F$. Hence $\Gamma_{cD} = n\Gamma_D + \Gamma_F$, where $n = [L : F]$. Also, we have $h_{cD}(\sigma) = n(h_D(\sigma)) + \Gamma_F$ for any $\sigma \in G' = H' = \text{Gal}(\overline{F}_{sep}/\overline{F})$, and $\ker(h_D) \subseteq \ker(h_{cD})$. So h_{cD} induces $\overline{h_{cD}}: \text{Gal}(Z(\overline{D})/\overline{F}) \rightarrow \Delta/\Gamma_F$ as $\text{Gal}(Z(\overline{D})/\overline{F}) = G'/\ker(h_D)$. Also, $\overline{h_{cD}}(\tau) = n\tilde{h}_D(\tau) + \Gamma_F = n\theta_D^{-1}(\tau) + \Gamma_F$ for any $\tau \in \text{Gal}(Z(\overline{D})/\overline{L}) = \text{Gal}(Z(\overline{D})/\overline{F})$. So as $\theta_D: \Gamma_D/\Gamma_L \rightarrow \text{Gal}(Z(\overline{D})/\overline{L})$ is an isomorphism by [JW, Lemma 5.1], $\ker(\overline{h_{cD}}) = \theta_D(\tilde{\Gamma})$, where $\tilde{\Gamma} = \{\alpha + \Gamma_L \in \Gamma_D/\Gamma_L \mid n\alpha \in \Gamma_F\}$. Hence $Z(\overline{cD}) = \mathcal{F}(\ker h_{cD}) = \mathcal{F}(\ker(\overline{h_{cD}})) = \mathcal{F}(\theta_D(\tilde{\Gamma}))$.

Note that $\tilde{\Gamma} \subseteq \tilde{\Gamma}_1$, where $\tilde{\Gamma}_1$ is the n -torsion subgroup of Γ_D/Γ_L . But, by [JW, Prop. 6.9], $Z(\overline{D^n}) = \mathcal{F}(\theta_D(\tilde{\Gamma}_1))$ where D^n is the underlying division algebra of $D^{\otimes n}$. So we have $Z(\overline{D^n}) \subseteq Z(\overline{cD})$. Also, as shown above, $\ker h_{cD} \supseteq \ker h_D$, so $Z(\overline{cD}) \subseteq Z(\overline{D})$. Therefore, $Z(\overline{D^n}) \subseteq Z(\overline{cD}) \subseteq Z(\overline{D})$.

The last assertion of the theorem follows from the definition of tame division algebra and the fact that ${}^cD \in \mathcal{D}_{is}(F)$ for $D \in \mathcal{D}_{is}(L)$. \square

3. The case when L/F is TRRT.

We begin this section by recalling the features of generalized crossed product algebras which will be needed. For further information on generalized crossed products (and proofs of the properties stated here), see [Ti₁, J], or [KY].

Let A be a central simple algebra over a field K , and suppose K is Galois over a subfield F ; let $G = \text{Gal}(K/F)$. A *generalized cocycle* of A with respect to G is a pair of functions (α, f) where $\alpha: G \rightarrow \text{Aut}_F(A)$ and $f: G \times G \rightarrow A^*$, such that for all $\sigma, \tau, \rho \in G$,

- (i) $\alpha(\sigma)|_K = \sigma$;
- (ii) $\alpha(\sigma) \circ \alpha(\tau) = \text{inn}(f(\sigma, \tau)) \circ \alpha(\sigma\tau)$, where $\text{inn}(f(\sigma, \tau))$ denotes conjugation by $f(\sigma, \tau)$;
- (iii) $f(\sigma, \tau)f(\sigma\tau, \rho) = [\alpha(\sigma)(f(\tau, \rho))]f(\sigma, \tau\rho)$.

(α, f) is said to be *normalized* if $\alpha(\text{id}_K) = \text{id}_A$ and $f(\text{id}_K, \sigma) = f(\sigma, \text{id}_K) = 1$ for all $\sigma \in G$. Given a normalized generalized cocycle (α, f) one forms the *generalized crossed product* $(A, G, (\alpha, f))$ as the free left A -module with base $\{x_\sigma \mid \sigma \in G\}$, which is made into a ring by the multiplication rule

$$(cx_\sigma)(dx_\tau) = [c\alpha(\sigma)(d)f(\sigma, \tau)]x_{\sigma\tau} \quad \text{for all } c, d \in A, \sigma, \tau \in G.$$

$(A, G, (\alpha, f))$ is a central simple F -algebra. Observe that if S is any central simple F -algebra containing K , then one sees using the Skolem-Noether theorem that there is a normalized generalized crossed product $(A, G, (\alpha, f))$ isomorphic to S , where $A = C_S(K)$. We will need the product theorem for generalized crossed products (cf. [Ti₁, Th. 4.6] or [J, (1.15)] or [KY, Th. 3]). This says if $(A, G, (\alpha, f))$ and $(B, G, (\beta, g))$ are generalized crossed products

of K over F , then with respect to the obvious induced generalized cocycle $(\alpha \otimes \beta, f \otimes g)$ of $A \otimes_K B$, we have in $\text{Br}(F)$,

$$(10) \quad (A, G, (\alpha, f)) \otimes_F (B, G, (\beta, g)) \sim (A \otimes_K B, G, (\alpha \otimes \beta, f \otimes g)).$$

Proposition 11. *For $i = 1, 2$, let $(T_i, v_i) \in \mathcal{D}_{\text{tr}}(F)$ with $v_1|_F = v_2|_F$. ($v_1|_F$ doesn't need be Heselien.) Suppose there is an extension field K of F of degree n such that $K \subseteq T_i$, $\Gamma_{T_1} \cap \Gamma_{T_2} = \Gamma_K$, and K is Galois over F with Galois group G . So there are normalized generalized crossed products $(C_i, G, (\alpha_i, f_i))$ isomorphic to T_i . Then, $C_1 \otimes_K C_2$ is a division ring with a unique valuation v such that $v|_{C_1} = v_1|_{C_1}$, $v|_{C_2} = v_2|_{C_2}$, $\Gamma_{C_1 \otimes_K C_2} = \Gamma_{C_1} + \Gamma_{C_2}$ and $(C_1 \otimes_K C_2, v) \in \mathcal{D}_{\text{tr}}(K)$. Also let $T = (C_1 \otimes_K C_2, G, (\alpha_1 \otimes \alpha_2, f_1 \otimes f_2))$. Then $T_1 \otimes_F T_2 \cong M_n(T)$ and T is a division ring with a valuation w such that $w|_{C_1 \otimes_K C_2} = v$ and $(T, w) \in \mathcal{D}_{\text{tr}}(F)$. So $K \subseteq T$ and $\Gamma_{C_1} + \Gamma_{C_2} \subseteq \Gamma_T$.*

Proof. Since $T_i \in \mathcal{D}_{\text{tr}}(F)$, the fundamental inequality (**S**, p. 21) gives $|\Gamma_{C_i} : \Gamma_K| = [C_i : K]$ and $\overline{C_1} = \overline{C_2} = \overline{K}$. Also $\Gamma_K \subseteq \Gamma_{C_1} \cap \Gamma_{C_2} \subseteq \Gamma_{T_1} \cap \Gamma_{T_2} = \Gamma_K$, so $\Gamma_{C_1} \cap \Gamma_{C_2} = \Gamma_K$. So by Prop. 4, $C_1 \otimes_K C_2$ is a division ring with a unique valuation v such that $v|_{C_1} = v_1|_{C_1}$ and $v|_{C_2} = v_2|_{C_2}$. Furthermore, $\Gamma_{C_1 \otimes_K C_2} = \Gamma_{C_1} + \Gamma_{C_2}$. So $|\Gamma_{C_1 \otimes_K C_2} : \Gamma_K| = [C_1 \otimes_K C_2 : K]$. Hence $(C_1 \otimes_K C_2, v) \in \mathcal{D}_{\text{tr}}(K)$ as $\text{char}(\overline{K}) \nmid [C_1 \otimes_K C_2 : K]$.

Let $T = (C_1 \otimes_K C_2, G, (\alpha_1 \otimes \alpha_2, f_1 \otimes f_2))$. Then by (10) in $\text{Br}(F)$, $T_1 \otimes_F T_2 \sim T$. As $[T : F] = [T_1 \otimes_F T_2 : F]/n^2$ where $n = [K : F]$, $T_1 \otimes_F T_2 \cong M_n(T)$. Our next goal is to define a valuation on T .

We have $T_1 \cong (C_1, G, (\alpha_1, f_1)) = \bigoplus_{\sigma \in G} C_1 x_\sigma$, i.e. the free left C_1 -module with base $\{x_\sigma \mid \sigma \in G\}$, which is made into a ring by the multiplication rule

$$(cx_\sigma)(dx_\tau) = [c\alpha_1(\sigma)(d)f_1(\sigma, \tau)]x_{\sigma\tau} \quad \text{for all } c, d \in C_1, \sigma, \tau \in G,$$

where (α_1, f_1) is a normalized generalized cocycle of C_1 with respect to G . Likewise, $T_2 \cong (C_2, G, (\alpha_2, f_2)) = \bigoplus_{\sigma \in G} C_2 y_\sigma$, with multiplication rule

$$(cy_\sigma)(\alpha y_\tau) = [c\alpha_2(\sigma)(d)f_2(\sigma, \tau)]y_{\sigma\tau} \quad \text{for all } c, d \in C_2, \sigma, \tau \in G,$$

where (α_2, f_2) is a normalized generalized cocycle of C_2 with respect to G .

We now claim that $v_1(x_\sigma) \in \Gamma_{C_1}$ if and only if $\sigma = \text{id}$. For, suppose $v_1(x_\sigma) = v_1(c)$ for some $c \in C_1$. Then by replacing x_σ by $c^{-1}x_\sigma$ we may assume that $v_1(x_\sigma) = 0$. Since $(T_1, v_1) \in \mathcal{D}_{\text{tr}}(F)$, there is a canonical (bilinear) pairing $\overline{C_{T_1}} : (\Gamma_{T_1}/\Gamma_F) \times (\Gamma_{T_1}/\Gamma_F) \rightarrow \mu_\ell(\overline{F})$ given by $(v_1(d) + \Gamma_F, v_1(e) + \Gamma_F) \mapsto \overline{ded^{-1}e^{-1}}$ where $\ell = \exp(\Gamma_{T_1}/\Gamma_F)$ and $\mu_\ell(\overline{F})$ is the set of ℓ distinct ℓ -th roots of unity in \overline{F} . (cf. [**TW**, Sec. 3]). As $v_1(x_\sigma) = 0 \in \Gamma_F$ and $x_\sigma k x_\sigma^{-1} = \sigma(k)$ for all $k \in K$, $\bar{1} = C_{T_1}(v_1(k) + \Gamma_F, v_1(x_\sigma) + \Gamma_F) =$

$\overline{kx_\sigma k^{-1}x_\sigma^{-1}} = \overline{k/\sigma(k)}$, so that $\overline{k/\sigma(k)} = \bar{1}$ for all $k \in K^*$. But since with respect to $v_1|_K$, K is tame and totally ramified over F and K is Galois over F with Galois group G , by [TW, Prop. 1.4] or [E, (20.11), pp. 161-162], there is a well-defined nondegenerate bilinear pairing

$$\gamma: (\Gamma_K/\Gamma_F) \times G \rightarrow \mu(\overline{F}) \quad \text{given by} \quad \gamma(v(k) + \Gamma_F, \tau) = \overline{k/\tau(k)} \in \overline{F}$$

for all $k \in K^*$, $\tau \in G$, where $\mu(\overline{F})$ is the set of roots of unity in \overline{F} . Thus $\sigma = \text{id}$ as claimed. Likewise, $v_2(y_\sigma) \in \Gamma_{C_2}$ if and only if $\sigma = \text{id}$.

It follows from the claim above that $v_1(x_\sigma)$ (resp. $v_2(y_\sigma)$), $\sigma \in G$, are distinct modulo Γ_{C_1} (resp. Γ_{C_2}). For, if $v_1(x_\sigma) - v_1(x_\tau) \in \Gamma_{C_1}$, then as $x_{\sigma\tau^{-1}}x_\tau = f(\sigma\tau^{-1}, \tau)x_\sigma$, we have $v(x_{\sigma\tau^{-1}}) = v(f(\sigma\tau^{-1}, \tau)) + [v(x_\sigma) - v(x_\tau)] \in \Gamma_{C_1}$. So, $\sigma\tau^{-1} = \text{id}$ by the claim above, i.e., $\sigma = \tau$.

We will now define a valuation w on $T = (C_1 \otimes_K C_2, G, (\alpha_1 \otimes \alpha_2, f_1 \otimes f_2)) = \bigoplus_{\sigma \in G} (C_1 \otimes_K C_2)z_\sigma$, such that $w|_{C_1 \otimes_K C_2} = v$ and $(T, w) \in \mathcal{D}_{\text{ttr}}(F)$. Define $w(z_\sigma) = v_1(x_\sigma) + v_2(y_\sigma)$ and

$$w\left(\sum_{\sigma \in G} c_\sigma z_\sigma\right) = \min_{\sigma \in G} (v(c_\sigma) + w(z_\sigma))$$

where $c_\sigma \in C_1 \otimes_K C_2$ for all $\sigma \in G$. If $w(z_\sigma) - w(z_\tau) \in \Gamma_{C_1 \otimes_K C_2} = \Gamma_{C_1} + \Gamma_{C_2}$, say $w(z_\sigma) - w(z_\tau) = \gamma_1 + \gamma_2$ for some $\gamma_i \in \Gamma_{C_i}$, then $v_1(x_\sigma) - v_1(x_\tau) - \gamma_1 = v_2(y_\tau) - v_2(y_\sigma) + \gamma_2 \in \Gamma_{T_1} \cap \Gamma_{T_2} = \Gamma_K$. So $v_1(x_\sigma) - v_1(x_\tau) \in \Gamma_{C_1}$, showing $\sigma = \tau$ as above. So the $w(z_\sigma)$, $\sigma \in G$, are distinct modulo $\Gamma_{C_1 \otimes_K C_2}$. Hence every element $\sum_{\sigma \in G} c_\sigma z_\sigma$ in T , with $c_\sigma \in C_1 \otimes_K C_2$, has a unique summand $c_\tau z_\tau$ with minimum value, which is called the *leading term* of $\sum_{\sigma \in G} c_\sigma z_\sigma$. We will show that w actually defines a valuation on T so that $(T, w) \in \mathcal{D}_{\text{ttr}}(F)$.

For $d = \sum_{\sigma \in G} d_\sigma z_\sigma \neq 0$ and $e = \sum_{\sigma \in G} e_\sigma z_\sigma \neq 0$ in T with $d \neq -e$, it is easy to see that $w(d + e) \geq \min(w(d), w(e))$; hence $w(d + e) = \min(w(d), w(e))$ if $w(d) \neq w(e)$. For the behavior of w under products, consider first $(d_\sigma z_\sigma) \cdot (e_\tau z_\tau)$ with $d_\sigma, e_\tau \neq 0 \in C_1 \otimes_K C_2$. Note that v is the unique extension of $v_1|_F = v_2|_F$ to $C_1 \otimes_K C_2$; so $v(z_\sigma e_\tau z_\sigma^{-1}) = v(e_\tau)$. Thus,

$$\begin{aligned} w((d_\sigma z_\sigma) \cdot (e_\tau z_\tau)) &= w([d_\sigma(\alpha_1 \otimes \alpha_2)(\sigma)(e_\tau)(f_1 \otimes f_2)(\sigma, \tau)]z_{\sigma\tau}) \\ &= v(d_\sigma \cdot (z_\sigma e_\tau z_\sigma^{-1}) \cdot f_1(\sigma, \tau) \cdot f_2(\sigma, \tau)) + w(z_{\sigma\tau}) \\ &= v(d_\sigma) + v(e_\tau) + v_1(f_1(\sigma, \tau)) + v_2(f_2(\sigma, \tau)) \\ &\quad + v_1(x_{\sigma\tau}) + v_2(y_{\sigma\tau}) \\ &= v(d_\sigma) + v(e_\tau) + v_1(x_\sigma x_\tau) + v_2(y_\sigma y_\tau) \\ &= [v(d_\sigma) + v_1(x_\sigma) + v_2(y_\sigma)] + [v(e_\tau) + v_1(x_\tau) + v_2(y_\tau)] \\ &= w(d_\sigma z_\sigma) + w(e_\tau z_\tau). \end{aligned}$$

Hence, if $de \neq 0$,

$$\begin{aligned} w(de) &= w\left(\left(\sum_{\sigma \in G} d_\sigma z_\sigma\right) \cdot \left(\sum_{\tau \in G} e_\tau z_\tau\right)\right) \\ &\geq \min\{w((d_\sigma z_\sigma)(e_\tau z_\tau)) \mid d_\sigma \neq 0, e_\tau \neq 0\} \\ &= \min\{w(d_\sigma z_\sigma) + w(e_\tau z_\tau) \mid d_\sigma \neq 0, e_\tau \neq 0\} = w(d) + w(e), \end{aligned}$$

i.e. $w(de) \geq w(d) + w(e)$. Now, let $d_{\sigma_0} z_{\sigma_0}$ be the leading term of d . Then $d = d_{\sigma_0} z_{\sigma_0} + d'$ where $d' = 0$ or $w(d') > w(d_{\sigma_0} z_{\sigma_0})$. Likewise, write $e = e_{\tau_0} z_{\tau_0} + e'$ where $e_{\tau_0} z_{\tau_0}$ is the leading term of e . Then $de = (d_{\sigma_0} z_{\sigma_0})(e_{\tau_0} z_{\tau_0}) + \rho$ where $\rho = d'(e_{\tau_0} z_{\tau_0}) + (d_{\sigma_0} z_{\sigma_0})e' + d'e'$. Now, if $\rho \neq 0$, $w(\rho) > w((d_{\sigma_0} z_{\sigma_0})(e_{\tau_0} z_{\tau_0}))$ by what has been already proved. Hence, $\rho \neq -(d_{\sigma_0} z_{\sigma_0})(e_{\tau_0} z_{\tau_0})$. Therefore, $de \neq 0$ and $w(de) = w((d_{\sigma_0} z_{\sigma_0})(e_{\tau_0} z_{\tau_0})) = w(d) + w(e)$. This shows that T has no zero divisors so T must be a division ring, since $\dim_F T < \infty$. The formula also shows w is a valuation on T . Clearly, $\Gamma_T = \{w(z_\sigma) \mid \sigma \in G\} + \Gamma_{C_1 \otimes_K C_2}$. As the $w(z_\sigma)$, $\sigma \in G$, are distinct modulo $\Gamma_{C_1 \otimes_K C_2}$ and $|G| = [K:F] = n$, we have $|\Gamma_T : \Gamma_F| = n \cdot |\Gamma_{C_1 \otimes_K C_2} : \Gamma_F| = n \cdot [C_1 \otimes_K C_2 : F] = [T:F]$. So $(T, w) \in \mathcal{D}_{\text{tr}}(F)$. \square

Next, when $(L, v) \supseteq (F, v)$ is a finite separable, TRRT extension of Henselian fields, we will give relations between $T \in \mathcal{D}_{\text{tr}}(L)$ (resp. $D \in \mathcal{D}_t(L)$) and cT (resp. cD) in Theorems 12 and 13. To prove these theorems, we need the following proposition.

Proposition 12. *Let (F, v) be a Henselian field and let p be a prime. Let $L = F(\gamma)$ with $\gamma^p \in F^*$ and $v(\gamma) \notin p\Gamma_F$. Let $\alpha, \beta \in L^*$.*

(1) *Every symbol algebra $(\alpha, \beta; L)_n$ with $\text{char}(\bar{L}) \nmid n$ is isomorphic to a symbol algebra of the form $(a, b\gamma^j; L)_n$ for some $a, b \in F^*$ and $0 \leq j \leq p-1$.*

(2) *If T is a symbol algebra in $\mathcal{D}_{\text{tr}}(L)$, then ${}^cT \in \mathcal{D}_{\text{tr}}(F)$, $p\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$, and $\exp(\Gamma_{{}^cT} / \Gamma_F) \mid \exp(\Gamma_T / \Gamma_L)$.*

Proof. Since $\gamma^p \in F^*$ and $v(\gamma) \notin p\Gamma_F$, $v(\gamma) + \Gamma_F$ has order p in $\frac{1}{p}\Gamma_F / \Gamma_F$, so L/F is totally ramified of degree p with $\Gamma_L = \langle v(\gamma) \rangle + \Gamma_F$. Hence L is a TRRT extension of F .

(1) Since $\{1, \gamma, \gamma^2, \dots, \gamma^{p-1}\}$ is an F -basis of L , $\alpha = a_0 + a_1\gamma + \dots + a_{p-1}\gamma^{p-1}$, and $\beta = b_0 + b_1\gamma + \dots + b_{p-1}\gamma^{p-1}$ with all $a_k, b_k \in F$. Since $v(\gamma^k)$, $0 \leq k \leq p-1$, are distinct modulo Γ_F , $\alpha = \sum_{k=0}^{p-1} a_k \gamma^k$ (resp. $\beta = \sum_{k=0}^{p-1} b_k \gamma^k$) has a unique summand, say $a_i \gamma^i$ (resp. $b_j \gamma^j$) with minimal value. Then $\alpha = a_i \gamma^i (1 + \alpha')$ and $\beta = b_j \gamma^j (1 + \beta')$ with $v(\alpha') > 0$ and $v(\beta') > 0$. Since (L, v) is Henselian and $\text{char}(\bar{L}) \nmid n$, $1 + \alpha' = \alpha_0^n$ and $1 + \beta' = \beta_0^n$ for some $\alpha_0, \beta_0 \in L^*$. So $(\alpha, \beta; L)_n \cong (a_i \gamma^i, b_j \gamma^j; L)_n$ for some $a_i, b_j \in F^*$ and i, j , $0 \leq i, j \leq p-1$.

Now recall that $(\alpha_1, \beta_1; L)_n \cong (\beta_1^{-1}, \alpha_1; L)_n$ by [D₁, p. 80, Lemma 5]. So if $j = 0$, $(\alpha, \beta; L)_n \cong (a_i \gamma^i, b_0; L)_n \cong (b_0^{-1}, a_i \gamma^i; L)_n$, as desired. Hence we may assume that $i > 0$ and $j > 0$. We argue by induction on $i + j$ that $(a_i \gamma^i, b_j \gamma^j; L)_n \cong (a, b \gamma^{j'}; L)_n$ for some a, b, j' . First, let's assume $i \leq j$. Then since $(a_i \gamma^i, -a_i \gamma^i; L)_n \sim L$ by [D₁, p. 82, Cor. 5], $(a_i \gamma^i, b_j \gamma^j; L)_n \cong (a_i \gamma^i, -a_i^{-1} b_j \gamma^{j-i}; L)_n$. As $i + (j - i) < i + j$, we have done by induction. We argue similarly when $i > j$. Note that this proof of (1) does not need the assumption that $[L : F]$ is prime.

(2) If T is a symbol division algebra of degree n in $\mathcal{D}_{tr}(L)$, by (1) above $T \cong (a, b \gamma^j; L)_n$ for some $a, b \in F^*$ and j , $0 \leq j \leq p - 1$. (So $\mu_n \subseteq L$.) Since T is tame and totally ramified over L , by [H₂, Prop. 3.1] in $\Gamma_L / n\Gamma_L$, $|\langle v(a) + n\Gamma_L, v(b \gamma^j) + n\Gamma_L \rangle| = n^2$ and $\Gamma_T = \langle \frac{1}{n}v(a), \frac{1}{n}v(b \gamma^j) \rangle + \Gamma_L = \langle \frac{1}{n}v(a), \frac{1}{n}v(b \gamma^j) \rangle + \langle v(\gamma) \rangle + \Gamma_F$. Let $K = L(\sqrt[n]{a})$. Then K is tame and totally ramified over L with $\Gamma_K / \Gamma_L = \langle \frac{1}{n}v(a) + \Gamma_L \rangle$ and $\exp(\Gamma_K / \Gamma_L) = n$ as $|\langle \frac{1}{n}v(a) + \Gamma_L : \Gamma_L \rangle| = n$. Also, since $\mu_n \subseteq L$, by [TW, Prop. 1.4 (iii)] or [S, p. 64, Th. 3], $\mu_n \subseteq \bar{L} = \bar{F}$, hence $\mu_n \subseteq F$ as (F, v) is Henselian.

(i) If $j = 0$, then $T \cong (a, b; L)_n$ for some $a, b \in F^*$ and

$$\Gamma_T = \left\langle \frac{1}{n}v(a), \frac{1}{n}v(b) \right\rangle + \langle v(\gamma) \rangle + \Gamma_F.$$

Since $|\Gamma_T : \Gamma_F| = n^2 p$, we must have $\Gamma_T / \Gamma_F = \langle \frac{1}{n}v(a) + \Gamma_F \rangle \times \langle \frac{1}{n}v(b) + \Gamma_F \rangle \times \langle v(\gamma) + \Gamma_F \rangle \cong \mathbb{Z}_n \times \mathbb{Z}_n \times \mathbb{Z}_p$. Since $\mu_n \subseteq F$, by [Ti₂, Th. 3.1] (Projection formula), as $N_{L/F}(b) = b^p$, ${}^cT \sim (a, b^p; F)_n$ in $\text{Br}(F)$.

If $p \nmid n$, then in $\frac{1}{n}\Gamma_F / \Gamma_F$, $|\langle \frac{1}{n}v(a) + \Gamma_F, \frac{1}{n}v(b^p) + \Gamma_F \rangle| = |\langle \frac{1}{n}v(a) + \Gamma_F, \frac{1}{n}v(b) + \Gamma_F \rangle| = n^2$. So by [H₂, Prop. 3.1], ${}^cT \in \mathcal{D}_{tr}(F)$ with

$$\Gamma_{{}^cT} = \left\langle \frac{1}{n}v(a), \frac{1}{n}v(b^p) \right\rangle + \Gamma_F,$$

whence $p\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$. If $p|n$, then ${}^cT \sim (a, b; F)_{n/p}$ and in $\frac{1}{(n/p)}\Gamma_F / \Gamma_F$,

$$\left| \left\langle \frac{1}{(n/p)}, v(a) + \Gamma_F, \frac{1}{(n/p)}v(b) + \Gamma_F \right\rangle \right| = (n/p)^2.$$

So by [H₂, Prop. 3.1], ${}^cT \cong (a, b; F)_{n/p} \in \mathcal{D}_{tr}(F)$ with $\Gamma_{{}^cT} = \langle \frac{p}{n}v(a), \frac{p}{n}v(b) \rangle + \Gamma_F$ whence $p\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$. In either case,

$$\exp(\Gamma_{{}^cT} / \Gamma_F) \mid \exp(\Gamma_T / \Gamma_L).$$

(ii) If $0 < j \leq p - 1$, then $T \cong (a, \beta; L)_n$ where $a \in F^*$ and $\beta = b \gamma^j$ with $b \in F^*$ and $1 \leq j \leq p - 1$. Since $v(\beta) = v(b) + jv(\gamma) \in \langle \frac{1}{n}v(\beta) \rangle + \Gamma_F$ and $jk \equiv 1 \pmod{p}$ for some integer k , and $\gamma^p \in F^*$, $v(\gamma) \equiv jkv(\gamma) \equiv kv(\beta)$

mod Γ_F so that $v(\gamma) \in \langle \frac{1}{n}v(\beta) \rangle + \Gamma_F$. Hence $\Gamma_T = \langle \frac{1}{n}v(a), \frac{1}{n}v(\beta) \rangle + \langle v(\gamma) \rangle + \Gamma_F = \langle \frac{1}{n}v(a), \frac{1}{n}v(\beta) \rangle + \Gamma_F$.

Since $\mu_n \subseteq F$, by [Ti₂, Th. 3.1] (Projection formula),

$${}^cT \sim (a, N_{L/F}(\beta); F)_n$$

in $\text{Br}(F)$. Note that $v(N_{L/F}(\beta)) = [L:F]v(\beta) = pv(\beta)$ by the argument in the proof of the theorem in [W₁]. So

$$\left| \Gamma_T : \left(\left\langle \frac{1}{n}v(a), \frac{1}{n}v(N_{L/F}(\beta)) \right\rangle + \Gamma_F \right) \right| \leq p.$$

As $|\Gamma_T/\Gamma_F| = |\langle \frac{1}{n}v(a) + \Gamma_F, \frac{1}{n}v(\beta) + \Gamma_F \rangle| = n^2p$,

$$\left| \left\langle \frac{1}{n}v(a) + \Gamma_F, \frac{1}{n}v(N_{L/F}(\beta)) + \Gamma_F \right\rangle \right| = n^2.$$

So by [H₂, Prop. 3.1],

$${}^cT \cong (a, N_{L/F}(\beta); F)_n \in \mathcal{D}_{ttr}(F)$$

with $\Gamma_{{}^cT} = \langle \frac{1}{n}v(a), \frac{p}{n}v(\beta) \rangle + \Gamma_F$. Hence $p\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$, and

$$\exp(\Gamma_{{}^cT}/\Gamma_F) \mid \exp(\Gamma_T/\Gamma_L).$$

□

Theorem 13. *Let $(L, v) \supseteq (F, v)$ be a finite separable TRRT extension of Henselian fields. If $T \in \mathcal{D}_{ttr}(L)$, then ${}^cT \in \mathcal{D}_{ttr}(F)$, $[L:F] \cdot \Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$, and $\exp(\Gamma_{{}^cT}/\Gamma_F) \mid \exp(\Gamma_T/\Gamma_L)$.*

Proof. If $[L:F]$ is not prime, then by [JW, Remark 4.2] there is a field F_1 such that $F \subsetneq F_1 \subsetneq L$ with $[F_1:F] = p$, prime, and $L/F_1, F_1/F$ TRRT. Assume that we can prove the theorem for F_1/F of prime degree. Since $[L:F_1] < [L:F]$, we can also assume that the theorem holds for L/F_1 by induction on $[L:F]$. Then the theorem is proved for L/F : Let T_1 be the underlying division algebra of $\text{cor}_{L/F_1}(T)$. Then by the assumption for L/F_1 , $T_1 \in \mathcal{D}_{ttr}(F_1)$, $[L:F_1] \cdot \Gamma_T \subseteq \Gamma_{T_1} \subseteq \Gamma_T$, and $\exp(\Gamma_{T_1}/\Gamma_{F_1}) \mid \exp(\Gamma_T/\Gamma_L)$. Also since ${}^cT \sim \text{cor}_{L/F}(T) \sim \text{cor}_{F_1/F}(T_1)$, by the assumption for F_1/F , ${}^cT \in \mathcal{D}_{ttr}(F)$ and $[F_1:F] \cdot \Gamma_{T_1} \subseteq \Gamma_{{}^cT} \subseteq \Gamma_{T_1}$, and $\exp(\Gamma_{{}^cT}/\Gamma_F) \mid \exp(\Gamma_{T_1}/\Gamma_{F_1})$. So $[L:F] \cdot \Gamma_T = [F_1:F][L:F_1]\Gamma_T \subseteq [F_1:F]\Gamma_{T_1} \subseteq \Gamma_{{}^cT} \subseteq \Gamma_{T_1} \subseteq \Gamma_T$, hence $[L:F]\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$. Also $\exp(\Gamma_{{}^cT}/\Gamma_F) \mid \exp(\Gamma_T/\Gamma_L)$.

So we may assume that $[L:F] = p$, prime, so that $L = F(\gamma)$ with $\gamma^p \in F^*$ and $v(\gamma) \notin p\Gamma_F$. Also by the primary decomposition we may assume that

$[T : F] = q^r$, a power of a prime number q . By $[\mathbf{D}_2, \text{Th. 1}]$ (Draxl's decomposition theorem for tame and totally ramified division algebras) $T \cong \bigotimes_{i=1}^k T_i$ with each T_i a symbol division algebra in $\mathcal{D}_{ttr}(L)$, and $\Gamma_T / \Gamma_L = \bigoplus_{i=1}^k (\Gamma_{T_i} / \Gamma_L)$. (So $(\Gamma_{T_1} + \cdots + \Gamma_{T_{j-1}}) \cap \Gamma_{T_j} = \Gamma_L$ for $2 \leq j \leq k$.) By Prop. 12 $T_i \cong (a_i, b_i \gamma^{e_i}; L)_{n_i}$ for some $a_i, b_i \in F^*$, $0 \leq e_i \leq p-1$, and n_i , a power of q , and $p\Gamma_{T_i} \subseteq \Gamma_{cT_i} \subseteq \Gamma_{T_i}$. So for $2 \leq j \leq k$, $(\Gamma_{cT_1} + \cdots + \Gamma_{cT_{j-1}}) \cap \Gamma_{cT_j} \subseteq (\Gamma_{T_1} + \cdots + \Gamma_{T_{j-1}}) \cap \Gamma_{T_j} = \Gamma_L$. As $|\Gamma_L : \Gamma_F| = p$, $\left| \left(\sum_{i=1}^{j-1} \Gamma_{cT_i} \right) \cap \Gamma_{cT_j} : \Gamma_F \right|$ divides p . Also since the (Schur) index of cT_j (i.e. $\sqrt{[{}^cT_j : F]}$) is a power of q , $|\Gamma_{cT_j} : \Gamma_F|$ is a power of q , so $\left| \left(\sum_{i=1}^{j-1} \Gamma_{cT_i} \right) \cap \Gamma_{cT_j} : \Gamma_F \right|$ is a power of q .

If $q \neq p$, then for $2 \leq j \leq k$, $\left(\sum_{i=1}^{j-1} \Gamma_{cT_i} \right) \cap \Gamma_{cT_j} = \Gamma_F$ as the index divides both p and a power of q . Also by Prop. 12 ${}^cT_i \in \mathcal{D}_{ttr}(F)$ for $1 \leq i \leq k$. So by applying Prop. 2 repeatedly, we have ${}^cT \cong \bigotimes_{i=1}^k {}^cT_i \in \mathcal{D}_{ttr}(F)$ and $\Gamma_{cT} = \sum_{i=1}^k \Gamma_{cT_i}$. So $p\Gamma_T = p \left(\sum_{i=1}^k \Gamma_{T_i} \right) \subseteq \sum_{i=1}^k \Gamma_{cT_i} = \Gamma_{cT} = \sum_{i=1}^k \Gamma_{cT_i} \subseteq \sum_{i=1}^k \Gamma_{T_i} = \Gamma_T$, as desired. So we may assume $q = p$, so that $p \mid n_i = \text{ind}(T_i)$ for $1 \leq i \leq k$.

Now we will prove the theorem by induction on k . If $k = 1$, the assertion is Prop. 12 (2).

If one of T_i , say T_{i_0} , is isomorphic to $(a, b; L)_n$ for some $a, b \in F^*$, then by reindexing we may assume $T_k \cong (a, b; L)_n$. Then $\Gamma_{T_k} / \Gamma_F = \langle \frac{1}{n}v(a) + \Gamma_F \rangle \oplus \langle \frac{1}{n}v(b) + \Gamma_F \rangle \oplus \langle v(\gamma) + \Gamma_F \rangle \cong \mathbb{Z}_n \oplus \mathbb{Z}_n \oplus \mathbb{Z}_p$. As ${}^cT_k = (a, b; F)_{n/p}$ with $p \mid n$, and $\Gamma_L = \langle v(\gamma) \rangle + \Gamma_F$, we have $\Gamma_{cT_k} \cap \Gamma_L = \Gamma_F$. This means that $(\Gamma_{cT_1} + \cdots + \Gamma_{cT_{k-1}}) \cap \Gamma_{cT_k} = \Gamma_F$ since $(\Gamma_{cT_1} + \cdots + \Gamma_{cT_{k-1}}) \cap \Gamma_{cT_k} \subseteq \Gamma_L$. Let $T_0 = \bigotimes_{i=1}^{k-1} T_i$. Then we have ${}^cT_0 \in \mathcal{D}_{ttr}(F)$ and $[L : F]\Gamma_{T_0} \subseteq \Gamma_{cT_0} \subseteq \Gamma_{T_0}$ by the inductive assumption on k . As ${}^cT_i \in \mathcal{D}_t(F)$, $\Gamma_{cT_0} \subseteq \Gamma_{cT_1} + \cdots + \Gamma_{cT_{k-1}}$ by $[\mathbf{JW}, \text{Cor. 6.7}]$, so $\Gamma_{cT_0} \cap \Gamma_{cT_k} = \Gamma_F$. Hence by Prop. 2, ${}^cT \cong {}^cT_0 \otimes_F {}^cT_k \in \mathcal{D}(F)$ and $\Gamma_{cT} = \Gamma_{cT_0} + \Gamma_{cT_k}$. So ${}^cT \in \mathcal{D}_{ttr}(F)$ and $p\Gamma_T \subseteq \Gamma_{cT} \subseteq \Gamma_T$, as claimed. Therefore we may assume $T = \bigotimes_{i=1}^k T_i$ where $T_i \cong (a_i, b_i \gamma^{e_i}; L)_{n_i}$, $0 < e_i \leq p-1$, and $a_i, b_i \in F^*$. As $T_i \in \mathcal{D}_{ttr}(L)$, $\Gamma_{T_i} = \langle \frac{1}{n_i}v(a_i), \frac{1}{n_i}v(b_i) + \frac{e_i}{n_i}v(\gamma) \rangle + \langle v(\gamma) \rangle + \Gamma_F$. Then ${}^cT \sim \bigotimes_{i=1}^k {}^cT_i$ in $\text{Br}(F)$. Also, ${}^cT_i \cong (a_i, b_i^p N_{L/F}(\gamma)^{e_i}; F)_{n_i} \in \mathcal{D}_{ttr}(F)$ and $p\Gamma_{T_i} \subseteq \Gamma_{cT_i} \subseteq \Gamma_{T_i}$ as shown in the last paragraph of the proof of Prop. 12 (2).

Let $K = F \left(\sqrt[p]{N_{L/F}(\gamma)^{e_1}} \right) = F \left(\sqrt[p]{N_{L/F}(\gamma)} \right)$. Note in passing that $K = L$

unless $p = 2$ and $\mu_4 \not\subseteq F$. Since each cT_i contains

$$F \left(\left(b_i^p N_{L/F}(\gamma)^{e_i} \right)^{1/n_i} \right)$$

as a maximal subfield, and n_i is a power of p , each cT_i contains K . Also $\Gamma_K = \Gamma_L$. Since $\mu_{n_i} \subseteq F$ as shown in the first paragraph of proof of Prop. 12 (2), K is Galois over F . Let $G = \text{Gal}(K/F)$, and let C_i be the centralizer of K in cT_i . Then ${}^cT_i \cong (C_i, G, (\alpha_i, f_i))$, a generalized crossed product of C_i with respect to G .

We show by induction on k that ${}^cT \in \mathcal{D}_{ttr}(F)$ and ${}^cT \cong (C_1 \otimes_K \cdots \otimes_K C_k, G, (\alpha_1 \otimes \cdots \otimes \alpha_k, f_1 \otimes \cdots \otimes f_k))$, a generalized crossed product. This holds for $k = 1$ by Prop. 12 (2). Assume that it is true for $k - 1$, so that if $T_0 = \bigotimes_{i=1}^{k-1} T_i$, then ${}^cT_0 \in \mathcal{D}_{ttr}(F)$ and ${}^cT_0 \cong (C_1 \otimes_K \cdots \otimes_K C_{k-1}, G, (\alpha_1 \otimes \cdots \otimes \alpha_{k-1}, f_1 \otimes \cdots \otimes f_{k-1}))$. Then cT is the underlying division algebra of ${}^cT_0 \otimes_F {}^cT_k$. As cT_i is tame over F for each i , $1 \leq i \leq k - 1$, and ${}^cT_0 \sim \bigotimes_{i=1}^{k-1} {}^cT_i$,

by [JW, Cor. 6.7] and Prop. 12 (2) we have $\Gamma_{{}^cT_0} \subseteq \sum_{i=1}^{k-1} \Gamma_{{}^cT_i} \subseteq \sum_{i=1}^{k-1} \Gamma_{T_i} = \Gamma_{T_0}$. So $\Gamma_{{}^cT_0} \cap \Gamma_{{}^cT_k} \subseteq \Gamma_{T_0} \cap \Gamma_{T_k} = \Gamma_L = \Gamma_K$, whence $\Gamma_{{}^cT_0} \cap \Gamma_{{}^cT_k} = \Gamma_K$. Since ${}^cT_0, {}^cT_k \in \mathcal{D}_{ttr}(F)$, $K \subseteq {}^cT_0$, $K \subseteq {}^cT_k$, and $\Gamma_{{}^cT_0} \cap \Gamma_{{}^cT_k} = \Gamma_K$, by Prop. 11 ${}^cT \in \mathcal{D}_{ttr}(F)$ and ${}^cT \cong (C_1 \otimes_K \cdots \otimes_K C_k, G, (\alpha_1 \otimes \cdots \otimes \alpha_k, f_1 \otimes \cdots \otimes f_k))$, as desired.

Since each C_i is the centralizer of K in ${}^cT_i \in \mathcal{D}_{ttr}(F)$, by [TW, Th. 3.8], $\Gamma_{C_i} / \Gamma_F = (\Gamma_K / \Gamma_F)^\perp$ relative to the canonical pairing $C_{{}^cT_i} : (\Gamma_{{}^cT_i} / \Gamma_F) \times (\Gamma_{{}^cT_i} / \Gamma_F) \rightarrow \mu(\overline{F})$ given by $(v(d) + \Gamma_F, v(e) + \Gamma_F) \mapsto \overline{ded^{-1}e^{-1}}$. Because the pairing is nondegenerate, $|\Gamma_{T_i} : \Gamma_{C_i}| = |\Gamma_{T_i} / \Gamma_F : (\Gamma_K / \Gamma_F)^\perp| = |\Gamma_K / \Gamma_F| = p$. So $p\Gamma_{T_i} \subseteq \Gamma_{C_i}$. Hence $\Gamma_{{}^cT} \supseteq \Gamma_{C_1 \otimes_K \cdots \otimes_K C_k} = \sum_{i=1}^k \Gamma_{C_i} \supseteq p \left(\sum_{i=1}^k \Gamma_{T_i} \right) = p\Gamma_T$ as $\Gamma_{C_1 \otimes_K \cdots \otimes_K C_k} = \sum_{i=1}^k \Gamma_{C_i}$ by Prop. 11. Also since ${}^cT_i \in \mathcal{D}_t(F)$ and $\Gamma_{{}^cT_i} \subseteq \Gamma_{T_i}$, by [JW, Cor. 6.7] $\Gamma_{{}^cT} \subseteq \sum_{i=1}^k \Gamma_{{}^cT_i} \subseteq \sum_{i=1}^k \Gamma_{T_i} = \Gamma_T$. Hence $p\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$ as desired.

Since $\Gamma_{{}^cT} \subseteq \sum_{i=1}^k \Gamma_{{}^cT_i}$, $\exp(\Gamma_{{}^cT} / \Gamma_F) \mid \exp \left(\left(\sum_{i=1}^k \Gamma_{{}^cT_i} \right) / \Gamma_F \right)$. But each $\exp(\Gamma_{{}^cT_i} / \Gamma_F) \mid \exp(\Gamma_{T_i} / \Gamma_L)$ by Prop. 12. So $\exp \left(\left(\sum_{i=1}^k \Gamma_{{}^cT_i} \right) / \Gamma_F \right) \mid \exp \left(\left(\sum_{i=1}^k \Gamma_{T_i} \right) / \Gamma_L \right) = \exp(\Gamma_T / \Gamma_L)$. Hence $\exp(\Gamma_{{}^cT} / \Gamma_F) \mid \exp(\Gamma_T / \Gamma_L)$. \square

While $[L : F] \cdot \Gamma_T \subseteq \Gamma_{{}^cT}$ holds for $T \in \mathcal{D}_{ttr}(L)$ when L/F is TRRT, $\exp(\Gamma_L / \Gamma_F) \cdot \Gamma_T \subseteq \Gamma_{{}^cT}$ need not hold, as the following example illustrates:

Example 14. Let p be a prime, and $m \geq k \geq 1$, integers. Let $n = p^m$. Let F_0 be any field with $\mu_{np^2} \subseteq F_0$. Let $F = F_0((x_1)) \cdots ((x_{k+2}))$ be the iterated Laurent power series field, with the usual Henselian valuation $v: F^* \rightarrow \mathbb{Z}^{k+2}$. That is,

$$v \left(\sum_{i_1} \cdots \sum_{i_{k+2}} c_{i_1 \dots i_{k+2}} x_1^{i_1} \cdots x_{k+2}^{i_{k+2}} \right) = \inf \{ (i_1, \dots, i_{k+2}) \mid c_{i_1 \dots i_{k+2}} \neq 0 \},$$

where \mathbb{Z}^{k+2} has the right-to-left lexicographical ordering, i.e.,

$$(i_1, \dots, i_{k+2}) < (j_1, \dots, j_{k+2})$$

if and only if there is a q with $i_q < j_q$ and $i_r = j_r$ for $q < r \leq k+2$ (cf. [Rb, p. 77, Prop. 4 and p. 198, Th. 4]). Let $L = F(\sqrt[k]{x_1}, \dots, \sqrt[k]{x_{k+2}})$. Then L is a TRRT extension of F with $[L: F] = p^{k+2}$.

Now, let $T = (\sqrt[k]{x_1}, \sqrt[k]{x_{k+2}}; L)_n$. Then by [H₂, Prop. 3.1], $T \in \mathcal{D}_{ttr}(L)$ and $\Gamma_T = \left\langle \frac{1}{np}v(x_1), \frac{1}{np}v(x_{k+2}) \right\rangle + \Gamma_L$. Since $T \sim (x_1, x_{k+2}; L)_{np^2}$, by [Ti₂, Th. 3.1] (Projection formula)

$$\text{cor}_{L/F}(T) \sim (x_1^{p^{k+2}}, x_2; F)_{np^2} \sim (x_1, x_2; F)_{n/p^k}$$

as $N_{L/F}(x_1) = x_1^{p^{k+2}}$. Then by [H₂, Prop. 3.1],

$${}^cT \cong (x_1, x_{k+2}; F)_{n/p^k} \in \mathcal{D}_{ttr}(F)$$

and

$$\Gamma_{{}^cT} = \left\langle \frac{p^k}{n}v(x_1), \frac{p^k}{n}v(x_{k+2}) \right\rangle + \Gamma_F.$$

So $p^{k+1}\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$. (Hence $[L: F]\Gamma_T = p^{k+2}\Gamma_T \subseteq \Gamma_{{}^cT}$.) But

$$\exp(\Gamma_L/\Gamma_F)\Gamma_T = p\Gamma_T \not\subseteq \Gamma_{{}^cT}.$$

Note that $\Gamma_T/\Gamma_F \cong (\mathbb{Z}_{np})^2 \times (\mathbb{Z}_p)^k$, and $\Gamma_{{}^cT}/\Gamma_F = p^{k+1}(\Gamma_T/\Gamma_F)$.

We end this section by giving relations between $D \in \mathcal{D}_t(L)$ and ${}^cD \in \mathcal{D}_t(F)$ when $(L, v) \supseteq (F, v)$ is a finite separable TRRT extension of Henselian fields.

Theorem 15. *Let $(L, v) \supseteq (F, v)$ be as above. If $D \in \mathcal{D}_t(L)$, then $[L: F]\Gamma_D \subseteq \Gamma_{{}^cD} \subseteq \Gamma_D$ and $Z(\overline{{}^cD}) \subseteq Z(\overline{D})$.*

Proof. (1) $[L: F]\Gamma_D \subseteq \Gamma_{{}^cD} \subseteq \Gamma_D$: By Prop. 3, in $\text{Br}(L)$, $D \sim S \otimes_L T$ for some $S \in \mathcal{D}_{is}(L)$ and $T \in \mathcal{D}_{ttr}(L)$, and $\Gamma_D = \Gamma_S + \Gamma_T$. Then ${}^cD \sim {}^cS \otimes_F {}^cT$

in $\text{Br}(F)$ where ${}^cS \in \mathcal{D}_{is}(F)$, with $\Gamma_{{}^cS} = [L : F]\Gamma_S + \Gamma_F$ by Th. 8, and ${}^cT \in \mathcal{D}_{tr}(F)$, with $[L : F]\Gamma_T \subseteq \Gamma_{{}^cT} \subseteq \Gamma_T$ by Th. 13 above. So by Prop. 3, $\Gamma_{{}^cD} = \Gamma_{{}^cS} + \Gamma_{{}^cT}$. Therefore, $[L : F]\Gamma_D = [L : F]\Gamma_S + [L : F]\Gamma_T \subseteq \Gamma_{{}^cS} + \Gamma_{{}^cT} = \Gamma_{{}^cD} \subseteq \Gamma_S + \Gamma_T = \Gamma_D$.

(2) $Z(\overline{{}^cD}) \subseteq Z(\overline{D})$: Since $D \in \mathcal{D}_i(L)$, $Z(\overline{D})$ is separable (so abelian Galois) over $\overline{L} = \overline{F}$ by [JW, Lemma 6.1]). Let Z be the inertial lift of $Z(\overline{D})$ over F . Then Z is Galois over F and $L \cap Z = F$ as L/F is TRRT and Z/F is inertial. So $L \otimes_F Z$ is the field $L \cdot Z$. Then by [D₁, p. 56, Ex.1] ${}^cD \otimes_F Z \sim \text{cor}_{LZ/Z}(D \otimes_L LZ)$ in $\text{Br}(Z)$. Let D_{LZ} and $({}^cD)_Z$ be the underlying division algebras of $D \otimes_L LZ$ and ${}^cD \otimes_F Z$, respectively. Since LZ/L is inertial, by [JW, Th. 3.1] $Z(\overline{D_{LZ}}) = Z(\overline{D}) \cdot \overline{LZ} = \overline{Z} \cdot \overline{LZ} = \overline{LZ}$. Then by [JW, Cor. 2.11], in $\text{Br}(LZ)$, $D_{LZ} \sim I \otimes_{LZ} T$ for some $I \in \mathcal{D}_i(LZ)$ and $T \in \mathcal{D}_{tr}(LZ)$. So $({}^cD)_Z \sim {}^cD \otimes_F Z \sim \text{cor}_{LZ/Z}(D_{LZ}) \sim \text{cor}_{LZ/Z}(I) \otimes_Z \text{cor}_{LZ/Z}(T)$.

Let I' and Z' be the underlying division algebras of $\text{cor}_{LZ/Z}(I)$ and $\text{cor}_{LZ/Z}(T)$, respectively. Since LZ/Z is TRRT, by Lemma 6 $I' \in \mathcal{D}_i(Z)$, and by Th. 13 $T' \in \mathcal{D}_{tr}(Z)$. So $Z(\overline{({}^cD)_Z}) = \overline{Z} = Z(\overline{D})$. But since Z/F is inertial, by Th. [JW, Th. 3.1] again $Z(\overline{({}^cD)_Z}) = Z(\overline{{}^cD}) \cdot \overline{Z}$ so we have $Z(\overline{{}^cD}) \cdot Z(\overline{D}) = Z(\overline{D})$, hence $Z(\overline{{}^cD}) \subseteq Z(\overline{D})$. \square

4. The case when L/F is tame.

Suppose $(L, v) \supseteq (F, v)$ is a finite separable extension of Henselian fields such that $\overline{L}/\overline{F}$ is separable and L/K is TRRT where K is the inertial lift of \overline{L} over F in L (i.e. the inertial extension of F with $\overline{K} = \overline{L}$). Then we can combine the previous results with ones of [H₂] to obtain relations between $D \in \mathcal{D}_i(L)$ and ${}^cD \in \mathcal{D}(F)$ since L/K is TRRT and K/F is inertial and ${}^cD \sim \text{cor}_{K/F}(\text{cor}_{L/K}(D))$ in $\text{Br}(F)$. Notably, if L is tame over F (i.e. $\text{char}(\overline{F}) \nmid [L : F]$), then L/F is such an extension: Note that $[L : F] = |\Gamma_L : \Gamma_F| \cdot [\overline{L} : \overline{F}] \cdot q^b$ for some nonnegative integer b , where $q = \text{char}(\overline{F})$ if $\text{char}(\overline{F}) \neq 0$, or $q = 1$ otherwise. (This is proved in [E, 20.21] when L is normal over F . By passing to the normal closure as done in the proof of [M, Cor. 3], this can be proved in general.) Since $\text{char}(\overline{F}) \nmid [L : F]$, necessarily $q^b = 1$, so $[L : F] = |\Gamma_L : \Gamma_F| \cdot [\overline{L} : \overline{F}]$. As $\text{char}(\overline{F}) \nmid [\overline{L} : \overline{F}]$, $\overline{L}/\overline{F}$ is separable. If K is the inertial lift of \overline{L} over F in L , then $[L : K] = [L : F] / [\overline{L} : \overline{F}] = |\Gamma_L : \Gamma_F| = |\Gamma_L : \Gamma_K|$, and $\text{char}(\overline{K}) = \text{char}(\overline{F}) \nmid [L : K]$. So L/K is tame and totally ramified. Since (K, v) is Henselian, L/K is TRRT by [S, p. 64, Th. 3].

Throughout this section, we assume that $(L, v) \supseteq (F, v)$ is a finite separable tame extension of Henselian fields and K is the inertial lift of \overline{L} over F in L . (So L/K is TRRT as we just showed and K/F is inertial.) Also,

for any $D \in \mathcal{D}_t(L)$, let ${}^cD \in \mathcal{D}(F)$ denote the underlying division algebra of $\text{cor}_{L/F}(D)$ as before.

Theorem 16. *Let $T \in \mathcal{D}_{\text{tr}}(L)$ and let T_1 be the underlying division algebra of $\text{cor}_{L/K}(T)$. Let $t = \exp(\Gamma_T/\Gamma_L)$ and $t_1 = \exp(\Gamma_{T_1}/\Gamma_K)$. (So $t_1 \mid t$ by Th. 13.) Let $e = \gcd([\bar{L} : \bar{F}], t)$ and $e_1 = \gcd([\bar{L} : \bar{F}], t_1)$. If $\mu_{t_1} \subseteq F$, then $Z({}^c\bar{T}) \subseteq \bar{F} \left((N_{\bar{L}/\bar{F}}(\bar{L}))^{1/e_1} \right) \subseteq \bar{F} \left((N_{\bar{L}/\bar{F}}(\bar{L}))^{1/e} \right)$, where $N_{\bar{L}/\bar{F}}$ is the norm map from \bar{L} to \bar{F} .*

Proof. Since L/K is TRRT, $T_1 \in \mathcal{D}_{\text{tr}}(K)$ by Th. 13. Also, since cT is the underlying division algebra of $\text{cor}_{K/F}(T_1)$ and K/F is inertial and $\mu_{t_1} \subseteq F$, by [H₂, Th. 3.9], $Z({}^c\bar{T}) \subseteq \bar{F} \left((\bar{N}_{\bar{K}/\bar{F}}(\bar{K}))^{1/e_1} \right) = \bar{F} \left((N_{\bar{L}/\bar{F}}(\bar{L}))^{1/e_1} \right)$ as $[K : F] = [\bar{L} : \bar{F}]$ and $\bar{K} = \bar{L}$.

The second inclusion is clear as $e_1 \mid e$. □

Recall that $\mathcal{N}(M/F)$ denotes the normal closure of M over F where M/F is an algebraic extension of fields.

Theorem 17. *Let L/F be tame.*

- (a) *If $I \in \mathcal{D}_i(L)$, then ${}^cI \in \mathcal{D}_i(F)$ and ${}^c\bar{I} \sim \text{cor}_{\bar{L}/\bar{F}}(\bar{I}^{\otimes |\Gamma_L : \Gamma_F|})$ in $\text{Br}(\bar{F})$.*
- (b) *If $D \in \mathcal{D}_{is}(L)$ then ${}^cD \in \mathcal{D}_{is}(F)$, $\Gamma_{{}^cD} \subseteq |\Gamma_L : \Gamma_F| \cdot \Gamma_D + \Gamma_F$, and $Z({}^c\bar{D}) \subseteq \mathcal{N}(\mathcal{F}(\theta_D(\tilde{\Gamma})) / \bar{F})$, where $\tilde{\Gamma} = \{\alpha + \Gamma_L \in \Gamma_D/\Gamma_L \mid |\Gamma_L : \Gamma_F| \cdot \alpha \in \Gamma_F\}$. (So $\Gamma_{{}^cD} \subseteq \Gamma_D$ and $Z({}^c\bar{D}) \subseteq \mathcal{N}(Z(\bar{D}) / \bar{F})$.)*
- (c) *If $D \in \mathcal{D}_t(L)$, then ${}^cD \in \mathcal{D}_t(F)$ and $\Gamma_{{}^cD} \subseteq \Gamma_D$.*

Proof. (a) Let I_1 be the underlying division algebra of $\text{cor}_{L/K}(I)$. Since $\bar{L} = \bar{K}$, by Lemma 4 $I_1 \in \mathcal{D}_i(K)$ and $\bar{I}_1 \sim \bar{I}^{\otimes |\Gamma_L : \Gamma_F|}$ in $\text{Br}(\bar{K})$ as $[L : K] = |\Gamma_L : \Gamma_F|$. Since ${}^cI \sim \text{cor}_{K/F}(I_1)$ in $\text{Br}(F)$, and K/F is inertial, by [H₂, Th. 2.4] ${}^cI \in \mathcal{D}_i(F)$ and ${}^c\bar{I} \sim \text{cor}_{\bar{K}/\bar{F}}(\bar{I}_1) \sim \text{cor}_{\bar{L}/\bar{F}}(\bar{I}^{\otimes |\Gamma_L : \Gamma_F|})$ in $\text{Br}(\bar{F})$.

(b) Let D_1 be the underlying division algebra of $\text{cor}_{L/K}(D)$. Since $\bar{L} = \bar{K}$, by Th. 8 $D_1 \in \mathcal{D}_{is}(K)$, $\Gamma_{D_1} = [L : K] \cdot \Gamma_D + \Gamma_K = |\Gamma_L : \Gamma_F| \cdot \Gamma_D + \Gamma_F$, and $Z(\bar{D}_1) = \mathcal{F}(\theta_D(\tilde{\Gamma}))$ where $\tilde{\Gamma} = \{\alpha + \Gamma_L \in \Gamma_D/\Gamma_L \mid [L : K] \cdot \alpha \in \Gamma_K\} = \{\alpha + \Gamma_L \in \Gamma_D/\Gamma_L \mid |\Gamma_L : \Gamma_F| \cdot \alpha \in \Gamma_F\}$. Then since K/F is inertial and ${}^cD \sim \text{cor}_{K/F}(D_1)$ in $\text{Br}(F)$, by [H₂, Th. 2.4] ${}^cD \in \mathcal{D}_{is}(F)$, $\Gamma_{{}^cD} \subseteq \Gamma_{D_1}$, and $Z({}^c\bar{D}) \subseteq \mathcal{N}(Z(\bar{D}_1) / \bar{F})$. Therefore, ${}^cD \in \mathcal{D}_{is}(F)$, $\Gamma_{{}^cD} \subseteq |\Gamma_L : \Gamma_F| \cdot \Gamma_D + \Gamma_F$, and $Z({}^c\bar{D}) \subseteq \mathcal{N}(\mathcal{F}(\theta_D(\tilde{\Gamma})) / \bar{F})$. So since $\mathcal{F}(\theta_D(\tilde{\Gamma})) \subseteq Z(\bar{D})$, we have $Z({}^c\bar{D}) \subseteq \mathcal{N}(Z(\bar{D}) / \bar{F})$.

(c) If $D \in \mathcal{D}_t(L)$, then ${}^cD \in \mathcal{D}_t(F)$ by definition of tameness and (b) above. Let D_1 be the underlying division algebra of $\text{cor}_{L/K}(D)$. Since L/K

is TRRT, $\Gamma_{D_1} \subseteq \Gamma_D$ by Th. 15. Since K/F is inertial and ${}^cD \sim \text{cor}_{K/F}(D_1)$ in $\text{Br}(F)$, $\Gamma_{{}^cD} \subseteq \Gamma_{D_1}$ by [H₂, Th.4.5]. So $\Gamma_{{}^cD} \subseteq \Gamma_D$. \square

Theorem 18. *Let L/F be tame and let K be the inertial lift of \bar{L} over F in L . Let $D \in \mathcal{D}_t(L)$ and let D_1 be the underlying division algebra of $\text{cor}_{L/K}(D)$. Let $t_1 = \exp(\ker \theta_{D_1}) = 2^{e_0} p_1^{e_1} \cdots p_r^{e_r}$ where p_i are odd primes, and $e_0 \geq 0$, $e_i > 0$ are integers, $1 \leq i \leq r$. Suppose $\mu_{p_i} \subseteq F$ for $1 \leq i \leq r_0$ and $\mu_{p_i} \not\subseteq F$ for $r_0 + 1 \leq i \leq r$. Let $s = 2^{e_0} p_1^{e_1} \cdots p_{r_0}^{e_{r_0}}$ and $s' = s/2^{e_0}$. Then*

(a) *if $\mu_4 \subseteq F$ or $4 \nmid t_1$, then*

$$Z({}^c\overline{D}) \subseteq \mathcal{N}(Z(\overline{D_1}) / \overline{F})^{1/n} \subseteq \mathcal{N}(Z(\overline{D}) / \overline{F})^{1/n},$$

where $n = (t_1/s) \cdot \gcd([\bar{L} : \bar{F}], s)$;

(b) *if $4 \mid t_1$ and $\mu_4 \not\subseteq F$, then*

$$Z({}^c\overline{D}) \subseteq \mathcal{N}(Z(\overline{D_1}) / \overline{F})^{1/n'} \subseteq \mathcal{N}(Z(\overline{D}) / \overline{F})^{1/n'},$$

where $n' = (t_1/s') \cdot \gcd([\bar{L} : \bar{F}], s')$. So in either case,

$$Z({}^c\overline{D}) \subseteq \mathcal{N}(Z(\overline{D_1}) / \overline{F})^{1/t_1} \subseteq \mathcal{N}(Z(\overline{D}) / \overline{F})^{1/t}$$

where $t = \exp(\ker \theta_D)$.

Proof. Since L/K is TRRT and $D_1 \sim \text{cor}_{L/K}(D)$, $D_1 \in \mathcal{D}_t(K)$ and $Z(\overline{D_1}) \subseteq Z(\overline{D})$ by Th. 8 and Th. 15. Also, since K/F is inertial and ${}^cD \sim \text{cor}_{K/F}(D_1)$, by [H₂, Th. 4.6] we have (a) and (b).

Since $n \mid t_1$ and $n' \mid t_1$, $Z({}^c\overline{D}) \subseteq \mathcal{N}(Z(\overline{D_1}) / \overline{F})^{1/t_1}$. By Prop. 3, we have $D \sim S \otimes_L T$ for some $S \in \mathcal{D}_{is}(L)$ and $T \in \mathcal{D}_{ttr}(L)$, and $\Gamma_T / \Gamma_L = \ker(\theta_D)$. So $t = \exp(\Gamma_T / \Gamma_L)$. Let S_1 and T_1 be the underlying division algebras of $\text{cor}_{L/K}(S)$ and $\text{cor}_{L/K}(T)$, respectively. Since L/K is TRRT, $D_1 \sim S_1 \otimes_K T_1$ where $S_1 \in \mathcal{D}_{is}(K)$ and $T_1 \in \mathcal{D}_{ttr}(K)$ and $\exp(\Gamma_{T_1} / \Gamma_K) \mid \exp(\Gamma_T / \Gamma_L)$ by Th. 8 and Th. 13. So by Prop. 3, $\ker(\theta_{D_1}) = \Gamma_{T_1} / \Gamma_K$, hence $t_1 = \exp(\Gamma_{T_1} / \Gamma_K) \mid \exp(\Gamma_T / \Gamma_L) = t$. So we have $\mathcal{N}(Z(\overline{D_1}) / \overline{F})^{1/t_1} \subseteq \mathcal{N}(Z(\overline{D}) / \overline{F})^{1/t_1} \subseteq \mathcal{N}(Z(\overline{D}) / \overline{F})^{1/t}$ as $Z(\overline{D_1}) \subseteq Z(\overline{D})$. \square

Remark. (The corestriction of central simple algebras with Dubrovin valuation rings.) There are generalizations of Theorems 17 and 18 above for central simple algebras S over a valued field (L, v) where v is not Henselian. We describe the generalizations here in Th. 20, but omit proofs, which can be found in [H₁, Chap. 5]. When v is not Henselian, it may not extend to a valuation on S , but there is always a unique (up to isomorphism) Dubrovin valuation ring B of S extending the valuation ring V of v on F , and B has

a value group Γ_B and a residue central simple algebra $\overline{B} = B/J(B)$, where $J(B)$ is the Jacobson radical of B (cf. [W₂]). The following proposition is used in proving the generalizations of Th. 17 and 18.

Proposition 19 [H₁, Th. 5.4]. *Let (F, v) be a Henselian field. If $D_i \in \mathcal{D}_t(F)$ for $1 \leq i \leq n$, and D is the underlying division algebra of $D_1 \otimes_F \cdots \otimes_F D_n$, then $Z(\overline{D}) \subseteq Z(\overline{D}_1)^{1/t_1} \cdots Z(\overline{D}_n)^{1/t_n}$ where $t_i = \exp(\ker(\theta_{D_i}))$.*

To state Th. 20, we introduce the following notation:

Let L be a finite separable extension of a field with valuation ring (F, V) and let W_1, \dots, W_k be all the valuation rings of L extending V . Let $L_i = (L, W_i)$, $(F^h, V^h) =$ the Henselization of (F, V) and $(L_i^h, W_i^h) =$ the Henselization of (L, W_i) for $1 \leq i \leq k$. Let S be a central simple L -algebra and let ${}^{\text{cor}}S = \text{cor}_{L/F}(S)$, the corestriction of S . Let A be a Dubrovin valuation ring of ${}^{\text{cor}}S$ with $A \cap F = V$ and let B_i be a Dubrovin valuation ring of S with $B_i \cap L = W_i$. Set $S_i = (S, B_i)$. Let S_i^h be the underlying division algebra of $S_i \otimes_L L_i^h$ and let $({}^{\text{cor}}S)^h$ be the underlying division algebra of ${}^{\text{cor}}S \otimes_F F^h$. Since $\Gamma_{B_i} = \Gamma_{S_i^h}$, $\Gamma_A = \Gamma_{({}^{\text{cor}}S)^h}$, $Z(\overline{B_i}) = Z(\overline{S_i^h})$ and $Z(\overline{A}) = Z(\overline{({}^{\text{cor}}S)^h})$ by [W₂, Th. B], we can obtain information about A by applying Th. 17 and 18 to the Henselizations. Thereby, we obtain the following theorem.

Theorem 20 [H₁, Th. 5.15]. *Assume all L_i^h/F^h are tame for $1 \leq i \leq k$.*

(1) *If S_i^h is inertially split over L_i^h for each i , $1 \leq i \leq k$, then $({}^{\text{cor}}S)^h$ is inertially split over F^h , $\Gamma_A \subseteq \sum_{i=1}^k \Gamma_{B_i}$ and*

$$Z(\overline{A}) \subseteq \mathcal{N} \left(\prod_{i=1}^k Z(\overline{B_i}) / \overline{F} \right).$$

(2) *If S_i^h is tame over L_i^h for each i , $1 \leq i \leq k$, then $({}^{\text{cor}}S)^h$ is tame over F^h and $\Gamma_A \subseteq \sum_{i=1}^k \Gamma_{B_i}$. Further, if $t_i = \exp(\ker(\theta_{S_i^h}))$ for $1 \leq i \leq k$, then $Z(\overline{A}) \subseteq \prod_{i=1}^k \mathcal{N} \left(Z(\overline{B_i}) / \overline{F} \right)^{1/m_i}$ where $m_i = t_i$ if $4 \nmid t_i$ or $\mu_4 \subseteq F$, or $m_i = 2t_i$ if $4 \mid t_i$ and $\mu_4 \not\subseteq F$. (The condition that S_i^h is inertially split (or tame) over L_i^h can be expressed in terms of B_i itself. See [H₁, Chap. 5, Sec. 2] for details.)*

References

- [Az] G. Azumaya, *On maximally central algebras*, Nagoya J. Math., **2** (1951), 119-150.
- [D₁] P. Draxl, *Skew Fields*, London Math. Soc. Lecture Note Series, Vol. 81, Cambridge Univ. Press, Cambridge, 1983.

- [D₂] P. Draxl, *Ostrowski's theorem for Henselian valued skew fields*, J. Reine Angew. Math., **354** (1984), 213-218.
- [E] O. Endler, *Valuation Theory*, Springer-Verlag, New York, 1972.
- [H₁] Y. Hwang, *The corestriction of valued division algebras over Henselian Fields*, Doctoral Dissertation, University of California, San Diego, 1992.
- [H₂] Y. Hwang, *The corestriction of valued division algebras over Henselian Fields I*, Pacific J. Math., **170** (1995), 53-81.
- [J] W. Jehne, *Separable adel algebras and a presentation of Weil groups*, J. Reine Angew. Math., **375/376** (1987), 211-237.
- [JW] B. Jacob and A. Wadsworth, *Division algebras over Henselian fields*, J. Algebra, **128** (1990), 126-179.
- [KY] V. V. Kursov and V. I. Yanchevskii, *Crossed products over simple algebras and their automorphism groups*, Akad. Nauk Belo. Inst. Math., preprint #35 (305), Minsk, No. 1987 [Russian].
- [M] P. Morandi, *The Henselization of a valued division algebra*, J. Algebra, **122** (1989), 232-243.
- [Rb] P. Ribenboim, *Théorie des Valuations*, Presses Univ. Montréal, Montréal, 1968.
- [S] O.F.G. Schilling, *The Theory of Valuations*, Math Surveys no. 4, Amer. Math. Soc., Providence, R.I., 1950.
- [Ti₁] J.-P. Tignol, *Generalized crossed products*, in *Seminaire Mathématique (nouvelle série)*, No. 106, Université Catholique de Louvain, Louvain-Neuve, Belgium, 1987.
- [Ti₂] J.-P. Tignol, *On the corestriction of central simple algebras*, Math. Zeitschrift, **194** (1987), 267-274.
- [TW] J.-P. Tignol and A. Wadsworth, *Totally ramified valuations on finite dimensional division algebras*, Trans. Amer. Math. Soc., **302** (1987), 223-250.
- [W₁] A. Wadsworth, *Extending valuations to finite dimensional division algebras*, Proc. Amer. Math. Soc., **98** (1986), 20-22.
- [W₂] A. Wadsworth, *Dubrovin valuation rings and Henselization*, Math. Ann., **283** (1989), 301-328.
- [We] E. Weiss, *Cohomology of Groups*, Academic Press, New York, 1969.

Received November 10, 1992 and revised April 8, 1993.

DEPARTMENT OF MATHEMATICS
 KOREA UNIVERSITY
 ANAM-DONG 5-1, SUNGBUK-KU
 SEOUL 136-701, KOREA
 E-MAIL ADDRESS: YHWANG@SEMI.KOREA.AC.KR

PACIFIC JOURNAL OF MATHEMATICS

Founded by E. F. Beckenbach (1906-1982) and F. Wolf (1904-1989)

EDITORS

Sun-Yung Alice Chang (Managing Editor)
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

F. Michael Christ
University of California
Los Angeles, CA 90095-1555
christ@math.ucla.edu

Thomas Enright
University of California
San Diego, La Jolla, CA 92093
tenright@ucsd.edu

Nicholas Ercolani
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

Robert Finn
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

Vaughan F. R. Jones
University of California
Berkeley, CA 94720
vfr@math.berkeley.edu

Steven Kerckhoff
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

Martin Scharlemann
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

Gang Tian
Courant Institute
New York University
New York, NY 10012-1100
tiang@taotao.cims.nyu.edu

V. S. Varadarajan
University of California
Los Angeles, CA 90095-1555
vsv@math.ucla.edu

SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
STANFORD UNIVERSITY
UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
UNIVERSITY OF UTAH
UNIVERSITY OF WASHINGTON
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$215.00 a year (10 issues). Special rate: \$108.00 a year to individual members of supporting institutions. Subscriptions, orders for back issues published within the last three years, and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,
the American Mathematical Society's TEX macro system.
Copyright © 1995 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 170 No. 1 September 1995

Generalized generalized spin models (four-weight spin models)	1
EIICHI BANNAI and ETSUKO BANNAI	
Fine structure of the Mackey machine for actions of abelian groups with constant Mackey obstruction	17
SIEGFRIED ECHTERHOFF and JONATHAN ROSENBERG	
The corestriction of valued division algebras over Henselian fields. I	53
YOON SUNG HWANG	
The corestriction of valued division algebras over Henselian fields. II	83
YOON SUNG HWANG	
The cohomology of expansive \mathbb{Z}^d -actions by automorphisms of compact, abelian groups	105
ANATOLE KATOK and KLAUS SCHMIDT	
The Anosov theorem for exponential solvmanifolds	143
EDWARD KEPPELMANN and CHRISTOPHER K. MCCORD	
Projections of measures on nilpotent orbits and asymptotic multiplicities of K -types in rings of regular functions. I	161
DONALD RAYMOND KING	
On almost-everywhere convergence of inverse spherical transforms	203
CHRISTOPHER MEANEY and ELENA PRESTINI	
Characters of supercuspidal representations of $SL(n)$	217
FIONA ANNE MURNAGHAN	
The cohomology of higher-dimensional shifts of finite type	237
KLAUS SCHMIDT	
On Gorenstein surface singularities with fundamental genus $p_f \geq 2$ which satisfy some minimality conditions	271
TADASHI TOMARU	