

*Pacific
Journal of
Mathematics*

**ON THE ODD PRIMARY COHOMOLOGY OF HIGHER
PROJECTIVE PLANES**

MARK FOSKEY AND MICHAEL DAVID SLACK

ON THE ODD PRIMARY COHOMOLOGY OF HIGHER PROJECTIVE PLANES

MARK FOSKEY AND MICHAEL SLACK

Let X be an n -fold loop space. Working with an auxiliary space $P_p^n X$ analogous to the projective plane $P_2 X$, we show that the existence of certain Steenrod connections in $H^*(P_p^n X; \mathbf{F}_p)$ (p odd) implies the vanishing of certain corresponding Dyer-Lashof operations in $H^*(X; \mathbf{F}_p)$, and vice versa.

1. Introduction.

A useful property of the projective plane of an H-space is that the vanishing of certain cup products in $H^*(P_2 X)$ implies the existence of certain related nonzero reduced coproducts in $H^*(X)$ (see [BT]). In [KSW], Kuhn, Slack, and Williams introduce the necessary theory to construct a space $P_p^n X$ with the property that the Steenrod action on its cohomology bears a similar relation to the Dyer-Lashof action on the homology of X . We develop this relation at odd primes in this paper. In particular, our main theorem can be summarized as follows:

Main Theorem. *Suppose X is an n -fold loop space, $1 \leq n \leq \infty$. Then there is a certain cofibration sequence*

$$\Sigma^n \tilde{C}_{n,p} X \xrightarrow{h} \Sigma^n X \xrightarrow{i} P_p^n X \xrightarrow{j} \Sigma^{n+1} \tilde{C}_{n,p} X$$

such that

1. For each $x \in \tilde{H}^q(X)$ there exist elements $\tilde{Q}_r x \in \tilde{H}^{pq+r}(\tilde{C}_{n,p} X)$ such that $Q_r^* w = x$ whenever $h^*(\sigma^n w) = \sigma^n \tilde{Q}_r x$, where σ is the suspension isomorphism and

$$Q_r^*: \tilde{H}^{pq+r}(X) \rightarrow H^q(X)$$

is an element of the opposite algebra to the Dyer-Lashof algebra.

2. If $\bar{x} \in \tilde{H}^{2s+1}(P_p^n X)$ and $i^*(\bar{x}) = \sigma^n x$ then

$$\begin{aligned} \mathcal{P}^s \bar{x} &= u \cdot j^* \left(\sigma^{n+1} \tilde{Q}_{(n-1)(p-1)-1} x \right), \\ \beta \mathcal{P}^s \bar{x} &= u \cdot j^* \left(\sigma^{n+1} \tilde{Q}_{(n-1)(p-1)} x \right), \end{aligned}$$

where u is some undetermined unit in \mathbf{F}_p .

The analogous result at the prime 2 was proved in [KSW].

Using this theorem and the long exact sequence of a cofibration it is easy to see how the triviality of certain Steenrod operations in $\tilde{H}^*(P_p^n X)$ can imply the existence of nontrivial Dyer-Lashof operations in $\tilde{H}_*(X)$, and vice versa. In [SI], the second author exploits this relation (in the stable case, see Corollary 4.2) to prove the following result.

Theorem (Slack). *If X is a connected infinite loop space (of finite type), and all of the Dyer-Lashof operations are trivial on the mod p homology of X , then X is mod p homotopy equivalent to a product of Eilenberg-MacLane spaces.*

The definitions of the spaces $P_p^n X$ and $\tilde{C}_{n,p} X$ are briefly summarized in Section 2. In Section 3 we define the elements $\tilde{Q}_{r,x}$ and show how they relate to the homology Dyer-Lashof operations, and in Section 4 we prove Part 2 of the Main Theorem. We conclude with an appendix giving the Nishida relations as they apply to the external operations \tilde{Q}_r , which are useful in applications.

We summarize here some of the notational conventions used in this paper. Let p be an odd prime, and take all homology and cohomology with coefficients in the field \mathbf{F}_p of p elements. Recall ([CLM]) that the homology of an n -fold loop space X admits certain Dyer-Lashof operations $Q_r: \tilde{H}_q(X) \rightarrow \tilde{H}_{p^q+r}(X)$. (We are using the lower notation of [CPS] in which $Q_{s(p-1)}v$ is defined to be $Q^{(s+\deg v)/2}v$.) In this paper, all spaces will be compactly generated Hausdorff spaces with non-degenerate basepoint $*$, and all maps will be based maps.

We would like to thank Nicholas Kuhn, Jim Lin, and Fred Cohen for many helpful notes and conversations.

2. The construction of the projective planes.

In this section we recall relevant information from [KSW]. Let $C_n X$ denote the standard approximation to $\Omega^n \Sigma^n X$ of [Ma2, Mi]. It admits a filtration

$$X \simeq C_{n,1} X \subset C_{n,2} X \subset \cdots \subset C_n X.$$

If there is a map $\theta_n^X: C_{n,p} X \rightarrow X$ extending the identity on X , then X is called an H_p^n -space, and there is a category \mathcal{H}_p^n for which the objects are H_p^n -spaces and the morphisms can be thought of as homotopy classes of maps preserving the H_p^n -structure.¹ From now on X will generally be assumed

¹Actually, they only preserve the H_p^n -structure up to homotopy, but they come equipped with prescribed classes of homotopies making the appropriate diagram commute.

to be an H_p^n -space, and, in the main theorem, the hypothesis that X is an n -fold loop space may be replaced by the somewhat weaker hypothesis that $X \in \text{Ob } \mathcal{H}_p^n$.

There is a functor $\Omega^n: \mathcal{H}_p^0 \rightarrow \mathcal{H}_p^n$ that, on objects, is the usual n -fold loop space functor. It admits a left adjoint $P_p^n: \mathcal{H}_p^n \rightarrow \mathcal{H}_p^0$ that is constructed (on objects) as follows. If X is an H_p^n -space, there are two natural retractions $\Sigma^n C_{n,p} X \rightarrow \Sigma^n X$; one is $\Sigma^n \theta_n^X$ and the other is the adjoint ε_n of the composition

$$C_{n,p} X \hookrightarrow C_n X \longrightarrow \Omega^n \Sigma^n X.$$

If $\tilde{C}_{n,p} X$ is the cofiber of the inclusion $X \hookrightarrow C_{n,p} X$, then there is a map $h: \Sigma^n \tilde{C}_{n,p} X \rightarrow \Sigma^n X$, making the diagram

$$\begin{array}{ccccc} \Sigma^n X & \longrightarrow & \Sigma^n C_{n,p} X & \longrightarrow & \Sigma^n \tilde{C}_{n,p} X \\ \downarrow & & \downarrow \Sigma^n \theta_n - \varepsilon_n & & \downarrow h \\ * & \longrightarrow & \Sigma^n X & \xlongequal{\quad} & \Sigma^n X \end{array}$$

homotopy commute; h is unique up to homotopy by the cofiber mapping sequence. The higher projective plane $P_p^n X$ is defined to be the cofiber of h , and we thus have the cofibration sequence

$$\Sigma^n \tilde{C}_{n,p} X \xrightarrow{h} \Sigma^n X \xrightarrow{i} P_p^n X \xrightarrow{j} \Sigma^{n+1} \tilde{C}_{n,p} X$$

of the Main Theorem.

By way of comparison, the ordinary projective plane $P_2 X$ can be defined as the cofiber of the map h determined by the commutative diagram

$$\begin{array}{ccccc} \Sigma(X \vee X) & \longrightarrow & \Sigma(X \times X) & \longrightarrow & \Sigma X \wedge X \\ \downarrow & & \downarrow -\Sigma\pi_1 + \Sigma\mu - \Sigma\pi_2 & & \downarrow h \\ * & \longrightarrow & \Sigma X & \xlongequal{\quad} & \Sigma^n X. \end{array}$$

3. The external operations.

We now start to define the external operations \tilde{Q}_r . We will begin by using the calculation of $H_*(C_n X)$ in [CLM] to observe that $H_*(\tilde{C}_{n,p} X)$ can be viewed as a particular direct summand of $H_*(C_n X)$. We then use this information to produce a specific basis for $H_*(\tilde{C}_{\infty,p} K(\mathbf{Z}/p, s + q - \gamma))$, in terms of which we define a map

$$G: \tilde{C}_{\infty,p} K(\mathbf{Z}/p, s + q - \gamma) \longrightarrow K(\mathbf{Z}/p, p(s + q) - 2\gamma).$$

(The roles of s , q , and γ , will be explained shortly.) Then, after recalling another map

$$\tilde{\varepsilon}_k: \Sigma^k \tilde{C}_{n+k,p} X \rightarrow \tilde{C}_{n,p} \Sigma^k X,$$

we define the external operations (Definition 3.3) in terms of $\tilde{\varepsilon}_k$, G , and a map f representing $x \in H^*(X)$. We end the section by demonstrating some of their properties.

In this section, r will denote $s(p-1) - \gamma$, where $\gamma \in \{0, 1\}$, so that $Q_r: \tilde{H}_q(X) \rightarrow \tilde{H}_{pq+r}(X)$ will be defined on an H_p^n -space provided $s < n$ and $q + s$ is even. Recall ([CLM, III]) that, along with the Dyer-Lashof operations Q_r , the homology of $C_n X$ admits Browder operations

$$\lambda_{n-1}: \tilde{H}_q(C_n X) \otimes \tilde{H}_m(C_n X) \longrightarrow \tilde{H}_{q+m+n-1}(C_n X).$$

We will say that a *Browder product of weight l* is a composition of Browder operations in which l (not necessarily distinct) variables appear; for instance, $\lambda_{n-1}(x, \lambda_{n-1}(x, y))$ is a Browder product of weight 3. We recall that the Browder operations are neither associative nor commutative.

Let $\eta = \eta_1: X \rightarrow C_n X$ and $\eta_k: C_{n,k} X \rightarrow C_n X$ be the respective inclusions. We then have the following proposition, which summarizes information contained in Cohen's proof of the structure theorem for $\tilde{H}_*(C_n X)$ ([CLM, III, Theorem 3.1]).

Proposition 3.1. *The map $(\eta_k)_*: \tilde{H}_*(C_{n,k} X) \rightarrow \tilde{H}_*(C_n X)$ is one-to-one, and the filtration on homology defined by $F_k \tilde{H}_*(C_n X) = \text{im}(\eta_k)_*$ is identical to the following algebraic filtration: If λ is a Browder product of weight l applied to elements of $\eta_*(\tilde{H}_*(X))$, then the filtration of $Q_{r_1} \cdots Q_{r_t} \lambda$ is $p^t l$, and, if $a \in F_j \tilde{H}_*(C_n X)$ and $b \in F_k \tilde{H}_*(C_n X)$ then $a * b \in F_{j+k} \tilde{H}_*(C_n X)$, where $a * b$ is the Pontryagin product of a and b .*

From this, along with the structure theorem for $\tilde{H}_*(C_n X)$, it follows that $\text{im}(\eta_p)_* \subset \tilde{H}_*(C_n X)$ can be written as $A \oplus B \oplus C$, where A , B , and C are defined as follows.

$$\begin{aligned} A &= \eta_* \tilde{H}_*(X), \\ B &= \{Q_{s(p-1)-\gamma} \eta_*(x) \mid s > 0, (s + \deg x) \text{ is even}\}, \end{aligned}$$

and C is the set $\{\eta_*(x_1) * \cdots * \eta_*(x_k) * \lambda_1 * \cdots * \lambda_t\}$, where $x_1, \dots, x_k \in \tilde{H}_*(X)$, $\lambda_1, \dots, \lambda_t$ are Browder products of weight l_1, \dots, l_t respectively, and $2 \leq k + l_1 + \cdots + l_t \leq p$. Since η_* and $(\eta_p)_*$ are one-to-one, we can by abuse of notation consider $B \oplus C$ as a decomposition of $\tilde{H}_*(\tilde{C}_{n,p} X)$. We can then make

Definition 3.2. If $v \in \tilde{H}_q(X)$ and $q + s$ is even then the external homology Dyer-Lashof operation $\tilde{Q}_{s(p-1)-\gamma} v \in \tilde{H}_*(\tilde{C}_{n,p} X) \subset \tilde{H}_*(C_{n,p} X)$ is defined as $Q_{s(p-1)-\gamma} \eta_*(v)$.

If $(X, \theta_n^X) \in \text{Ob } \mathcal{H}_p^n$, then the structure map θ_n^X characterizes the internal Dyer-Lashof operations in $\tilde{H}_*(X)$ by the formula

$$Q_r v = (\theta_n^X)_*(\overline{Q}_r v).$$

We also have

$$h_*(\sigma^n \overline{Q}_r v) = \sigma^n Q_r v.$$

To see this, first note that $(\varepsilon_n)_*(\overline{Q}_r v) = 0$ for otherwise the formula $(\varepsilon_n)_*(\overline{Q}_r v)$ would define a natural positive-dimensional homology operation on an arbitrary space, which is impossible. The formula then follows by a simple diagram chase on the diagram from Section 2 defining h .

Now we come to the maps G and $\tilde{\varepsilon}_k$. From the point of view of the theory of spectra, one way to think about G is that, when $\gamma = 0$, the existence of the G we want is essentially equivalent to the fact that the mod p Eilenberg-MacLane spectrum admits the structure of an H_∞^2 ring spectrum (as defined by Bruner et. al. [BMMS]). In [KSW], the analogous key fact is the existence of an H_∞^1 ring spectrum structure on the mod 2 Eilenberg-MacLane spectrum. In fact, using the methods of this paper and of [KSW] it should be possible to generalize our main theorem to a theorem on arbitrary H_∞^d ring spectrums E , which would relate power operations in the E^* cohomology of a generalized projective plane to power operations in the E_* homology of the original space.

For our purposes, it is convenient to have an explicit description of G . The main technical difficulty in the odd primary case which does not cause a problem in the 2 primary case in [KSW] is the fact that $\tilde{C}_{\infty,p} X$ is not homotopy equivalent to the p -adic construction, $D_p X$, when p is odd (stably it contains $D_p X$ as a wedge summand), whereas it is when $p = 2$. This leads to many possible choices for the definition of G in the odd primary case, and it is important to carefully define the ‘‘correct’’ one.

To simplify notation let $K = K(\mathbf{Z}/p, s + q - \gamma)$. Then the fundamental class $\iota = \iota_{s+q-\gamma}$, its images under the Steenrod algebra, and their respective products form a standard basis $\mathcal{B}\tilde{H}^*(K)$ for $\tilde{H}^*(K)$; let its dual basis be $\mathcal{B}\tilde{H}_*(K)$, and let the element of $\mathcal{B}\tilde{H}_*(K)$ dual to ι be ν .

By Theorem 3.1, $\tilde{H}_*(\tilde{C}_{\infty,p} K) \cong F_p \tilde{H}_*(C_\infty K) / F_1 \tilde{H}_*(C_\infty K)$. If we then apply the structure theorem for $\tilde{H}_*(C_\infty X)$ ([CLM, I, 4.1]), it is not hard to show that one gets a basis for $\tilde{H}_*(\tilde{C}_{\infty,p} K)$ by totally ordering the elements of $\mathcal{B}\tilde{H}_*(K)$ and then defining the set $\mathcal{B}\tilde{H}_*(\tilde{C}_{\infty,p} K)$ to be $A \cup B$, where

$$\begin{aligned} A &= \{\eta_*(v_1) * \cdots * \eta_*(v_k) | v_1, \dots, \\ &\quad v_k \in \mathcal{B}\tilde{H}_*(K), 2 \leq k \leq p, v_1 \leq v_2 \leq \cdots \leq v_k\}, \\ B &= \{\overline{Q}_{s(p-1)-\gamma} v | v \in \mathcal{B}\tilde{H}_*(K), (s + \deg v) \text{ even}, s > 0, \gamma = 0 \text{ or } 1\}. \end{aligned}$$

We emphasize that the notation $\eta_*(v_1) * \cdots * \eta_*(v_k)$ for an element of $\tilde{H}_*(\tilde{C}_{\infty,p}K)$ only makes sense via the isomorphism with

$$F_p \tilde{H}_*(C_\infty K) / F_1 \tilde{H}_*(C_\infty K),$$

since $\tilde{C}_{\infty,p}K$ itself is not an H-space and thus admits no Pontryagin products.

We now let $\mathcal{B}\tilde{H}^*(\tilde{C}_{\infty,p}K)$ be the dual basis to $\mathcal{B}\tilde{H}_*(\tilde{C}_{\infty,p}K)$, and, for each v in $\mathcal{B}\tilde{H}_*(\tilde{C}_{\infty,p}K)$, we denote the dual basis element by $(v)^{\text{dual}}$. If $\gamma = 0$ we then define $G: \tilde{C}_{\infty,p}K(\mathbf{Z}/p, s+q) \rightarrow K(\mathbf{Z}/p, p(s+q))$ to be the map representing the cohomology class

$$\sum (\eta_*(\nu^{t_1}) * \cdots * \eta_*(\nu^{t_k}))^{\text{dual}}$$

where the sum runs over sequences (t_1, \dots, t_k) such that

1. $k \leq p$,
2. $t_1 + \cdots + t_k = p$, and
3. $\nu^{t_1} \leq \nu^{t_2} \leq \cdots \leq \nu^{t_k}$ in the ordering on $\mathcal{B}\tilde{H}_*(K)$.

This definition makes sense since we have required that $s+q$ be even.

If $\gamma = 1$ then we define $G: \tilde{C}_{\infty,p}K(\mathbf{Z}/p, s+q-1) \rightarrow K(\mathbf{Z}/p, p(s+q)-2)$ to be the map representing the cohomology class

$$(\overline{Q}_{p-2}\nu)^{\text{dual}}.$$

Now we consider the map $\tilde{\varepsilon}_k$. It is constructed in [KS \mathbf{W}], and it fits into a commutative diagram

$$\begin{array}{ccccc} \Sigma^k \tilde{C}_{n+k,p}X & \longleftarrow & \Sigma^k C_{n+k,p}X & \longrightarrow & \Sigma^k \Omega^{n+k} \Sigma^{n+k} X \\ \downarrow \tilde{\varepsilon}_k & & \downarrow \varepsilon_k & & \downarrow \varepsilon \\ \tilde{C}_{n,p} \Sigma^k X & \longleftarrow & C_{n,p} \Sigma^k X & \longrightarrow & \Omega^n \Sigma^{n+k} X \end{array}$$

where ε_k is a map constructed in [Ma $\mathbf{2}$] and $\varepsilon: \Sigma^k \Omega^k Y \rightarrow Y$ is the evaluation map. If X is connected, so that $C_n X \simeq \Omega^n \Sigma^n X$, then ε_k may be seen as “filtering” ε .

One important fact about $\tilde{\varepsilon}_k$ is that it commutes with the (homology) Dyer-Lashof operations in the sense that

$$(\tilde{\varepsilon}_k)_* \left(\sigma^k \overline{Q}_{s(p-1)-\gamma} v \right) = \begin{cases} \overline{Q}_{(s-k)(p-1)-\gamma} \sigma^k v & \text{if } s-k > \gamma; \\ 0 & \text{otherwise.} \end{cases}$$

This follows from the above commutative diagram combined with the standard fact that the Dyer-Lashof operations commute with the loop homomorphism induced by ε (see, e.g., [CLM, III, Theorem 1.4]). We also note that $\tilde{\varepsilon}_k$ is still defined when $n = \infty$.

We now define the external cohomology operations.

Definition 3.3. Suppose γ is an element of the set $\{0, 1\}$, and $s \geq \gamma$ and $q \geq 0$ are integers such that $s + q$ is even. Let $r = s(p - 1) - \gamma$. For every $1 \leq n \leq \infty$ and any space X , let

$$\tilde{Q}_r: \tilde{H}^q(X) \longrightarrow \tilde{H}^{pq+r}(\tilde{C}_{n,p}X)$$

be defined as follows. If $x \in \tilde{H}^q(X)$ is represented by $f: X \rightarrow K(\mathbf{Z}/p, q)$, let $\sigma^{s-\gamma}(\tilde{Q}_r x)$ be represented by the composite

$$\begin{aligned} \Sigma^{s-\gamma} \tilde{C}_{n,p}X &\rightarrow \Sigma^{s-\gamma} \tilde{C}_{\infty,p}X \xrightarrow{\tilde{\varepsilon}_{s-\gamma}} \tilde{C}_{\infty,p} \Sigma^{s-\gamma} X \xrightarrow{\tilde{C}_{\infty,p} \Sigma^{s-\gamma} f} \tilde{C}_{\infty,p} \Sigma^{s-\gamma} K(\mathbf{Z}/p, q) \\ &\xrightarrow{\tilde{C}_{\infty,p} \epsilon} \tilde{C}_{\infty,p} K(\mathbf{Z}/p, s+q-\gamma) \xrightarrow{G} K(\mathbf{Z}/p, p(s+q)-2\gamma). \end{aligned}$$

The element $\tilde{Q}_r x$ for $r > 0$ is sometimes denoted $e_r \otimes x^p$ or $e_r \int x$ in the literature (the way our operations are defined, $\tilde{Q}_0 x \neq e_0 \int x$). We remark for future reference that when $s \geq n$ above, the operation \tilde{Q}_r is trivial.

The next proposition relates our external cohomology operations \tilde{Q}_r to the external homology operations, via the Kronecker pairing $\langle \cdot, \cdot \rangle$. This will make it reasonable to see \tilde{Q}_r as a sort of “dual external Dyer-Lashof operation”.

Proposition 3.4. *Suppose $x \in \tilde{H}^q(X)$ and $v \in \tilde{H}_q(X)$. Then*

$$\langle \tilde{Q}_r x, \overline{Q}_{r'} v \rangle = \begin{cases} \langle x, v \rangle & \text{if } r = r'; \\ 0 & \text{if } r \neq r'. \end{cases}$$

Furthermore, if $r = s(p - 1) - \gamma > 0$, then the element $\tilde{Q}_r x$ paired with any Browder or Pontryagin product operation yields zero.

Proof. If $r > 0$ we assume by naturality that $n = \infty$ and make the following calculation. By the definition of $\tilde{Q}_r x$,

$$\begin{aligned} \langle \tilde{Q}_r x, \overline{Q}_{r'} v \rangle &= \langle \sigma^{s-\gamma} \tilde{Q}_{s(p-1)-\gamma} x, \sigma^{s-\gamma} \overline{Q}_{s'(p-1)-\gamma'} v \rangle \\ &= \langle \tilde{\varepsilon}_{s-\gamma}^* (\tilde{C}_{\infty,p} \Sigma^{s-\gamma} f)^* (\tilde{C}_{\infty,p} \epsilon)^* [G], \sigma^{s-\gamma} \overline{Q}_{s'(p-1)-\gamma'} v \rangle \\ &= \langle [G], (\tilde{C}_{\infty,p} \epsilon)_* (\tilde{C}_{\infty,p} \Sigma^{s-\gamma} f)_* (\tilde{\varepsilon}_{s-\gamma})_* \sigma^{s-\gamma} \overline{Q}_{s'(p-1)-\gamma'} v \rangle. \end{aligned}$$

Now, we have already observed how $(\tilde{\varepsilon}_{s-\gamma})_*$ commutes with the $\overline{Q}_{r'}$. In addition, for any map $g: X \rightarrow Y$, we have $(C_{\infty g})_*(Q_r \eta_* v) = Q_r g_*(\eta_* v)$ [CLM], and hence

$$(\tilde{C}_{\infty,p} g)_*(\overline{Q}_{r'} v) = \overline{Q}_{r'} g_*(v).$$

Thus,

$$\begin{aligned} \langle [G], (\tilde{C}_{\infty,p}\epsilon)_*(\tilde{C}_{\infty,p}\Sigma^{s-\gamma}f)_*(\tilde{\epsilon}_{s-\gamma})_*\sigma^{s-\gamma}\overline{Q}_{s'(p-1)-\gamma'}v \rangle \\ = \langle [G], \overline{Q}_{(s'-s+\gamma)(p-1)-\gamma'}\epsilon_*\sigma^{s-\gamma}f_*(v) \rangle. \end{aligned}$$

Here we let $\overline{Q}_r x = 0$ if $r < 0$. Now, by the construction of G , the last quantity can be nonzero only if $\overline{Q}_{(s'-s+\gamma)(p-1)-\gamma'}\epsilon_*\sigma^{s-\gamma}f_*(v)$ equals either $u \cdot \overline{Q}_0\nu_{s+q}$ or $u \cdot \overline{Q}_{p-2}\nu_{s+q-1}$, where $u \in \mathbf{F}_p$ is some unit. In the former case, $\gamma = \gamma' = 0$ and $s - s' = 0$, and in the latter case, $\deg(\epsilon_*\sigma^{s-\gamma}f_*(v)) = s + q - \gamma$ must be odd for the Dyer-Lashof operation to be nontrivial, implying (since $s + q$ is even) that $\gamma = 1$. Thus $\gamma = \gamma' = 1$ and $s - s' = 0$, and so in either case $s' = s$ and $\gamma' = \gamma$, i.e., $r' = r$.

Now let us write r'' for $(s' - s + \gamma)(p - 1) - \gamma' = \gamma(p - 1) - \gamma$. Then $\langle [G], u \cdot \overline{Q}_{r''}\nu_{s+q-\gamma} \rangle = u$, so we need to show that

$$\overline{Q}_{r''}\epsilon_*\sigma^{s-\gamma}f_*(v) = \langle x, v \rangle \cdot \overline{Q}_{r''}\nu_{s+q-\gamma}.$$

By linearity of the Dyer-Lashof operations it suffices to show that $\epsilon_*\sigma^{s-\gamma}f_*(v) = \langle x, v \rangle \cdot \nu_{s+q-\gamma}$, and, since $H_{s+q-\gamma}(K(\mathbf{Z}/p, s + q - \gamma))$ is one-dimensional, this amounts to showing that $\langle \nu_{s+q-\gamma}, \epsilon_*\sigma^{s-\gamma}f_*(v) \rangle = \langle x, v \rangle$. This is easily seen to be true.

For the other formulas we are assuming $r > 0$, which means that $\tilde{Q}_r x$ is in the kernel of the map $\tilde{H}^*(\tilde{C}_{n,p}X) \rightarrow \tilde{H}^*(\tilde{C}_{1,p}X)$ induced by inclusion. Thus the pairing of $\tilde{Q}_r x$ with any Pontryagin product operation applied to (v, w) must be zero. Similarly, since $\tilde{Q}_r x$ comes from the cohomology of $\tilde{C}_{\infty,p}X$, it must pair trivially with any Browder operation. \square

It follows from Proposition 3.4, along with the formula for $h_*(\sigma^n \overline{Q}_r v)$, that if $h^*(\sigma^n y) = \sigma^n \tilde{Q}_r x$ for some $x, y \in \tilde{H}^*(X)$, then $\langle x, v \rangle = \langle y, \overline{Q}_r v \rangle$ for all $v \in \tilde{H}_*(X)$. Equivalently, if $h^*(\sigma^n y) = \sigma^n \tilde{Q}_r x$, then $Q_r^* y = x$ as in Part 1 of the Main Theorem.

We close this section with the following lemma, which shows that the \tilde{Q}_r commute with $\tilde{\epsilon}_k^*$ just as one would want.

Lemma 3.5. *Let $k > 0$ be an integer, let X be an H_p^{n+k} -space, and choose $x \in \tilde{H}^q(X)$. Then*

$$\tilde{\epsilon}_k^*(\tilde{Q}_{s(p-1)-\gamma}\sigma^k x) = \sigma^k \tilde{Q}_{(s+k)(p-1)-\gamma} x$$

for any $\gamma \leq s < n + k$.

Proof. As with Proposition 3.4, we assume $n = \infty$ by naturality, and we will show that

$$\sigma^{q-\gamma} \tilde{\varepsilon}_k^* (\tilde{Q}_{s(p-1)-\gamma} \sigma^k x) = \sigma^{q-\gamma+k} \tilde{Q}_{(s+k)(p-1)-\gamma} x.$$

Let x be represented by the map $f: X \rightarrow K(\mathbf{Z}/p, q)$, in which case $\sigma^k x$ will be represented by the map $\epsilon \circ \Sigma^k f: \Sigma^k X \rightarrow K(\mathbf{Z}/p, q+k)$. Then, by definition

$$\sigma^{q-\gamma} \tilde{\varepsilon}_k^* (\tilde{Q}_{s(p-1)-\gamma} \sigma^k x) = (\Sigma^{s-\gamma} \tilde{\varepsilon}_k)^* \tilde{\varepsilon}_{s-\gamma}^* (\tilde{C}_{\infty,p} \Sigma^{s-\gamma} (\epsilon \circ \Sigma^k f))^* (\tilde{C}_{\infty,p} \epsilon)^* [G].$$

Now, $\tilde{\varepsilon}_j \circ \Sigma^j \tilde{\varepsilon}_k = \tilde{\varepsilon}_{j+k}$ by construction (see [KSW]), and

$$\tilde{C}_{\infty,p} \Sigma^{s-\gamma} (\epsilon \circ \Sigma^k f) = \tilde{C}_{\infty,p} \Sigma^{s-\gamma} \epsilon \circ \tilde{C}_{\infty,p} \Sigma^{k+s-\gamma} f$$

by functoriality, so we actually have

$$\sigma^{q-\gamma} \tilde{\varepsilon}_k^* (\tilde{Q}_{s(p-1)-\gamma} \sigma^k x) = \tilde{\varepsilon}_{k+s-\gamma}^* (\tilde{C}_{\infty,p} \Sigma^{k+s-\gamma} f)^* (\tilde{C}_{\infty,p} \Sigma^{s-\gamma} \epsilon)^* (\tilde{C}_{\infty,p} \epsilon)^* [G].$$

But then $\epsilon \circ \Sigma^{s-\gamma} \epsilon = \epsilon$, so in fact

$$\begin{aligned} \sigma^{q-\gamma} \tilde{\varepsilon}_k^* (\tilde{Q}_{s(p-1)-\gamma} \sigma^k x) &= \tilde{\varepsilon}_{k+s-\gamma}^* (\tilde{C}_{\infty,p} \Sigma^{k+s-\gamma} f)^* (\tilde{C}_{\infty,p} \epsilon)^* [G] \\ &= \sigma^{q-\gamma+k} \tilde{Q}_{(s+k)(p-1)-\gamma} x. \end{aligned}$$

□

4. The Main Theorem.

Part 1 of the Main Theorem has already been proved; here we restate Part 2 in slightly altered form. First we introduce the following convention. Suppose x_1 and x_2 are two elements of an \mathbf{F}_p -algebra (e.g. the mod p cohomology of a space). Then we say $x_1 \doteq x_2$ if $x_1 = u \cdot x_2$ where $u \in \mathbf{F}_p$ is a unit. This is easily seen to be an equivalence relation.

Theorem 4.1. *Suppose p is an odd prime and $X \in \text{Ob } \mathcal{H}_p^n$, with basic cofibration sequence*

$$\Sigma^n \tilde{C}_{n,p} X \xrightarrow{h} \Sigma^n X \xrightarrow{i} P_p^n X \xrightarrow{j} \Sigma^{n+1} \tilde{C}_{n,p} X.$$

Let $n = 2s + 1 - \delta$ for some $\delta \in \{0, 1\}$ and $s \geq \delta$. If $\bar{x} \in \tilde{H}^{2q+\delta+n}(P_p^n X)$, then

$$\begin{aligned} \mathcal{P}^{q+s} \bar{x} &\doteq j^* \left(\sigma^{n+1} \tilde{Q}_{(n-1)(p-1)-1} x \right), \\ \beta \mathcal{P}^{q+s} \bar{x} &\doteq j^* \left(\sigma^{n+1} \tilde{Q}_{(n-1)(p-1)} x \right), \end{aligned}$$

where $i^*\bar{x} = \sigma^n x$ defines $x \in \tilde{H}^{2q+\delta}(X)$.

We first prove Theorem 4.1 for the universal example $X = K(\mathbf{Z}/p, 2q + \delta)$ and $x = \iota_{2q+\delta}$; the universal example is then used to prove the theorem for general X . We will prove the formulas for $\beta\mathcal{P}^{q+s}\bar{\iota}_{2q+\delta}$ and $\mathcal{P}^{q+s}\bar{\iota}_{2q+\delta}$ separately.

For the $\beta\mathcal{P}^{q+s}\bar{\iota}_{2q+\delta}$ formula we first consider the special case in which $s = \delta = 0$. Let $\bar{\iota}_{2q} \in \tilde{H}^{2q+1}(P_p^1 K(\mathbf{Z}/p, 2q + \delta))$ be (the unique class) such that $i^*(\bar{\iota}_{2q}) = \iota_{2q}$; we know such a class exists because ι_{2q} is representable by an infinite loop map, and hence $h^*(\iota_{2q}) = 0$. We use the following commutative diagram, where $K(\mathbf{Z}/p, t)$ is written K_t , κ represents $\beta\mathcal{P}^q \iota_{2q+1}$, and E is the homotopy fiber of κ .

$$\begin{array}{ccccccc} K_{2pq+1} & \xrightarrow{\lambda} & E & \xrightarrow{\pi} & K_{2q+1} & \xrightarrow{\kappa} & K_{2pq+2} \\ \uparrow f_2 & & \uparrow f_1 & & \uparrow f & & \uparrow f'_2 \\ \Sigma\tilde{C}_{1,p}K_{2q} & \xrightarrow{h} & \Sigma K_{2q} & \xrightarrow{i} & P_p^1 K_{2q} & \xrightarrow{j} & \Sigma^2\tilde{C}_{1,p}K_{2q} \end{array} .$$

Here the map f represents the class $\bar{\iota}_{2q}$, and the map f_1 exists by the lifting property of a fibration since $\kappa f i$ represents

$$\beta\mathcal{P}^q i^*(\bar{\iota}_{2q}) = \beta\mathcal{P}^q \iota_{2q} = 0$$

and is thus homotopically trivial. There is of course some indeterminacy in the choice of f_1 ; we will begin with an arbitrary choice and then alter it as necessary by maps factoring through the fiber.

We will give a specific construction of f_2 making the leftmost square commute, and then we can choose f'_2 to be the adjoint of f_2 . This will make the rightmost square commute as well. (We will in general let g' denote the adjoint of g , where g may either be a map $X \rightarrow \Omega Y$ or $\Sigma X \rightarrow Y$.) The construction of f_2 will in turn be as the adjoint of a map $\tilde{C}_{1,p}K_{2q} \rightarrow K_{2pq}$ factoring through ΩE , and the key to the proof is that the image of ι_{2pq} in $H^*(\Omega E)$ is connected by the opposite Dyer-Lashof operation Q_0^* to $(\Omega\pi)^*(\iota_{2q})$, and that this connection is preserved by the map $\tilde{C}_{1,p}K_{2q} \rightarrow \Omega E$.

First let us construct f_2 . Because $\Omega\kappa$ is homotopically trivial, $\Omega E \simeq K_{2pq} \times K_{2q}$. Hence there is a map $\rho: \Omega E \rightarrow K_{2pq}$ such that $\rho\Omega\lambda$ is homotopic to the identity. We then define f_2 to be the adjoint of the composition

$$\tilde{C}_{1,p}K_{2q} \xrightarrow{h'} \Omega\Sigma K_{2q} \xrightarrow{\Omega f_1} \Omega E \xrightarrow{\rho} K_{2pq}.$$

Thus, if we write $\rho^*(\iota_{2pq})$ as just $\iota_{2pq} \in \tilde{H}^*(\Omega E)$, then f_2 represents $\sigma(h')^*(\Omega f_1)^* \iota_{2pq}$. Using the definitions of the various maps and the relationships between them, we then see that

$$\beta\mathcal{P}^q \bar{\iota} = j^*(\sigma^2(h')^*(\Omega f_1)^* \iota_{2pq}).$$

Hence to prove the theorem in this case we need only show that

$$(h')^*(\Omega f_1)^* \iota_{2pq} \doteq \tilde{Q}_0 \iota_{2q}.$$

We will do this by showing that

$$\langle (h')^*(\Omega f_1)^* \iota_{2pq}, v \rangle \doteq \begin{cases} 1 & \text{if } v = \eta_*(\nu_{2q}^{t_1}) * \cdots * \eta_*(\nu_{2q}^{t_k}) \\ & \text{with } t_1 + \cdots + t_k = p, \\ 0 & \text{otherwise,} \end{cases}$$

for any v in the basis

$$\begin{aligned} & \mathcal{B}\tilde{H}_*(\tilde{C}_{1,p}K_{2q}) \\ &= \left\{ \eta_*(v_1) * \cdots * \eta_*(v_k) \mid v_1, \dots, v_k \in \mathcal{B}\tilde{H}_*(K_{2q}), 2 \leq k \leq p \right\}. \end{aligned}$$

It is easy to see that this characterizes $\tilde{Q}_0 \iota_{2q}$.

We will be using the fact that $\langle (h')^*(\Omega f_1)^* \iota_{2pq}, v \rangle = \langle \iota_{2pq}, (\Omega f_1)_*(h')_* v \rangle$, and so we begin by considering ι_{2pq} . We claimed above that ι_{2pq} was connected to ι_{2q} by Q_0^* . More precisely, let $\mathcal{B}\tilde{H}^*(\Omega E)$ be the obvious basis consisting of products of elements of $\mathcal{B}\tilde{H}^*(K(\mathbf{Z}/p, 2q))$ with elements of $\mathcal{B}\tilde{H}^*(K(\mathbf{Z}/p, 2pq))$, and let $\mathcal{B}\tilde{H}_*(\Omega E)$ be the dual basis. Then

$$(\nu_{2q}^p)^{\text{dual}} \doteq \iota_{2pq}.$$

This follows from the well-known formula for the coproduct of ι_{2pq} in $H^*(E)$; see [Za, §3]. We cannot simply choose a representation of ΩE as $K_{2q} \times K_{2pq}$ to make the equality exact as does Zabrodsky, since the projection $\rho: \Omega E \rightarrow K_{2pq}$ is constrained by the requirement that it be a retraction of the map $\Omega\lambda$.

Since $(\nu_{2q}^p)^{\text{dual}} \doteq \iota_{2pq}$, it will suffice to show, for $v \in \mathcal{B}\tilde{H}_*(\tilde{C}_{1,p}K_{2q})$, that $(\Omega f_1)_*(h')_* v \doteq \nu_{2q}^p$ if and only if $v \doteq \eta_*(\nu_{2q}^{t_1}) * \cdots * \eta_*(\nu_{2q}^{t_k})$ with $t_1 + \cdots + t_k = p$. We will prove the following formula:

$$(\Omega f_1)_*(h')_*(\eta_*(v_1) * \cdots * \eta_*(v_k)) = -v_1 * \cdots * v_k,$$

which does the trick.

We begin with $(h')_*$. By considering how h was constructed (Section 2, see [KSW] for details), we see that its adjoint h' fits into the following commutative diagram.

$$\begin{array}{ccccc} K_{2q} & \longrightarrow & C_{1,p}K_{2q} & \longrightarrow & \tilde{C}_{1,p}K_{2q} \\ \downarrow & & \downarrow (\Sigma\theta_p^1)' - \varepsilon'_1 & & \downarrow h' \\ * & \longrightarrow & \Omega\Sigma K_{2q} & \xlongequal{\quad} & \Omega\Sigma K_{2q} \end{array}.$$

Here we note that ε'_1 is the map

$$C_{1,p}K_{2q} \hookrightarrow C_1K_{2q} \xrightarrow{\simeq} \Omega\Sigma K_{2q},$$

where the second arrow is the usual equivalence of May [Ma2].

Let $\eta_*(v_1) * \cdots * \eta_*(v_k) \in \mathcal{B}\tilde{H}_*(\tilde{C}_{1,p}K_{2q})$ be given, and observe, since $(\varepsilon'_1)_*$ is operation preserving, that

$$(h')_*(\eta_*(v_1) * \cdots * \eta_*(v_k)) = (\Sigma\theta_p^1)'_*(\eta_*(v_1) * \cdots * \eta_*(v_k)) - \eta_*(v_1) * \cdots * \eta_*(v_k).$$

Now $(\Sigma\theta_p^1)'$ is the composition

$$C_{1,p}K_{2q} \xrightarrow{\eta} \Omega\Sigma C_{1,p}K_{2q} \xrightarrow{\Omega\Sigma\theta_p^1} \Omega\Sigma K_{2q},$$

implying that $(\Sigma\theta_p^1)'_*(\eta_*(v_1) * \cdots * \eta_*(v_k)) = \eta_*(v_1 * \cdots * v_k)$ and hence

$$(h')_*(\eta_*(v_1) * \cdots * \eta_*(v_k)) = \eta_*(v_1 * \cdots * v_k) - \eta_*(v_1) * \cdots * \eta_*(v_k).$$

We will handle the two terms on the right hand separately, first altering f_1 by a map factoring through K_{2pq} so that $\langle \iota_{2pq}, (\Omega f_1)_*\eta_*(v_1 * \cdots * v_k) \rangle = 0$. For each $v \in \mathcal{B}\tilde{H}_{2pq}(K_{2q})$, let $f_v: \Sigma K_{2q} \rightarrow K_{2pq+1}$ be the map representing the cohomology class $\langle (\Omega f_1)^*\iota_{2pq}, \eta_*(v) \rangle \cdot \sigma(v^{\text{dual}})$, and take f_1 now to be the map obtained from the original choice of f_1 by subtracting the sum

$$\sum_{v \in \mathcal{B}\tilde{H}_{2pq}(K_{2q})} \lambda \circ f_v$$

using the H-structure of E . Redefine f_2 accordingly. Then

$$\langle \iota_{2pq}, (\Omega f_1)_*\eta_*(v) \rangle = 0$$

for any v in $\mathcal{B}\tilde{H}_{2pq}(K_{2q})$ by the construction of f_1 , and in particular,

$$\langle \iota_{2pq}, (\Omega f_1)_*\eta_*(v_1 * \cdots * v_k) \rangle = 0.$$

We now have to consider $\langle \iota_{2pq}, (\Omega f_1)_*(-\eta_*(v_1) * \cdots * \eta_*(v_k)) \rangle$. By construction,

$$(\Omega\pi)(\Omega f_1)\eta \simeq (\Omega f)(\Omega i)\eta \simeq \text{id},$$

and hence $(\Omega f_1)_*\eta_*(v) = v$ for any $v \in H_*(K_{2q})$, where, on the right hand side, v denotes $1 \otimes v$ in $H_*(\Omega E)$. Since Ωf_1 is an H-map, we see that $(\Omega f_1)_*(-\eta_*(v_1) * \cdots * \eta_*(v_k)) = -v_1 * \cdots * v_k \in \tilde{H}^{2pq}(\Omega E)$. Thus, putting together our calculations of $(\Omega f_1)_*$ and $(h')_*$, we have $(\Omega f_1)_*(h')_*(\eta_*(v_1) * \cdots * \eta_*(v_k)) = -v_1 * \cdots * v_k$ for any $\eta_*(v_1) * \cdots * \eta_*(v_k) \in \mathcal{B}\tilde{H}_*(\tilde{C}_{1,p}K_{2q})$, as we wanted.

Consider now the case in which $\delta = 0$ or 1 and $s > 1$. Theorem 4.1 of [KSW] implies that the following diagram commutes up to homotopy.

$$\begin{array}{ccc}
 P_p^n K(\mathbf{Z}/p, 2q + \delta) & \xrightarrow{j} & \Sigma^{n+1} \tilde{C}_{n,p} K(\mathbf{Z}/p, 2q + \delta) \\
 \downarrow \epsilon & & \downarrow \Sigma^2 \tilde{\epsilon}_{n-1} \\
 & & \Sigma^2 \tilde{C}_{1,p} \Sigma^{n-1} K(\mathbf{Z}/p, 2q + \delta) \\
 & & \downarrow \Sigma^2 \tilde{C}_{1,p} \epsilon \\
 P_p^1 K(\mathbf{Z}/p, 2(q+s)) & \xrightarrow{j} & \Sigma^2 \tilde{C}_{2,p} K(\mathbf{Z}/p, 2(q+s))
 \end{array}$$

Here, ϵ represents one of two natural transformations, running either

$$P_p^{n+k} \Omega^n \rightarrow P_p^k \quad \text{or} \quad \Sigma^{n+k} \Omega^n \rightarrow \Sigma^k,$$

which are determined by the adjointness of P_p^n and Σ^n with Ω^n , respectively. Since the choices of $\bar{\iota}_{2q+1}$ and $\bar{\iota}_{2(q+s-1)+1}$ are unique, one sees that $\epsilon^*(\bar{\iota}_{2(q+s)}) = \bar{\iota}_{2q+\delta}$, and hence $\epsilon^*(\beta \mathcal{P}^{q+s} \bar{\iota}_{2(q+s)}) = \beta \mathcal{P}^{q+s} \bar{\iota}_{2q+\delta}$. By Lemma 3.5,

$$\left(\Sigma^2 (\tilde{C}_{1,p} \epsilon \circ \tilde{\epsilon}_{n-1}) \right)^* (\sigma^2 \tilde{Q}_0 \iota_{2(q+s)}) = \sigma^{n+1} \tilde{Q}_{(n-1)(p-1) \iota_{2q+\delta}};$$

thus

$$\begin{aligned}
 j^* \left(\sigma^{n+1} \tilde{Q}_{(n-1)(p-1) \iota_{2q+\delta}} \right) &= j^* \left(\Sigma^2 (\tilde{C}_{1,p} \epsilon \circ \tilde{\epsilon}_{n-1}) \right)^* \left(\sigma^2 \tilde{Q}_0 \iota_{2(q+s)} \right) \\
 &= \epsilon^* j^* \left(\sigma^2 \tilde{Q}_0 \iota_{2(q+s)} \right) \\
 &\doteq \epsilon^* (\beta \mathcal{P}^{q+s} \bar{\iota}_{2(q+s)}) \\
 &= \beta \mathcal{P}^{q+s} \bar{\iota}_{2q+\delta}.
 \end{aligned}$$

In order to prove the formula $\mathcal{P}^{q+s} \bar{\iota}_{2q+\delta} \doteq j^* \left(\sigma^{n+1} \tilde{Q}_{(n-1)(p-1) \iota_{2q+\delta}} \right)$ for $s > 0$, we need only show the case in which $n = 2$ (and hence $s = \delta = 1$). Then the case for general n will follow by exactly the same argument as for $\beta \mathcal{P}^{q+s} \bar{\iota}_{2q+\delta}$.

Because of the commutative diagram

$$\begin{array}{ccc}
 \Sigma^{-2} \Sigma^\infty P_p^2 K(\mathbf{Z}/p, 2q+1) & \xrightarrow{j} & \Sigma \Sigma^\infty \tilde{C}_{2,p} K(\mathbf{Z}/p, 2q+1) \\
 \downarrow & & \downarrow \\
 P_p K(\mathbf{Z}/p, 2q+1) & \xrightarrow{j} & \Sigma \Sigma^\infty \tilde{C}_{\infty,p} K(\mathbf{Z}/p, 2q+1)
 \end{array}$$

it suffices to show $\mathcal{P}^{q+1} \bar{\iota}_{2q+1} \doteq j^* (\sigma \tilde{Q}_{p-2} \iota_{2q+1})$ holds in $\tilde{H}^*(P_p K(\mathbf{Z}/p, 2q+1))$.

The class $\bar{\iota}_{2q+1} \in \tilde{H}^{2q+1}(P_p K(\mathbf{Z}/p, 2q+1))$ is the unique element for which $i^* \bar{\iota}_{2q+1} = \iota_{2q+1}$. Since we have computed that $\beta \mathcal{P}^{q+1} \bar{\iota}_{2q+1} \neq 0$ in

$\tilde{H}^*(P_p^2 K(\mathbf{Z}/p, 2q+1))$, the above diagram shows us that $\mathcal{P}^{q+1} \bar{\iota}_{2q+1} \neq 0$. However, $i^*(\mathcal{P}^{q+1} \bar{\iota}_{2q+1}) = \mathcal{P}^{q+1} \iota_{2q+1} = 0$ by the unstable condition; therefore, $0 \neq \mathcal{P}^{q+1} \bar{\iota}_{2q+1} \in \text{im } j^*$. On the other hand, because the space $P_p^1 K(\mathbf{Z}/p, 2q+1)$ has L-S category 2, all threefold cup products vanish in its cohomology, and hence the image of

$$\mathcal{P}^{q+1} \bar{\iota}_{2q+1} = \bar{\iota}_{2q+1}^p \quad \text{in} \quad \tilde{H}^{2p(q+1)}(P_p^1 K(\mathbf{Z}/p, 2q+1))$$

must be zero.

In order to simplify notation, let $K = K(\mathbf{Z}/p, 2q+1)$. Consider the following commutative diagram.

$$\begin{array}{ccc} \Sigma^\infty K & \longrightarrow & \Sigma^{-1} \Sigma^\infty P_p^1 K \\ \downarrow i & & \downarrow \\ P_p K & \xlongequal{\quad} & P_p K \\ \downarrow j & & \downarrow \\ \Sigma \Sigma^\infty \tilde{C}_{\infty,p} K & \xrightarrow{p} & \Sigma \Sigma^\infty (\tilde{C}_{\infty,p} K / \tilde{C}_{1,p} K) \end{array} .$$

The discussion above implies that there is an element

$$v \in \tilde{H}^{2p(q+1)-1}(\tilde{C}_{\infty,p} K / \tilde{C}_{1,p} K)$$

for which $j^* p^*(v) = \mathcal{P}^{q+1} \bar{\iota}_{2q+1} \neq 0$. Since all (external) Pontryagin product elements in the mod p homology of $\tilde{C}_{\infty,p} K$ must come from the mod p homology of $\tilde{C}_{1,p} K$, $p^*(v)$ must be equal (up to a unit in \mathbf{F}_p) to $\tilde{Q}_{p-2} \iota_{2q+1}$, since it is the only element in that degree which is in the image of p^* . Thus $\mathcal{P}^{q+1} \bar{\iota}_{2q+1} \doteq j^*(\tilde{Q}_{p-2} \iota_{2q+1})$ as desired.

For the general case, let $f: X \rightarrow K(\mathbf{Z}/p, 2q+\delta)$ represent x . Then Theorem 6.1 of [KSW] implies that, for every choice of element \bar{x} such that $i^*(\bar{x}) = \sigma^n x$, there is a corresponding (unique) H_p^n -structure on f (note that if $i^*(\bar{x}) = 0$, then $\bar{x} \in \text{im } j^*$, and $\mathcal{P}^{q+s} \bar{x} = 0$ by the unstable condition). Now consider the following commutative diagram.

$$\begin{array}{ccccc} \Sigma^n X & \xrightarrow{i} & P_p^n X & \xrightarrow{j} & \Sigma^{n+1} \tilde{C}_{n,p} X \\ \downarrow f & & \downarrow P_p^n f & & \downarrow \Sigma^{n+1} \tilde{C}_{n,p} f \\ \Sigma^n K(\mathbf{Z}/p, 2q+\delta) & \xrightarrow{i} & P_p^n K(\mathbf{Z}/p, 2q+\delta) & \xrightarrow{j} & \Sigma^{n+1} \tilde{C}_{n,p} K(\mathbf{Z}/p, 2q+\delta) \end{array}$$

By construction, $f^*(\iota_{2q+\delta}) = x$, $(P_p^n f)^*(\bar{\iota}_{2q+\delta}) = \bar{x}$ and $(\tilde{C}_{n,p} f)^*(\tilde{Q}_r \iota_{2q+\delta}) \doteq \tilde{Q}_r x$, and the result follows. QED

The following corollary is the stable version of Theorem 4.1. Recall from [KSW] that there is a natural map $\Sigma^{-n} \Sigma^\infty P_p^n X \rightarrow P_p X$ for every

$X \in \text{Ob } \mathcal{H}_p^\infty$. The geometric filtration determined by these maps yields an algebraic filtration of $\tilde{H}^*(P_p X)$, and we define

$$F^{k+1} = \ker\{\tilde{H}^*(P_p X) \rightarrow \tilde{H}^*(\Sigma^{-k}\Sigma^\infty P_p^k X)\}.$$

Corollary 4.2. *Suppose p is an odd prime and $X \in \text{Ob } \mathcal{H}_p^\infty$, with basic cofibration sequence*

$$\Sigma^\infty \tilde{C}_{\infty,p} X \xrightarrow{h} \Sigma^\infty X \xrightarrow{i} P_p X \xrightarrow{j} \Sigma \Sigma^\infty \tilde{C}_{\infty,p} X.$$

Let $n = 2s + 1 - \delta$ for some $\delta \in \{0, 1\}$ and $s \geq \delta$. If $\bar{x} \in \tilde{H}^{2q+\delta}(P_p X)$, and $s > 0$, then

$$\begin{aligned} \mathcal{P}^{q+s} \bar{x} &\doteq j^*(\sigma^\infty \tilde{Q}_{(n-1)(p-1)-1} x) \pmod{F^{n+1}}, \\ \beta \mathcal{P}^{q+s} \bar{x} &\doteq j^*(\sigma^\infty \tilde{Q}_{(n-1)(p-1)} x) \pmod{F^{n+1}}, \end{aligned}$$

where $i^* \bar{x} = \sigma^\infty x$ defines $x \in \tilde{H}^{2q+\delta}(X)$.

Appendix: The Nishida relations.

The Nishida relations [NI, CLM], are formulas which relate the action of Dyer-Lashof operations to the (adjoint) action of the Steenrod algebra on $\tilde{H}_*(\tilde{C}_{n,p} X)$. In computations using the main theorem it is helpful to have the analogous formulas in cohomology. They are given in the next proposition.

Recall from Section 3 that $\tilde{H}_*(\tilde{C}_{n,p} X) \cong B \oplus C$, for certain submodules B and C . Let $\text{Ann}(B) \subset \tilde{H}^*(\tilde{C}_{n,p} X)$ be the submodule of elements which pair trivially with every element of B under the Kronecker pairing.

Let $\delta \in \{0, 1\}$ and define a function λ by $\lambda(2j + \delta) = \delta$. Finally, let $u \in \mathbf{F}_p$ denote an undetermined unit.

Proposition A.1. *The following formulas (where (i) is taken mod $\text{Ann}(B)$ in the case that $r = 0$) hold for every $x \in \tilde{H}^q(X)$.*

(i)

$$\begin{aligned} \mathcal{P}^k \tilde{Q}_r x &\doteq \sum_{i=0}^{\lfloor k/p \rfloor} \binom{\lfloor r/2 \rfloor + (p-1)(q/2 - i)}{k - pi} \tilde{Q}_{r+2(p-1)(k-pi)} \mathcal{P}^i x \\ &+ u \lambda(r-1) \sum_{i=0}^{\lfloor k/p \rfloor} \binom{\lfloor r/2 \rfloor + (p-1)(q/2 - i) - 1}{k - pi - 1} \tilde{Q}_{r+2(p-1)(k-pi)-p} \beta \mathcal{P}^i x. \end{aligned}$$

(ii)

$$\beta \tilde{Q}_{2r-1} x \doteq \tilde{Q}_{2r} x.$$

Proof. It suffices to prove the formulas in $\tilde{H}^*(D_p K(\mathbf{Z}/p, q))$, with $x = \iota_q$, where they are dual to the usual Nishida formulas. \square

References

- [BT] W. Browder and E. Thomas, *On the projective plane of an H-space*, Ill. J. Math., **7** (1963), 493–501.
- [BMMS] R. Bruner, J.P. May, J. McClure, and M. Steinberger, *H_∞ Ring Spectra and their Applications*, Springer Verlag, LNM, **1176** (1986).
- [CLM] F.R. Cohen, T.J. Lada, and J.P. May, *The Homology of Iterated Loop Spaces*, Springer Verlag, LNM, **533** (1976).
- [CPS] H. Campbell, F. Peterson, and P. Selick, *Self-Maps of Loop Spaces*, I, Trans. Amer. Math. Soc., **293** (1986), 1-39.
- [KSW] N.J. Kuhn, M. Slack, and F. Williams, *A Hopf type construction for iterated loop spaces*, preprint.
- [Ma2] J.P. May, *The Geometry of Iterated Loop Spaces*, Springer Verlag, LNM **271** (1970).
- [Mi] R.J. Milgram, *Iterated loop spaces*, Ann. Math., **11** (1967), 242-250.
- [Ni] G. Nishida, *Cohomology operations in iterated loop spaces*, Proc. J. Acad., **44** (1968), 104-109.
- [Sl] M. Slack, *Infinite loop spaces with trivial Dyer-Lashof operations*, Math. Proc. Camb. Phil. Soc., **113** (1993), 311-328.
- [Th] E. Thomas, *On functional cup-products and the transgression operator*, Arch. Math., **12** (1961), 435-444.
- [Za] A. Zabrodsky, *Secondary operations in the cohomology of H-spaces*, Illinois J. Math., **15** (1971), 648-655.

Received September 27, 1993.

JACKSONVILLE UNIVERSITY
 JACKSONVILLE, FL 32211
E-mail address: mfoskey@junix.ju.edu

AND

WESTERN MICHIGAN UNIVERSITY
 KALAMAZOO, MI 49008
E-mail address: slack@math-stat.wmich.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded by

E. F. Beckenbach (1906-1982) F. Wolf (1904-1989)

EDITORS

Sun-Yung A. Chang (Managing Editor)
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

F. Michael Christ
University of California
Los Angeles, CA 90095-1555
christ@math.ucla.edu

Nicholas Ercolani
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

Robert Finn
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

Steven Kerckhoff
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

Martin Scharlemann
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

Gang Tian
Massachusetts Institute of Technology
Cambridge, MA 02139
tian@math.mit.edu

V. S. Varadarajan
University of California
Los Angeles, CA 90095-1555
vsv@math.ucla.edu

Dan Voiculescu
University of California
Berkeley, CA 94720
dvv@math.berkeley.edu

SUPPORTING INSTITUTIONS

CALIFORNIA INSTITUTE OF TECHNOLOGY
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
STANFORD UNIVERSITY
UNIVERSITY OF ARIZONA
UNIVERSITY OF BRITISH COLUMBIA
UNIVERSITY OF CALIFORNIA
UNIVERSITY OF HAWAII

UNIVERSITY OF MONTANA
UNIVERSITY OF NEVADA, RENO
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA
UNIVERSITY OF UTAH
UNIVERSITY OF WASHINGTON
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The table of contents and the abstracts of the papers in the current issue, as well as other information about the Pacific Journal of Mathematics, may be found on the Internet at <http://www.msri.org>.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$245.00 a year (10 issues). Special rate: \$123.00 a year to individual members of supporting institutions. Subscriptions, back issues published within the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143 Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,
the American Mathematical Society's TEX macro system.

Copyright © 1995 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 173 No. 1 March 1996

Isometric immersions of H_1^n into H_1^{n+1}	1
KINETSU ABE	
Rotationally symmetric hypersurfaces with prescribed mean curvature	29
MARIE-FRANÇOISE BIDAUT-VÉRON	
The covers of a Noetherian module	69
JIAN-JUN CHUAI	
On the odd primary cohomology of higher projective planes	77
MARK FOSKEY and MICHAEL DAVID SLACK	
Unit indices of some imaginary composite quadratic fields. II	93
MIKIHITO HIRABAYASHI	
Mixed automorphic vector bundles on Shimura varieties	105
MIN HO LEE	
Trace ideal criteria for Toeplitz and Hankel operators on the weighted Bergman spaces with exponential type weights	127
PENG LIN and RICHARD ROCHBERG	
On quadratic reciprocity over function fields	147
KATHY DONOVAN MERRILL and LYNNE WALLING	
(A_2) -conditions and Carleson inequalities in Bergman spaces	151
TAKAHIKO NAKAZI and MASAHIRO YAMADA	
A note on a paper of E. Boasso and A. Larotonda: "A spectral theory for solvable Lie algebras of operators"	173
C. OTT	
Tensor products with anisotropic principal series representations of free groups	181
CARLO PENSVALLE and TIM STEGER	
On Ricci deformation of a Riemannian metric on manifold with boundary	203
YING SHEN	
The Weyl quantization of Poisson $SU(2)$	223
ALBERT JEU-LIANG SHEU	
Weyl's law for $SL(3, \mathbb{Z}) \backslash SL(3, \mathbb{R}) / SO(3, \mathbb{R})$	241
ERIC GEORGE STADE and DOROTHY IRENE WALLACE (ANDREOLI)	
Minimal hyperspheres in two-point homogeneous spaces	263
PER TOMTER	
Subalgebras of little Lipschitz algebras	283
NIKOLAI ISAAC WEAVER	