

# *Pacific Journal of Mathematics*

**ON THE CAUCHY PROBLEM FOR A SINGULAR PARABOLIC  
EQUATION**

XIANGSHENG XU

# ON THE CAUCHY PROBLEM FOR A SINGULAR PARABOLIC EQUATION

XIANGSHENG XU

The existence of a renormalized solution is established for the Cauchy problem for the parabolic P-Laplacian equation in which  $p$  is allowed to be close to 1 and the initial data are only assumed to be locally integrable.

## 1. Introduction.

We shall be concerned with the existence of a solution to the following problem

$$(1.1a) \quad \frac{\partial}{\partial t} u - \operatorname{div} (|\nabla u|^{p-2} \nabla u) = 0 \quad \text{in } \Sigma_T \equiv \mathbf{R}^N \times (0, T),$$

$$(1.1b) \quad u(x, 0) = u_0(x) \quad \text{on } \mathbf{R}^N$$

in the case where  $T > 0$ ,  $1 < p < 2$ , and  $u_0 \in L^1_{\text{loc}}(\mathbf{R}^N)$ . The restriction on  $p$  makes the equation (1.1a) singular because the term  $|\nabla u|^{p-2}$ , which measures the modulus of ellipticity of the principal part of (1.1a), is unbounded at points where  $|\nabla u|$  is 0. Thus we are dealing with a singular parabolic problem.

It is observed in [DH] that in the generality considered here an estimate of the form

$$(1.2) \quad |\nabla u| \in L^q_{\text{loc}}(\Sigma_T), \quad q \geq 1$$

is no longer possible. This suggests that solutions of (1.1a) display new phenomena that cannot be incorporated into the classical weak formulation. To define our notion of a weak solution, we follow the approach adopted in [X1]. Let  $\mathcal{A} = \{\theta \in C(\mathbf{R}) : \theta \text{ is a Lipschitz function whose derivative } \theta'(s) \text{ exists except at finitely many points and } \theta'(s) = 0 \text{ for } |s| \text{ sufficiently large}\}$ . If a measurable function  $v$  on  $\Sigma_T$  is such that  $\theta(v) \in L^p(0, T; W^{1,p}_{\text{loc}}(\mathbf{R}^N))$  for all  $\theta \in \mathcal{A}$ , then we can define a measurable function  $g : \Sigma_T \rightarrow \mathbf{R}^N$  so that

$$g = \nabla P_M(v) \quad \text{almost everywhere on } \{|v| < M\}$$

for all  $M > 0$ , where  $P_M(s) = \min\{|s|, M\} \operatorname{sign}(s)$ . The function  $g$  is viewed as the spatial gradient of  $v$ , and is also denoted by  $\nabla v$ . We are ready to present our definition of a solution.

**Definition.** A measurable function  $u$  on  $\Sigma_T$  is said to be a renormalized solution of (1.1) if:

1.  $u \in C([0, T]; L^1_{\text{loc}}(\mathbf{R}^N));$

2. For each  $\theta \in \mathcal{A}$ ,  $\theta(u) \in L^p(0, T; W_{\text{loc}}^{1,p}(\mathbf{R}^N))$  and  $\nabla\theta(u) = \theta'(u)\nabla u$  almost everywhere on  $\Sigma_T$ , where  $\theta'(u)$  is understood to be 0 if  $u \in B_\theta \equiv \{s \in \mathbf{R} : \theta'(s) \text{ does not exist}\}$ ;
3.  $|\nabla u|^{p-1} \in L^1(0, T; L_{\text{loc}}^1(\mathbf{R}^N))$  and

$$\begin{aligned}
& - \int_{\Sigma_T} \int_0^u \theta(s) ds \varphi_t dx dt + \int_{\Sigma_T} |\nabla u|^{p-2} \nabla u (\nabla\theta(u)\varphi + \theta(u)\nabla\varphi) dx dt \\
& = \int_{\mathbf{R}^N} \varphi(x, 0) \int_0^{u_0(x)} \theta(s) ds dx
\end{aligned}$$

for all  $\theta \in \mathcal{A}$  and all  $\varphi \in C_0^\infty(\mathbf{R}^N \times (-\infty, T))$ .

The idea of a renormalized solution was originated in the study of the Boltzmann equation; see [DL1, DL2] for details. An elliptic version of this idea appears in [BGDM]. The definition here is a slight modification of that in [X1]; also see [X2] where it is evident that the notion of a renormalized solution is the correct notion of solution for p-Laplacian problems. The objective of this paper is to show that there exists a renormalized solution to (1.1).

If  $u_0 \geq 0$ , the existence and uniqueness of a solution to (1.1) are established in [DH]. In [X1], the sign restriction on  $u_0$  is removed, but  $\mathbf{R}^N$  is replaced with a bounded domain  $\Omega$ . The stationary problem is considered in [X2] and references therein. The question of existence and uniqueness of a solution to (1.1) in the case where  $u_0$  may change sign was proposed as an open problem in [DH]. In this paper, we solve the question of existence, while the question of uniqueness remains open.

It is interesting to note that we obtain a renormalized solution to (1.1) without imposing any growth condition on  $u_0$ . This is in sharp contrast with the case  $p > 2$  [D]. Also, it is easy to infer from the argument in [D, p. 188-192] that if  $u_0 \in L^s(\mathbf{R}^N)$ ,  $s = N(2-p)/p$ ,  $1 < p < 2N/(N+1)$ , and  $N \geq 2$ , then the renormalized solution  $u$  constructed here will extinct in finite time, i.e., there exists a positive number  $T^*$  such that  $u(x, t) = 0$  for all  $t > T^*$ .

The main gap between the case  $u_0 \geq 0$ , and the case where  $u_0$  may change sign, is that in the latter case an estimate of the type

$$\int_s^T \int_{\{|x| < R\}} \frac{u_t^2}{(1 + |u|)^{1+\varepsilon}} dx dt < \infty, \quad s \in (0, T), \varepsilon > 0, R > 0$$

is no longer available. To overcome this difficulty, we develop an analysis that combines the best features of the arguments in [DH] and [X1] with a compactness theorem of Simon [S].

This work is organized as follows. In Section 2, we prove a comparison principle for classical weak solutions of (1.1a). This result is used in Section 3 to prove the existence of a renormalized solution.

We conclude this section by making some remarks on notation. Let  $R > 0$ , and we denote by  $B_R$  the ball centered at the origin with radius  $R$ . Fix  $R > r > 0$ . We say that  $\xi$  is a cut-off function associated with  $R$  and  $r$  if  $\xi \in C_0^\infty(B_R)$ ,  $0 \leq \xi \leq 1$ ,  $\xi = 1$  on  $B_r$ , and  $|\nabla \xi| \leq \frac{2}{R-r}$ . Let  $E$  be a measurable set in  $\mathbf{R}^{N+1}$ . We use  $|E|$  to denote the Lebesgue measure of  $E$ .

## 2. Preliminaries.

In this section we consider the problem

$$(2.1a) \quad \frac{\partial}{\partial t} u - \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0 \quad \text{in } \Sigma_T,$$

$$(2.1b) \quad u(x, 0) = u_0(x) \quad \text{on } \mathbf{R}^N$$

in the case where  $u_0 \in L_{\text{loc}}^2(\mathbf{R}^N)$  and  $1 < p < 2$ . A function  $u$  on  $\Sigma_T$  is said to be a classical weak solution of (2.1) if:

- (i)  $u \in C([0, T]; L_{\text{loc}}^2(\mathbf{R}^N)) \cap L^p(0, T; W_{\text{loc}}^{1,p}(\mathbf{R}^N))$ ;
- (ii)  $-\int_{\Sigma_T} u \varphi_t dx dt + \int_{\Sigma_T} |\nabla u|^{p-2} \nabla u \nabla \varphi dx dt = \int_{\mathbf{R}^N} \varphi(x, 0) u_0(x) dx$  for all  $\varphi \in C_0^\infty(\mathbf{R}^N \times (-\infty, T))$ .

Let  $u$  be a classical weak solution to (2.1). Then we can easily deduce from (ii) that for each  $\rho > 0$ ,

$$(2.2) \quad u_t \in L^{p'}(0, T; W^{-1,p'}(B_\rho))$$

$$(2.3) \quad u_t - \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0 \quad \text{in } W^{-1,p'}(B_\rho) \quad \text{for almost every } t \in (0, T).$$

Here and in what follows  $p' = p/(p-1)$ .

**Lemma 2.1.** *Let  $u$  be a classical weak solution of (2.1). Then  $u_0 \in L_{\text{loc}}^\infty(\mathbf{R}^N)$  implies  $u \in L^\infty(0, T; L_{\text{loc}}^\infty(\mathbf{R}^N))$ .*

**Remark.** If  $u_0 \geq 0$ , then this lemma is a direct consequence of Theorem III.6.2 in [DH].

*Proof of Lemma 2.1.* We modify a device in [DH]. Fix  $R > 0$ . For  $n = 0, 1, 2, \dots$ , define

$$\rho_n = R(1 + 2^{-n}), B_n = B_{\rho_n}, k_n = M(2 - 2^{-n}),$$

where  $M \geq \|u_0\|_{L^\infty(B_{2R})}$  will be selected later. Let  $\xi_n$  be a cut-off function associated with  $\rho_n$  and  $\rho_{n+1}$ . Then we can derive from the chain rule [X1] that the function  $t \rightarrow \frac{1}{2} \int_{B_n} [(u - k_n)^+]^2 \xi_n^p dx$  is absolutely continuous on  $[0, T]$ , and

$$(2.4) \quad \frac{d}{dt} \frac{1}{2} \int_{B_n} [(u - k_n)^+]^2 \xi_n^p dx = (u_t, (u - k_n)^+ \xi_n^p)$$

almost everywhere on  $(0, T)$ ,

where  $(\cdot, \cdot)$  denotes the duality pairing between  $W^{-1,p'}(B_n)$  and  $W_0^{1,p}(B_n)$ . Keep this in mind, use  $(u - k_n)^+ \xi_n^p$  as a test function in (2.3), thereby obtain

$$\begin{aligned} & \frac{d}{dt} \frac{1}{2} \int_{B_n} [(u - k_n)^+]^2 \xi_n^p dx + \int_{B_n} |\nabla (u - k_n)^+|^p \xi_n^p dx \\ &= - \int_{B_n} |\nabla (u - k_n)^+|^{p-2} \nabla (u - k_n)^+ (u - k_n)^+ p \xi_n^{p-1} \nabla \xi_n dx \\ &\leq \frac{1}{2} \int_{B_n} |\nabla (u - k_n)^+|^p \xi_n^p dx + 2^{p-1} \left(\frac{p}{R}\right)^p 2^{p(n+1)} \int_{B_n} [(u - k_n)^+]^p dx. \end{aligned}$$

Consequently,

$$\begin{aligned} (2.5) \quad & \max_{0 \leq t \leq T} \int_{B_n} [(u - k_n)^+]^2 \xi_n^p dx + \int_{B_n \times (0, T)} |\nabla (u - k_n)^+|^p \xi_n^p dx dt \\ &\leq \left(\frac{p}{R}\right)^p 2^{p(n+2)} \int_{B_n \times (0, T)} [(u - k_n)^+]^p dx dt. \end{aligned}$$

This, in conjunction with the Gagliardo-Nirenberg-Sobolev inequality, implies

$$\begin{aligned} & \int_{B_n \times (0, T)} [(u - k_n)^+ \xi_n]^{p \frac{N+2}{N}} dx dt \\ &\leq c_0 \left( \sup_{0 \leq t \leq T} \int_{B_n} [(u - k_n)^+ \xi_n]^2 dx \right)^{\frac{N}{2}} \\ &\quad \cdot \int_{B_n \times (0, T)} |\nabla ((u - k_n)^+ \xi_n)|^p dx dt \\ &\leq c_1 \frac{2^{\left(\frac{p(N+p)}{N}\right)n}}{R^{\frac{p(N+p)}{N}}} \left( \int_{B_n \times (0, T)} [(u - k_n)^+]^p dx dt \right)^{\frac{(N+p)}{N}}. \end{aligned}$$

Here, and in what follows,  $c_i, i \in \{0, 1, 2, \dots\}$ , denote positive constants depending only upon  $p, N$ . We estimate

$$\int_{B_{n+1} \times (0, T)} [(u - k_{n+1})^+]^p dx dt$$

$$\begin{aligned}
 (2.6) \quad & \leq \int_{B_n \times (0, T)} \left[ (u - k_{n+1})^+ \xi_n \right]^p dxdt \\
 & \leq |B_n \times (0, T) \cap \{u > k_{n+1}\}|^{\frac{2}{N+2}} \\
 & \quad \cdot \left( \int_{B_n \times (0, T)} \left[ (u - k_{n+1})^+ \xi_n \right]^{p \frac{N+2}{N}} dxdt \right)^{\frac{N}{N+2}} \\
 & \leq c_2 \frac{2^{\left(\frac{p(N+p)}{(N+2)}\right)n}}{R^{\frac{p(N+p)}{(N+2)}}} |B_n \times (0, T) \cap \{u > k_{n+1}\}|^{\frac{2}{N+2}} \\
 & \quad \cdot \left( \int_{B_n \times (0, T)} \left[ (u - k_n)^+ \right]^p dxdt \right)^{\frac{N+p}{N+2}}.
 \end{aligned}$$

Observe that

$$\begin{aligned}
 & \int_{B_n \times (0, T)} \left[ (u - k_n)^+ \right]^p dxdt \\
 & \geq \int_{B_n \times (0, T) \cap \{u > k_{n+1}\}} (k_{n+1} - k_n)^p dxdt \\
 & = M^p 2^{-p(n+1)} |B_n \times (0, T) \cap \{u > k_{n+1}\}|.
 \end{aligned}$$

This, together with (2.6) shows that

$$\begin{aligned}
 & \int_{B_{n+1} \times (0, T)} \left[ (u - k_{n+1})^+ \right]^p dxdt \\
 & \leq c_3 \frac{2^{\left[\frac{p(N+p)}{(N+2)} + \frac{2p}{(N+2)}\right]n}}{R^{\frac{p(N+p)}{(N+2)}} M^{\frac{2p}{(N+2)}}} \left( \int_{B_n \times (0, T)} \left[ (u - k_n)^+ \right]^p dxdt \right)^{1 + \frac{p}{N+2}}.
 \end{aligned}$$

According to a result in [LSU, p. 95],  $\lim_{n \rightarrow \infty} \int_{B_n \times (0, t)} \left[ (u - k_n)^+ \right]^p dxdt = 0$ , provided we can select  $M \geq \|u_0\|_{L^\infty(B_{2R})}$  so that

$$\begin{aligned}
 (2.7) \quad & \int_{B_{2R} \times (0, T)} \left[ (u - M)^+ \right]^p dxdt \leq \left( \frac{c_3}{R^{\frac{p(N+p)}{(N+2)}} M^{\frac{2p}{(N+2)}}} \right)^{-\frac{N+2}{p}} \\
 & \quad \cdot \left( 2^{\frac{(pN+p^2+2p)}{(N+2)}} \right)^{-\left(\frac{N+2}{p}\right)^2} \\
 & \leq c_4 R^{(N+p)} M^2.
 \end{aligned}$$

This can be easily done, and hence

$$\int_{B_R \times (0, T)} \left[ (u - 2M)^+ \right]^p dxdt \leq \lim_{n \rightarrow \infty} \int_{B_R \times (0, T)} \left[ (u - k_n)^+ \right]^p dxdt = 0.$$

To see that  $u$  is also bounded below, note that  $v = -u$  is a classical weak solution of the following problem

$$\begin{aligned} \frac{\partial v}{\partial t} - \operatorname{div}(|\nabla v|^{p-2} \nabla v) &= 0 \quad \text{in } \Sigma_T, \\ v(x, 0) &= -u_0(x) \quad \text{in } \mathbf{R}^N. \end{aligned}$$

This completes the proof of the lemma.

Before we continue, let us recall the following lemma from [O, pp. 145–147].

**Lemma 2.2.** *Let  $x, y$  be any two vectors in  $\mathbf{R}^N$  and  $p \in (1, 2]$ . Then,*

- (a)  $(|x|^{p-2}x - |y|^{p-2}y)(x - y) \geq (p - 1) \frac{|x - y|^2}{(|x| + |y|)^{2-p}};$
- (b)  $||x|^{p-2}x - |y|^{p-2}y| \leq \sqrt{5}|x - y|^{p-1}.$

**Lemma 2.3.** *Let  $u_0, v_0$  be two functions in  $L_{\text{loc}}^\infty(\mathbf{R}^N)$ . Assume that  $u$  and  $v$  are classical weak solutions of (2.1a) with initial conditions  $u_0$  and  $v_0$ , respectively. Then  $u_0 \leq v_0$  implies  $u \leq v$ .*

*Proof.* Fix  $R > r > 0$ . Let  $\xi$  be a cut-off function associated with  $R$  and  $r$ . By Lemma 2.2,  $u, v \in L^\infty(0, T; L_{\text{loc}}^\infty(\mathbf{R}^N))$ . Thus for each  $q > 1$ ,  $[(u - v)^+]^q \xi^2 \in L^p(0, T; W_0^{1,p}(B_R))$ . We can conclude from (2.3) and the chain rule [X1] that

$$\begin{aligned} (2.8) \quad & \frac{d}{dt} \frac{1}{q+1} \int_{B_R} [(u - v)^+]^{q+1} \xi^2 dx \\ & + \int_{B_R} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) q [(u - v)^+]^{q-1} \nabla(u - v) \xi^2 dx \\ & = - \int_{B_R} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) [(u - v)^+]^q 2\xi \nabla \xi dx \\ & \leq \frac{2}{R - r} \int_{B_R} ||\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v| [(u - v)^+]^q \xi dx. \end{aligned}$$

Set

$$A_t = \left\{ x : (u(x, t) - v(x, t))^+ \frac{2}{R - r} \leq \frac{1}{2} q \left| \nabla(u(x, t) - v(x, t))^+ \right| \xi(x) \right\}.$$

We compute, with the aid of Lemma 2.2, that

$$\frac{2}{R - r} \int_{B_R} ||\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v| [(u - v)^+]^q \xi dx$$

$$\begin{aligned}
&\leq \frac{1}{2} \int_{B_R \cap A_t} \left| |\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v \right| |\nabla(u-v)^+| q [(u-v)^+]^{q-1} \xi^2 dx \\
&\quad + \frac{2}{R-r} \int_{B_R \setminus A_t} \sqrt{5} |\nabla u - \nabla v|^{p-1} [(u-v)^+]^q \xi dx \\
&\leq \frac{1}{2} \int_{B_R} (|\nabla u|^{p-2} \nabla u - |\nabla v|^{p-2} \nabla v) \nabla(u-v) q [(u-v)^+]^{q-1} \xi^2 dx \\
&\quad + \frac{2}{R-r} \int_{B_R} \sqrt{5} \left( \frac{4}{q(R-r)} (u-v)^+ \right)^{p-1} [(u-v)^+]^q dx.
\end{aligned}$$

Use this in (2.8) to obtain

$$(2.9) \quad \int_{B_r} [(u-v)^+]^{q+1} dx \leq \frac{\sqrt{5}(q+1)2^{2p-1}}{q^{p-1}(R-r)^p} \int_{B_R \times (0,t)} [(u-v)^+]^{q+p-1} dx d\tau.$$

Now we are ready to employ an argument in [DH]. Fix  $\rho > 0$ , and set

$$\begin{aligned}
\rho_n &= \left( \sum_{i=0}^n 2^{-i} \right) \rho, \quad B_n = B_{\rho_n}, \\
\Lambda_n &= \sup_{0 \leq T \leq t} \int_{B_n} [(u-v)^+]^{q+1} dx \quad (n = 0, 1, 2, \dots).
\end{aligned}$$

We can infer from (2.9) that

$$\begin{aligned}
\Lambda_n &\leq c \frac{2^{p(n+1)}}{\rho^p} \int_{B_{n+1} \times (0,t)} [(u-v)^+]^{q+p-1} dx d\tau \\
&\leq c t^{\frac{2-p}{q+1}+1} (2p)^N \frac{2^{\frac{q+p-1}{q+1}}}{\rho^p} \Lambda_{n+1} \\
&= c_1 t^{\frac{3-p+q}{q+1}} \frac{2^{pn}}{\rho^{p-\frac{(2-p)N}{(q+1)}}} \Lambda_{n+1}^{\frac{(q+p-1)}{(q+1)}} \\
&\leq \delta \Lambda_{n+1} + \left( 2^{p\frac{q+1}{2-p}} \right)^n c(\delta) \left( \frac{t^{\frac{(3-p+q)}{(q+1)}}}{\rho^{p-\frac{(2-p)N}{(q+1)}}} \right)^{\frac{q+1}{2-p}}.
\end{aligned}$$

Here  $\delta > 0$  is arbitrary. This implies

$$(2.10) \quad \Lambda_0 \leq \delta^n \Lambda_n + \frac{1}{\delta} c(\delta) \left( \frac{t^{\frac{(3-p+q)}{(q+1)}}}{\rho^{p-\frac{(2-p)N}{(q+1)}}} \right)^{\frac{q+1}{2-p}} \sum_{i=0}^{n+1} \left( \delta 2^{p\frac{q+1}{2-p}} \right)^i.$$

Now we select  $\delta > 0$  and  $q > 0$  so that

$$2^{p\frac{q+1}{2-p}} = \frac{1}{2} \quad \text{and} \quad (q+1)p - (2-p)N > 0.$$



We conclude from (2.10) that

$$\sup_{0 \leq \tau \leq t} \int_{B_\rho} [(u - v)^+]^{q+1} dx \leq c \left( \frac{t^{\frac{(3-p+q)}{(q+1)}}}{\rho^{\frac{(q+1)p - (2-p)N}{q+1}}} \right)^{\frac{q+1}{2-p}}$$

$$\rightarrow 0 \quad \text{as} \quad \rho \rightarrow \infty.$$

This proves the lemma. □

An easy consequence of Lemma 2.1 and Lemma 2.3 is that

$$\|u(\cdot, t)\|_{L^\infty(\mathbf{R}^N)} \leq \|u_0\|_{L^\infty(\mathbf{R}^N)}$$

for each  $t > 0$ .

### 3. Existence.

The main result of this section is:

**Theorem 3.1.** *Assume that  $u_0 \in L^1_{\text{loc}}(\mathbf{R}^N)$ , and  $1 < p < 2$ . Then there exists a renormalized solution to (1.1).*

*Proof.* If  $k \in \{1, 2, \dots\}$ , define

$$(3.1) \quad f_k(x) = \min \{u_0^+(x), k\},$$

$$(3.2) \quad g_k(x) = \min \{u_0^-(x), k\}.$$

For each  $k$ , consider the approximating problem

$$(3.3a) \quad \frac{\partial u_k}{\partial t} - \operatorname{div} (|\nabla u_k|^{p-2} \nabla u_k) = 0 \quad \text{on } \Sigma_T,$$

$$(3.3b) \quad u(x, 0) = u_{0k}(x) = f_k - g_k \quad \text{in } \mathbf{R}^N.$$

The existence of a classical weak solution to (3.3) can be inferred from a result in [DH, D]. Since  $u_{0k} \in L^\infty(\mathbf{R}^N)$ , Lemma 2.3 asserts the uniqueness. The remaining proof is divided into several lemmas. □

**Lemma 3.1.** *For each  $\rho > 0$ , there exists a  $c(\rho) > 0$  such that*

$$(3.4) \quad \max_{0 \leq t \leq T} \int_{B_\rho} |u_k(x, t)| dx \leq c(\rho),$$

$$(3.5) \quad \int_{B_\rho \times (0, T)} |\nabla u_k|^{p-1} dx dt \leq c(\rho) \quad (k = 1, 2, \dots).$$

*Proof.* For each  $k$ , let  $v_k$  be the classical weak solution of the following problem

$$(3.6a) \quad \frac{\partial}{\partial t} v_k - \operatorname{div} \left( |\nabla v_k|^{p-2} \nabla v_k \right) = 0 \quad \text{in } \Sigma_T,$$

$$(3.6b) \quad v_k(x, 0) = f_k(x) \quad \text{on } \mathbf{R}^N,$$

and  $w_k$  be the classical weak solution of the following problem

$$(3.7a) \quad \frac{\partial}{\partial t} w_k - \operatorname{div} \left( |\nabla w_k|^{p-2} \nabla w_k \right) = 0 \quad \text{in } \Sigma_T,$$

$$(3.7b) \quad w_k(x, 0) = -g_k(x) \quad \text{on } \mathbf{R}^N.$$

In light of Lemma 2.3, we have

$$(3.8) \quad w_k \leq u_k \leq v_k \quad \text{almost everywhere on } \Sigma_T$$

for all  $k$ . Since  $f_k \geq 0$  on  $\mathbf{R}^N$ , we can invoke a result in [DH, p. 260] to obtain that there exists a  $c_1(\rho) > 0$  such that

$$(3.9) \quad \max_{0 \leq t \leq T} \int_{B_\rho} v_k(x, t) dx \leq c_1(\rho) \quad (k = 1, 2, \dots).$$

Note that  $z_k = -w_k$  is the classical weak solution of the problem

$$\begin{aligned} \frac{\partial}{\partial t} z_k - \operatorname{div} \left( |\nabla z_k|^{p-2} \nabla z_k \right) &= 0 \quad \text{in } \Sigma_T, \\ z_k(x, 0) &= g_k(x) \quad \text{on } \mathbf{R}^N. \end{aligned}$$

Thus, we can find  $c_2(\rho) > 0$  with

$$(3.10) \quad \max_{0 \leq t \leq T} \int_{B_\rho} |w_k(x, t)| dx \leq c_2(\rho) \quad (k = 1, 2, \dots).$$

We see that (3.4) is a consequence of (3.8), (3.9), and (3.10). To see (3.5), for each  $\varepsilon > 0$  define

$$(3.11) \quad \phi_\varepsilon(s) = \begin{cases} 1 - \frac{1}{(1+s)^\varepsilon} & \text{if } s \geq 0, \\ -\phi_\varepsilon(-s) & \text{if } s < 0. \end{cases}$$

Let  $\xi$  be a cut-off function associated with  $2\rho$  and  $\rho$ . Then using  $\phi_\varepsilon(u_k)\xi^p$  as a test function in (3.3a), we derive from a standard argument [X1] that

$$(3.12) \quad \frac{d}{dt} \int_{B_{2\rho}} \int_0^{u_k(x,t)} \phi_\varepsilon(s) ds \xi^p(x) dx + \int_{B_{2\rho}} \phi'_\varepsilon(u_k) |\nabla u_k|^p \xi^p dx$$

$$= - \int_{B_{2\rho}} |\nabla u_k|^{p-2} \nabla u_k \phi_\varepsilon(u_k) p \xi^{p-1} \nabla \xi dx.$$

Note that

$$\phi'_\varepsilon = \frac{\varepsilon}{(1 + |s|)^{1+\varepsilon}} \quad \text{and} \quad |\phi_\varepsilon| \leq 1$$

and that

$$(3.13) \quad ab \leq \sigma a^p + \sigma^{-\frac{p'}{p}} b^{p'}, \quad a > 0, \quad b > 0, \quad \sigma > 0.$$

We deduce from (3.12) that

$$(3.14) \quad \begin{aligned} & \int_{B_{2\rho}} \int_0^{u_k(x,t)} \phi_\varepsilon(s) ds \xi^p(x) dx + \frac{\varepsilon}{2} \int_{B_{2\rho} \times (0,t)} \frac{|\nabla u_k|^p \xi^p}{(1 + |u_k|)^{1+\varepsilon}} dx d\tau \\ & \leq \int_{B_{2\rho}} \int_0^{u_{0k}(x)} \phi_\varepsilon(s) ds \xi^p(x) dx \\ & \quad + \left(\frac{\varepsilon}{2}\right)^{1-p} \left(\frac{p}{\rho}\right)^p \int_{B_{2\rho} \times (0,t)} (1 + |u_k|)^{(1+\varepsilon)(p-1)} dx d\tau. \end{aligned}$$

Observe that  $\int_0^{u_k(x,t)} \phi_\varepsilon(s) ds \geq 0$  on  $\Sigma_T$ . Then select  $\varepsilon_0 > 0$  so that

$$(1 + \varepsilon_0)(p - 1) = 1.$$

It follows from (3.14) and (3.4) that there exists a  $c(\rho) > 0$  with

$$\int_{B_\rho \times (0,T)} \frac{|\nabla u_k|^p}{(1 + |u_k|)^{1+\varepsilon_0}} dx dt \leq c(\rho).$$

We estimate that

$$\begin{aligned} \int_{B_\rho \times (0,T)} |\nabla u_k|^{p-1} dx dt &= \int_{B_\rho \times (0,T)} \frac{|\nabla u_k|^{p-1}}{(1 + |u_k|)^{\frac{(1+\varepsilon_0)}{p'}}} (1 + |u_k|)^{\frac{(1+\varepsilon_0)}{p'}} dx dt \\ &\leq \frac{\varepsilon_0}{2} \int_{B_\rho \times (0,T)} \frac{|\nabla u_k|^p}{(1 + |u_k|)^{1+\varepsilon_0}} dx dt \\ &\quad + \left(\frac{\varepsilon_0}{2}\right)^{1-p} \int_{B_\rho \times (0,T)} (1 + |u_k|)^{(1+\varepsilon_0)(p-1)} dx dt. \end{aligned}$$

This implies (3.5). □

**Lemma 3.2.** For  $k \in \{1, 2, \dots\}$ , there hold

$$(3.15) \quad \int_{B_\rho \times (0,T)} \frac{1}{(1 + |u_k|)^{1+\varepsilon}} |\nabla u_k|^p dx dt \leq \frac{c(\rho)}{\varepsilon} \quad (\varepsilon > 0),$$

$$(3.16) \quad \int_{B_\rho \times (0,T) \cap \{|u_k| \leq M\}} |\nabla u_k|^p dxdt \leq Mc(\rho) \quad (M > 0)$$

for some  $c(\rho) > 0$ .

*Proof.* Let  $\rho > 0$  and  $\xi$  be a cut-off function associated with  $2\rho$  and  $\rho$ . Use  $\phi_\varepsilon(u_k)\xi$  as a test function in (3.3a) to obtain

$$\begin{aligned} & \int_{B_\rho \times (0,T)} \frac{\varepsilon}{(1 + |u_k|)^{1+\varepsilon}} |\nabla u_k|^p dxdt \\ & \leq \int_{B_{2\rho}} |u_0(x)| dx + \frac{1}{\rho} \int_{B_{2\rho} \times (0,T)} |\nabla u_k|^{p-1} dxdt. \end{aligned}$$

This, together with (3.5) implies (3.15). To see (3.16), for  $M > 0$  let  $P_M(s)$  be given as before. Then use  $P_M(u_k)\xi$  as a test function in (3.3a) to get

$$\int_{B_\rho \times (x,T)} P'_M(u_k) |\nabla u_k|^p dxdt \leq M \int_{B_{2\rho}} |u_0| dx + \frac{M}{\rho} \int_{B_{2\rho} \times (0,T)} |\nabla u_k|^{p-1} dxdt.$$

This completes the proof.  $\square$

**Lemma 3.3.** *There exists a subsequence of  $\{u_k\}$ , still denoted by  $\{u_k\}$ , and a function  $u \in L^1_{\text{loc}}(\mathbf{R}^N \times (0, T))$  with*

$$(3.17) \quad u_k \rightarrow u \text{ almost everywhere on } \Sigma_T.$$

*Proof.* Fix  $\rho > 0$ , and let  $\xi$  be given as in the proof of Lemma 3.2. We conclude from (3.3a) that

$$\begin{aligned} (3.18) \quad & \int_0^T \left( \frac{\partial}{\partial t} u_k, \frac{1}{1 + u_k^2} \xi \varphi \right) dt + \int_{B_{2\rho} \times (0,T)} |\nabla u_k|^{p-2} \nabla u_k \nabla \xi \varphi dxdt \\ & + \int_{B_{2\rho} \times (0,T)} \frac{1}{1 + u_k^2} |\nabla u_k|^{p-2} \nabla u_k \xi \nabla \varphi dxdt \\ & - \int_{B_{2\rho} \times (0,T)} \frac{2u_k}{(1 + u_k^2)^2} |\nabla u_k|^p \xi \varphi dxdt = 0 \end{aligned}$$

for all  $\varphi \in C_0^\infty(B_{2\rho} \times (0, T))$ . Here,  $(\cdot, \cdot)$  denotes the duality pairing between  $W^{-1,p'}(B_{2\rho})$  and  $W_0^{1,p}(B_{2\rho})$ . We infer from an argument in [X1] that

$$\left( \frac{\partial}{\partial t} u_k, \frac{1}{1 + u_k^2} \xi \varphi \right) = \left( \frac{\partial}{\partial t} (\xi \arctan u_k), \varphi \right) \quad \text{almost everywhere on } (0, T).$$

This, combined with (3.18) indicates that

$$\begin{aligned} (3.19) \quad & \frac{\partial}{\partial t} (\xi \arctan u_k) - \operatorname{div} \left( \frac{1}{1 + u_k^2} \xi |\nabla u_k|^{p-2} \nabla u_k \right) \\ & + |\nabla u_k|^{p-2} \nabla u_k \nabla \xi - \frac{2u_k}{(1 + u_k^2)^2} \xi |\nabla u_k|^p = 0 \\ & \text{in} \quad \mathcal{D}'(B_{2\rho} \times (0, T)). \end{aligned}$$

Now set

$$\begin{aligned} F_k &= \operatorname{div} \left( \frac{1}{1 + u_k^2} \xi |\nabla u_k|^{p-2} \nabla u_k \right), \\ G_k &= -|\nabla u_k|^{p-2} \nabla u_k \nabla \xi - \frac{2u_k}{(1 + u_k^2)^2} \xi |\nabla u_k|^p. \end{aligned}$$

It is easy to verify from (3.5) and (3.15) that

$$\begin{aligned} \{G_k\} &\text{ is bounded in } L^1(B_{2\rho} \times (0, T)), \\ \{F_k\} &\text{ is bounded in } L^{p'}(0, T; W^{-1,p'}(B_{2\rho})), \\ \{\xi \arctan u_k\} &\text{ is bounded in } L^p(0, T; W_0^{1,p}(B_{2\rho})). \end{aligned}$$

This puts us in a position to invoke Lemma 4.2 in [BM] to conclude that

$$\{\xi \arctan u_k\} \text{ is precompact in } L^p_{\text{loc}}(B_{2\rho} \times (0, T)).$$

In particular, we can extract a subsequence of  $\{u_k\}$ , still denoted by  $\{u_k\}$ , such that

$$\arctan u_k \text{ converges almost everywhere on } B_\rho \times (0, T).$$

Note that  $u_k = \tan(\arctan u_k)$ . We may define

$$u(x, t) = \lim_{k \rightarrow \infty} u_k(x, t) \quad \text{for almost everywhere } (x, t) \in B_\rho \times (0, T).$$

To conclude that  $\{u_k\}$  converges almost everywhere on  $B_\rho \times (0, T)$ , we must show that  $|u| < \infty$  almost everywhere on  $B_\rho \times (0, T)$ . However, this is an easy consequence of Fatou’s lemma and (3.4). Since  $\rho > 0$  is arbitrary, we can appeal to the classical diagonal argument to conclude the proof.  $\square$

**Lemma 3.4.** *There exists a subsequence of  $\{u_k\}$ , still denoted by  $\{u_k\}$ , and a measurable function  $F(x, t)$  on  $\Sigma_T$  such that*

$$(3.20) \quad \nabla u_k \rightarrow F \quad \text{almost everywhere on } \Sigma_T.$$

*Proof.* Fix  $\rho > 0$ , and let  $\xi$  be given as in the proof of Lemma 3.3. Assume (3.17) holds. According to Egorov's theorem, for each  $\eta > 0$  there exists a measurable set  $E_\eta \subset B_\rho \times (0, T)$  such that

$$|B_\rho \times (0, T) \setminus E_\eta| < \eta \quad \text{and} \quad u_k \rightarrow u \quad \text{uniformly on } E_\eta.$$

We may assume that  $\{u_k\}$  is bounded in  $L^\infty(E_\eta)$ , and thus by (3.16),

$$(3.21) \quad \int_{E_\eta} |\nabla u_k|^p dx dt \leq c(\eta, \rho).$$

For  $\delta > 0$ , we can find a  $K(\delta)$  with

$$(3.22) \quad |u_k - u_m| < \delta \quad \text{on } E_\eta \quad \text{for all } m, k > K(\delta).$$

Let  $P_\delta$  be defined as before. We can derive from (3.3a) that

$$\begin{aligned} & \frac{d}{dt} \int_{B_{2\rho}} \int_0^{u_k(x,t) - u_m(x,t)} P_\delta(s) ds \xi(x) dx + \\ & \int_{B_{2\rho}} \left( |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right) (\nabla u_k - \nabla u_m) \xi(x) P'_\delta(u_k - u_m) dx \\ & = - \int_{B_{2\rho}} \left( |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right) \nabla \xi(x) P_\delta(u_k - u_m) dx \\ & \leq \frac{\delta}{\rho} \int_{B_{2\rho}} \left( |\nabla u_k|^{p-1} + |\nabla u_m|^{p-1} \right) dx, \end{aligned}$$

for  $k, m$  sufficiently large. Thus,

$$\begin{aligned} (3.23) \quad & \int_{B_{2\rho} \times (0, T)} \left( |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right) \\ & \cdot (\nabla u_k - \nabla u_m) \xi(x) P'_\delta(u_k - u_m) dx dt \\ & \leq \int_{B_{2\rho}} \int_0^{u_{0k} - u_{0m}} P_\delta(s) ds dx + \frac{\delta}{\rho} \int_{B_{2\rho} \times (0, T)} \left( |\nabla u_k|^{p-1} + |\nabla u_m|^{p-1} \right) dx dt \\ & \leq c(\rho) \delta \end{aligned}$$

for  $k, m$  sufficiently large. We estimate, with the aid of (3.21), (3.22), and (3.23) that

$$(3.24) \quad \int_{E_\eta} |\nabla u_k - \nabla u_m|^p dx dt$$

$$\begin{aligned}
&= \int_{E_\eta} \frac{|\nabla u_k - \nabla u_m|^p}{(|\nabla u_k| + |\nabla u_m|)^{\frac{(2-p)p}{2}}} (|\nabla u_k| + |\nabla u_m|)^{\frac{(2-p)p}{2}} dxdt \\
&\leq \left( \int_{E_\eta} \frac{|\nabla u_k - \nabla u_m|^2}{(|\nabla u_m| + |\nabla u_k|)^{2-p}} dxdt \right)^{\frac{p}{2}} \left( \int_{E_\eta} (|\nabla u_m| + |\nabla u_k|)^p dxdt \right)^{\frac{(2-p)}{2}} \\
&\leq c(\eta, \rho) \left( \int_{B_{2\rho} \times (0, T)} (|\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m) \right. \\
&\quad \cdot (\nabla u_k - \nabla u_m) \xi(x) P'_\delta(u_k - u_m) dxdt \Big)^{\frac{p}{2}} \\
&\leq c_1(\eta, \rho) \delta^{\frac{p}{2}}
\end{aligned}$$

for  $k, m$  sufficiently large. We see that  $\{\nabla u_k\}$  is a Cauchy sequence in  $(L^p(E_\eta))^N$ . In particular, we can select a subsequence of  $\{u_k\}$ , still denoted by  $\{u_k\}$ , so that

$$\nabla u_k \quad \text{converges almost everywhere on } E_\eta.$$

This is true for each  $\eta > 0$ , and so  $\{\nabla u_k\}$  converges almost everywhere on  $B_\rho \times (0, T)$ . The lemma follows from the classical diagonal argument.  $\square$

**Lemma 3.5.**  $\{|\nabla u_k|^{p-2} \nabla u_k\}$  is precompact in  $L^1(B_\rho \times (0, T))$  for each  $\rho > 0$ .

*Proof.* Note that the function  $G(x) \equiv |x|^{p-2}x$  is continuous because  $\lim_{|x| \rightarrow 0} |x|^{p-2}x = 0 \equiv G(0)$ . Thus, we may assume that

$$(3.25) \quad \{|\nabla u_k|^{p-2} \nabla u_k\} \quad \text{converges almost everywhere on } B_\rho \times (0, T).$$

Now for each  $q \in \left(0, \frac{p}{2}\right)$ , we can choose  $\varepsilon_0 > 0$  so that  $q = \frac{1}{2 + \varepsilon_0}p$ . We deduce from (3.4) and (3.15) that

$$\begin{aligned}
(3.26) \quad &\int_{B_\rho \times (0, T)} |\nabla u_k|^q dxdt \\
&= \int_{B_\rho \times (0, T)} \frac{1}{(1 + |u_k|)^{(1+\varepsilon_0)\frac{q}{p}}} |\nabla u_k|^q (1 + |u_k|)^{(1+\varepsilon_0)\frac{q}{p}} dxdt \\
&\leq \left( \int_{B_\rho \times (0, T)} \frac{1}{(1 + |u_k|)^{1+\varepsilon_0}} |\nabla u_k|^p dxdt \right)^{\frac{q}{p}}
\end{aligned}$$

$$\begin{aligned} & \cdot \left( \int_{B_\rho \times (0, T)} (1 + |u_k|)^{(1+\varepsilon_0)\frac{q}{(p-q)}} dx dt \right)^{\frac{(p-q)}{p}} \\ & \leq c(\rho) \left( \int_{B_\rho \times (0, T)} (1 + |u_k|) dx dt \right)^{\frac{(p-q)}{p}} \leq c(\rho). \end{aligned}$$

Since  $0 < p - 1 < \frac{p}{2}$ , there exists a  $q \in \left(p - 1, \frac{p}{2}\right)$  such that

$$\int_{B_\rho \times (0, T)} |\nabla u_k|^q dx dt \leq c(\rho),$$

at least for  $k$  large enough. This implies that  $\{|\nabla u_k|^{p-2} \nabla u_k\}$  is uniformly integrable. This, in conjunction with (3.32) and Vitali's theorem, yields the lemma.  $\square$

**Lemma 3.6.**  *$\{u_k\}$  is precompact in  $C([0, T]; L^1(B_\rho))$  for each  $\rho > 0$ .*

*Proof.* For  $\delta > 0$  let

$$\theta_\delta(s) = \begin{cases} 1 & \text{if } s > \delta \\ s & \text{if } |s| < \delta \\ -1 & \text{if } s < -\delta \end{cases},$$

and  $\xi$  be given as in the proof of Lemma 3.2. We can conclude from (3.3a) that

$$\begin{aligned} (3.27) \quad & \int_{B_{2\rho}} \int_0^{u_k(x,t) - u_m(x,t)} \theta_\delta(s) ds \xi(x) dx \\ & + \int_{B_{2\rho} \times (0, t)} (|\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m) \\ & \cdot (\nabla u_k - \nabla u_m) \xi(x) \theta'_\delta(u_k - u_m) dx d\tau \\ & = \int_{B_{2\rho}} \int_0^{u_{0k}(x) - u_{0m}(x)} \theta_\delta(s) ds \xi(x) dx \\ & - \int_{B_{2\rho} \times (0, T)} (|\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m) \theta_\delta(u_k - u_m) \nabla \xi dx d\tau. \end{aligned}$$

Observe that the second integral in (3.27) is nonnegative. Hence, we obtain

$$\int_{B_\rho} |u_k(x, t) - u_m(x, t)| dx$$



$$\leq \int_{B_{2\rho}} |u_{0k} - u_{0m}| dx + \frac{1}{\rho} \int_{B_{2\rho} \times (0, T)} \left| |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right| dx dt.$$

Then the lemma follows from Lemma 3.5.  $\square$

**Lemma 3.7.** *Let  $E \subset \mathbf{R}^N \times (0, T)$  be bounded and measurable. Assume that there exists an  $M > 0$  such that*

$$|u_k| \leq M \quad \text{almost everywhere on } E \quad \text{for } k \text{ sufficiently large.}$$

*Then  $\{\nabla u_k\}$  is precompact in  $(L^p(E))^N$ .*

*Proof.* Let  $\rho > 0$  be such that

$$B_\rho \times (0, T) \supset E,$$

and let  $\xi$  be given as in the proof of Lemma 3.2. We conclude from (3.39) that

$$\begin{aligned} & \int_{B_{2\rho}} \xi(x) \int_0^{u_k(x,t) - u_m(x,t)} P_{2M}(s) ds dx \\ & + \int_{B_{2\rho} \times (0, T)} \left( |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right) \\ & \quad \cdot (\nabla u_k - \nabla u_m) P'_{2M}(u_k - u_m) \xi(x) dx d\tau \\ & = \int_{B_{2\rho}} \xi(x) \int^{u_{0k} - u_{0m}} P_{2M}(s) ds dx \\ & \quad - \int_{B_{2\rho} \times (0, T)} \left( |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right) P_{2M}(u_k - u_m) \nabla \xi(x) dx d\tau. \end{aligned}$$

Subsequently,

$$\begin{aligned} & \int_E \left( |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right) (\nabla u_k - \nabla u_m) dx d\tau \\ & \leq 2M \int_{B_{2\rho}} |u_{0k} - u_{0m}| dx \\ & \quad + \frac{2M}{\rho} \int_{B_{2\rho} \times (0, T)} \left| |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right| dx dt. \end{aligned}$$

A calculation similar to (3.24) yields

$$\begin{aligned} & \int_E |\nabla u_k - \nabla u_m|^p dx dt \\ & \leq c(M, \rho) \left( \int_{B_{2\rho}} |u_{0k} - u_{0m}| dx \right. \end{aligned}$$

$$+ \int_{B_{2\rho}} \left| |\nabla u_k|^{p-2} \nabla u_k - |\nabla u_m|^{p-2} \nabla u_m \right| dx dt \Big)^{\frac{p}{2}}.$$

This implies the desired result.  $\square$

Now we are ready to conclude the proof of Theorem 3.1. Let  $\{v_k\}$ ,  $\{u_k\}$  be given as before. Note from Lemma 2.3 that

$$\begin{aligned} v_k &\leq v_{k+1} && \text{on } \Sigma_T && \text{for all } k, \\ w_k &\geq w_{k+1} && \text{on } \Sigma_T && \text{for all } k. \end{aligned}$$

Define

$$\begin{aligned} v(x, t) &= \lim_{k \rightarrow \infty} v_k(x, t), \\ w(x, t) &= \lim_{k \rightarrow \infty} w_k(x, t). \end{aligned}$$

Consequently,

$$(3.28) \quad w \leq u_k \leq v \quad \text{almost everywhere.}$$

By a result in [DH], there holds

$$\int_s^T \int_{B_\rho} \frac{(z_t)^2}{(|z| + 1)^{1+\varepsilon}} dx dt \leq c(\varepsilon, s, p), T > s > 0, \varepsilon > 0, \rho > 0,$$

where  $z = w$  or  $v$ . The remaining proof is entirely similar to that in [X1]. The only difference is that in (3.23) of [X1] we require

$$\varphi \in C_0^\infty(\mathbf{R}^N \times (-\infty, T)).$$

This completes the proof.

**Acknowledgment:** This work was supported in part by the Arkansas Science & Technology Authority 94-B-18. The author would like to thank Prof. E. DiBenedetto for his interest in this work.

## References

- [BM] L. Boccardo and F. Murat, *Almost everywhere convergence of the gradients of solutions to elliptic and parabolic equations*, Nonlinear Anal., **19** (1992), 581–597.
- [BGDM] L. Boccardo, T. Gallouet, J.I. Diaz and F. Murat, *Existence and regularity of renormalized solutions for some elliptic problems involving derivatives of nonlinear terms*, J. Differential equations, to appear.
- [D] E. DiBenedetto, *Degenerate Parabolic Equations*, Springer-Verlag, New York, 1993.

- [DH] E. DiBenedetto and M.A. Herrero, *Non-negative solutions of the evolution  $p$ -Laplacian equation, initial traces, and Cauchy problem when  $1 < p < 2$* , Arch. Rat. Mech. Anal., **111** (1990), 225–290.
- [DL1] R.J. DiPerna and P.L. Lions, *Globale existence for the Fokker-Planck-Boltzmann equations*, Comm. Pure Appl. Math., **11** (1989), 729–758.
- [DL2] ———, *On the Cauchy problem for Boltzmann equations: globale existence and weak stability*, Ann. Math., **130** (1989), 321–366.
- [LSU] O.A. Ladyzenskaya, V.A. Solonnikov and N.N. Uralceva, *Linear and Quasilinear equations of Parabolic Type*, AMS, Rhode Island, 1968.
- [O] J.T. Oden, *Qualitative Methods in Nonlinear Mechanics*, Prentice-Hall, Inc., New Jersey, 1986.
- [S] J. Simon, *Compact sets in the space  $L^p(0, T, B)$* , Am. Mat. Pura Appl., **146** (1987), 65–96.
- [X1] X. Xu, *On the initial-boundary-value problem for  $u_t - \operatorname{div}(|\nabla u|^{p-2} \nabla u) = 0$* , Arch. Rational Mech. Anal., **127** (1994), 337–360.
- [X2] T. Kilpeläinen and X. Xu, *On the uniqueness problem for quasilinear elliptic equations involving measures*, Revista Mat. Iberoamericana, **12** (1996), to appear.

Received November 23, 1993.

MISSISSIPPI STATE UNIVERSITY  
 MISSISSIPPI STATE, MS 39762  
*E-mail address:* xxu@math.msstate.edu



# PACIFIC JOURNAL OF MATHEMATICS

Volume 174

No. 1

May 1996

---

|   |     |
|---|-----|
| A distance formula for algebras on the disk   | 1   |
| CHRISTOPHER J. BISHOP   |     |
| Rigidity of isotropic maps  | 29  |
| FERNANDO CUKIERMAN  |     |
| The Schwartz space of a general semisimple Lie group. V. Schwartz<br>class wave packets                           | 43  |
| REBECCA A. HERB   |     |
| Rational polynomials with a $C^*$ -fiber  | 141 |
| SHULIM KALIMAN  |     |
| Linear combinations of logarithmic derivatives of entire functions with<br>applications to differential equations | 195 |
| JOSEPH B. MILES and JOHN ROSSI  |     |
| Factorization problems in the invertible group of a homogeneous<br>$C^*$ -algebra                                 | 215 |
| N. CHRISTOPHER PHILLIPS   |     |
| Higher order estimates in complex interpolation theory  | 247 |
| RICHARD ROCHBERG  |     |
| Braid commutators and Vassiliev invariants  | 269 |
| TED STANFORD  |     |
| On the Cauchy problem for a singular parabolic equation   | 277 |
| XIANGSHENG XU   |     |