

*Pacific
Journal of
Mathematics*

**DIMENSIONS OF NILPOTENT ALGEBRAS OVER FIELDS OF
PRIME CHARACTERISTIC**

CORA M. STACK

DIMENSIONS OF NILPOTENT ALGEBRAS OVER FIELDS OF PRIME CHARACTERISTIC

CORA STACK

In this short paper we consider the conjecture that for a finite dimensional commutative nilpotent algebra M over a perfect field of prime characteristic p , $\dim M \geq p \dim M^{(p)}$ where $M^{(p)}$ is the subalgebra of M generated by $x^p, x \in M$. We prove that for any finite dimensional nilpotent algebra M (not necessarily commutative) over any field of prime characteristic p , $\dim M \geq p \dim M^{(p)}$ for $\dim M^{(p)} \leq 2$.

1. Introduction.

It is generally accepted that the structure of nilpotent algebras is not well understood. In this paper we discuss some questions which could lead to significant further development of this structure theory.

Let M be an algebra over a field K . The algebra M is said to be nilpotent if $M^n = 0$ for some $n \geq 1$. (Recall that for $j \geq 1$, M^j is the subalgebra of M generated by all monomials of degree j in the elements of M .) If n is the least such integer, then n is called the nilpotency index of M . If $M^j \neq 0$, then it follows from the nilpotency of M that $M^j \supset M^{j+1}$. Let $d_i = \dim_K(M^i/M^{i+1})$, where $\dim_K V$ or $\dim V$ denotes of course the dimension of V as a vector space over K . Then clearly $\dim M = \sum_{i=1}^{n-1} d_i$.

In [1] N. Eggert conjectures that for a finite commutative nilpotent algebra over a field K of characteristic p ,

$$(1.1) \quad \dim_K M \geq p \dim_K M^{(p)}$$

where $M^{(p)}$ is the subalgebra of M generated by $x^p, x \in M$. In [1] Eggert proves (1.1) when $\dim M^{(p)} \leq 2$. His proof however does not appear to be easy to generalise. In view of this we provide a simpler proof in Section 2. Since these proofs do not require the nilpotent algebra to be finite, it is believed that all that is required in (1.1) is that M be finite dimensional.

Now (1.1) obviously holds when $\dim M^{(p)} = 0$. If $\dim M^{(p)} = 1$ then it follows from the remarks above, that $\dim M \geq \sum_{i=1}^p d_i \geq p$, and so (1.1) holds in this case. In the next section we prove that for any finite dimensional nilpotent algebra M over any field of characteristic p , $\dim M \geq p \dim M^{(p)}$ if $\dim M^{(p)} = 2$. This yields a second proof of Eggert's theorem. Our proof however does not require the field K to be perfect, nor indeed the algebra M to be commutative, and in this sense will be a slight extension.

2. The Two Dimensional Case.

Some Notation. Let V be a vectorspace over a field K of characteristic p . If $X \subseteq V$ by $\langle X \rangle$ we mean the linear span (over K) of X . If X is a singleton say $\{x\}$ we write Kx for $\langle X \rangle$. Let $N^{(p)}$ denote the subspace of M generated by x^p , $x \in M$. Of course if M is commutative, then $M^{(p)} = N^{(p)}$.

We now prove the following theorem, of which our main theorem will be just a simple corollary.

Theorem 1. *Let M be any nilpotent algebra over the field K of prime characteristic, and let $d_k = 1$ for some $1 \leq k \leq p-1$, then if $N^{(p)} = \langle z^p, z \in M \rangle$, then $N^{(p)} = \langle y^{p^i}, i \geq 1 \rangle$ for some $y \in M$.*

Proof. If $M^p = 0$, then $N^{(p)} = 0$ and there is nothing to prove. Hence we may assume that $M^p \neq 0$. Since $d_k = 1$, there exists $m = m_1 m_2 \dots m_k$ for which $\langle m + M^{k+1} \rangle = M^k / M^{k+1}$. Write $m = y\sigma$ where $y = m_1 \in M$ and $\sigma = m_2 \dots m_k \in M^{k-1}$ (if $k = 1$, regard σ as an empty product). Now

$$(2.1) \quad M^k = Ky\sigma + M^{k+1},$$

and so using the algebra properties of M , $yM^k \subseteq Ky^2\sigma + M^{k+2}$. Consequently since $M^{k+1} = M^k M$ we have that

$$(2.2) \quad M^{k+1} = Ky\sigma M + M^{k+2} \subseteq yM^k + M^{k+2} \subseteq Ky^2\sigma + M^{k+2}.$$

Assuming $d_k = 1$, choose now $1 \leq i \leq k$ maximal with $M^k = Ky^i\sigma + M^{k+1}$, with $\sigma = a_1 a_2 \dots a_{k-i}$ a product of $k-i$ elements of M . From (2.1) it is clear that such an i exists. We now show that $i = k$. If not, then σ is not an empty product and by applying (2.2) to this formula we get that $M^{k+1} = Ky^{i+1}\sigma + M^{k+2}$. But $M^{k+1} \neq M^{k+2}$, since otherwise $M^{k+1} = 0$ and $k+1 \leq p$ would imply that $M^p = 0$ - a contradiction. Hence $y^{i+1}\sigma \notin M^{k+2}$, and therefore $y^{i+1}a_1 \dots a_{k-i-1} \notin M^{k+1}$. But then $y^{i+1}a_1 \dots a_{k-i-1} \in M^k \setminus M^{k+1}$, and so since $d_k = 1$, $M^k = Ky^{i+1}a_1 a_2 \dots a_{k-i-1} + M^{k+1}$, contradicting the maximality of i . Thus $i = k$ and $M^k = Ky^k + M^{k+1}$.

Now equation (2.2) and induction imply that $M^{k+j} = Ky^{k+j} + M^{k+j+1}$ for all $j \geq 0$ and hence $M^{k+j} = \langle y^{k+j}, y^{k+j+1}, \dots \rangle$. Since $k \leq p-1$, we have in particular that

$$(2.3) \quad M^{p-1} = \langle y^{p-1}, y^p, y^{p+1}, \dots \rangle.$$

Now let $Y = \langle y, y^2, \dots \rangle$. If $p = 2$ then by (2.3), $M = Y$. Hence M is commutative and $N^{(p)} = \langle y^p, y^{2p}, \dots \rangle$ as required. We may assume therefore that $p \geq 3$. Let $A = \text{Annih}_M(y^{p-2})$. The map $m \rightarrow my^{p-2}$ (of additive

groups) sends M into M^{p-1} , has kernel A and by (2.3) maps Y onto M . It is easy to verify that this forces $Y + A = M$.

We next claim that any product of the elements of Y and A which involves p factors, with at least one from A must equal 0. For consider the first A term in such a product $y_1 \dots y_j a m_1 \dots m_t$ where $j + t = p - 1$, $y_i \in Y$ and $a_i \in A$. Notice that $y_2 \dots y_j a m_1 \dots m_y \in M^{p-1}$. Since Y clearly centralises M^{p-1} , we may slide the factor y_1 above to the far right. Thus $x = y_2 \dots y_j a m_1 \dots m_t y_1$, and continuing in this manner we get that $x = a m_1 \dots m_t y_1 \dots y_j \in a M^{p-1} = 0$. But then $x = 0$ proving our claim.

Finally, if z is an arbitrary element of M then since $M = Y + A$, we may write $z = b + a$, ($b \in Y$, $a \in A$.) Then $z^p = b^p +$ a sum of terms involving p factors of Y^s and of A^s with at least one A factor occurring. Thus, by the above this last sum is zero and so $z^p = b^p$ and so is a K -linear sum of the y^{p^i} 's, $i \geq 1$. Thus $N^{(p)} \subseteq \langle y^{p^i}, i \geq 1 \rangle$ completing the proof of the theorem. □

We now have our main theorem as a corollary of this:

Theorem 2. *Let M be a finite dimensional nilpotent algebra over a field K of characteristic p , and let $M^{(p)}$ be the subalgebra of M generated by the p^{th} powers of elements of M . If $\dim M^{(p)} = 2$ then $\dim M \geq 2p$.*

Proof. Assume the theorem is false and let M be a counter example of least dimension. Then $\dim M < 2p$ and so $M^{2p} = 0$. In particular we have that $m^n n^p = 0$ for every $m, n \in M$ and so $M^{(p)}$ will just be the linear span of elements of the form $x^p, x \in M$, i.e $M^{(p)} = N^{(p)}$. Since $\dim M < 2p$ and $\dim M^p \geq 2$, it follows that there exists $k, 1 \leq k \leq p - 1$ for which $d_k = 1$. But then, by Theorem 1, $M^{(p)} = N^{(p)} = \langle y^{p^i}, i \geq 1 \rangle$. But since $M^{2p} = 0$, we must have that $M^{(p)} = K y^p$, and so $\dim M^{(p)} \leq 1$ a contradiction. This completes the proof of the theorem. □

We remark that the nonnecessity of the assumptions of commutativity and perfectness of the field in our previous theorem suggests perhaps (and believed to be the case by the author) and (2.1) may hold for finite dimensional noncommutative nilpotent algebras and perhaps also over fields of prime characteristic.

Acknowledgement. The author would like to thank Dr. B. Corbas and the referee for suggesting a number of improvements to this paper.

References

[1] N. Eggert, *Quasi Regular Groups of Finite Commutative Nilpotent Algebras.*

- [2] I. Kaplansky, *Fields and Rings*, Chicago Lectures in Mathematics, The University of Chicago Press, Chicago and London, 1969.
- [3] N.H. McCoy, *The Theory of Rings*, The MacMillan company, New York, 1966.
- [4] B.R. Mac Donald, *Finite Rings with identity*, Marcel Dekker, New York, 1974.
- [5] D. Passman, *The Algebraic Structure of Group Rings*, John Wiley and Sons, New York and London, 1977.
- [6] Zariski and Samuel, *Commutative Algebra*, D. Van Nostrand and company, 1 (1966).

Received July 5, 1994 and revised December 15, 1994.

UNIVERSITY COLLEGE GALWAY
GALWAY, IRELAND
E-mail address: 0003998S@bodkin.ucg.ie

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by

E. F. Beckenbach (1906-1982)

F. Wolf (1904-1989)

EDITORS

Sun-Yung A. Chang (Managing Editor)
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

F. Michael Christ
University of California
Los Angeles, CA 90095-1555
christ@math.ucla.edu

Nicholas Ercolani
University of Arizona
Tucson, AZ 85721
ercolani@math.arizona.edu

Robert Finn
Stanford University
Stanford, CA 94305
finn@gauss.stanford.edu

Steven Kerckhoff
Stanford University
Stanford, CA 94305
spk@gauss.stanford.edu

Martin Scharlemann
University of California
Santa Barbara, CA 93106
mgscharl@math.ucsb.edu

Gang Tian
Massachusetts Institute of Technology
Cambridge, MA 02139
tian@math.mit.edu

V. S. Varadarajan
University of California
Los Angeles, CA 90095-1555
vsv@math.ucla.edu

Dan Voiculescu
University of California
Berkeley, CA 94720
dvv@math.berkeley.edu

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIF. INST. OF TECHNOLOGY
CHINESE UNIV. OF HONG KONG
HONG KONG UNIV. OF SCI. & TECH.
KEIO UNIVERSITY
MACQUARIE UNIVERSITY
MATH. SCI. RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.
PEKING UNIVERSITY
RITSUMEIKAN UNIVERSITY
STANFORD UNIVERSITY

TOKYO INSTITUTE OF TECHNOLOGY
UNIVERSIDAD DE LOS ANDES
UNIV. OF ARIZONA
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIF., BERKELEY
UNIV. OF CALIF., DAVIS
UNIV. OF CALIF., IRVINE
UNIV. OF CALIF., LOS ANGELES
UNIV. OF CALIF., RIVERSIDE
UNIV. OF CALIF., SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF HAWAII
UNIV. OF MELBOURNE
UNIV. OF MONTANA
UNIV. NACIONAL AUTONOMA DE MEXICO
UNIV. OF NEVADA, RENO
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

The supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

Manuscripts must be prepared in accordance with the instructions provided on the inside back cover.

The table of contents and the abstracts of the papers in the current issue, as well as other information about the Pacific Journal of Mathematics, may be found on the Internet at <http://www.math.uci.edu/pjm.html>.

The *Pacific Journal of Mathematics* (ISSN 0030-8730) is published monthly except for July and August. Regular subscription rate: \$245.00 a year (10 issues). Special rate: \$123.00 a year to individual members of supporting institutions.

Subscriptions, back issues published within the last three years and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. Prior back issues are obtainable from Kraus Periodicals Co., Route 100, Millwood, NY 10546.

The Pacific Journal of Mathematics at the University of California, c/o Department of Mathematics, 981 Evans Hall, Berkeley, CA 94720 (ISSN 0030-8730) is published monthly except for July and August. Second-class postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 6143, Berkeley, CA 94704-0163.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS at University of California,
Berkeley, CA 94720, A NON-PROFIT CORPORATION

This publication was typeset using AMS-LATEX,
the American Mathematical Society's TEX macro system.
Copyright © 1995 by Pacific Journal of Mathematics

PACIFIC JOURNAL OF MATHEMATICS

Volume 176 No. 1 November 1996

Moduli spaces of isometric pluriharmonic immersions of Kähler manifolds into indefinite Euclidean spaces	1
HITOSHI FURUHATA	
On a theorem of Koch	15
FARSHID HAJIR	
Degree-one maps onto lens spaces	19
CLAUDE HAYAT-LEGRAND, SHICHENG WANG and HEINER ZIESCHANG	
Unitary representation induced from maximal parabolic subgroups for split F_4	33
CHENG CHON HU	
New constructions of models for link invariants	71
FRANÇOIS JAEGER	
Solvability of Dirichlet problems for semilinear elliptic equations on certain domains	117
ZHIREN JIN	
Hadamard-Frankel type theorems for manifolds with partially positive curvature	129
KATSUEI KENMOTSU and CHANGYU XIA	
Boundary behavior of the Bergman curvature in strictly pseudoconvex polyhedral domains	141
KANG-TAE KIM and JIYE YU	
Existence and behavior of the radial limits of a bounded capillary surface at a corner	165
KIRK LANCASTER and DAVID SIEGEL	
Triangle subgroups of hyperbolic tetrahedral groups	195
COLIN MACLACHLAN	
Chern classes of vector bundles on arithmetic varieties	205
TOHRU NAKASHIMA and YUICHIRO TAKEDA	
Haar measure on $E_q(2)$	217
ARUP KUMAR PAL	
Domains of partial attraction in noncommutative probability	235
VITTORINO PATA	
Partitioning products of $\mathcal{P}(\omega)/\text{fin}$	249
OTMAR SPINAS	
Dimensions of nilpotent algebras over fields of prime characteristic	263
CORA M. STACK	
Tensor products of structures with interpolation	267
FRIEDRICH WEHRUNG	
Fourier multipliers for $L_p(\mathbb{R}^n)$ via q -variation	287
QUANHUA XU	