

*Pacific
Journal of
Mathematics*

BOTT FORMULA OF THE MASLOV-TYPE INDEX
THEORY

YIMING LONG

Volume 187 No. 1

January 1999

BOTT FORMULA OF THE MASLOV-TYPE INDEX THEORY

YIMING LONG

In this paper the integer valued ω -index theory parameterized by all ω on the unit circle for paths in the symplectic group $\text{Sp}(2n)$ is established. Based on this index theory, the Bott formula of the Maslov-type index theory for iterated paths in $\text{Sp}(2n)$ is established, the mean index for periodic solutions of Hamiltonian systems is defined, and the increasing estimate of the iterated Maslov-type index in terms of the mean index is proved.

1. Introduction and main results.

For $n \in \mathbf{N}$, the set of natural numbers, as usual we define the symplectic group

$$\text{Sp}(2n) \equiv \text{Sp}(2n, \mathbf{R}) = \{ M \in \mathcal{L}(\mathbf{R}^{2n}) \mid M^T J M = J \},$$

where $J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$, I_n is the identity matrix on \mathbf{R}^n , and $\mathcal{L}(\mathbf{R}^{2n})$ is the space of $2n \times 2n$ real matrices. When there is no confusion, we shall omit the subindex of the identity matrices. The metric on $\text{Sp}(2n)$ is inherited through $\mathcal{L}(\mathbf{R}^{2n})$ from that of \mathbf{R}^{4n^2} . For $\tau > 0$, suppose $H \in C^2(S_\tau \times \mathbf{R}^{2n}, \mathbf{R})$, where $S_\tau \equiv \mathbf{R}/(\tau\mathbf{Z})$. Let x be a τ -periodic solution of the nonlinear Hamiltonian system

$$(1.1) \quad \dot{x}(t) = JH'(t, x(t)), \quad x \in \mathbf{R}^{2n}.$$

Then the fundamental solution of the linearized Hamiltonian system

$$(1.2) \quad \dot{y} = JB(t)y, \quad y \in \mathbf{R}^{2n},$$

with $B(t) = H''(t, x(t))$, is a path $\gamma_x \in C^1([0, +\infty), \text{Sp}(2n))$ with $\gamma_x(0) = I$. In order to study the property of periodic solutions of (1.1), C. Conley and E. Zehnder defined their index theory for nondegenerate paths in $\text{Sp}(2n)$ starting from the identity matrix in their celebrated work [CZ] for the case of $n \geq 2$. This index theory was extended by E. Zehnder and the author in [LZ] to nondegenerate paths in $\text{Sp}(2)$, by the author in [Lo1] and C. Viterbo in [Vi2] independently via different methods to degenerate paths which are fundamental solutions of linear Hamiltonian systems (1.2) for any $B \in C(S_\tau, \mathcal{L}_s(\mathbf{R}^{2n}))$, where $\mathcal{L}_s(\mathbf{R}^{2n})$ is the set of all real symmetric

$2n \times 2n$ matrices, and recently by the author in [Lo8] to any degenerate paths in $\text{Sp}(2n)$ using ideas of [Lo1] and [Lo5] together with an axiom characterization of this index theory. In this paper we call it the Maslov-type index theory. Denote by

$$\mathcal{P}_\tau(2n) = \{\gamma \in C([0, \tau], \text{Sp}(2n)) \mid \gamma(0) = I\}.$$

This index theory assigns a pair of integers $(i_\tau(\gamma), \nu_\tau(\gamma)) \in \mathbf{Z} \times \{0, \dots, 2n\}$ to each $\gamma \in \mathcal{P}_\tau(2n)$. For any $\gamma \in \mathcal{P}_\tau(2n)$, define the iteration path $\tilde{\gamma} \in C([0, +\infty), \text{Sp}(2n))$ of γ by

$$\tilde{\gamma}(t) = \gamma(t - j\tau)\gamma(\tau)^j, \quad \text{for } j\tau \leq t \leq (j + 1)\tau \text{ and } j \in \{0\} \cup \mathbf{N}.$$

Let $\gamma^k = \tilde{\gamma}|_{[0, k\tau]}$. Then we can associate to γ through $\tilde{\gamma}$ a sequence of Maslov-type indices $\{(i_{k\tau}(\tilde{\gamma}), \nu_{k\tau}(\tilde{\gamma}))\}_{k \in \mathbf{N}}$. When $\gamma : [0, +\infty) \rightarrow \text{Sp}(2n)$ is the fundamental solution of (1.2) with $B \in C(S_\tau, \mathcal{L}_s(\mathbf{R}^{2n}))$, there holds $(\gamma|_{[0, \tau]})^\sim = \gamma$. Thus for $B(t) = H''(t, x(t))$ this index sequence with $\gamma = \gamma_x$ reflects important properties of the τ -periodic solution x of (1.1).

In order to gain more information on the iterated Maslov-type indices, in Section 2 of this paper, the Maslov-type index theory is generalized to the ω -index theory for all $\omega \in \mathbf{U}$, where \mathbf{U} is the unit circle in the complex plane \mathbf{C} . This ω -index theory assigns a pair of integers $(i_{\tau, \omega}(\gamma), \nu_{\tau, \omega}(\gamma)) \in \mathbf{Z} \times \{0, \dots, 2n\}$ to each $\gamma \in \mathcal{P}_\tau(2n)$. When $\omega = 1$, the ω -index theory coincides with the above mentioned Maslov-type index theory. The following are our main results on the ω -index theory in this paper.

Definition 1.1. For any $M \in \text{Sp}(2n)$, define the **homotopy set** of M in $\text{Sp}(2n)$ by

$$\begin{aligned} \Omega(M) &= \{N \in \text{Sp}(2n) \mid \sigma(N) \cap \mathbf{U} = \sigma(M) \cap \mathbf{U}, \text{ and} \\ &\quad \dim_{\mathbf{C}} \ker_{\mathbf{C}}(N - \lambda I) = \dim_{\mathbf{C}} \ker_{\mathbf{C}}(M - \lambda I), \forall \lambda \in \sigma(M) \cap \mathbf{U}\}. \end{aligned}$$

We denote by $\Omega^0(M)$ the path connected component of $\Omega(M)$ which contains M , and call it the **homotopy component** of M in $\text{Sp}(2n)$. For $\omega \in \mathbf{U}$, define

$$\Omega_\omega(M) = \{N \in \text{Sp}(2n) \mid \dim_{\mathbf{C}} \ker_{\mathbf{C}}(N - \omega I) = \dim_{\mathbf{C}} \ker_{\mathbf{C}}(M - \omega I)\}.$$

We denote by $\Omega_\omega^0(M)$ the path connected component of $\Omega_\omega(M)$ which contains M , and call it the ω -**homotopy component** of M in $\text{Sp}(2n)$.

For any $M \in \text{Sp}(2n)$, define $[M] = \{N \in \text{Sp}(2n) \mid N = P^{-1}MP \text{ for some } P \in \text{Sp}(2n)\}$. Then $[M] \subset \Omega^0(M) \subset \Omega_\omega^0(M)$ for all $\omega \in \mathbf{U}$.

Theorem 1.2. For any $\tau > 0$, $\gamma \in \mathcal{P}_\tau(2n)$, and $\omega \in \mathbf{U}$, the ω -index pair $(i_{\tau, \omega}(\gamma), \nu_{\tau, \omega}(\gamma))$ of γ is completely determined by the homotopy class of γ in $\mathcal{P}_\tau(2n)$ in the sense of Definition 2.2 below. More specially, $i_{\tau, \omega}(\gamma)$ is completely determined by the ω -homotopy component $\Omega_\omega^0(\gamma(\tau))$ of $\gamma(\tau)$ in

$\mathrm{Sp}(2n)$ up to an additive constant, and $\nu_{\tau,\omega}(\gamma)$ is completely determined by $\Omega_\omega^0(\gamma(\tau))$.

The idea of our ω -index theory is closely related to those of [CZ], [LZ], [Lo1, Lo2, Lo5, Lo8], and [DL], but is very different from those of [Bo], [Ed], [CD], [Ek], and [Vil]. Our method is rather elementary and intuitive. The establishment of the ω -index theory and the proof of Theorem 1.2 are based on the understanding of the topological structures of the singular hypersurface $\mathrm{Sp}(2n)_\omega^0$, which contains all symplectic matrices possessing the eigenvalue ω , and its complement $\mathrm{Sp}(2n)_\omega^*$ in $\mathrm{Sp}(2n)$ obtained in the papers [CZ], [LZ], [Lo1]-[Lo6], and [LD]. Fix $\omega \in \mathbf{U}$. For any $\gamma \in \mathcal{P}_\tau(2n)$ we define $\nu_{\tau,\omega}(\gamma)$ to be $\dim_{\mathbf{C}} \ker_{\mathbf{C}}(\gamma(\tau) - \omega I)$. Since $\mathrm{Sp}(2n)$ is homeomorphic to a product of \mathbf{U} and a simply connected space, any path $\gamma \in \mathcal{P}_\tau(2n)$ can be viewed as rotating around the unit circle part in $\mathrm{Sp}(2n)$. Because $\mathrm{Sp}(2n)_\omega^*$ possesses precisely two path connected components and $\mathrm{Sp}(2n)_\omega^*$ is simply connected in $\mathrm{Sp}(2n)$, we can give a way to count this rotation suitably for every non-degenerate path γ , i.e. $\omega \notin \sigma(\gamma(\tau))$, which defines the ω -index $i_{\tau,\omega}(\gamma) \in \mathbf{Z}$. For any degenerate path $\gamma \in \mathcal{P}_\tau(2n)$, i.e. $\omega \in \sigma(\gamma(\tau))$, using the topology of $\mathrm{Sp}(2n)$ near $\mathrm{Sp}(2n)_\omega^0$, we construct topologically two families of non-degenerate paths γ_{-s} and γ_s near γ for $s \in (0, 1]$ satisfying $i_{\tau,\omega}(\gamma_s) - i_{\tau,\omega}(\gamma_{-s}) = \nu_{\tau,\omega}(\gamma)$. We also prove analytically in Section 4 that γ_{-s} and γ_s takes the minimum and maximum values of $i_{\tau,\omega}$ among all non-degenerate path in $\mathcal{P}_\tau(2n)$ near γ . Then we define this minimum to be $i_{\tau,\omega}(\gamma)$. In Theorem 2.11 below, we give an axiom characterization of the ω -index theory as an integer valued map defined on $\cup_{n \geq 1} \mathcal{P}_\tau(2n)$ via the homotopy invariance, symplectic additivity, and the values of it on $\mathcal{P}_\tau(2)$.

In Section 4 we introduce the concept of splitting numbers:

Theorem 1.3. *For any $M \in \mathrm{Sp}(2n)$ and $\omega \in \mathbf{U}$, choose $\tau > 0$ and $\gamma \in \mathcal{P}_\tau(2n)$ with $\gamma(\tau) = M$, and define*

$$(1.3) \quad S_M^\pm(\omega) = \lim_{\epsilon \rightarrow 0^+} i_{\tau, \exp(\pm \epsilon \sqrt{-1})\omega}(\gamma) - i_{\tau,\omega}(\gamma).$$

*Then these two integers are independent of the choice of the path γ . They are called the **splitting numbers** of M at ω .*

We also prove that the splitting numbers coincide with the **ultimate type** of $\omega \in \mathbf{U}$ for M , a new concept introduced in the following Definition 4.8. Specially we prove that splitting numbers of M at ω are constants on $\Omega^0(M)$, and give a complete characterization of splitting numbers.

Based on the ω -index theory and the idea of R. Bott in [Bo], in Section 5 we are able to establish the following Bott-type formulae of the Maslov-type index theory for iterations of any path in $\mathcal{P}_\tau(2n)$.

Theorem 1.4. *For any $\tau > 0$, $\gamma \in \mathcal{P}_\tau(2n)$, and $k \in \mathbf{N}$, there hold*

$$(1.4) \quad i_{k\tau}(\tilde{\gamma}) = i_{k\tau,1}(\tilde{\gamma}) = \sum_{\omega^k=1} i_{\tau,\omega}(\gamma),$$

$$(1.5) \quad \nu_{k\tau}(\tilde{\gamma}) = \nu_{k\tau,1}(\tilde{\gamma}) = \sum_{\omega^k=1} \nu_{\tau,\omega}(\gamma).$$

As a direct consequence of Theorem 1.4, the mean index per τ for any path $\gamma \in \mathcal{P}_\tau(2n)$ can be defined and an estimate on $i_{k\tau}$ is obtained.

Theorem 1.5. *For any $\tau > 0$ and $\gamma \in \mathcal{P}_\tau(2n)$ there hold*

$$(1.6) \quad \hat{i}_\tau(\gamma) \equiv \lim_{k \rightarrow +\infty} \frac{i_{k\tau}(\tilde{\gamma})}{k} = \frac{1}{2\pi} \int_{\mathbf{U}} i_{\tau,\omega}(\gamma) d\omega,$$

$$(1.7) \quad \hat{\nu}_\tau(\gamma) \equiv \lim_{k \rightarrow +\infty} \frac{\nu_{k\tau}(\tilde{\gamma})}{k} = \frac{1}{2\pi} \int_{\mathbf{U}} \nu_{\tau,\omega}(\gamma) d\omega = 0.$$

Specially, $\hat{i}_\tau(\gamma)$ is always a finite real number, which is called the **mean index** per τ of γ .

Theorem 1.6. *For any $\tau > 0$ and $\gamma \in \mathcal{P}_\tau(2n)$, there holds*

$$(1.8) \quad |k\hat{i}_\tau(\gamma) - i_{k\tau}(\tilde{\gamma})| \leq n, \quad \forall k \in \mathbf{N}.$$

Then through the fundamental solution γ_x of (1.2) with $B(t) = H''(t, x(t))$ the mean index per period τ of a τ -periodic solution x of the nonlinear system (1.1) can be defined by $\hat{i}_\tau(x) = \hat{i}_\tau(\gamma_x)$. When τ is the minimal period of x , we denote by $\hat{i}(x) = \hat{i}_\tau(x)$. This yields a new invariant to each periodic solution of the system (1.1).

Since Bott's celebrated work [Bo] in 1956, iteration properties of various Morse and Maslov type index theories have been studied by [Ed], [CD], [KI], [Ek], [Vi1], [DL], [Lo5], and others. But upto the author's knowledge, the Bott-type formula were established only by H. Edwards for the Sturm form with finite Morse indices (cf. [Ed]), by I. Ekeland (cf. [Ek]) for convex Hamiltonian systems, and by C. Viterbo (cf. [Vi1]) for star-shaped Hamiltonian systems under certain nondegeneracy conditions. In Section 6, we prove that our ω -index theory generalizes the Bott functions $\Lambda(\omega)$ and $N(\omega)$ defined in [Bo] and Ekeland's index functions defined in [Ek]. Therefore from Theorem 1.4, we recover their formulae for iterations. Our results also generalizes those in [Vi1]. Results obtained in the current paper have been applied to the study of the existence, multiplicity, and stability of periodic solutions of nonconvex Hamiltonian systems in our other papers.

In Section 7, for reders convenience, we briefly introduce results obtained in [LD] on the basic normal forms of symplectic matrices, in [Lo6] on the topological structures of the hypersurface $\text{Sp}(2n)_\omega^0$ and its complement in $\text{Sp}(2n)$, and study properties of homotopy components of symplectic matrices, which are fundamental for our discussion in this paper.

2. The ω -index theory.

In this section, we define the ω -index theory as a generalization of the Maslov-type index theory, and study its basic properties. In the following we fix $n \in \mathbf{N}$ and use notations introduced in Section 7.

For any $\tau > 0$, we define the metric of $\mathcal{P}_\tau(2n)$ to be the one induced from the usual metric of $C([0, \tau], \text{Sp}(2n))$. For $\omega \in \mathbf{U}$, we define two subsets of $\mathcal{P}_\tau(2n)$ by

$$\mathcal{P}_{\tau,\omega}^*(2n) = \{\gamma \in \mathcal{P}_\tau(2n) \mid \gamma(\tau) \in \text{Sp}(2n)_\omega^*\}, \quad \mathcal{P}_{\tau,\omega}^0(2n) = \mathcal{P}_\tau(2n) \setminus \mathcal{P}_{\tau,\omega}^*(2n).$$

We define the following subfamilies of paths in $\mathcal{P}_\tau(2n)$ which are fundamental solutions of Hamiltonian systems (1.2) with continuous symmetric and τ -periodic coefficients.

$$\begin{aligned} \hat{\mathcal{P}}_\tau(2n) &= \{\gamma \in \mathcal{P}_\tau(2n) \cap C^1([0, \tau], \text{Sp}(2n)) \mid \dot{\gamma}(\tau) = \dot{\gamma}(0)\gamma(\tau)\}, \\ \hat{\mathcal{P}}_{\tau,\omega}^*(2n) &= \hat{\mathcal{P}}_\tau(2n) \cap \mathcal{P}_{\tau,\omega}^*(2n), \quad \hat{\mathcal{P}}_{\tau,\omega}^0(2n) = \hat{\mathcal{P}}_\tau(2n) \setminus \hat{\mathcal{P}}_{\tau,\omega}^*(2n). \end{aligned}$$

Let the metric of $\hat{\mathcal{P}}_\tau(2n)$ be the one induced from the usual metric of $C^1([0, \tau], \text{Sp}(2n))$. Note that for any $\gamma \in \mathcal{P}_\tau(2n)$, we can choose a C^1 -path $\beta \in \mathcal{P}_\tau(2n)$ possessing the same end points as γ and as C^0 -close to γ as we want. Now composing β with a suitable function $\rho \in C^2([0, \tau], [0, 1])$ satisfying $\rho(0) = 0$, $\dot{\rho}(0) = 0$, $\rho(\tau) = 1$, $\dot{\rho}(\tau) = 0$, and $\dot{\rho}(t) \geq 0$ for all $0 \leq t \leq \tau$, without loss of generality, we can assume $\beta \in \hat{\mathcal{P}}_\tau(2n)$. Thus $\hat{\mathcal{P}}_\tau(2n)$ is dense in $\mathcal{P}_\tau(2n)$.

Definition 2.1. For any $\tau > 0$ and $\gamma \in \mathcal{P}_\tau(2n)$, we define

$$(2.1) \quad \nu_{\tau,\omega}(\gamma) = \dim_{\mathbf{C}} \ker_{\mathbf{C}}(\gamma(\tau) - \omega I), \quad \forall \omega \in \mathbf{U}.$$

Definition 2.2 (cf. [Lol]). For $\tau > 0$ and $\omega \in \mathbf{U}$, given two paths γ_0 and $\gamma_1 \in \mathcal{P}_\tau(2n)$, if there exists a map $\delta \in C([0, 1] \times [0, \tau], \text{Sp}(2n))$ such that $\delta(0, \cdot) = \gamma_0(\cdot)$, $\delta(1, \cdot) = \gamma_1(\cdot)$, $\delta(s, 0) = I$, and $\nu_{\tau,\omega}(\delta(s, \cdot))$ is constant for $0 \leq s \leq 1$, then γ_0 and γ_1 are **ω -homotopic on $[0, \tau]$ along $\delta(\cdot, \tau)$** and we write $\gamma_0 \sim_\omega \gamma_1$. If $\gamma_0 \sim_\omega \gamma_1$ on $[0, \tau]$ along $\delta(\cdot, \tau)$ for all $\omega \in \mathbf{U}$, then γ_0 and γ_1 are **homotopic on $[0, \tau]$ along $\delta(\cdot, \tau)$** and we write $\gamma_0 \sim \gamma_1$.

As well known, every $M \in \text{Sp}(2n)$ has its unique polar decomposition $M = AU$, where $A = (MM^T)^{1/2}$ is symmetric, positive definite, and symplectic, U is orthogonal and symplectic. Therefore U has the form

$$U = \begin{pmatrix} u_1 & -u_2 \\ u_2 & u_1 \end{pmatrix}, \quad \text{where } u = u_1 + \sqrt{-1}u_2 \in \mathcal{L}(\mathbf{C}^n) \text{ is a unitary matrix.}$$

So for every path $\gamma \in \mathcal{P}_\tau(2n)$ we can associate to γ a path $u : [0, \tau] \rightarrow U(n)$ in the unitary group $U(n)$. If $\Delta(t)$ is any continuous real function satisfying $\det u(t) = \exp(\sqrt{-1}\Delta(t))$, the difference $\Delta(\tau) - \Delta(0)$ depends only on γ but

not on the choice of the function $\Delta(t)$. Therefore we may define the rotation number of γ on $[0, \tau]$ by

$$(2.2) \quad \Delta_\tau(\gamma) = \Delta(\tau) - \Delta(0) \in \mathbf{R}.$$

Note that $\Delta_\tau : \mathcal{P}_\tau(2n) \rightarrow \mathbf{R}$ is continuous.

Lemma 2.3. *If γ_0 and $\gamma_1 \in \mathcal{P}_\tau(2n)$ possess common end point $\gamma_0(\tau) = \gamma_1(\tau)$, then $\Delta_\tau(\gamma_0) = \Delta_\tau(\gamma_1)$ if and only if $\gamma_0 \sim_\omega \gamma_1$ on $[0, \tau]$ for some $\omega \in \mathbf{U}$, if and only if $\gamma_0 \sim \gamma_1$ on $[0, \tau]$.*

This is a well known result as mentioned in [YS]. It can also be proved by using the function $\rho_n : \text{Sp}(2n) \rightarrow \mathbf{U}$ of [SZ] and the topological structure of the symplectic groups.

For any $\gamma \in \mathcal{P}_{\tau,\omega}^*(2n)$, by Theorem 7.1 we can connect $\gamma(\tau)$ to M_n^- or M_n^+ by a path β within $\text{Sp}(2n)_\omega^*$ and get a product path $\beta * \gamma$ defined by

$$\beta * \gamma(t) = \begin{cases} \gamma(2t), & \text{if } 0 \leq t \leq \frac{\tau}{2}, \\ \beta(2t - \tau), & \text{if } \frac{\tau}{2} \leq t \leq \tau. \end{cases}$$

Define

$$(2.3) \quad k \equiv \frac{1}{\pi} \Delta_\tau(\beta * \gamma).$$

Then k is an integer, and by Theorem 7.1, it is independent of the choice of the path β . In this case we write $\gamma \in \mathcal{P}_{\tau,\omega}^*(2n, k)$. These $\mathcal{P}_{\tau,\omega}^*(2n, k)$'s give a homotopy classification of $\mathcal{P}_{\tau,\omega}^*(2n)$.

Definition 2.4. We define

$$(2.4) \quad i_{\tau,\omega}(\gamma) = k, \quad \text{if } \gamma \in \mathcal{P}_{\tau,\omega}^*(2n, k).$$

Thus for $\beta, \gamma \in \mathcal{P}_{\tau,\omega}^*(2n)$, $i_{\tau,\omega}(\beta) = i_{\tau,\omega}(\gamma)$ if and only if $\beta \sim_\omega \gamma$.

Next we study the degenerate paths. Fix $\omega \in \mathbf{U}$ and $\gamma \in \mathcal{P}_{\tau,\omega}^0(2n)$. Using Theorem 7.5, there exists $P \in \text{Sp}(2n)$ such that (7.8) holds for $M = \gamma(\tau)$. Let $\Sigma_0(\omega, \gamma(\tau))$ be the subset of $\text{Sp}(2n)_\omega^0$ which contains all matrices M satisfying

$$\dim_{\mathbf{C}} \ker_{\mathbf{C}}(M - \omega I) > \nu_{\tau,\omega}(\gamma).$$

Let $\Sigma_1(\omega, \gamma(\tau))$ be the path connected component of $\text{Sp}(2n)_\omega^0 \setminus \Sigma_0(\omega, \gamma(\tau))$ which contains $\gamma(\tau)$. For $\epsilon > 0$, let $B_\epsilon(\gamma(\tau))$ be the open ball in $\text{Sp}(2n)$ centered at $\overline{\gamma(\tau)}$ with radius ϵ . Choose $\epsilon > 0$ to be sufficiently small so that $\overline{B_\epsilon(\gamma(\tau))}$ is contractible and possesses no intersection with $\text{Sp}(2n)_\omega^0 \setminus \Sigma_1(\omega, \gamma(\tau))$.

Let $\theta_0 \in (0, \frac{\pi}{8n})$ and the integers $\{m_1, \dots, m_{p+2q}\}$ be the numbers defined in (7.9). For $s_i \in [-1, 1]$ with $1 \leq i \leq p + 2q$, we define

$$(2.5) \quad Q(s_1, \dots, s_{p+2q}) \equiv \gamma(\tau)P^{-1}R_{m_1}(s_1\theta_0) \cdots R_{m_{p+2q}}(s_{p+2q}\theta_0)P.$$

Then for all $s_i \in [-1, 1] \setminus \{0\}$ with $1 \leq i \leq p + 2q$, there hold

$$\begin{cases} Q(s_1, \dots, s_{p+2q}) \in \mathrm{Sp}(2n)_\omega^* \cap B_\epsilon(\gamma(\tau)), \\ Q(s_1, \dots, s_{p+2q})P^{-1}R_{m_k}(-s_k\theta_0)P \in \mathrm{Sp}(2n)_\omega^0, & 1 \leq k \leq p + 2q, \\ \dim_{\mathbf{C}} \ker_{\mathbf{C}}(Q(s_1, \dots, s_{p+2q})P^{-1}R_{m_k}(-s_k\theta_0)P - \omega I) = a_k, & 1 \leq k \leq p + 2q, \end{cases}$$

where the constant $a_k = 1$ or 2 .

For $t_0 \in (0, \tau)$, let $\rho \in C^2([0, \tau], [0, 1])$ such that $\rho(t) = 0$ for $0 \leq t \leq t_0$, $\rho(t) \geq 0$ for $0 \leq t \leq \tau$, $\rho(\tau) = 1$, and $\dot{\rho}(\tau) = 0$. As in [Lo1], whenever $t_0 \in (0, \tau)$ is sufficiently close to τ , there holds $\gamma([t_0, \tau]) \subset B_\epsilon(\gamma(\tau))$. For any $(s, t) \in [-1, 1] \times [0, \tau]$, we define the rotational perturbation paths

$$(2.6) \quad \gamma_s(t) = \gamma(t)P^{-1}R_{m_1}(s\rho(t)\theta_0) \cdots R_{m_{p+2q}}(s\rho(t)\theta_0)P.$$

They satisfy $\gamma_s \in \mathcal{P}_{\tau, \omega}^*(2n)$ for $s \neq 0$, $\gamma_0 = \gamma$, $\gamma_s(t) = \gamma(t)$ for $0 \leq t \leq t_0$, $\gamma_{\pm 1}(\tau) = Q(\pm 1, \dots, \pm 1)$, γ_s is continuous in $s \in [-1, 1]$ and converges to γ in C^0 as $s \rightarrow 0$, and

$$(2.7) \quad \begin{cases} \gamma_s(t) \in B_\epsilon(\gamma(\tau)), & \forall (s, t) \in [0, 1] \times [t_0, \tau], \\ \nu_{\tau, \omega}(\gamma_s) = 0, & \forall s \in [-1, 1] \setminus \{0\}, \\ i_{\tau, \omega}(\gamma_s) = i_{\tau, \omega}(\gamma_{s'}), & \forall s, s' \in [-1, 1] \text{ with } ss' > 0. \end{cases}$$

Note that if $\gamma \in \hat{\mathcal{P}}_{\tau, \omega}^0(2n)$, we have $\gamma_s \in \hat{\mathcal{P}}_{\tau, \omega}^*(2n)$ for $s \neq 0$ and $\gamma_s \rightarrow \gamma$ as $s \rightarrow 0$ in the topology of $\hat{\mathcal{P}}_\tau(2n)$.

Theorem 2.5. *For $\gamma \in \mathcal{P}_{\tau, \omega}^0(2n)$, the rotational perturbation paths of γ defined above by (2.6) satisfy*

$$(2.8) \quad i_{\tau, \omega}(\gamma_1) - i_{\tau, \omega}(\gamma_{-1}) = \nu_{\tau, \omega}(\gamma).$$

Proof. Without loss of generality, we assume $\tau = 1$. Note that this theorem has been proved in [Lo8] for $\omega = 1$, and the case of $\omega = -1$ follows directly from the case of $\omega = 1$ by the fact $D_{-1}(M) = D_1(-M)$ for all $M \in \mathrm{Sp}(2n)$ with $D_{\pm 1}$ defined in Section 7. Here we only prove the theorem for $\omega \in \mathbf{U} \setminus \mathbf{R}$. Fix $\omega \in \mathbf{U} \setminus \mathbf{R}$, $\epsilon > 0$, and $\gamma \in \mathcal{P}_{1, \omega}^0(2n)$. Define γ_s for $s \in [-1, 1]$ by (2.6). In the following we fix an $s \in (0, 1]$.

For $0 \leq k \leq p + 2q$ define

$$(2.9) \quad \alpha_{s, k}(t) = \gamma(t)\gamma(1)^{-1}Q(\mu_1(s, t), \dots, \mu_{p+2q}(s, t)), \quad \forall t \in [0, 1],$$

where $\mu_i(s, t) = \rho(t)s$ for $1 \leq i \leq k$ and $\mu_i(s, t) = -\rho(t)s$ for $k + 1 \leq i \leq p + 2q$. Then by a direct verification, we have $\alpha_{s, k} \in \mathcal{P}_{1, \omega}^*(2n)$ for $0 \leq k \leq p + 2q$, $\alpha_{s, 0} = \gamma_{-s}$, $\alpha_{s, p+2q} = \gamma_s$. We obtain $\nu_{1, \omega}(\gamma) = \sum_{k=1}^{p+2q} a_k$. Assume the following equalities hold,

$$(2.10) \quad i_{1, \omega}(\alpha_{s, k}) - i_{1, \omega}(\alpha_{s, k-1}) = a_k, \quad \text{for } 1 \leq k \leq p + 2q.$$

Summing (2.10) up from $k = 1$ to $k = p + 2q$, by Remark 7.4 and Theorem 7.5 we obtain (2.8). Therefore the proof of Theorem 2.5 is reduced to that of (2.10). Write α_k for $\alpha_{s,k}$ for notational simplicity.

Fix a $k \in \{1, \dots, p + 2q\}$. Let $\eta(t) = \alpha_{k-1}(1)P^{-1}R_{m_k}(2ts\theta_0)P$ for $t \in [0, 1]$. Then by definition we have $\eta(0) = \alpha_{k-1}(1)$, $\eta(1) = \alpha_k(1)$, $\eta(1/2) = \alpha_k(1)P^{-1}R_{m_k}(-s\theta_0)P \in \text{Sp}(2n)_\omega^0$, $\dim \ker(\eta(1/2) - \omega I) = c_k$, and $\eta(t) \in \text{Sp}(2n)_\omega^*$ for $t \in [0, 1] \setminus \{1/2\}$. By definition we obtain that $\alpha_k|_{[t_0, 1]}$ is homotopic to $\eta * \alpha_{k-1}|_{[t_0, 1]}$ with fixed end points. Therefore we obtain $\alpha_k \sim \eta * \alpha_{k-1}$ with fixed end points. Then (2.10) becomes

$$(2.11) \quad i_{1,\omega}(\eta * \alpha_{k-1}) - i_{1,\omega}(\alpha_{k-1}) = a_k.$$

When $a_k = 2$, (2.11) follows from the corresponding proof for Lemma 3 of [Lo1] or Theorem 3.2 of [Lo8] with the aid of Lemma 7.10. When $a_k = 1$, (2.11) follows from the corresponding proof of Theorem 3.2 in [Lo8] with the aid of Theorem 7.2. Then the proof is complete. \square

Theorem 2.6. *For $\gamma \in \mathcal{P}_{\tau,\omega}^0(2n)$, there exists $\epsilon > 0$ sufficiently small such that for any $\alpha \in \mathcal{P}_{\tau,\omega}^*(2n) \cap B_\epsilon^0(\gamma)$ there holds*

$$(2.12) \quad i_{\tau,\omega}(\gamma_{-s}) \leq i_{\tau,\omega}(\alpha) \leq i_{\tau,\omega}(\gamma_s),$$

where the paths γ_{-s} and γ_s are defined by (2.6) for $s \in (0, 1]$, $B_\epsilon^0(\gamma)$ is the open ball centered at γ with radius ϵ in $\mathcal{P}_1(2n)$.

The proof of the Theorem 2.6 is postponed to Section 3. Then Theorem 2.6 yields:

Corollary 2.7. *For $\gamma \in \mathcal{P}_{\tau,\omega}^0(2n)$ and $s \in (0, 1]$, the perturbation paths of γ defined by (2.6) satisfy*

$$(2.13) \quad i_{\tau,\omega}(\gamma_{-s}) = \inf\{i_{\tau,\omega}(\alpha) \mid \alpha \in \mathcal{P}_\tau(2n) \text{ is } C^0 \text{ sufficiently close to } \gamma\}.$$

Specially for $s, s' \in [-1, 1] \setminus \{0\}$ there holds

$$(2.14) \quad i_{\tau,\omega}(\gamma_s) = i_{\tau,\omega}(\gamma_{s'}) \quad \text{if } ss' > 0.$$

Definition 2.8. For any $\gamma \in \mathcal{P}_{\tau,\omega}^0(2n)$, we define

$$(2.15) \quad i_{\tau,\omega}(\gamma) = i_{\tau,\omega}(\gamma_{-s}), \quad \forall 0 < s \leq 1.$$

Proposition 2.9. *Definition 2.8 of $i_{\tau,\omega}(\gamma)$ for any $\gamma \in \mathcal{P}_{\tau,\omega}^0(2n)$ is well defined.*

Proof. Without loss of generality, suppose $\tau = 1$. Fix an $\omega \in \mathbf{U}$, a path $\gamma \in \mathcal{P}_{1,\omega}^0(2n)$, and an $s \in (0, 1]$. Define $\zeta_s(t) = Q(-st, \dots, -st)$ for $t \in [0, 1]$ by (2.5). Then $\zeta_s(1) = \gamma_{-s}(1)$. Let $\psi_{-s} : [0, 1] \rightarrow \text{Sp}(2n)_\omega^*$ be a path connecting $\gamma_{-s}(1)$ to M_n^+ or M_n^- . Then the path γ_{-s} is homotopic to the

joint path $\zeta_s * \gamma$. Thus these two paths possess the same rotation numbers. Then we obtain

$$(2.16) \quad i_{1,\omega}(\gamma_s) = \Delta_1(\psi_{-s} * \zeta_s * \gamma)/\pi = \Delta_1(\psi_{-s})/\pi + \Delta_1(\zeta_s)/\pi + \Delta_1(\gamma)/\pi.$$

Thus $i_{1,\omega}(\gamma)$ is well defined. □

Definition 2.10. For every path $\gamma \in \mathcal{P}_\tau(2n)$, Definitions 2.1, 2.4, and 2.8 assign a pair of integers to γ :

$$(i_{\tau,\omega}(\gamma), \nu_{\tau,\omega}(\gamma)) \in \mathbf{Z} \times \{0, \dots, 2n\}.$$

This pair of integers is called the ω -**index** of γ . We call $i_{\tau,\omega}(\gamma)$ the ω -**rotation index** of γ and $\nu_{\tau,\omega}(\gamma)$ the ω -**nullity** of γ .

Using \diamond -product defined in Section 7, for any paths $\gamma_j \in \mathcal{P}_\tau(2n_j)$ with $j = 0$ and 1, define $\gamma_0 \diamond \gamma_1(t) = \gamma_0(t) \diamond \gamma_1(t)$ for all $t \in [0, \tau]$. For any $\tau > 0$, $n \in \mathbf{N}$, and $k \in \mathbf{Z}$ using notations in Section 7 we define a sequence of **zigzag standard paths** $\alpha_{n,k,\tau}$ in $\mathcal{P}_\tau(2n)$ as follows.

$$\begin{aligned} \phi_{\theta,\tau}(t) &= R(\theta t/\tau), \quad \forall t \in [0, \tau], \\ \alpha_{1,0,\tau}(t) &= D(1 + t/\tau), \quad \forall t \in [0, \tau], \\ \alpha_{n,k,\tau} &= [(D(2)\phi_{k\pi,\tau}) * \hat{\alpha}_{1,0,\tau}] \diamond (\hat{\alpha}_{1,0,\tau})^{\diamond(n-1)}. \end{aligned}$$

Then there hold $\alpha_{n,k,\tau} \in \mathcal{P}_{\tau,\omega}^*(2n)$, $i_{\tau,\omega}(\alpha_{n,k,\tau}) = k$ for all $\omega \in \mathbf{U}$ and $\alpha_{n,k,\tau}(\tau) = M_n^\pm$ if $(-1)^k = \pm 1$. The following theorem gives a characterization of our ω -index theory.

Theorem 2.11. For any $\tau > 0$ and $\omega = \exp(\theta\sqrt{-1}) \in \mathbf{U}$, there exists a unique function $i_{\tau,\omega} : \cup_{n \in \mathbf{N}} \mathcal{P}_\tau(2n) \rightarrow \mathbf{Z}$ satisfying the following five axioms:

1° (**Homotopy invariant**). For γ_0 and $\gamma_1 \in \mathcal{P}_\tau(2n)$, if $\gamma_0 \sim_\omega \gamma_1$ on $[0, \tau]$, then

$$i_{\tau,\omega}(\gamma_0) = i_{\tau,\omega}(\gamma_1).$$

2° (**Symplectic additivity**). For any $\gamma_i \in \mathcal{P}_\tau(2n_i)$ with $i = 0$ and 1, there holds

$$i_{\tau,\omega}(\gamma_0 \diamond \gamma_1) = i_{\tau,\omega}(\gamma_0) + i_{\tau,\omega}(\gamma_1).$$

3° (**Clockwise continuity**). For any $\gamma \in \mathcal{P}_\tau(2n)$ and $\omega \in \mathbf{U}$ satisfying $\gamma(\tau) = N_1(\omega, B)$ defined in Section 7, there exists a $\theta_0 > 0$ such that

$$i_{\tau,\omega}((\gamma(\tau)\phi_{-\theta,\tau}) * \gamma) = i_{\tau,\omega}(\gamma), \quad \forall 0 < \theta \leq \theta_0.$$

4° (**Counterclockwise jumping**). For any $\gamma \in \mathcal{P}_\tau(2n)$ and $\omega \in \mathbf{U}$ satisfying $\gamma(\tau) = N_1(\omega, B)$ defined in Section 7, there exists a $\theta_0 > 0$ such that

$$i_{\tau,\omega}((\gamma(\tau)\phi_{\theta,\tau}) * \gamma) = i_{\tau,\omega}(\gamma) + 1, \quad \forall 0 < \theta \leq \theta_0.$$

5° (**Normality**). There holds

$$i_{\tau,\omega}(\alpha_{1,0,\tau}) = 0.$$

Proof. When $\omega = 1$, this theorem is proved in [Lo8]. From the fact $D_{-1}(M) = D_1(-M)$ for all $M \in \text{Sp}(2n)$, the case of $\omega = -1$ follows from that of $\omega = 1$. In the following we fix an $\omega \in \mathbf{U} \setminus \mathbf{R}$ with $\omega = \exp(\theta\sqrt{-1})$ and only prove the theorem for this case. Without loss of generality we suppose $\tau = 1$. By our above discussions, similar to the proof in [Lo8], the index function $i_{1,\omega} : \cup_{n \in \mathbf{N}} \mathcal{P}_1(2n) \rightarrow \mathbf{Z}$ defined by Definition 2.10 satisfies these five axioms.

Suppose we have another index function $\mu : \cup_{n \in \mathbf{N}} \mathcal{P}_1(2n) \rightarrow \mathbf{Z}$ satisfying these five conditions. It suffices to prove $\mu = i_{1,\omega}$ on $\mathcal{P}_1(2n)$. Let $M = \gamma(1)$ in Corollary 7.11, we obtain a path $f \in \Omega_\omega^0(\gamma(1))$ with $f(0) = \gamma(1)$ and $f(1)$ is given by (7.14). Choose $\gamma_j \in \mathcal{P}_1(2)$ with $\gamma_j = M_j$ for $0 \leq j \leq n$ such that $\sum_{j=1}^n i_{1,\omega}(\gamma_j) = i_{1,\omega}(\gamma)$. Then by 2°, it suffices to prove $\mu = i_{1,\omega}$ on every path γ in $\mathcal{P}_1(2)$ with $\gamma(1)$ possessing the form $R(\theta)$, $R(2\pi - \theta)$, $D(2)$ or $D(-2)$. Thus by 3°, the value of μ on $\mathcal{P}_1(2n)$ is completely determined by the value of μ on $\mathcal{P}_{1,\omega}^*(2)$. By 1° these values are completely determined by μ on the standard paths $\alpha_{1,k,1}$'s. By 3°, 4° and 5° we obtain $\mu(\alpha_{1,k,1}) = k$ for all $k \in \mathbf{Z}$. Thus $\mu(\gamma) = i_{1,\omega}(\gamma)$ holds for every $\gamma \in \mathcal{P}_1(2n)$. This completes the proof. □

Remark 2.12. It is easy to construct examples to show that the five axioms in Theorem 2.11 are independent from each other. Theorem 1.2 is a direct consequences of Theorem 2.11. Note that when $\omega = 1$ the ω -index theory defined above coincides with the Maslov-type index theory defined in [CZ], [LZ], [Lo1], [Lo8], and [Vi2]. Thus when $\omega = 1$, we simply write $(i_\tau(\gamma), \nu_\tau(\gamma)) = (i_{\tau,1}(\gamma), \nu_{\tau,1}(\gamma))$.

Theorem 2.13. (Inverse homotopy invariant.) *For any two paths γ_0 and $\gamma_1 \in \mathcal{P}_\tau(2n)$ with $i_{\tau,\omega}(\gamma_0) = i_{\tau,\omega}(\gamma_1)$, suppose that there exists a continuous path $h : [0, 1] \rightarrow \text{Sp}(2n)$ such that $h(0) = \gamma_0(\tau)$, $h(1) = \gamma_1(\tau)$, and $\dim_{\mathbf{C}} \ker_{\mathbf{C}}(h(s) - \omega I) = \nu_{\tau,\omega}(\gamma_0)$ for all $s \in [0, 1]$. Then $\gamma_0 \sim_\omega \gamma_1$ on $[0, \tau]$ along h .*

The proof of this theorem is similar to that of Theorem 6.4 in [Lo8] and thus is omitted.

Remark 2.14. For any end point free curve $f \in C([a, b], \text{Sp}(2n))$, choose $\gamma \in \mathcal{P}_1(2n)$ so that $\gamma(1) = f(a)$. Define $\eta(t) = \gamma(2t)$ for $t \in [0, 1/2]$, and $\eta(t) = f(a + (2t - 1)(b - a))$ for $t \in [1/2, 1]$. Define

$$i_\omega(f) = i_{1,\omega}(\eta) - i_{1,\omega}(\gamma).$$

By our above study of the index $i_{\tau,\omega}$ it is easy to see that $i_\omega(f)$ only depends on f itself. Similar to Theorem 2.11, an axiom characterization of this index can be given.

3. Morse indices and ω -indices.

In this section we study the relationship between the ω -indices of the linear Hamiltonian system (1.2) and the Morse indices of the corresponding functionals defined on the truncated loop space, and prove Theorem 2.6.

For $\tau > 0$, define L_τ to be the Hilbert space $L^2([0, \tau], \mathbf{C}^{2n})$ with the usual L^2 inner product $\langle \cdot, \cdot \rangle_{L_\tau}$, and $E_\tau = W^{1,2}(S_\tau, \mathbf{C}^{2n})$ with the usual $W^{1,2}$ norm as a subspace of L_τ . For $\omega = \exp(\theta\sqrt{-1}) \in \mathbf{U}$, define E_τ^ω to be the subspace $\{y \in E_\tau \mid y(\tau) = \omega y(0)\}$ of L_τ . Any $y \in E_\tau^\omega$ has the form

$$y(t) = \sum_{k \in \mathbf{Z}} e^{\sqrt{-1}(\theta+2k\pi)t/\tau} \xi_k, \quad \xi_k \in \mathbf{C}^{2n},$$

where there holds

$$\sum_{k \in \mathbf{Z}} ((\theta + 2k\pi)^2 + 1) |\xi_k|^2 < +\infty.$$

Define $E_{\tau,k}^\omega = E_{\tau,k}^{\omega,+} \oplus E_{\tau,k}^{\omega,-}$ with

$$E_{\tau,k}^{\omega,\pm} = e^{\sqrt{-1}(\theta+2k\pi)t/\tau} (J \pm \sqrt{-1}I) \mathbf{R}^{2n}.$$

Then there holds $E_\tau^\omega = \bigoplus E_{\tau,k}^\omega$.

Define $A = -J \frac{d}{dt} : E_\tau^\omega \rightarrow L_\tau$. Then A is continuous and symmetric, i.e. it satisfies

$$(3.1) \quad \langle Ax, y \rangle_{L_\tau} = \langle x, Ay \rangle_{L_\tau}, \quad \forall x, y \in E_\tau^\omega.$$

Viewing A as from the subspace E_τ^ω of L_τ to L_τ , we have

$$\sigma(A) = \{\lambda_k^\pm \mid k \in \mathbf{Z}\}, \quad \lambda_k^\pm = \pm \frac{\theta + 2k\pi}{\tau},$$

each eigenvalue λ_k^\pm of A has multiplicity $2n$ and the corresponding eigenspace is $E_{\tau,k}^{\omega,\pm}$.

Given $B \in C(S_\tau, \mathcal{L}_s(\mathbf{R}^{2n})) \subset C(S_\tau, \mathcal{L}_s(\mathbf{C}^{2n}))$, it induces a symmetric operator on L_τ by

$$(3.2) \quad \langle Bx, y \rangle_{L_\tau} = \int_0^\tau B(t)x(t) \cdot \overline{y(t)} dt, \quad \forall x, y \in L_\tau.$$

We consider the linear Hamiltonian system (1.2) for $y \in E_\tau^\omega$. Then τ -periodic solutions of (1.2) are one to one correspondent to critical points of the functional

$$(3.3) \quad f_{\tau,\omega}(y) = \frac{1}{2} \langle (A - B)y, y \rangle_{L_\tau}, \quad \forall y \in E_\tau^\omega \subset L_\tau.$$

Then $f_{\tau,\omega} : E_\tau^\omega \rightarrow \mathbf{R}$ is smooth in the topology of L_τ .

Using the saddle point reduction method (cf. [Ch]) we obtain a subspace

$$(3.4) \quad Z_\tau^\omega = \bigoplus_{|m| \leq m_0} E_{\tau,k}^\omega$$

with a sufficiently large $m_0 \in \mathbf{N}$ (cf. Section 4.2.1 of [Ch]), an injection map $u_{\tau,\omega} \in C^\infty(Z_\tau^\omega, L_\tau)$, and a smooth functional $a_{\tau,\omega} \in C^\infty(Z_\tau^\omega, \mathbf{R})$ defined by

$$(3.5) \quad a_{\tau,\omega}(z) = f_{\tau,\omega}(u_{\tau,\omega}(z)), \quad \forall z \in Z_\tau^\omega.$$

Note that the origin of Z_τ^ω as a critical point of $a_{\tau,\omega}$ corresponds to the origin of E_τ^ω as a critical point of $f_{\tau,\omega}$. Denote by $m_{\tau,\omega}^*$ for $* = +, 0,$ and $-$, the positive, null, and negative Morse indices of the functional $a_{\tau,\omega}$ at the origin respectively. Denote by $2d_{\tau,\omega} = \dim_{\mathbf{C}} Z_\tau^\omega$.

Theorem 3.1. *Let $\gamma \in \hat{\mathcal{P}}_\tau(2n)$ be the fundamental solution of (1.2) with $B(t)$ given above. For any $\omega \in \mathbf{U}$, there hold*

$$(3.6) \quad m_{\tau,\omega}^- = d_{\tau,\omega} + i_{\tau,\omega}(\gamma),$$

$$(3.7) \quad m_{\tau,\omega}^0 = \nu_{\tau,\omega}(\gamma),$$

$$(3.8) \quad m_{\tau,\omega}^+ = d_{\tau,\omega} - i_{\tau,\omega}(\gamma) - \nu_{\tau,\omega}(\gamma).$$

Proof. For the case of $\omega = 1$, this theorem was proved in [CZ], [LZ], [Lo1], and [Lo8]. The case of $\omega = -1$ can be proved similarly and is left to the readers. Thus we fix an $\omega \in \mathbf{U} \setminus \mathbf{R}$ and prove the theorem for this case only. Without loss of generality, we suppose $\tau = 1$.

Step 1. The case of $\gamma \in \hat{\mathcal{P}}_{\tau,\omega}^*(2n)$.

For $k \in \mathbf{Z}$ define

$$(3.9) \quad B_k = (k\pi I_2) \diamond O_{2n-2}, \quad \beta_k(t) = \exp(tk\pi J) \diamond I_{2n-2},$$

where O_m is the $m \times m$ zero matrix. Then $\beta_k \sim_\omega \hat{\alpha}_{n,k,1}$ for all $k \in \mathbf{Z}$. Therefore there exists a $k \in \mathbf{Z}$ such that $\beta_k \sim_\omega \gamma$. By an analogue of Lemma 4.2 of [Lo8] for Z_1^ω and Definition 2.4, it suffices to prove (3.6)-(3.8) for $\gamma = \beta_k$ with any $k \in \mathbf{Z}$.

Fix $k \in \mathbf{Z}$. Define $B(t) \equiv B_k$ of (3.9) in (1.2). Since B_k is a symmetric constant matrix, the corresponding functional $a_{\tau,\omega}$ satisfies (cf. Theorem 3.2.1 of [Lo4])

$$(3.10) \quad a_{\tau,\omega}(z) = \frac{1}{2} \langle (A - B_k)z, z \rangle_{L_\tau}, \quad \forall z \in Z^\omega.$$

Thus $a_{\tau,\omega}''(0) = A - B_k$. Denote its Morse indices by m_k^* for $* = +, 0,$ and $-$. We obtain (3.7) by the definition of β_k .

In order to verify (3.6) and (3.8), we consider properties of $(A - B_k)|_{E_{\tau,m}^\omega}$. Write $\omega = e^{\sqrt{-1}\theta}$ with $\theta \in (0, \pi) \cup (\pi, 2\pi)$. For any $m \in \mathbf{Z}$, let $\lambda_m = \theta + 2m\pi$. For $\zeta^\pm = (\xi^\pm, \eta^\pm) \in \mathbf{R}^2 \times \mathbf{R}^{2n-2}$ we have

$$z^\pm = e^{\sqrt{-1}\lambda_m t} (J \pm \sqrt{-1}I) \zeta^\pm \in E_m^{\omega,\pm}.$$

Then there holds

$$(3.11) \quad \begin{aligned} & \langle (A - B_k)(z^+ + z^-), (z^+ + z^-) \rangle_{L_\tau} \\ &= 2(\lambda_m - k\pi)|\xi^+|^2 + 2\lambda_m|\eta^+|^2 - 2(\lambda_m + k\pi)|\xi^-|^2 - 2\lambda_m|\eta^-|^2. \end{aligned}$$

Thus $A - B_k$ possesses the following four eigenvalues:

$$(3.12) \quad \begin{cases} \Lambda_1 = 2(\lambda_m - k\pi) = 4\pi(m + \frac{\theta}{2\pi} - \frac{k}{2}), \\ \Lambda_2 = 2\lambda_m = 4\pi(m + \frac{\theta}{2\pi}), \\ \Lambda_3 = -2(\lambda_m + k\pi) = -4\pi(m + \frac{\theta}{2\pi} + \frac{k}{2}), \\ \Lambda_4 = -2\lambda_m = -4\pi(m + \frac{\theta}{2\pi}), \end{cases}$$

where Λ_1 and Λ_3 are double eigenvalues, Λ_2 and Λ_4 are $(2n - 2)$ -fold eigenvalues. Since β_k is nondegenerate, there hold

$$-\frac{\theta}{2\pi} \pm \frac{k}{2} \quad \text{and} \quad -\frac{\theta}{2\pi} \notin \mathbf{Z}.$$

Now we distinguish the study in three cases. Suppose $k > 0$. By (3.12) there hold

$$\begin{aligned} \Lambda_1 < 0, \quad \Lambda_2 < 0, \quad \Lambda_3 > 0, \quad \Lambda_4 > 0, & \quad \text{if } m < -\frac{\theta}{2\pi} - \frac{k}{2}, \\ \Lambda_1 < 0, \quad \Lambda_2 < 0, \quad \Lambda_3 < 0, \quad \Lambda_4 > 0, & \quad \text{if } -\frac{\theta}{2\pi} - \frac{k}{2} < m < -\frac{\theta}{2\pi}, \\ \Lambda_1 < 0, \quad \Lambda_2 > 0, \quad \Lambda_3 < 0, \quad \Lambda_4 < 0, & \quad \text{if } -\frac{\theta}{2\pi} < m < -\frac{\theta}{2\pi} + \frac{k}{2}, \\ \Lambda_1 > 0, \quad \Lambda_2 > 0, \quad \Lambda_3 < 0, \quad \Lambda_4 < 0, & \quad \text{if } -\frac{\theta}{2\pi} + \frac{k}{2} < m. \end{aligned}$$

From $-\theta/(2\pi) \pm (k/2) \notin \mathbf{Z}$ we obtain

$$\# \left\{ \mathbf{Z} \cap \left(-\frac{\theta}{2\pi} - \frac{k}{2}, -\frac{\theta}{2\pi} + \frac{k}{2} \right) \right\} = k.$$

Summing up total numbers of positive and negative eigenvalues respectively for all $m \in [-m_0, m_0]$ defined in (3.4) we obtain

$$(3.13) \quad m_k^- = d_{1,\omega} + k, \quad m_k^+ = d_{1,\omega} - k.$$

By similar computations, when $k < 0$, we get (3.13), and when $k = 0$, we obtain $m_0^- = d_{1,\omega} = m_0^+$. This completes the proof of Step 1.

Step 2. The case of $\gamma \in \hat{\mathcal{P}}_{\tau,\omega}^0(2n)$.

Similar to the proof in [Lo8] for $\omega = 1$ we obtain

$$(3.14) \quad m^0 = \dim_{\mathbf{C}} \ker_{\mathbf{C}} a''_{1,\omega}(0) = \dim_{\mathbf{C}} \ker_{\mathbf{C}}(\gamma(1) - \omega I) = \nu_{1,\omega}(\gamma).$$

Define the perturbation paths γ_s for $s \in [-1, 1]$ by (2.6). Define $B_s(t) = -J\dot{\gamma}_s(t)\gamma_s(t)^{-1}$ for $(s, t) \in [-1, 1] \times [0, 1]$. Denote the functionals and maps on E_1^ω and the truncated space Z_1^ω corresponding to B_s by $a_s = f_s \circ u_s$. Here $a_0 = a_{1,\omega}$. Since $\gamma_s \rightarrow \gamma$ in C^1 as $s \rightarrow 0$, we obtain $B_s \rightarrow B$ in C^0

as $s \rightarrow 0$. Therefore by choosing $2d_{1,\omega} = \dim_{\mathbb{C}} Z_1^\omega$ to be large enough, the space Z_1^ω can be defined uniformly for all $s \in [-1, 1]$. Denote by m_s^* for $*$ = +, 0, and -. The origin $z = 0$ is a nondegenerate isolated critical point of a_s when $s \neq 0$. By Step 1, Theorem 3.1 holds for B_s when $s \neq 0$, i.e.

$$m_s^- = d_{1,\omega} + i_{1,\omega}(\gamma_s), \quad m_s^0 = 0, \quad m_s^+ = d_{1,\omega} - i_{1,\omega}(\gamma_s), \quad \text{if } s \in [-1, 1] \setminus \{0\}.$$

We also have a_s converges to a in C^2 as $s \rightarrow 0$. When $|s| > 0$ is sufficiently small, the matrix $a_s''(0)$ is a small perturbation of $a''(0)$. Thus in this case there hold $m^+ \leq m_s^+$ and $m^- \leq m_{-s}^-$. So when $0 < s \leq 1$ and sufficiently close to 0 there hold

$$(3.15) \quad m^+ \leq m_s^+ = d_{1,\omega} - i_{1,\omega}(\gamma_s),$$

$$(3.16) \quad m^- \leq m_{-s}^- = d_{1,\omega} + i_{1,\omega}(\gamma_{-s}).$$

By Theorem 2.5, the above (3.15) can be rewritten into

$$(3.17) \quad m^+ \leq d_{1,\omega} - i_{1,\omega}(\gamma_s) = d_{1,\omega} - i_{1,\omega}(\gamma_{-s}) - \nu_{1,\omega}(\gamma).$$

Note that $\dim Z_1^\omega = 2d_{1,\omega}$. We then obtain

$$(3.18)$$

$$d_{1,\omega} - i_{1,\omega}(\gamma_{-s}) \leq 2d_{1,\omega} - m^- = m^+ + m^0 = m^+ + \nu_{1,\omega}(\gamma) \leq d_{1,\omega} - i_{1,\omega}(\gamma_{-s}).$$

Thus equalities must hold in (3.18) and they yield

$$(3.19) \quad m^- = d_{1,\omega} + i_{1,\omega}(\gamma_{-s}) \quad \text{and} \quad m^+ = d_{1,\omega} - i_{1,\omega}(\gamma_{-s}) - \nu_{1,\omega}(\gamma).$$

Now (3.14) and (3.19) finish Step 2.

The proof of Theorem 3.1 is complete. \square

Now we can give:

Proof of Theorem 2.6. Without loss of generality we suppose $\tau = 1$. We prove Theorem 2.6 in two steps.

Step 1. Firstly we suppose $\gamma \in \hat{\mathcal{P}}_1(2n)$. Fix $s \in (0, 1]$. We claim that there exists $\epsilon = \epsilon(\gamma) > 0$ depending on γ such that (2.12) holds for any path $\alpha \in \hat{\mathcal{P}}_{1,\omega}^*(2n) \cap B_\epsilon^1(\gamma)$, where $B_\epsilon^1(\gamma)$ is the open ball centered at γ with radius ϵ in $\hat{\mathcal{P}}_1(2n)$.

We prove the left inequality of (2.12) indirectly by assuming that there exist $\alpha_k \in \hat{\mathcal{P}}_{1,\omega}^*(2n)$ for $k \in \mathbb{N}$ such that α_k converges to γ in C^1 as $k \rightarrow \infty$, and there holds

$$(3.20) \quad i_{1,\omega}(\gamma_{-s}) > i_{1,\omega}(\alpha_k).$$

Denote by $B_k(t) = -J\dot{\alpha}_k(t)\alpha_k^{-1}(t)$. Then $B_k \rightarrow B$ in C^0 . Therefore we can choose the truncated space Z_1^ω to be large enough for all the B and B_k to carry out the saddle point reduction. Let $2d_{1,\omega} = \dim Z_1^\omega$. Denote by m_k^- , m_k^0 , and m_k^+ the Morse indices of the functional a_k on Z_1^ω corresponding to α_k , and by m^- , m^0 , and m^+ the Morse indices of the functional $a_{1,\omega}$ on Z_1^ω

corresponding to γ . Then whenever k is large enough, by Theorem 3.1 and (3.20) we obtain

$$m^- \leq m_k^- = d_{1,\omega} + i_{1,\omega}(\alpha_k) < d_{1,\omega} + i_{1,\omega}(\gamma_{-s}).$$

Together with Theorem 3.1, we get the following contradiction

$$d_{1,\omega} - i_{1,\omega}(\gamma_{-s}) = m^+ + m^0 = 2d_{1,\omega} - m^- > d_{1,\omega} - i_{1,\omega}(\gamma_{-s}).$$

This proves the left inequality in (2.12). The right inequality in (2.12) follows by a similar proof.

Step 2. Fix $\gamma \in \mathcal{P}_{1,\omega}^0(2n)$, choose $\hat{\gamma} \in \hat{P}_1(2n)$ such that $\hat{\gamma}(1) = \gamma(1)$ and $\hat{\gamma} \sim \gamma$. Specially this implies $\Delta_1(\hat{\gamma}) = \Delta_1(\gamma)$. Thus by the construction in the proof of Theorem 2.5, we obtain $\hat{\gamma}_s(1) = \gamma_s(1)$ and $\hat{\gamma}_s \sim \gamma_s$ for all $s \in [-1, 1]$ provided $\hat{\gamma}$ being sufficiently C^0 close to γ . By the homotopy invariant of the ω -index for non-degenerate paths, we obtain

$$(3.21) \quad i_{1,\omega}(\hat{\gamma}_s) = i_{1,\omega}(\gamma_s), \quad \nu_{1,\omega}(\hat{\gamma}_s) = \nu_{1,\omega}(\gamma_s), \quad \forall s \in [-1, 1] \setminus \{0\}.$$

For the constant $\epsilon = \epsilon(\hat{\gamma}) > 0$ given by the step 1, let $B_{\epsilon/2}^0(\gamma)$ be the open ball centered at γ with radius $\epsilon/2$ in $\mathcal{P}_1(2n)$ under the $C([0, 1], \text{Sp}(2n))$ topology. By further requiring this $\epsilon > 0$ to be much smaller, we can assume

$$\begin{aligned} |\Delta_1(\hat{\gamma}) - \Delta_1(\hat{\alpha})| &\leq 1/2, & \forall \hat{\alpha} \in B_\epsilon^1(\hat{\gamma}), \\ |\Delta_1(\gamma) - \Delta_1(\alpha)| &\leq 1/2, & \forall \alpha \in B_{\epsilon/2}^0(\gamma). \end{aligned}$$

Fix a path $\alpha \in \mathcal{P}_{1,\omega}^*(2n) \cap B_{\epsilon/2}^0(\gamma)$. Choose $\hat{\alpha} \in \hat{P}_{1,\omega}^*(2n) \cap B_\epsilon^1(\hat{\gamma})$ such that $\hat{\alpha}(1) = \alpha(1)$. Then we obtain $|\Delta_1(\hat{\alpha}) - \Delta_1(\alpha)| \leq 1$. This then implies $\Delta_1(\hat{\alpha}) = \Delta_1(\alpha)$. By Lemma 2.3, this implies $\hat{\alpha} \sim \alpha$. Then by the homotopy invariant for the non-degenerate paths, we obtain $i_{1,\omega}(\hat{\alpha}) = i_{1,\omega}(\alpha)$. Thus by (2.12) proved in Step 1 for $\hat{\gamma}$, we obtain (2.12) for $\gamma, \gamma_{-s}, \gamma_s$, and α . The proof is complete. \square

Corollary 3.2. *Fix $\omega \in \mathbf{U}$ and $\gamma \in \mathcal{P}_{\tau,\omega}^0(2n)$. For any paths α and $\beta \in \mathcal{P}_{\tau,\omega}^*(2n)$ which are sufficiently C^0 -close to γ , there hold*

$$(3.22) \quad |i_{\tau,\omega}(\beta) - i_{\tau,\omega}(\alpha)| \leq \nu_{\tau,\omega}(\gamma).$$

4. Splitting numbers.

Fix $n \in \mathbf{N}$, $\tau > 0$ and a path $\gamma \in \mathcal{P}_\tau(2n)$. In this section, we study the properties of the ω -indices of γ ,

$$(4.1) \quad i_\tau(\omega) \equiv i_{\tau,\omega}(\gamma) \quad \text{and} \quad \nu_\tau(\omega) \equiv \nu_{\tau,\omega}(\gamma),$$

as functions of $\omega \in \mathbf{U}$. By definition we obtain

$$(4.2) \quad i_\tau(\omega) = i_\tau(\bar{\omega}) \quad \text{and} \quad \nu_\tau(\omega) = \nu_\tau(\bar{\omega}), \quad \forall \omega \in \mathbf{U}.$$

Lemma 4.1. i_τ is locally constant on $\mathbf{U} \setminus \sigma(\gamma(\tau))$, and thus is constant on each connected component of $\mathbf{U} \setminus \sigma(\gamma(\tau))$. There holds

$$(4.3) \quad \nu_\tau(\omega) = 0, \quad \forall \omega \in \mathbf{U} \setminus \sigma(\gamma(\tau)).$$

Proof. Note that $\mathbf{U} \cap \sigma(\gamma(\tau))$ contains at most $2n$ points. For any $\omega_0 \in \mathbf{U} \setminus \sigma(\gamma(\tau))$, let $\mathcal{N}(\omega_0)$ be an open connected neighborhood of ω_0 in $\mathbf{U} \setminus \sigma(\gamma(\tau))$. By definition we obtain $\nu_\tau(\omega) = 0$ for all $\omega \in \mathcal{N}(\omega_0)$. Thus (4.3) holds and $\gamma \in \mathcal{P}_{\tau, \omega}^*(2n)$ for all $\omega \in \mathcal{N}(\omega_0)$. By Theorem 7.1, there is a path $\beta : [0, 1] \rightarrow \mathcal{P}_{\tau, \omega_0}^*(2n)$ connecting $\gamma(\tau)$ to M_n^+ or M_n^- . By the compactness of the image of β and the openness of $\text{Sp}(2n)_{\omega_0}^*$ in $\text{Sp}(2n)$, we can further require $\mathcal{N}(\omega_0)$ to be so small that β is completely located within $\text{Sp}(2n)_{\omega}^*$ for all $\omega \in \mathcal{N}(\omega_0)$. Then by definition, this implies $i_\tau(\omega) = i_\tau(\omega_0)$ for all $\omega \in \mathcal{N}(\omega_0)$, and proves the lemma. \square

Corollary 4.2. The discontinuity points of $i_\tau(\cdot)$ and $\nu_\tau(\cdot)$ are contained in $\mathbf{U} \cap \sigma(\gamma(\tau))$.

Definition 4.3 (cf. [YS] and Section I.2 of [Ek]). Recall that for $M \in \text{Sp}(2n)$ and $\omega \in \mathbf{U} \cap \sigma(M)$ being an m -fold eigenvalue, the Hermitian form $\langle \sqrt{-1}J \cdot, \cdot \rangle$, which is called the **Krein form**, is always nondegenerate on the invariant root vector space $E_\omega(M) = \ker_{\mathbf{C}}(M - \omega I)^m$, where $\langle \cdot, \cdot \rangle$ denotes the inner product in \mathbf{C}^{2n} . Then ω is of **Krein type** (p, q) with $p + q = m$ if the restriction of the Krein form on $E_\omega(M)$ has signature (p, q) . ω is Krein positive if it has Krein type $(p, 0)$, is Krein negative if it has Krein type $(0, q)$. If $\omega \in \mathbf{U} \setminus \sigma(M)$, we define the Krein type of ω by $(0, 0)$.

It is well known that whenever 1 or $-1 \in \sigma(M)$, it must be a $2p$ -fold eigenvalue for some $p \in \mathbf{N}$ and have Krein type (p, p) . By direct computations we obtain:

Lemma 4.4. For any $M \in \text{Sp}(2n)$ and $\lambda \in \sigma(M)$, the algebraic and geometric multiplicities and the Krein type of $\lambda \in \sigma(N)$ for any $N \in \Omega^0(M)$ are the same as those of M .

We study next the splitting numbers $S_M^\pm(\omega)$ defined in Theorem 1.3.

Lemma 4.5. The splitting numbers $S_M^\pm(\omega)$ are well defined, i.e. is independent of the path $\gamma \in \mathcal{P}_1(2n)$ appeared in (1.3). For $\omega \in \mathbf{U}$ and $M \in \text{Sp}(2n)$, $S_N^\pm(\omega)$ are constant for any $N \in \Omega^0(M)$ of Definition 1.1.

Proof. We prove the theorem in two steps.

Step 1. Claim 1. $S_M^\pm(\omega)$ is independent of the path γ in its definition.

In fact, for γ in (1.3) using (2.5) and a sufficiently small $a > 0$, we define

$$\xi_a(t) = \gamma(1)Q(-ta, \dots, -ta), \quad \forall t \in [0, 1].$$

Then we obtain

$$i_{1, \omega}(\gamma) = i_{1, \omega}(\xi_a * \gamma),$$

and by Theorem 7.1 there is a path $\beta_0 : [0, 1] \rightarrow \text{Sp}(2n)_\omega^*$ which connects $\xi_\alpha(1)$ to M_n^+ or M_n^- defined in Section 7. Choose $\epsilon > 0$ to be small enough such that $\lambda(s) = \exp(s\epsilon\sqrt{-1})\omega \notin \sigma(M)$ for $0 < s \leq 1$. By Lemma 4.1 there holds $i_{1,\lambda(s)}(\gamma) = \text{constant}$ for all $s \in (0, 1]$. Thus by Theorem 7.1, there exists a path $\beta_\epsilon : [0, 1] \rightarrow \text{Sp}(2n)_{\lambda(1)}^*$ such that $\beta_\epsilon(0) = M$ and $\beta_\epsilon(1) = M_n^+$ or M_n^- . Thus by Definition 2.4 we obtain

$$\begin{aligned} S_M^+(\omega) &= i_{1,\lambda(1)}(\gamma) - i_{1,\omega}(\gamma) \\ &= \frac{1}{\pi} \Delta_1(\beta_\epsilon * \gamma) - \frac{1}{\pi} \Delta_1(\beta_0 * \xi_a * \gamma) \\ &= \frac{1}{\pi} [\Delta_1(\beta_\epsilon) - \Delta_1(\beta_0) - \Delta_1(\xi_a)]. \end{aligned}$$

Thus $S_M^+(\omega)$ is independent of the choice of γ . By Theorem 7.1, this number is also independent of the choices of the above $a > 0$ and paths β_ϵ and β_0 , and then is completely determined by M and ω .

Step 2. Claim 2. $S_M^\pm(\omega) = S_N^\pm(\omega)$ for any $N \in \Omega^0(M)$.

In fact, by the definition of $N \in \Omega^0(M)$, there exists a path $\xi : [0, 1] \rightarrow \Omega(M)$ such that $\xi(0) = M$ and $\xi(1) = N$. Choose any $\gamma \in \mathcal{P}_1(2n)$ such that $\gamma(1) = M$. Then by the definition of $\Omega(M)$, there holds $\gamma \sim \xi * \gamma$ on $[0, 1]$ along ξ . Thus by Theorem 2.11, we obtain

$$\begin{aligned} S_M^\pm(\omega) &= \lim_{\epsilon \rightarrow 0^+} i_{1,\exp(\pm\epsilon\sqrt{-1})\omega}(\gamma) - i_{1,\omega}(\gamma) \\ &= \lim_{\epsilon \rightarrow 0^+} i_{1,\exp(\pm\epsilon\sqrt{-1})\omega}(\xi * \gamma) - i_{1,\omega}(\xi * \gamma) \\ &= S_N^\pm(\omega). \end{aligned}$$

The proof is complete. □

Note that Theorem 1.3 is contained in Lemma 4.5.

Lemma 4.6. For $M \in \text{Sp}(2n)$ and $\omega \in \mathbf{U}$, there hold

$$(4.4) \quad S_M^\pm(\omega) = 0, \quad \text{if } \omega \notin \sigma(M).$$

$$(4.5) \quad S_M^+(\omega) = S_M^-(\bar{\omega}).$$

For any $M_i \in \text{Sp}(2n_i)$ with $i = 0$ and 1 , there holds

$$(4.6) \quad S_{M_0 \diamond M_1}^\pm(\omega) = S_{M_0}^\pm(\omega) + S_{M_1}^\pm(\omega), \quad \forall \omega \in \mathbf{U}.$$

Proof. This is a direct consequence of Definition (1.3), Theorem 2.11, and Lemma 4.5. □

In order to give a complete explanation of the splitting number, we introduce the concept of ultimate type of any $\omega \in \mathbf{U}$ as an eigenvalue of

$M \in \mathrm{Sp}(2n)$ as follows. We use the notation of basic normal forms introduced in Section 7.

Definition 4.7. A basic normal form matrix M given by Definition 7.7 is **trivial**, if for sufficiently small $a > 0$, $MR((t-1)a)^{\diamond n}$ possesses no eigenvalue on \mathbf{U} for $t \in [0, 1)$, and is **nontrivial** otherwise.

Note that by direct computations, $N_1(1, -1)$, $N_1(-1, 1)$, $N_2(\omega, B)$ and $N_2(\bar{\omega}, B) \in \mathcal{M}_\omega^1(4)$ with $\omega = \exp(\theta\sqrt{-1}) \in \mathbf{U} \setminus \mathbf{R}$ and $(b_2 - b_3) \sin \theta > 0$ are trivial, and any other basic normal form matrix is nontrivial.

Definition 4.8. For any basic normal form $M \in \mathrm{Sp}(2n)$ given by Definition 7.7 and $\omega \in \mathbf{U} \cap \sigma(M)$, we define the **ultimate type** (p, q) of ω for M to be its usual Krein type if M is nontrivial, and to be $(0, 0)$ if M is trivial. For any $M \in \mathrm{Sp}(2n)$, we define the ultimate type of ω for M to be $(0, 0)$ if $\omega \in \mathbf{U} \setminus \sigma(M)$. For any $M \in \mathrm{Sp}(2n)$, by Theorem 7.8 there exists a \diamond -product expansion (7.12) in the homotopy component $\Omega^0(M)$ of M where each M_i is a basic normal form for $1 \leq i \leq k$ and $\sigma(M_0) \cap \mathbf{U} = \emptyset$. Denote the ultimate type of ω for M_i by (p_i, q_i) for $0 \leq i \leq k$. Let $p = \sum_{i=0}^k p_i$ and $q = \sum_{i=0}^k q_i$. We define the **ultimate type** of ω for M by (p, q) .

Proposition 4.9. *The ultimate type of $\omega \in \mathbf{U}$ for M is uniquely determined by ω and M , therefore is well defined. It is constant on $\Omega^0(M)$ for fixed $\omega \in \mathbf{U}$.*

Proof. It suffices to notice that the \diamond -product (7.12) is uniquely determined by M module the arranging order of $\{M_1, \dots, M_k, M_0\}$ and the choice of the symplectic matrix M_0 satisfying $\omega \notin \sigma(M_0)$, and (7.12) holds for any $N \in \Omega^0(M)$. But these two factors make no change on the ultimate type defined above. \square

Lemma 4.10. *For $\omega \in \mathbf{U}$ and $M \in \mathrm{Sp}(2n)$, denote the Krein type and the ultimate type of ω for M by (P, Q) and (p, q) . Then there holds*

$$(4.7) \quad P - p = Q - q \geq 0.$$

Proof. Note that (4.7) holds for all basic normal forms of eigenvalues in \mathbf{U} . By the Krein Theorem (Theorem III.1.3 of [YS]), whenever the eigenvalue ω leaves \mathbf{U} by a perturbation on M , both the Krein positive and negative type numbers of ω must decrease by the same integer. Therefore by the proof of Theorem 7.8, (4.7) must hold. \square

Theorem 4.11. *For any $\omega \in \mathbf{U}$ and $M \in \mathrm{Sp}(2n)$, there hold*

$$(4.8) \quad S_M^+(\omega) = p \quad \text{and} \quad S_M^-(\omega) = q,$$

where (p, q) is the ultimate type of ω for M .

Proof. By Lemmas 4.5, 4.6 and Theorem 7.8, it suffices to prove (4.8) for M being the one of basic normal forms $N_1(\pm 1, B)$ with $B = 1, 0, -1$, $N_1(\omega, B)$

and $N_2(\omega, B) \in \mathcal{M}_\omega^1(2n)$ for $\omega \in \mathbf{U} \setminus \mathbf{R}$ defined in Section 7. We continue the proof according to different basic normal forms.

Case 1. $M = N_1(1, B)$ with $B = \pm 1$ or 0.

Define $f_B(t) = N_1(1, tB)$ for $t \in [0, 1]$. The matrix $M = N_1(1, B)$ is non-trivial when $B = 1$ or 0 and possesses both the Krein type and the ultimate type (1, 1) of the eigenvalue 1. The matrix $M = N(1, -1)$ is trivial, and possesses the Krein type (1, 1) and the ultimate type (0, 0) of the eigenvalue 1. Then using the \mathbf{R}^3 cylindrical coordinate representation introduced in [Lo2], it is easy to see that for any $\epsilon > 0$ sufficiently small there hold

$$\begin{aligned} i_{1,1}(f_B) &= -1, & i_{1,\exp(\pm\epsilon\sqrt{-1})}(f_B) &= 0, & \text{if } B &= 1 \text{ or } 0, \\ \text{and } i_{1,1}(f_{-1}) &= 0, & i_{1,\exp(\pm\epsilon\sqrt{-1})}(f_{-1}) &= 0. \end{aligned}$$

Thus (4.8) holds when $\omega = 1$.

Case 2. $M = N_1(-1, B)$ with $B = \pm 1$ or 0.

The proof of the case of $\omega = -1$ is similar by using the \mathbf{R}^3 -cylindrical coordinate representation of [Lo2] and is left to the readers.

Case 3. $M = R(\theta)$ or $R(2\pi - \theta)$ with $\omega = e^{\theta\sqrt{-1}} \in \mathbf{U} \setminus \mathbf{R}$.

Fix $\omega = e^{\theta\sqrt{-1}} \in \mathbf{U} \setminus \mathbf{R}$. By Lemma 4.6 it suffices to prove (4.8) for $\theta \in (0, 2\pi)$ and $M = R(\theta)$. Fix a $\gamma \in \mathcal{P}_\tau(2)$ in (1.3).

When $\theta \in (0, \pi)$, note that $\sigma(R(\theta))$ consists of two simple eigenvalues ω and $\bar{\omega}$. By direct computation, $R(\theta)$ possess $(p, q) = (0, 1)$ as the Krein type and ultimate type of ω .

On the other hand, let $\lambda(\pm\epsilon) = \exp(\pm\epsilon\sqrt{-1})\omega$ with $\epsilon > 0$ sufficiently small. When $\theta \in (0, \pi)$, the matrix $\gamma_{-1}(1)$ can be connected to $D(2)$ within $\text{Sp}(2)_\omega^+$, where γ_{-1} is the perturbation path of γ defined by (2.6). Note also that the matrix $R(\theta)$ can be connected to $D(2)$ within $\text{Sp}(2)_{\lambda(\epsilon)}^+$, and the matrix $R(\theta)$ can be connected to $D(-2)$ within $\text{Sp}(2)_{\lambda(-\epsilon)}^-$. Thus there hold

$$(4.9) \quad i_1(\lambda(-\epsilon)) = i_1(\omega) + 1, \quad i_1(\lambda(\epsilon)) = i_1(\omega).$$

Therefore we obtain (4.8).

When $\theta \in (\pi, 2\pi)$, similarly the matrix $\gamma_{-1}(1)$ can be connected to $D(-2)$ within $\text{Sp}(2)_{\bar{\omega}}^-$, the matrix $R(\theta)$ can be connected to $D(-2)$ within $\text{Sp}(2)_{\lambda(\epsilon)}^-$, and the matrix $R(\theta)$ can be connected to $D(2)$ within $\text{Sp}(2)_{\lambda(-\epsilon)}^+$. Thus (4.8) still hold.

Next we continue our study for $M = N_2(\omega, B) \in \mathcal{M}_\omega^1(4)$ with $\omega = e^{\theta\sqrt{-1}} \in \mathbf{U} \setminus \mathbf{R}$ in two cases. By Lemma 4.6, it suffices to consider the case of $\omega \in \sigma(N_2(\omega, B))$ with $\theta \in (0, \pi) \cup (\pi, 2\pi)$. In the following we always suppose $\theta \in (0, \pi)$. The case of $\theta \in (\pi, 2\pi)$ can be studied similarly and is left to the readers.

Case 4. $M = N_2(\omega, B) \in \mathcal{M}_\omega^1(4)$ and $(b_2 - b_3) \sin \theta < 0$.

Since $N_2(\omega, B)$ is nontrivial, by direct computation we obtain that ω as an eigenvalue of M possesses both the Krein type and the ultimate type $(1, 1)$.

Define

$$(4.10) \quad \xi(t) = MR((t - 1)\alpha)^{\diamond 2}, \quad \forall t \in [0, 1],$$

for some small $\alpha \in (0, \theta/2]$. Note that there holds

$$(4.11) \quad \det(\xi(t) - \lambda I_4) = \lambda^4 - 4A_\xi(t)\lambda^3 + B_\xi(t)\lambda^2 - 4A_\xi(t)\lambda + 1.$$

Then denoting $s = \sin \theta$, $c = \cos \theta$, $s_1 = \sin((t - 1)\alpha)$ and $c_1 = \cos((t - 1)\alpha)$ for $j = 1$ and 2 there, we obtain that for sufficiently small $\alpha > 0$ and $t \in [0, 1)$,

$$(4.12) \quad D_\omega(\xi(t)) = 2s_1c_1s(b_2 - b_3) + o(|s_1|) > 0,$$

$$(4.13) \quad 4A_\xi^2(t) + 2 - B_\xi(t) = 2s_1c_1s(b_2 - b_3) + o(|s_1|) > 0,$$

where $s_1 = \sin((t - 1)\alpha) < 0$. Here the condition $s(b_2 - b_3) < 0$ is crucial. Thus by Lemma 7.9, for $0 \leq t < 1$ the matrix $\xi(t)$ possesses two pairs of simple eigenvalues $\{\lambda_1(t), \overline{\lambda_1(t)}, \lambda_2(t), \overline{\lambda_2(t)}\}$ such that for $j = 1$ and 2 , $\lambda_j \in C([0, 1], \mathbf{U})$, $\lambda_j(1) = \omega$, $\lambda_j(t) \neq \omega$ and $|\lambda_j(t) - \omega|$ is sufficiently small provided $\alpha > 0$ is small enough for $t \in [0, 1)$. Denote by $\lambda_j(t) = \exp(\beta_j(t)\sqrt{-1})$ with $\beta_j \in C([0, 1], \mathbf{R})$ for $j = 1$ and 2 , and $\beta_1(1) = \beta_2(1) = \theta$. Note that $\xi(t)$ is the fundamental solution of the system (1.2) with $B(t) = -J\dot{\xi}(t)\xi(t)^{-1}$ being negative definite in the sense of [Ek]. So When $0 \leq t < 1$, $\lambda_1(t)$ and $\lambda_2(t)$ must have different Krein type. Without loss of generality, for $0 \leq t < 1$ we assume that $\lambda_1(t)$ is Krein positive and $\lambda_2(t)$ is Krein negative. By the proof of Propositions I.3.2 of [Ek], we obtain

$$(4.14) \quad \beta_1(t) > \theta > \beta_2(t), \quad \forall 0 \leq t < 1,$$

and $\beta_1(t)$ and $\beta_2(t)$ are strictly decreasing and increasing respectively when t increases to 1.

By (4.12), we have $\xi(0) \in \text{Sp}(4)_\omega^-$. Thus there is a path $\eta : [0, 1] \rightarrow \text{Sp}(2n)_\omega^-$ connecting $\eta(0) = D(-2) \diamond D(2)$ to $\eta(1) = \xi(0)$. Denote by $\psi = \alpha_{2,1,1}$ the standard zigzag path defined in Section 2. Then ψ connects I_4 to $D(-2) \diamond D(2)$ and there holds $i_{1,\lambda}(\psi) = 1$ for all $\lambda \in \mathbf{U}$. Thus

$$(4.15) \quad i_{1,\omega}(\eta * \psi) = 1.$$

Define

$$(4.16) \quad \gamma = \xi * \eta * \psi.$$

Denote by s and c as in (4.12). For any $b \in [-\alpha, 0]$, let $s_1 = \sin(-\alpha)$, $c_1 = \cos(-\alpha)$, $s_2 = \sin b$, $c_2 = \cos b$. By Lemma 7.10 we further require

$\alpha > 0$ to be so small such that for all $b \in [-\alpha, 0]$ there holds

$$(4.17) \quad D_\omega(M[R(-\alpha) \diamond R(b)]) = 2s^2(2-c_1-c_2) + 2s^2s_1s_2 - 2(1+c^2)(1-c_1)(1-c_2) > 0.$$

Thus $M[R(-\alpha) \diamond I_2]$ and $M[R(-\alpha) \diamond R(-\alpha)]$ can be connected within $\text{Sp}(4)_\omega^*$. This proves $\gamma_{-1} \sim_\omega \eta * \psi$, where γ_{-1} is the perturbation path of γ defined by (2.6). Thus

$$(4.18) \quad i_{1,\omega}(\gamma) = i_{1,\omega}(\gamma_{-1}) = i_{1,\omega}(\eta * \psi) = 1.$$

Recall that we have supposed $\theta \in (0, \pi)$. So we can fix $\epsilon > 0$ sufficiently small so that for $\chi(\pm\epsilon) = e^{\pm\epsilon\sqrt{-1}}\omega$ there hold

$$(4.19) \quad i_{1,\chi(\pm\epsilon)}(\gamma) = \lim_{s \rightarrow 0^+} i_{1,\exp(\pm s\sqrt{-1})\omega}(\gamma),$$

$$(4.20) \quad S(\omega) \equiv \{\exp(a\sqrt{-1})\omega \in \mathbf{U} \mid |a| \leq \epsilon/2\} \subset \mathbf{U} \setminus \mathbf{R}.$$

Choose $r \in (0, 1)$ close to 1 enough so that

$$(4.21) \quad \sigma(\xi(t)) \subset S(\omega) \cup S(\bar{\omega}), \quad \forall t \in [r, 1].$$

Define $\hat{\xi}(t) = \xi(tr)$ for $t \in [0, 1]$. Note that (4.21) implies

$$(4.22) \quad i_{1,\chi(\pm\epsilon)}(\gamma) = i_{1,\chi(\pm\epsilon)}(\hat{\xi} * \eta * \psi).$$

By (4.14) and the discussion there, $\lambda_1(r)$ and $\lambda_2(r)$ are Krein positive and negative respectively. By (4.20) and (4.21), both $\beta_1(r)$ and $\beta_2(r)$ are in $(0, \pi)$. Thus by Theorem 7.6 there exists $P \in \text{Sp}(4)$ such that $P\xi(r)P^{-1} = R(2\pi - \beta_1(r)) \diamond R(\beta_2(r))$. Let $p \in \mathcal{P}_1(4)$ satisfy $p(1) = P$. Define $\zeta(t) = p(t)\xi(r)p(t)^{-1}$ for all $t \in [0, 1]$. Then the conjugate path ζ connects $\xi(r)$ to $R(2\pi - \beta_1(r)) \diamond R(\beta_2(r))$. Define

$$(4.23) \quad \begin{aligned} f_1(t) &= R(t(2\pi - \beta_1(r))), & f_2(t) &= R(t\beta_2(r)), \\ f(t) &= f_1 \diamond f_2(t), & \forall t &\in [0, 1]. \end{aligned}$$

By (4.14), there holds $\beta_1(r) > \theta > \beta_2(r)$. Thus by Theorem 2.11 and a direct computation for paths in $\text{Sp}(2)$, we obtain

$$(4.24) \quad i_{1,\omega}(f) = 1 + 0 = 1,$$

$$(4.25) \quad i_{1,\chi(\epsilon)}(f) = 2 + 0 = 2, \quad i_{1,\chi(-\epsilon)}(f) = 1 + 1 = 2.$$

By (4.15) and the definitions of $\hat{\xi}$ and ζ we obtain

$$(4.26) \quad i_{1,\omega}(\zeta * \hat{\xi} * \eta * \psi) = i_{1,\omega}(\eta * \psi) = 1.$$

In Figure 4.1, this computation is intuitively explained via the \mathbf{R}^3 -cylindrical representation of $\text{Sp}(2)$ of [Lo2] on the plane $\{z = 0\}$, where f_1 and f_2 are represented by two nearby paths with the same topological properties. Since $\zeta * \hat{\xi} * \eta * \psi(1) = f(1)$, by Theorem 2.13, (4.24), and (4.26), we obtain $\zeta * \hat{\xi} * \eta * \psi \sim f$ on $[0, 1]$ with fixed end points. Since ζ is a conjugation

path, this implies $\hat{\xi} * \eta * \psi \sim f$ on $[0, 1]$ along ζ . Then by Theorem 2.11 and (4.25) we obtain

$$(4.27) \quad i_{1,\chi(\pm\epsilon)}(\hat{\xi} * \eta * \psi) = i_{1,\chi(\pm\epsilon)}(f) = 2.$$

Together with (4.22) we obtain

$$(4.28) \quad i_{1,\chi(\pm\epsilon)}(\gamma) = 2.$$

By (4.18), (4.19), and (4.28), we obtain

$$(4.29) \quad S_M^+(\omega) = S_M^-(\omega) = 1,$$

i.e. (4.8) holds in this case.




Figure 4.1.

Case 5. $M = N_2(\omega, B) \in \mathcal{M}_\omega^1(4)$ and $(b_2 - b_3) \sin \theta > 0$.

Since in this case M is trivial, by direct computation we obtain that ω as an eigenvalue of M possesses the Krein type $(1, 1)$ and the ultimate type $(0, 0)$.

Define the path $\xi : [0, 1] \rightarrow \text{Sp}(4)$ by (4.10) for some small $\alpha \in (0, \theta/2]$. By the notations of Case 4, we obtain that for sufficiently small $\alpha > 0$ and $t \in [0, 1)$ as in (4.12) and (4.13) there hold:

$$(4.30) \quad D_\omega(\xi(t)) < 0, \quad 4A_\xi^2(t) + 2 - B_\xi(t) < 0,$$

where $s_1 = \sin((t - 1)\alpha) < 0$. Here the condition $s(b_2 - b_3) < 0$ is crucial. By Lemma 7.9, this implies

$$(4.31) \quad \sigma(\xi(t)) \cap (\mathbf{U} \cup \mathbf{R}) = \emptyset, \quad \forall t \in [0, 1).$$

Similar to our study in Case 1, by (4.30) we obtain that M_2^+ can be connected to $\xi(0)$ by a path $\eta : [0, 1] \rightarrow \text{Sp}(2n)$ satisfying

$$(4.32) \quad \sigma(\eta(t)) \cap \mathbf{U} = \emptyset, \quad \forall t \in [0, 1].$$

Define $\psi = \alpha_{2,0,1}$ of Section 2. Then ψ connects I_4 to M_2^+ and satisfies $i_{1,\lambda}(\psi) = 0$ for all $\lambda \in \mathbf{U}$. Together with (4.32) there holds

$$(4.33) \quad i_{1,\omega}(\eta * \psi) = 0.$$

Similar to the proof of (4.18), using Lemma 7.10, we get that for $\gamma = \xi * \eta * \psi$ there holds

$$(4.34) \quad i_{1,\omega}(\gamma) = i_{1,\omega}(\eta * \psi) = 0.$$

Fix $\epsilon > 0$ so small that (4.19) holds for $\chi(\pm\epsilon) = e^{\pm\epsilon\sqrt{-1}}\omega \in \mathbf{U} \setminus \mathbf{R}$. By (4.33) and (4.34), $\chi(\pm\epsilon) \notin \sigma(\gamma(t))$ for $0 \leq t \leq 1$. Since γ passes through M_2^+ , we obtain

$$(4.35) \quad i_{1,\chi(\pm\epsilon)}(\gamma) = 0.$$

Then together with (4.19) and (4.34) we obtain

$$(4.36) \quad S_M^+(\omega) = S_M^-(\omega) = 0,$$

i.e. (4.8) holds in this case.

The proof is complete. □

Now we obtain the following results immediately.

Corollary 4.12. *If $\omega \in \mathbf{U} \cap \sigma(\gamma(\tau))$ is of Krein type (p, q) , there holds*

$$(4.37) \quad \lim_{\epsilon \rightarrow 0^+} \left(i_\tau(e^{\epsilon\sqrt{-1}}\omega) - i_\tau(e^{-\epsilon\sqrt{-1}}\omega) \right) = p - q.$$

Corollary 4.13. *For any $\omega \in \mathbf{U}$ and $M \in \text{Sp}(2n)$, there holds*

$$(4.38) \quad 0 \leq S_M^\pm(\omega) \leq \dim_{\mathbf{C}} \ker_{\mathbf{C}}(M - \omega I).$$

Corollary 4.14. *The integer valued splitting number pair $(S_M^+(\omega), S_M^-(\omega))$ defined for all $(\omega, M) \in \mathbf{U} \times \cup_{n \geq 1} \text{Sp}(2n)$ are uniquely determined by the following axioms:*

1° (Homotopy invariant). $S_M^\pm(\omega) = S_N^\pm(\omega)$ for all $N \in \Omega^0(M)$.

2° (Symplectic additivity). $S_{M_1 \diamond M_2}^\pm(\omega) = S_{M_1}^\pm(\omega) + S_{M_2}^\pm(\omega)$ for all $M_i \in \text{Sp}(2n_i)$ with $i = 1$ and 2 .

3° (Vanishing). $S_M^\pm(\omega) = 0$ if $\omega \notin \sigma(M)$.

4° (Normality). $(S_M^+(\omega), S_M^-(\omega))$ coincides with the ultimate type of ω for M when M is any basic normal form given by Definition 7.7.

Remark 4.15. Our Theorem 4.11 and Corollary 4.14 give a complete understanding of the splitting numbers for all $M \in \text{Sp}(2n)$. Consequently, the ultimate type is also characterized by Corollary 4.14. Note that Example II on page 181 of [Bo] coincides with our Theorem 4.11. Since [Bo] uses the same J as we do, the Theorem IV of [Bo] should be corrected by exchanging the positive and the negative Krein numbers.

5. Bott-type formulae and the mean index.

Based on the results of the previous sections, following the idea of [Bo], we now prove Theorems 1.4, 1.5 and 1.6 for the Maslov-type index theory.

Proof of Theorem 1.4. Fix $\tau > 0$ and $B \in C(S_\tau, \mathcal{L}_s(\mathbf{R}^{2n}))$. Let $\gamma : [0, +\infty) \rightarrow \text{Sp}(2n)$ be the fundamental solution of (1.2). Fix $k \in \mathbf{N}$. The bilinear form $\phi_{k\tau}$ on $E_{k\tau}^1$ corresponding to $f_{k\tau,1}$ in Section 3 is given by

$$\phi_{k\tau}(x, y) = \frac{1}{2} \langle (A - B)x, y \rangle_{L_\tau}, \quad \forall x, y \in \text{dom}A = E_{k\tau}^1 \subset L_{k\tau}^1.$$

Two subspaces F_1 and F_2 of $E_{k\tau}^1$ are $\phi_{k\tau}$ -orthogonal and is denoted by $F_1 \perp F_2$, if there holds $\phi_{k\tau}(x, y) = 0$ for all $x \in F_1$ and $y \in F_2$. We denote by $F_1 \oplus F_2$ the $\phi_{k\tau}$ -orthogonal product of F_1 and F_2 . For $\omega \in \mathbf{U}$ define

$$E_{k\tau}^1(\tau, \omega) = \{y \in E_{k\tau}^1 \mid y(t + \tau) = \omega y(t), \forall t\}.$$

In the following for notational simplicity we identify $E_{k\tau}^1(\tau, \omega)$ with E_τ^ω . For $0 \leq p \leq k$, define $\omega_p = \exp(2p\pi/k\sqrt{-1})$. Then $\omega_p^k = 1$. We claim that there hold

$$(5.1) \quad E_{k\tau}^1(\tau, \omega_p) \perp E_{k\tau}^1(\tau, \omega_q), \quad \text{if } 0 \leq p \neq q \leq k,$$

$$(5.2) \quad E_{k\tau}^1 = \bigoplus_{\omega^k=1} E_{k\tau}^1(\tau, \omega).$$

In fact, for $x \in E_{k\tau}^1(\tau, \omega_p)$ and $y \in E_{k\tau}^1(\tau, \omega_q)$ with $0 \leq p \neq q \leq k$, we have

$$\phi_{k\tau}(x, y) = \phi_\tau(x, y) \sum_{m=0}^{k-1} (\omega_p \bar{\omega}_q)^m = 0.$$

Thus (5.1) holds, and then (5.2) follows.

We consider the functional $f_{k\tau,1}$ defined by (3.3). For any $\alpha_i \in \mathbf{C}$ and $y_i \in E_{k\tau}^1(\tau, \omega_i)$ with $\omega_i^k = 1$ and $0 \leq i \leq k-1$, there holds

$$f_{k\tau,1} \left(\sum_{i=0}^{k-1} \alpha_i y_i \right) = \sum_{i=0}^{k-1} f_{k\tau,1}(\alpha_i y_i) = k \sum_{i=0}^{k-1} f_{\tau,1}(\alpha_i y_i).$$

This yields

$$(5.3) \quad f_{k\tau,1} \Big|_{E_{k\tau}^1} = \sum_{i=0}^{k-1} f_{k\tau,1} \Big|_{E_{k\tau}^1(\tau, \omega_i)} = k \sum_{i=0}^{k-1} f_{\tau, \omega_i} \Big|_{E_{\tau}^{\omega_i}}.$$

Now we carry out the saddle point reduction for $f_{k\tau,1}$ on $E_{k\tau}^1$, and obtain the functional $a_{k\tau,1} = f_{k\tau,1} \circ u_{k\tau,1}$ defined on $Z_{k\tau}^1$. Simultaneously this induces saddle point reductions for f_{τ, ω_i} on $E_{k\tau}^1(\tau, \omega_i)$ for $0 \leq i \leq k-1$, and yields the functional $a_{\tau,1}(\tau, \omega_i) = f_{\tau,1} \circ u_{\tau,1}(\tau, \omega_i)$ defined on $Z_{k\tau}^1(\tau, \omega)$. Denote the Morse indices of $a_{\tau,1}(\tau, \omega_i)$ at the origin by $\hat{m}_{\tau}^*(\omega_i)$ for $*$ = +, 0, and -. Then these Morse indices coincide with the Morse indices $m_{\tau}^*(\omega_i)$ of the functional a_{τ, ω_i} on Z_{τ}^{ω} at the origin defined next to (3.5). Therefore from (5.2) and (5.3) we obtain

$$(5.4) \quad m_{k\tau}^*(1) = \sum_{\omega^k=1} \hat{m}_{\tau}^*(\omega) = \sum_{\omega^k=1} m_{\tau}^*(\omega), \quad \text{for } * = +, 0, -.$$

Note that there holds

$$(5.5) \quad d_{k\tau,1} = \sum_{\omega^k=1} d_{\tau, \omega}.$$

Combining (5.4), (5.5), and Theorem 3.1, we have proved the special case of the Bott formula Theorem 1.4 for $\gamma \in \hat{\mathcal{P}}_{\tau}(2n)$.

Now fix $\gamma \in \mathcal{P}_{\tau}(2n)$. Choose $\beta \in \hat{\mathcal{P}}_{\tau}(2n)$ such that $\beta(\tau) = \gamma(\tau)$ and $\beta \sim \gamma$. We obtain $i_{\tau, \omega}(\beta) = i_{\tau, \omega}(\gamma)$ for all $\omega \in \mathbf{U}$. From $\beta \sim \gamma$ with fixed end points, this homotopy can be extended to $[0, 1] \times [0, k\tau]$. By Theorem 2.11, we then obtain $\beta^k \sim \gamma^k$. Thus $i_{k\tau}(\beta^k) = i_{k\tau}(\gamma^k)$ holds. Then the Bott-type formula for β implies that for γ . This completes the proof of Theorem 1.4. \square

Remark 5.1. Theorem 1.4 can also be proved based on Theorem 2.11 of the ω -index theory by reducing the path $\gamma \in \text{Sp}(2n)$ to paths in $\text{Sp}(2)$, and then by direct verifications without using the above analytic method of R. Bott.

Fix $\tau > 0$ and $\gamma \in \mathcal{P}_{\tau}$ in the rest of this section. Let $i_{k\tau}(\omega) = i_{k\tau, \omega}(\tilde{\gamma})$ and $\nu_{k\tau}(\omega) = \nu_{k\tau, \omega}(\tilde{\gamma})$ for $\omega \in \mathbf{U}$ and $k \in \mathbf{N}$.

Proof of Theorem 1.5. By Theorem 1.4, there hold

$$(5.6) \quad \frac{2\pi}{k} i_{k\tau} = \frac{2\pi}{k} \sum_{\omega^k=1} i_\tau(\omega), \quad \frac{2\pi}{k} \nu_{k\tau} = \frac{2\pi}{k} \sum_{\omega^k=1} \nu_\tau(\omega).$$

By Lemma 4.1, the function $i_\tau(\omega)$ is locally constant and $\nu_\tau(\omega)$ is locally zero on \mathbf{U} except at finitely many points. Therefore the right hand sides of (5.6) are Riemannian sums, and converge to the corresponding integrals as $k \rightarrow \infty$. \square

As a direct consequence of Theorem 1.5, we obtain

$$(5.7) \quad \hat{i}_{k\tau} = k\hat{i}_\tau, \quad \forall k \in \mathbf{N}.$$

Next we give:

Proof of Theorem 1.6. Based on the results of splitting numbers in Section 4, the proof is completely parallel to that of Theorem 1.4 of [Rd], and therefore is omitted. \square

Remark 5.2. Recently based on our above results, the following increasing estimate for any $\gamma \in \mathcal{P}_\tau(2n)$ is proved by C. Liu and the author in [LL],

$$(5.8) \quad k\hat{i}_\tau(\gamma) - n \leq i_{k\tau}(\gamma^k) \leq k\hat{i}_\tau(\gamma) + n - \nu_{k\tau}(\gamma^k), \quad \forall k \in \mathbf{N}.$$

This estimate is optimal on both sides of (5.8) as shown in [LL].

6. Relations with other index functions.

In this section we study the relationship between our ω -index theory and the index functions of R. Bott defined in [Bo] and I. Ekeland defined in [Ek].

Firstly we consider the real periodic Sturm system:

$$(6.1) \quad -\frac{d}{dt} \left(P \frac{d}{dt} x + Qx \right) + Q^T \frac{d}{dt} x + Rx = 0,$$

where $P, Q,$ and $R \in C^1(S_\tau, \mathcal{L}(\mathbf{R}^n))$, $P(t)$ and $R(t)$ are symmetric, and $P(t)$ is positive definite for all t . R. Bott defined his index function $\Lambda(\omega)$ and $N(\omega)$ in Section 1 of [Bo] for such systems and $\omega \in \mathbf{U}$. Define

$$(6.2) \quad B(t) \equiv B_x(t) = \begin{pmatrix} P^{-1}(t) & -P^{-1}(t)Q(t) \\ -Q(t)^T P^{-1}(t) & Q(t)^T P^{-1}(t)Q(t) - R(t) \end{pmatrix}.$$

Then our ω -index theory is defined for the linear Hamiltonian system (1.2) with this $B(t)$, which we denote by $(i_\tau(\omega), \nu_\tau(\omega))$ for $\omega \in \mathbf{U}$.

Theorem 6.1. *Under the above assumptions, there hold*

$$(6.3) \quad i_\tau(\omega) = \Lambda(\omega), \quad \nu_\tau(\omega) = N(\omega), \quad \forall \omega \in \mathbf{U}.$$

Proof. Note that the second equality holds by definition of the two sides. To prove the first equality, denote by $BS_M^\pm(\omega)$ the splitting numbers defined on the page 180 of [Bo]. Then by a direct verification we obtain

that $(BS_M^+(\omega), BS_M^-(\omega))$ satisfies the four axioms in Corollary 4.14 for all $M \in \cup_{n \geq 1} \text{Sp}(2n)$. Therefore it coincides with our splitting numbers $(S_M^+(\omega), S_M^-(\omega))$. Thus we obtain

$$(6.4) \quad \Lambda(\omega) - i_\tau(\omega) = \Lambda(1) - i_\tau(1).$$

By [Vi2] or [LA] we obtain that the right hand side of (6.4) is zero. Thus the first equality in (6.3) holds. \square

For $\tau > 0$ denote by $\mathcal{E}_\tau(2n)$ the set of $\gamma \in \hat{\mathcal{P}}_\tau(2n)$ which are fundamental solutions of the systems (1.2) with negative definite $B(t)$ for all $t \in [0, \tau]$. In his celebrated work [Ek], I. Ekeland systematically studied such Hamiltonian systems and defined his index theory via dual action principle. For any $\gamma \in \mathcal{E}_\tau(2n)$, denote the Ekeland index by $(j_{\tau,\omega}(\gamma), n_{\tau,\omega}(\gamma))$ for all $\omega \in \mathbf{U}$ via Definition I.5.3 of [Ek]. Note that $i_\tau^E(\gamma) = j_{\tau,1}(\gamma)$ and $\nu_\tau^E(\gamma) = n_{\tau,1}(\gamma)$ is the Ekeland index defined by Definition I.4.3 of [Ek]. The following result give the relation between our index theory and Ekeland's.

Theorem 6.2. *For any $\gamma \in \mathcal{E}_\tau(2n)$, there hold*

$$(6.5) \quad \nu_{\tau,\omega}(\gamma) = n_{\tau,\omega}(\gamma), \quad \forall \omega \in \mathbf{U},$$

$$(6.6) \quad i_{\tau,1}(\gamma) + \nu_{\tau,1}(\gamma) = -j_{\tau,1}(\gamma) - n,$$

$$(6.7) \quad i_{\tau,\omega}(\gamma) + \nu_{\tau,\omega}(\gamma) = -j_{\tau,\omega}(\gamma), \quad \forall \omega \in \mathbf{U} \setminus \{1\}.$$

We need:

Lemma 6.3. *For $\tau > 0$ and $\gamma \in \mathcal{E}_\tau(2n)$, denote by (p, p) the ultimate type of $1 \in \sigma(\gamma(\tau))$, by m_0 the total multiplicity of $1 \in \sigma(\gamma(\tau))$ and by $2r_0^-$ the number of Floquet multipliers which arrive on the unit circle at 1 along γ as defined in the Definition I.3.6 of [Ek]. Then there holds*

$$(6.8) \quad \nu_\tau(\gamma) - p = \frac{m_0}{2} - r_0^-.$$

Proof. Set $\tau = 1$. Firstly note that the definition of r_0^- in [Ek] can be localized, i.e. every short curve $\beta : [0, 1] \rightarrow \text{Sp}(2n)$ satisfying $\beta(1) = \gamma(1)$ and $-J\dot{\beta}(t)\beta(t)^{-1}$ being negative definite for all $t \in [0, 1]$ possesses the same r_0^- , whose proof is left to the readers. By Theorem 4.11, $p = S_{\gamma(1)}^+(1)$ satisfies Corollary 4.14. Thus together with Theorems 2.11, 2.13, and Theorem 7.8 for $M = \gamma(1)$, the proof of (6.8) is reduced to the case of $\gamma(1) = N_1(1, B)$ with $B = \pm 1$ or 0. Then (6.8) of these three cases follows by direct computation. Note that the perturbation path $D(1 + t\epsilon)$ in the cases 1 and 2 of the proof of Theorem 7.8 should be replaced by $D(1 + t\epsilon)R(t\theta)$ with a small $-\theta > 0$ and a much smaller $\epsilon > 0$. \square

Proof of Theorem 6.2. Without loss of generality, we suppose $\tau = 1$. Note that (6.5) follows from the Floquet theory.

Next we consider the proof of (6.6) when $\nu_1(\gamma) = 0$. Similar to our discussions in Section 2, by a sufficiently small rotational perturbation, γ is homotopic to a nearby \diamond -product path β of paths β_j in $\mathcal{E}_1(2) \cup \mathcal{E}_1(4)$ for $1 \leq j \leq k$ with some $k \in [1, n]$ and each $\beta_j(1)$ possessing only simple eigenvalues. If $\beta_j \in \mathcal{E}_1(4)$, by extending it within $\mathcal{E}_1(4)$ to reach a matrix possessing a double eigenvalue pair on $\mathbf{R} \setminus \{0, \pm 1\}$, then by a small perturbation, similar to our discussions in Sections 2 and 4 we can use a \diamond -product of two paths in $\mathcal{E}_1(2)$ to replace this β_j . Thus without loss of generality, we can assume all $\beta_j \in \mathcal{E}_1(2)$. Note that Ekeland index for nondegenerate paths is constant under small perturbations and is symplectic additive. Therefore the proof of (6.6) for γ is reduced to that for each β_j . Here we leave the details of the above discussion to the readers. By direct computation we obtain

$$(6.9) \quad i_\tau(\beta_j) = -1 - \sum_{0 < t < 1} \nu_t(\beta_j|_{[0,t]}), \quad \text{for } \beta_j \in \mathcal{E}_1(2).$$

Thus (6.6) follows from Theorem I.4.6 of [Ek] and (6.9).

When $\nu_1(\gamma) > 0$, let γ_s for $s \in [-1, 1]$ be the perturbation paths of γ defined by (2.6). Since $\gamma \in \mathcal{E}_1(2n)$, we can choose $\theta_0 > 0$ in (2.6) so small that $\gamma_s \in \mathcal{E}_1(2n)$, and there holds

$$i_1^E(\gamma_{-1}) - \nu_1^E(\gamma) \leq i_1^E(\gamma) \leq i_1^E(\gamma_1).$$

Thus by Theorem 2.5 and (6.6) for γ_s with $s \neq 0$, we obtain

$$i_1^E(\gamma_1) = i_1^E(\gamma_{-1}) - \nu_1^E(\gamma), \quad i_1^E(\gamma) = i_1^E(\gamma_1).$$

Then together with (6.6) for γ_1 and Theorem 2.5, we obtain (6.6) for γ .

To prove (6.7), by Theorem 4.11, we obtain $\lim_{\epsilon \rightarrow 0^+} i_{1, \exp(\epsilon\sqrt{-1})}(\gamma) = i_{1,1}(\gamma) + p$, where (p, p) is the ultimate type of $1 \in \sigma(\gamma(1))$. By Proposition I.3.5 of [Ek], there holds $\lim_{\epsilon \rightarrow 0^+} j_{1, \exp(\epsilon\sqrt{-1})}(\gamma) = j_{1,1}(\gamma) + n + \frac{m_0}{2} - r_0^-$. Thus together with Proposition I.5.11 of [Ek], we obtain

$$\begin{aligned} \lim_{\epsilon \rightarrow 0^+} i_{1, \exp(\epsilon\sqrt{-1})}(\gamma) &= -j_{1,1}(\gamma) - n_{1,1}(\gamma) - n + p \\ &= - \left[j_{1,1}(\gamma) + n + \frac{m_0}{2} - r_0^- \right] \\ &= - \lim_{\epsilon \rightarrow 0^+} j_{1, \exp(\epsilon\sqrt{-1})}(\gamma). \end{aligned}$$

Using (I.5.54) of [Ek] and our (4.2) we then obtain

$$\lim_{\epsilon \rightarrow 0^+} i_{1, \exp(-\epsilon\sqrt{-1})}(\gamma) = - \lim_{\epsilon \rightarrow 0^+} j_{1, \exp(-\epsilon\sqrt{-1})}(\gamma).$$

Then (6.7) holds for all $\omega \in \mathbf{U} \setminus (\sigma(\gamma(1)) \cup \{1\})$ by Proposition I.5.9 of [Ek] and Theorem 4.11.

Suppose $\omega \in \sigma(\gamma(1)) \cap (\mathbf{U} \setminus \{1\})$. Choose the perturbation paths γ_s with $s \in [-1, 1]$ of γ defined by (2.6) sufficiently close to γ so that $\gamma_s \in \mathcal{E}_1(2n)$

and there holds

$$j_{1,\omega}(\gamma_{-1}) - \nu_{1,\omega}(\gamma) \leq j_{1,\omega}(\gamma) \leq j_{1,\omega}(\gamma_1).$$

Similar to the proof of (6.6), using (6.7) for paths in $\mathcal{P}_{1,\omega}^*(2n)$ and Theorem 2.5 we obtain

$$j_{1,\omega}(\gamma_1) = -i_{1,\omega}(\gamma_1) = -i_{1,\omega}(\gamma_{-1}) - \nu_{1,\omega}(\gamma) = j_{1,\omega}(\gamma_{-1}) - \nu_{1,\omega}(\gamma).$$

Then we obtain

$$j_{1,\omega}(\gamma) = j_{1,\omega}(\gamma_1).$$

Then together with (6.7) for γ_1 and Theorem 2.5 we obtain

$$\begin{aligned} i_{1,\omega}(\gamma) &= i_{1,\omega}(\gamma_{-1}) = i_{1,\omega}(\gamma_1) - \nu_{1,\omega}(\gamma) \\ &= -j_{1,\omega}(\gamma_1) - \nu_{1,\omega}(\gamma) = -j_{1,\omega}(\gamma) - \nu_{1,\omega}(\gamma). \end{aligned}$$

This proves (6.7) for $\omega \in \sigma(\gamma(1)) \cap (\mathbf{U} \setminus \{1\})$. □

Corollary 6.4. *For $\tau > 0$ and $\gamma \in \mathcal{E}_\tau(2n)$, denote by (p^0, q^0) and (p, q) the Krein type and the ultimate type of $\omega \in \sigma(\gamma(\tau)) \cap (\mathbf{U} \setminus \{1\})$ respectively, and by $2r_0^-$ the number of Floquet multipliers which arrive on the unit circle at ω along γ as defined in Definition I.3.6 of [Ek]. Then there hold*

$$(6.10) \quad \nu_{\tau,\omega}(\gamma) - p = q^0 - r_0^-, \quad \nu_{\tau,\omega}(\gamma) - q = p^0 - r_0^-.$$

Proof. This corollary follows from Proposition I.5.11 of [Ek] and Theorem 6.2. □

As Lemma 6.3, (6.10) can also be proved directly.

Now from Theorems 1.4 and 6.1, we recover the Bott formula for the Morse index theory of closed geodesics. From Theorems 1.4 and 6.2 we recover the Bott-type formula Corollary I.5.4 of [Ek] for the Ekeland index theory of convex Hamiltonian systems. By Theorems 1.5 and 6.2 the corresponding mean indices satisfy $\hat{i}_\tau(x) = -\hat{i}_\tau^E(x)$.

Remark 6.5. Our choice of the standard symplectic matrix J on \mathbf{R}^{2n} in Section 1 coincides with those in [Bo], [Ra], [CZ], [LZ], [Lo1]-[Lo9], [DL] and [LD]. Note that there is a sign difference of this matrix between our choice and [YS] and [Ek]. Because of this reason, there is a sign difference between our Corollary 4.12 and Proposition I.5.9 of [Ek]. If we take the same J as defined in Section 1 to be the standard symplectic matrix, then (6.6) should be replaced by $i_\tau(\gamma) = i_\tau^E(\gamma) + n$ for all $\gamma \in \mathcal{E}_\tau(2n)$, which was proved by Theorem II.C.1 of [Br] when $\nu_\tau(\gamma) = 0$ and by Lemma 1.3 of [Lo7] when $\nu_\tau(\gamma) > 0$. In [Br], the Conley-Zehnder index theory i_T defined in [CZ] is extended with J being replaced by $-J$ to the degenerate linear Hamiltonian systems by Definition II.D.1 of [Br], and two extensions $i_T^\pm = m_{1/2,T}^\pm$ of i_T are obtained. In Theorem III.C.1 of [Br], the indices i_T^\pm are extended to functions $j^\pm : \mathbf{U} \rightarrow \mathbf{Z}$, and the corresponding Bott-type formulae (III.C.5.1) and (III.C.5.2) of [Br] for i_T^\pm in terms of j^\pm are claimed without any proofs.

These two formulae are incorrect, since they make no distinction between trivial and nontrivial normal forms. To see this for (III.C.5.1) of [Br], let $T = 1$ and $H(t) = \text{diag}(\pi/2, \pi/2)$ for all $t \in [0, 1]$ in the notations of [Br]. Then there holds $i_4^+ = 3 \neq 1 = \sum_{\omega^4=1} j^+(\omega)$. For (III.C.5.2) of [Br], one can consider $H(t) = (h_{i,j})$ with $h_{2,2} = 1$ or -1 , and all other $h_{i,j} = 0$ with $1 \leq i, j \leq 2$.

7. Normal forms and homotopy components.

In this section we briefly recall results obtained in [Lo6] on topological structures of ω -subsets, in [LD] on normal forms for eigenvalues in \mathbf{U} , and study the homotopy components for symplectic matrices.

Given any two symplectic matrices of square block form:

$$M_1 = \begin{pmatrix} A_1 & B_1 \\ C_1 & D_1 \end{pmatrix}_{2i \times 2i}, \quad M_2 = \begin{pmatrix} A_2 & B_2 \\ C_2 & D_2 \end{pmatrix}_{2j \times 2j},$$

we define an operation \diamond -product of M_1 and M_2 to be the $2(i+j) \times 2(i+j)$ matrix $M_1 \diamond M_2$ given by

$$M_1 \diamond M_2 = \begin{pmatrix} A_1 & 0 & B_1 & 0 \\ 0 & A_2 & 0 & B_2 \\ C_1 & 0 & D_1 & 0 \\ 0 & C_2 & 0 & D_2 \end{pmatrix}.$$

Denote by $M^{\diamond k}$ the k -fold \diamond -product $M \diamond \cdots \diamond M$. Note that the \diamond -multiplication is associative, and the \diamond -product of any two symplectic matrices is symplectic. For $\theta \in \mathbf{R}$ and $a \in \mathbf{R} \setminus \{0\}$ denote by $D(a) = \text{diag}(a, 1/a)$, $R(\theta) = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$, and $R_k(\theta) = I_{2k-2} \diamond R(\theta) \diamond I_{2n-2k}$.

For any $\omega \in \mathbf{U}$ and $M \in \text{Sp}(2n)$ define

$$(7.1) \quad D_\omega(M) = (-1)^{n-1} \omega^{-n} \det(M - \omega I).$$

Then $D_\omega = D_{\bar{\omega}}$ for all $\omega \in \mathbf{U}$ and $D \in C^\infty(\mathbf{U} \times \text{Sp}(2n), \mathbf{R})$. For $\omega \in \mathbf{U}$, define $\text{Sp}(2n)_\omega^0 = \{M \in \text{Sp}(2n) \mid D_\omega(M) = 0\}$, $\text{Sp}(2n)_\omega^\pm = \{M \in \text{Sp}(2n) \mid \pm D_\omega(M) < 0\}$, and $\text{Sp}(2n)_\omega^* = \text{Sp}(2n)_\omega^+ \cup \text{Sp}(2n)_\omega^-$. There hold $\text{Sp}(2n)_\omega^0 = \text{Sp}(2n)_{\bar{\omega}}^0$ and $\text{Sp}(2n)_\omega^* = \text{Sp}(2n)_{\bar{\omega}}^*$. Define $M_n^+ = D(2)^{\diamond n}$, $M_n^- = D(-2) \diamond D(2)^{\diamond(n-1)}$, and

$$(7.2) \quad \mathcal{M}_\omega^k(2n) = \{M \in \text{Sp}(2n)_\omega^0 \mid \dim_{\mathbf{C}} \ker_{\mathbf{C}}(M - \omega I) = k\}, \quad \text{for } k = 1, \dots, 2n.$$

An intuitive picture of these sets can be found in [Lo2] and Figure 4.1 by the \mathbf{R}^3 -cylindrical representation of $\text{Sp}(2)$ introduced in [Lo2].

Theorem 7.1 (cf. [Lo6]). *For any $\omega \in \mathbf{U}$, the ω -nonsingular set $\text{Sp}(2n)_\omega^*$ contains exactly two path-connected components $\text{Sp}(2n)_\omega^+$ and $\text{Sp}(2n)_\omega^-$ and*

there holds $M_n^\pm \in \text{Sp}(2n)_\omega^\pm$. $\text{Sp}(2n)_\omega^*$ is simply connected in $\text{Sp}(2n)$ for any $\omega \in \mathbf{U}$.

Theorem 7.2 (cf. [Lo6]). For any $M \in \mathcal{M}_\omega^1(2n)$ with $\omega \in \mathbf{U} \setminus \mathbf{R}$, by Theorem 7.1 there exists $P \in \text{Sp}(2n)$ such that

$$PMP^{-1} = N_k(\hat{\omega}, B) \diamond M_1 \equiv N,$$

with $\hat{\omega} \equiv e^{\theta\sqrt{-1}} = \omega$ or $\bar{\omega}$, and $M_1 \in \text{Sp}(2h)_\omega^*$ with $h + k = n$. Then there exist $M_0 \in \text{Sp}(2n - 2)_\omega^*$ and a continuous path $\sigma : [0, 1] \rightarrow \mathcal{M}_\omega^1(2n)$ such that

$$\sigma(0) = M, \quad \sigma(1) = R(\theta) \diamond M_0.$$

Moreover for any sufficiently small $\alpha > 0$, there exist continuous paths $\sigma^\pm : [0, 1] \rightarrow \text{Sp}(2n)_\omega^*$ such that

$$\begin{aligned} \sigma^-(0) &= MP^{-1}R_1(-\alpha)P, & \sigma^-(1) &= [R(\theta)R(-\alpha)] \diamond M_0, \\ \sigma^+(0) &= MP^{-1}R_1(\alpha)P, & \sigma^+(1) &= [R(\theta)R(\alpha)] \diamond M_0, \end{aligned}$$

and the distance between σ^\pm and σ is not greater than twice of the distance between N and $NR_1(\alpha)$.

Next we consider normal forms of symplectic matrices for eigenvalues on \mathbf{U} studied in [LD].

Case 1. The normal form $N_k(\pm 1, B) \in \mathcal{M}_{\pm 1}^1(2k) \cup \mathcal{M}_{\pm 1}^2(2k)$.

For $\lambda = \pm 1$, we define

$$(7.3) \quad N_1(\lambda, B) = \begin{pmatrix} \lambda & B \\ 0 & \lambda \end{pmatrix}, \quad \text{for } B = \pm 1, \text{ or } 0.$$

For $k \geq 2$, the matrix $N_k(\lambda, b)$ is of the following form with $k \times k$ real matrices $A(\lambda)$, $B(\lambda, b)$, $C(\lambda)$, and $B = (b_1, \dots, b_k) \in \mathbf{R}^k$:

$$(7.4) \quad N_k(\lambda, B) = \begin{pmatrix} A_k(\lambda) & B_k(\lambda, B) \\ 0 & C_k(\lambda) \end{pmatrix},$$

where $A_k(\lambda) = (A_{i,j})$ is a Jordan block of λ , i.e. $A_{i,i} = \lambda$, $A_{i,i+1} = 1$, and all other $A_{i,j} = 0$; $B_k(\lambda, B) = (B_{i,j})$ is a lower triangle matrix satisfying $B_{i,j} = 0$ if $i < j$ and $B_{i,j} = (-\lambda)^{j-1}b_i$ if $i \geq j$; $C_k(\lambda) = (C_{i,j})$ is a lower triangle matrix satisfying $C_{i,j} = 0$ if $i < j$ and $C_{i,j} = -(-\lambda)^{i-j+1}$ if $i \geq j$.

Case 2. The normal form $N_{2k}(\omega, B) \in \mathcal{M}_\omega^1(4k) \cup \mathcal{M}_\omega^2(4k)$ for the eigenvalue pair $\{\omega, \bar{\omega}\}$ with $\omega = e^{\theta\sqrt{-1}} \in \mathbf{U} \setminus \mathbf{R}$.

The matrix $N_{2k}(\omega, B)$ is of the following form with $2k \times 2k$ real matrices $A_{2k}(\omega)$, B , and $C_{2k}(\omega)$:

$$(7.5) \quad N_{2k}(\omega, B) = \begin{pmatrix} A_{2k}(\omega) & B \\ 0 & C_{2k}(\omega) \end{pmatrix},$$

where $A_{2k}(\omega) = (A_{i,j})$ and $C_{2k}(\omega) = (C_{i,j})$ are 2×2 blockwise blockwise lower triangular matrices and satisfy $A_{i,i} = R(\theta)$, $A_{i,i+1} = I_2$, all other $A_{i,j} = 0$; $C_{i,j} = 0$ if $i < j$, and $C_{i,j} = -(-1)^{i-j}R((i-j+1)\theta)$ if $i \geq j$.

Case 3. The normal form $N_k(\omega, B) \in \text{Sp}(2k)$ for the eigenvalue pair $\{\omega, \bar{\omega}\}$ with $\omega = e^{\theta\sqrt{-1}} \in \mathbf{U} \setminus \mathbf{R}$ and $k \in 2\mathbf{N} - 1$.

When $k = 1$, $N_1(\omega, 0) = R(\theta)$ and $N_1(\bar{\omega}, 0) = R(2\pi - \theta) \in \mathcal{M}_\omega^1(2)$.

When $k = 2m + 1 \geq 2$, the following matrix $N_k(\omega, B) \in \mathcal{M}_\omega^1(2k)$ with $2m \times 2m$ matrices $A_{2m}(\theta)$, B , $C_{2m}(\theta)$, defined in (7.5), and $2m \times 1$ matrices D , E , F , G :

$$(7.6) \quad N_k(\omega, B) = \begin{pmatrix} A_{2m}(\omega) & D & B & E \\ 0 & \cos \hat{\theta} & F^T & -\sin \hat{\theta} \\ 0 & 0 & C_{2m}(\omega) & 0 \\ 0 & \sin \hat{\theta} & G^T & \cos \hat{\theta} \end{pmatrix},$$

and

$$\begin{cases} \hat{\theta} = \theta, & D = (0, \dots, 0, 1, 0)^T, & E = (0, \dots, 0, 0, 1)^T, & \text{or} \\ \hat{\theta} = -\theta, & D = (0, \dots, 0, 0, 1)^T, & E = (0, \dots, 0, 1, 0)^T. \end{cases}$$

The case of $k = 2m \geq 2$: The matrix $N_k(\omega, B) \in \mathcal{M}_\omega^2(2k)$ is given by (7.5), or by

$$(7.7) \quad N_k(\omega, B) = \begin{pmatrix} N_m(\omega, B_1) & B_0 \\ 0 & N_m(\omega, B_2) \end{pmatrix},$$

with $N_m(\omega, B_i)$ for $i = 1$ and 2 given by (7.6).

Definition 7.3. The normal form $N_n(\omega, B) \in \text{Sp}(2n)$ belonging to the eigenvalue $\omega \in \mathbf{U}$ possesses the **type number** $t(N_n(\omega, B)) = 1$ if $N_n(\omega, B)[R(\alpha) \diamond I_{2n-2}] \in \text{Sp}(2n)_\omega^*$ for all sufficiently small $|\alpha| > 0$, and $t(N_n(\omega, B)) = 2$ if $N_n(\omega, B)[R(\alpha) \diamond I_{2n-2}] \in \text{Sp}(2n)_\omega^0$ for all $\alpha \in \mathbf{R}$.

Remark 7.4. Note that when $t(N_n(\omega, B)) = 2$, there must hold $N_n(\omega, B) \in \mathcal{M}_\omega^2(2n)$, $N_n(\omega, B)[R(\alpha) \diamond I_{2n-2}] \in \mathcal{M}_\omega^1(2n)^0$, $N_n(\omega, B)[I_{2n-2} \diamond R(\alpha)] \in \mathcal{M}_\omega^1(2n)^0$, and

$$N_n(\omega, B)[R(\alpha) \diamond I_{2n-4} \diamond R(\alpha)] \in \text{Sp}(2n)_\omega^*$$

for all sufficiently small $|\alpha| > 0$. Note also that $t(N_n(\omega, B)) = 2$ if and only if $\omega = \pm 1$ and $n \geq 2$, or $\omega \in \mathbf{U} \setminus \mathbf{R}$ and $n \geq 3$.

Theorem 7.5 (cf. [LD]). *For any $M \in \text{Sp}(2n)$ and $\omega = e^{\theta\sqrt{-1}} \in \mathbf{U}$, there exist $P \in \text{Sp}(2n)$ and two nonnegative integers p and q satisfying $1 \leq p + q \leq n$, first type normal forms $H_i \in \text{Sp}(2h_i)$ and second type normal forms $K_j \in \text{Sp}(2k_j)$ belonging to the eigenvalue ω for $1 \leq i \leq p$ and $1 \leq$*

$j \leq q$ defined above, and $M_0 \in \text{Sp}(2n - 2m)_\omega^*$ with $m = \sum_{i=1}^p h_i + \sum_{i=1}^q k_i$, such that there holds

$$(7.8) \quad PMP^{-1} = H_1 \diamond \cdots \diamond H_p \diamond K_1 \diamond \cdots \diamond K_q \diamond M_0.$$

In this case setting $k_0 = h_0 = 0$, we define

$$(7.9) \quad \begin{cases} m_i = \sum_{s=0}^{i-1} h_s + 1, & \text{for } 1 \leq i \leq p, \\ m_{p+2j-1} = \sum_{s=0}^p h_s + \sum_{s=0}^{j-1} k_s + 1, & \text{for } 1 \leq j \leq q, \\ m_{p+2j} = \sum_{s=0}^p h_s + \sum_{s=0}^{j-1} k_s + k_j, & \text{for } 1 \leq j \leq q. \end{cases}$$

Theorem 7.6 (cf. [LD]). *For any $M \in \text{Sp}(2n)$, there exist $P \in \text{Sp}(2n)$, integer $p \in [0, n]$, and normal forms $M_j \in \text{Sp}(2k_j)$ for $1 \leq j \leq p$ belonging to eigenvalues on \mathbf{U} , $M_0 \in \text{Sp}(2n - \sum_{j=1}^p k_j)$ satisfying $\sigma(M_0) \cap \mathbf{U} = \emptyset$, such that there holds*

$$(7.10) \quad PMP^{-1} = M_1 \diamond \cdots \diamond M_p \diamond M_0.$$

Next using normal forms, we study the homotopy components. For any $M \in \text{Sp}(2n)$, to find much simpler elements in $\Omega^0(M)$ or $\Omega_\omega^0(M)$ than (7.10).

Case 1. $M = N_n(1, B) \in \mathcal{M}_1^1(2n)$ with $n \geq 2$.

For $\epsilon > 0$ small enough, we define

$$f(t) = M[I_2 \diamond D(1 + t\epsilon)^{\diamond(n-1)}], \quad \forall t \in [0, 1].$$

Applying Theorem 7.6 to $f(1)$, we obtain $P \in \text{Sp}(2n)$ such that $Pf(1)P^{-1} = N_1(\lambda, c) \diamond M_0$ with $\sigma(M_0) \cap \mathbf{U} = \emptyset$. Together with Theorem 7.1, M can be connected to $N_1(1, c) \diamond M_0$ within $\Omega^0(M)$.

Case 2. $M = N_n(1, B) \in \mathcal{M}_\lambda^2(2n)$ with $n \geq 2$.

We define $f(t) = N_n(1, (1 - t)B)$ for $t \in [0, 1]$ and

$$g(t) = f(1)[I_2 \diamond D(1 + t)^{\diamond(n-2)} \diamond I_2], \quad \forall t \in [0, 1].$$

Applying Theorem 7.6 to $g(1)$, we obtain a path h in $\Omega_1(g(1))$ which connects $g(1)$ to $N_2(1, c) \diamond M_0$ with $\sigma(M_0) \cap \mathbf{U} = \emptyset$. Define 4×4 matrix paths $\phi(t) = N_2(1, (1 - t)c)$ and $\psi(t) = (\psi_{i,j}(t))$ with $\psi_{1,1}(t) = \psi_{3,3}(t) = 1$, $\psi_{2,2}(t) = \psi_{4,4}(t)^{-1} = 1 + t$, $\psi_{1,2}(t) = 1 - t^2$, $\psi_{4,3}(t) = t - 1$, and all other $\psi_{i,j}(t) = 0$ for all $t \in [0, 1]$. Then the path $[(\psi * \phi) \diamond M_0] * h * g * f$ connects M to $I_2 \diamond D(2) \diamond M_0$ in $\Omega^0(M)$.

Case 3. $M = N_n(\omega, B) \in \mathcal{M}_\omega^1(2n)$ is a normal form of the eigenvalue $\omega \in \mathbf{U} \setminus \mathbf{R}$.

For $\epsilon > 0$ small enough, we define

$$f(t) = N_n(\omega, B)[D(1 + t\epsilon)^{\diamond(n-k)} \diamond I_{2k}], \quad \forall t \in [0, 1],$$

where $k = 2$ if M is given by (7.5) or (7.7), and $k = 1$ if M is given by (7.6). Then $f(t) \in \Omega^0(M)$ for all $t \in [0, 1]$. Thus together with Theorem

7.6, M can be connected to $N_k(\hat{\omega}, c) \diamond M_0$ within $\Omega^0(M)$ with $\hat{\omega} = \omega$ or $\bar{\omega}$, $\sigma(M_0) \cap \mathbf{U} = \emptyset$, and some c .

Case 4. $M = N_n(\omega, B) \in \mathcal{M}_\omega^2(2n)$ is a normal form of the eigenvalue $\omega \in \mathbf{U} \setminus \mathbf{R}$ given by (7.5).

We define $f(t) = N_n(\omega, (1-t)B)$ and

$$g(t) = f(1)[D(1+t)^{\diamond(n-2)} \diamond I_4], \quad \forall t \in [0, 1].$$

Then there exists a path h in $\mathcal{M}_\omega^2(2n)$ which connects $g(1)$ to $N_2(\hat{\omega}, 0) \diamond M_0$ with $\hat{\theta} = \theta$ or $2\pi - \theta$ and $M_0 \in \text{Sp}(2n-4)$ satisfies $\sigma(M_0) \cap \mathbf{U} = \emptyset$. We define

$$(7.11) \quad k(t) = \begin{pmatrix} c & -s(1-t) & -st & s\sqrt{2t-2t^2} \\ s(1-t) & c & s\sqrt{2t-2t^2} & st \\ st & -s\sqrt{2t-2t^2} & c & -s(1-t) \\ -s\sqrt{2t-2t^2} & -st & s(1-t) & c \end{pmatrix}, \quad \forall t \in [0, 1],$$

where $c = \cos \hat{\theta}$ and $s = \sin \hat{\theta}$. By direct computation we obtain $k(t) \in \mathcal{M}_\omega^2(4)$ for all $t \in [0, 1]$. Note that the path k connects $N_2(\hat{\omega}, 0)$ to $R(\hat{\theta}) \diamond R(-\hat{\theta})$ within $\mathcal{M}_\omega^2(4)$. Thus there is a path in $\Omega^0(M)$ which connects M to $R(\hat{\theta}) \diamond R(-\hat{\theta}) \diamond M_0$.

Note that $N_n(\omega, B)$ for $\omega \in \mathbf{U}$ in other cases can also be connected to \diamond -product of basic normal forms defined below and matrices possessing no eigenvalues on \mathbf{U} . Since the proofs are similar, they are omitted.

Definition 7.7. A normal form matrix $M = N_n(\omega, B) \in \text{Sp}(2n)$ defined above is a **basic normal form** if $n = 1$ and $\omega \in \mathbf{U}$, or $M = N_2(\omega, B) \in \mathcal{M}_\omega^1(4)$ and $\omega \in \mathbf{U} \setminus \mathbf{R}$.

Together with Theorem 7.6 and our above discussion, we obtain the following result:

Theorem 7.8. For any $M \in \text{Sp}(2n)$, there is a path $f \in C^\infty([0, 1], \Omega^0(M))$ such that $f(0) = M$ and

$$(7.12) \quad f(1) = M_1 \diamond \cdots \diamond M_p \diamond M_0,$$

where the integer $p \in [0, n]$, each M_i is a basic normal form of eigenvalues on \mathbf{U} for $1 \leq i \leq p$, and the symplectic matrix M_0 satisfies $\sigma(M_0) \cap \mathbf{U} = \emptyset$.

Lemma 7.9 (cf. [Lo6]). For A and $B \in \mathbf{R}$ suppose the polynomial equation,

$$(7.13) \quad \lambda^4 - 4A\lambda^3 + B\lambda^2 - 4A\lambda + 1 = 0,$$

possesses no real roots. Then (7.13) possesses two pairs of conjugate simple roots on \mathbf{U} if and only if $4A^2 + 2 > B$. (7.13) possesses one pair of conjugate double roots on \mathbf{U} if and only if $4A^2 + 2 = B$. (7.13) possesses four simple roots away from $\mathbf{U} \cup \mathbf{R}$ if and only if $4A^2 + 2 < B$.

Proof. By direct computation. □

Lemma 7.10 (cf. [Lo6]). For $\omega = e^{\theta\sqrt{-1}}$ with $\theta \in (0, 2\pi) \setminus \{\pi\}$ and $B = \begin{pmatrix} b_1 & b_2 \\ b_3 & b_4 \end{pmatrix} \in \mathcal{L}(\mathbf{R}^2)$, consider the matrix $M = N_2(\omega, B)$.

1° $M \in \text{Sp}(4)$ if and only if $(b_2 - b_3) \cos \theta + (b_1 + b_4) \sin \theta = 0$. In the following we always suppose $M \in \text{Sp}(4)$.

2° $\omega = \cos \theta + \sqrt{-1} \sin \theta$ and $\bar{\omega}$ are double eigenvalues of M . $M \in \mathcal{M}_\omega^1(4)$ if and only if $b_2 - b_3 \neq 0$, and $M \in \mathcal{M}_\omega^2(4)$ if and only if $b_2 - b_3 = 0$.

3° For $M \in \mathcal{M}_\omega^1(4)$, there exists $\alpha_0 > 0$ small enough such that $M(R(\alpha) \diamond I_2) \in \text{Sp}(4)_\omega^*$ if $0 < |\alpha| \leq \alpha_0$.

4° Any $M \in \mathcal{M}_\omega^1(4)$ can be connected within $\mathcal{M}_\omega^1(4)$ to $N_2(\omega, D)$ such that for any $\epsilon > 0$ there exists a perturbation path $\gamma : [0, 1] \rightarrow \mathcal{M}_\omega^1(4)$ such that $\gamma(0) = N_2(\omega, D)$, $\sigma(\gamma(t))$ possesses two pairs of simple eigenvalues $\{\omega, \bar{\omega}, \lambda(t), \overline{\lambda(t)}\}$ and $\lambda \in C([0, 1], \mathbf{U})$ satisfies $\lambda(0) = \omega$, $\lambda(t) \neq \omega$ and $|\lambda(t) - \omega| \leq \epsilon$ for $0 < t \leq 1$.

5° For $M \in \mathcal{M}_\omega^2(4)$, there holds $D_\omega(M[R(\alpha) \diamond I_2]) = 0$ if and only if $\alpha = 0 \pmod{2\pi}$.

Corollary 7.11. For $M \in \text{Sp}(2n)$ and $\omega \in \mathbf{U}$, there exist a path $f : [0, 1] \rightarrow \Omega_\omega^0(M)$ such that $f(0) = M$ and

$$(7.14) \quad f(1) = M_1 \diamond \cdots \diamond M_n.$$

Here for $1 \leq i \leq n$, each M_i is of the form $N_1(\omega, B)$, $N_1(\bar{\omega}, B)$, $D(2)$, or $D(-2)$ for $1 \leq i \leq n$.

Proof. This corollary follows from Theorems 7.1, 7.8, and Lemma 7.10. □

Acknowledgements. The first manuscript [Lo9] of this paper appeared in Fall 1995, which has contained the main idea of the current version of this paper. Since then, specially during September to December 1996, I discussed with Professor Nancy Hingston on this manuscript by email and mail many times, and tried to explain to her my idea and certain details of this paper. I would like to sincerely thank her for her careful reading and pointing out some mistakes in the manuscript of this paper.

References

- [BTZ] W. Ballmann, G. Thobergsson and W. Ziller, *Closed geodesics on positively curved manifolds*, Ann. of Math., **116** (1982), 213-247.
- [Be] V. Benci, *A new approach to the Morse-Conley theory and some applications*, Ann. di Mat. pura ed appl., **CLVIII** (1991), 231-305.
- [Bo] R. Bott, *On the iteration of closed geodesics and the Sturm intersection theory*, Comm. Pure Appl. Math., **9** (1956), 171-206.
- [Br] V. Brousseau, *Espaces de Krein et index des systemes hamiltoniens*, Ann. Inst. H. Poincaré Anal non linéaire, **7** (1990), 525-560.

- [Ch] K.C. Chang, *Infinite Dimensional Morse Theory and Multiple Solution Problems*, Birkhäuser, Boston, 1993.
- [CZ] C. Conley and E. Zehnder, *Morse-type index theory for flows and periodic solutions for Hamiltonian equations*, Comm. Pure Appl. Math., **37** (1984), 207-253.
- [CD] R. Cushman and J.J. Duistermaat, *The behavior of the index of a periodic linear Hamiltonian system under iteration*, Adv. in Math., **23** (1977), 1-21.
- [DL] D. Dong and Y. Long, *The iteration formula of the Maslov-type index theory with applications to nonlinear Hamiltonian systems*, Nankai Inst. of Math., Nankai Univ., preprint, 1993; ICTP. Report. No. IC/94/318, 1994, Trans. Amer. Math. Soc., **349** (1997), 2619-2661.
- [Ed] H.M. Edwards, *A generalized Sturm theorem*, Ann. of Math., **80** (1964), 22-57.
- [Ek] I. Ekeland, *Convexity Methods in Hamiltonian Mechanics*, Springer, Berlin, 1990.
- [Kl] W. Klingenberg, *Lectures on closed geodesics*, Springer, Berlin, 1978.
- [LL] C. Liu and Y. Long, *An optimal increasing estimate for Maslov-type indices of iterations*, Nankai Inst. of Math., Nankai Univ., preprint, 1997; Chinese Bull. Sci., **42** (1997) 2275-2278.
- [Lo1] Y. Long, *Maslov-type index, degenerate critical points, and asymptotically linear Hamiltonian systems*, Science in China (Scientia Sinica), Series A, **33** (1990), 1409-1419.
- [Lo2] ———, *The structure of the singular symplectic matrix set*, Science in China (Scientia Sinica), Series A, **5** (1991), 457-465 (Chinese edition); **34** (1991), 897-907 (English edition).
- [Lo3] ———, *A Maslov-type index theory and asymptotically linear Hamiltonian systems*, in 'Dyna. Syst. and Rel. Topics', K. Shiraiwa ed., World Sci., (1991), 333-341.
- [Lo4] ———, *The Index Theory of Hamiltonian Systems with Applications*, (in Chinese), Science Press, Beijing, 1993.
- [Lo5] ———, *Periodic points of Hamiltonian diffeomorphisms on tori and a conjecture of C. Conley*, ETH-Zürich Preprint, Dec. 1994.
- [Lo6] ———, *The topological structures of ω -subsets of symplectic groups*, Nankai Inst. of Math., Nankai Univ., preprint, 1997; Acta Math. Sinica, to appear.
- [Lo7] ———, *Hyperbolic closed characteristics on compact convex smooth hypersurfaces in \mathbf{R}^{2n}* , Nankai Inst. of Math., Nankai Univ., preprint, 1996; J. Diff. Equa., to appear.
- [Lo8] ———, *A Maslov-type index theory for symplectic paths*, Nankai Inst. of Math., Nankai Univ., preprint, 1997, Top. Meth. Nonl. Anal., **10** (1997), 47-78.
- [Lo9] ———, *The Bott formula of the Maslov-type index theory*, Research Report, Nankai Inst. of Math., Nankai Univ., (1995, 1996).
- [LA] Y. Long and T. An, *Indexing domains of instability for Hamiltonian systems*, Nodea, to appear.
- [LD] Y. Long and D. Dong, *Normal forms of symplectic matrices*, Nankai Inst. of Math., Nankai Univ., preprint, (1995, 1997); Acta Math. Sinica, to appear.
- [LZ] Y. Long and E. Zehnder, *Morse theory for forced oscillations of asymptotically linear Hamiltonian systems*, in 'Stoc. Proc. Phys. and Geom., S. Albeverio et al. ed., World Sci., (1990), 528-563.

- [Ra] P. Rabinowitz, *Periodic solutions of Hamiltonian systems*, Comm. Pure Appl. Math., **31** (1978), 157-184.
- [Rd] H.-B. Rademacher, *On the average indices of closed geodesics*, J. Diff. Geod., **29** (1989), 65-83.
- [SZ] D. Salamon and E. Zehnder, *Morse theory for periodic solutions of Hamiltonian systems and the Maslov index*, Comm. Pure and Appl. Math., **45** (1992), 1303-1360.
- [Vi1] C. Viterbo, *Equivariant Morse theory for starshaped Hamiltonian systems*, Trans. Amer. Math. Soc., **311** (1989), 621-655.
- [Vi2] ———, *A new obstruction to embedding Lagrangian tori*, Invent. Math., **100** (1990), 301-320.
- [YS] V. Yakubovich and V. Starzhinskii, *Linear Differential Equations with Periodic Coefficients*, New York, John Wiley & Sons, 1975.

Received August 25, 1997 and revised December 14, 1997. The author was partially supported by the NNSF and MCSEC of China and the Qiu Shi Sci. Tech. Foundation. The author is also an associate member of ICTP.

NANKAI INSTITUTE OF MATHEMATICS
NANKAI UNIVERSITY
TIANJIN 300071
PEOPLE'S REPUBLIC OF CHINA
E-mail address: longym@sun.nankai.edu.cn