

Pacific Journal of Mathematics

ENTROPY OF CUNTZ'S CANONICAL ENDOMORPHISM

MARIE CHODA

ENTROPY OF CUNTZ'S CANONICAL ENDOMORPHISM

MARIE CHODA

Let $\{S_i\}_{i=1}^n$ be generators of the Cuntz algebra \mathcal{O}_n and let Φ be the $*$ -endomorphism of \mathcal{O}_n defined by $\Phi(x) = \sum_{i=1}^n S_i x S_i^*$. Then both of Connes–Narnhofer–Thirring's entropy $h_\phi(\Phi)$ and Voiculescu's topological entropy $ht(\Phi)$ are $\log n$, where ϕ is the unique $\log n$ -KMS state of \mathcal{O}_n . Also Longo's canonical endomorphism for $N \subset M$ have the same entropy $\log n$, where the inclusion $N \subset M$ comes from \mathcal{O}_n .

1. Introduction.

Connes–Størmer entropy $H(\cdot)$ extended the entropy invariant of Kolmogorov–Sinai to trace preserving automorphisms of finite von Neumann algebras ([CS]). Replacing a finite trace to an invariant state ϕ , Connes–Narnhofer–Thirring entropy $h_\phi(\cdot)$ is defined for automorphisms of C^* -algebras as a generalization of $H(\cdot)$ ([CNT]). These entropies depend on an invariant state under a given automorphism.

The first typical interesting example to compute the entropy is the Bernoulli shift β_n on the infinite product space of n -point sets.

In the context of operator algebras (von Neumann algebras or C^* -algebras), the non-commutative Bernoulli shift α_n takes the place of the Bernoulli shift β_n . It is the shift automorphism on the infinite tensor product $A = \bigotimes_{i=-\infty}^{\infty} A_i$ (where A_i is the $n \times n$ -matrix algebra) and $H(\alpha_n) = \log n = h_\tau(\alpha_n)$ ([CS], [CNT]), where τ is the unique tracial state of A .

Let γ be an aperiodic automorphism of an algebra B . Then there exists an implementing unitary operator u for γ in the crossed product $M = B \rtimes_\gamma \mathbb{Z}$. The inner automorphism Ad_u , ($Ad_u(x) = uxu^*$) of M is an extension of γ to M . In general, the entropy of γ is less than the entropy of Ad_u . Størmer [S] asked if the equality between the entropies of γ and Ad_u holds.

Voiculescu [V] defined topological entropy $ht(\cdot)$ for automorphisms of nuclear C^* -algebras (cf. [Hu], [T]), which does not depend on any state but is based on approximations. As an application, he showed that his topological entropy satisfies the equality for the Bernoulli shift β_n , so that Connes–Narnhofer–Thirring entropy does too.

In this paper, we show the equality for both of the automorphism α_n and the unital $*$ -endomorphism of the type of the non-commutative Bernoulli shift.

In §3, we denote only the fact that

$$H(Ad_u) = h_\tau(Ad_u) = ht(Ad_u) = \log n,$$

where τ is the unique tracial state of the reduced crossed product $A \rtimes_{\alpha_n} \mathbb{Z}$. These are proved by similar method as in §4 and §5.

The definition of Connes-Størmer entropy is available to trace preserving *-endomorphisms on finite von Neumann algebras. Similarly, we can apply the definition of Connes-Narnhofer-Thirring entropy to unital and state preserving *-endomorphisms of C^* -algebras, and also Voiculescu's topological entropy to unital *-endomorphisms of nuclear C^* -algebras. We apply here, in particular, to the unital *-endomorphism which is an extension of the *-endomorphism coming from the non-commutative Bernoulli shift α_n as follows.

If we restrict our algebra A to the *half side* infinite C^* -tensor product (or von Neumann tensor product) $B = \bigotimes_{i=0}^{\infty} A_i$ of matrix algebras, then the restriction of α_n to B defines a unit preserving *-endomorphism σ_n of B , which is canonical in the sense of [Ch2, Ch3]. Then we have the extension algebra $\langle B, \sigma_n \rangle$ of B by σ_n ([Ch2, Ch3]). In the case of C^* algebras, $\langle B, \sigma_n \rangle$ is the crossed product $B \rtimes_{\rho} \mathbb{N}$ of B by the corner endomorphism ρ in [R, I2], which is defined by σ_n using the canonical property of σ_n . Further, the canonical extension $\hat{\sigma}_n$ (in the sense of [Ch2, Ch3]) of σ_n to $\langle B, \sigma_n \rangle$ is obtained. The *-endomorphism $\hat{\sigma}_n$ of $\langle B, \sigma_n \rangle$ is defined by a modification of the automorphism Ad_u of $A \rtimes_{\alpha_n} \mathbb{Z}$ and has the property like the canonical extension in the sense of [I1, HS]. In the case of C^* -algebras, the extension algebra $\langle B, \sigma_n \rangle$ is the Cuntz algebra \mathcal{O}_n and $\hat{\sigma}_n$ is nothing but Cuntz's canonical inner endomorphism Φ of \mathcal{O}_n defined by $\Phi(x) = \sum_{i=1}^n S_i x S_i^*$, ($x \in \mathcal{O}_n$) for generators $\{S_1, \dots, S_n\}$ of \mathcal{O}_n . In the case of von Neumann algebras, $\langle B, \sigma_n \rangle$ is the unique injective type III_{1/n} factor and $\hat{\sigma}_n$ is Longo's canonical endomorphism for the subfactor of $\langle B, \sigma_n \rangle$, which appears naturally in the construction of the extension algebra $\langle B, \sigma_n \rangle$ by the canonical *-endomorphism σ_n ([Ch3]).

In §4, we show that

$$ht(\Phi) = \log n = ht(\sigma_n).$$

Applying to Connes-Narnhofer-Thirring's entropy $h_\phi(\cdot)$ relative to the unique $\log n$ -KMS state ϕ of \mathcal{O}_n , we have

$$h_\phi(\Phi) = \log n = h_\phi(\sigma_n).$$

This relation implies the same relation for Longo's canonical endomorphism. Thus the canonical extension of the non-commutative Bernoulli shift has the same entropy with the original one in the case of *-endomorphisms too.

The author thanks F. Hiai for his interest in this work and encouragement during the preparation.

2. Preliminaries.

2.1. Let H_0 be a Hilbert space of dimension $n < \infty$. Put $H_i = H_0$, $i \in \mathbb{Z}$. For two integers i and j with $i < j$, we put

$$H_{[i,j]} = H_i \otimes H_{i+1} \otimes \cdots \otimes H_j.$$

Let $\{\delta(i) : i = 1, \dots, n\}$ be an orthonormal basis of H_0 . The embedding $H_{[i,j]} \hookrightarrow H_{[i-1,j+1]}$ is given by $\xi \in H_{[i,j]} \rightarrow \delta(1) \otimes \xi \otimes \delta(1) \in H_{[i-1,j+1]}$. We denote by \mathcal{H}_i the inductive limit of $\{H_{[i,i+j]} : j = 0, 1, \dots\}$ and by \mathcal{H} the inductive limit of the increasing sequence $\{\mathcal{H}_i : i = 0, -1, \dots\}$.

Given $k, l \in \mathbb{Z}$ $k < l$, let

$$W_{[k,l]}^n = \{\mu = (\mu_k, \dots, \mu_l) : \mu_i \in \{1, \dots, n\}, (k \leq i \leq l)\}.$$

Let $\mu \in W_{[k,l]}^n$ and $\nu \in W_{[l+1,m]}^n$. We put

$$\mu \cdot \nu = (\mu_k, \dots, \mu_l, \nu_{l+1}, \dots, \nu_m).$$

Further, let

$$W_0^n = \{0\}, \quad W_{[0,\infty]}^n = \cup_{k=0}^{\infty} W_{[0,k]}^n \quad \text{and} \quad W_{\infty}^n = \cup_{k=0}^{\infty} W_{[-k,k]}^n.$$

The shift $\alpha : i \in \mathbb{Z} \rightarrow i + 1$ induces the mapping on W_{∞}^n , which we denote by the same notation α .

For $\mu \in W_{[k,l]}^n$, we put

$$\delta(\mu) = \delta(\mu_k) \otimes \cdots \otimes \delta(\mu_l) \in H_{[k,l]}.$$

Then $\{\delta(\mu) : \mu \in W_{[k,l]}^n\}$ is an orthonormal basis in $H_{[k,l]}$.

Let $A_0 = B(H_0)$ and $\{e(i, j) : i, j = 1, \dots, n\}$ be the matrix unit of A_0 with respect to the orthonormal basis $\{\delta(i) : i = 1, \dots, n\}$. We denote the trace $(1/n)\text{Tr}$ of A_0 by τ_0 . Put $A_i = A_0$, ($i \in \mathbb{Z}$) and $\tau_i = \tau_0$. For two integers $i < j$, let

$$A_{[i,j]} = A_i \otimes A_{i+1} \otimes \cdots \otimes A_j.$$

For $\mu, \nu \in W_{[k,l]}^n$, we put

$$e(\mu, \nu) = e(\mu_k, \nu_k) \otimes \cdots \otimes e(\mu_l, \nu_l) \in A_{[k,l]}.$$

Then $\{e(\mu, \nu) : \mu, \nu \in W_{[k,l]}^n\}$ is a matrix units of $A_{[k,l]}$.

2.2. We apply the entropy of Connes-Narnhofer-Thirring and Voiculescu's topological entropy to both of automorphisms and unital *-endomorphisms on C^* -algebras. To fix notations, we recall the definition of the topological entropy. Let B be a nuclear C^* -algebra with unity. Let $CAP(B)$ be triples (ρ, η, C) , where C is a finite dimensional C^* -algebra, and $\rho : B \rightarrow C$ and $\eta : C \rightarrow B$ are unital completely positive maps. Let Ω be the set of finite subsets of B . For an $\omega \in \Omega$, put

$$rcp(\omega; \delta) = \inf\{\text{rank } C : (\rho, \eta, C) \in CAP(B), \|\eta \cdot \rho(a) - a\| < \delta, a \in B\},$$

where $\text{rank } C$ means the dimension of a maximal abelian self-adjoint subalgebra of C . For a unital $*$ -endomorphism β of B , put

$$ht(\beta, \omega; \delta) = \overline{\lim}_{N \rightarrow \infty} \frac{1}{N} \log rcp(\omega \cup \beta(\omega) \cup \cdots \cup \beta^{N-1}(\omega); \delta)$$

and

$$ht(\beta, \omega) = \sup_{\delta > 0} ht(\beta, \omega; \delta).$$

Then the topological entropy $ht(\beta)$ of β is defined by

$$ht(\beta) = \sup_{\omega \in \Omega} ht(\beta, \omega).$$

Assume that there exists an increasing sequence $(\omega_j)_{j \in \mathbb{N}}$ of finite subsets of B such that the linear span of $\cup_{j \in \mathbb{N}} \omega_j$ is dense in B . Even in the case of $*$ -endomorphisms which are not automorphisms, by the obvious analogues of [V, Proposition 4.3], $ht(\cdot)$ is obtained as the following form which we use later:

$$ht(\beta) = \sup_{j \in \mathbb{N}} ht(\beta, \omega_j).$$

Let ϕ be a state of B with $\phi \cdot \beta = \phi$. The essential relation between $ht(\beta)$ and Connes-Narnhofer-Thirring entropy $h_\phi(\beta)$ is by [V, Proposition 4.6]

$$h_\phi(\beta) \leq ht(\beta).$$

3. Entropy of Ad_u for non-commutative Bernoulli shift.

In this section, we only state results without proof. We remark that these are proved by similar methods as in §4 and §5.

3.1. Let $n(2 \leq n < \infty)$ be an integer. Let $A_i, \tau_i(i \in \mathbb{Z})$ be as in §2.1 and let A be the infinite C^* -tensor product $A = \bigotimes_{i \in \mathbb{Z}} A_i$. We denote the unique tracial state of A by τ . The non-commutative Bernoulli shift α_n is the automorphism of the C^* -algebra A induced by the shift $\alpha : i(\in \mathbb{Z}) \rightarrow i + 1$. Let u be the implementing unitary in the reduced C^* -crossed product $A \rtimes_{\alpha_n} \mathbb{Z}$ for α_n . Let E be the conditional expectation of $A \rtimes_{\alpha_n} \mathbb{Z}$ onto A with $E(u^j) = 0, (j \neq 0)$. Then $\tau \cdot E$ is a tracial state of $A \rtimes_{\alpha_n} \mathbb{Z}$ which is invariant under Ad_u . We denote by the same notation α_n the extension of α_n to the hyperfinite II_1 factor $\bigotimes_{i \in \mathbb{Z}} (A_i, \tau_i) \rtimes_{\alpha_n} \mathbb{Z}$.

Theorem 3.2. *Under the above notations,*

$$ht(\alpha_n) = ht(Ad_u) = h_{\tau \cdot E}(Ad_u) = h_\tau(\alpha_n) = \log n = H(\alpha_n) = H(Ad_u).$$

4. Entropy of Cuntz's canonical endomorphism.

In this section, we apply the definition of Connes-Narnhofer-Thirring entropy and Voiculescu's topological entropy for unital $*$ -endomorphisms of nuclear C^* -algebras. All facts for automorphisms, which we need here, work for unital $*$ -endomorphisms by the analogues of definitions and proofs in [CNT] and [V].

Let n ($2 \leq n < \infty$) be an integer. Given n isometries $\{S_i\}$ on a Hilbert space such that $\sum_{i=1}^n S_i S_i^* = 1$, Cuntz defined the Cuntz algebra \mathcal{O}_n as the C^* -algebra generated by $\{S_i\}_i$ ([Cu1]). So called Cuntz's canonical endomorphism Φ of \mathcal{O}_n is defined by

$$\Phi(x) = \sum_{i=1}^n S_i x S_i^*, \quad (x \in \mathcal{O}_n).$$

The \mathcal{O}_n has exactly one log n -KMS state ϕ ([OP]). In this section we compute Voiculescu's topological entropy of Φ and Connes-Narnhofer-Thirring's entropy $h_\phi(\Phi)$. Applying to the factor generated by $\pi_\phi(\mathcal{O}_n)$, we get the entropy of Longo's canonical endomorphism.

4.1. To compute the entropy of Φ , we recall some of the representation for the Cuntz algebra \mathcal{O}_n as a crossed product in [Cu1], (cf., [Ch2, I2, P, R]). Let $A_i, \tau_i, (i \in \mathbb{Z})$ and $e(i, j), (i, j \in \mathbb{N})$ be the same as in §2.1. For a $j \in \mathbb{Z}$, \mathcal{A}_j is given as the infinite tensor product:

$$\mathcal{A}_j = \bigotimes_{i=j}^{\infty} A_i.$$

Define embeddings

$$\mathcal{A}_j \hookrightarrow \mathcal{A}_{j-1} \hookrightarrow \mathcal{A}_{j-2} \hookrightarrow \cdots$$

by $x \in \mathcal{A}_j \rightarrow e_{j-1}(1, 1) \otimes x \in \mathcal{A}_{j-1}$, where $e_{j-1}(i, l)$ is a copy of $e(i, l)$ in \mathcal{A}_{j-1} . The inductive limit of this sequence is denoted by \mathcal{A} . Since the embedding $\mathcal{A}_j \hookrightarrow \mathcal{A}_{j-1}$ and the embedding $\mathcal{H}_i \hookrightarrow \mathcal{H}_{i-1}$ in §2.1 are compatible, we can consider \mathcal{A} acting faithfully on \mathcal{H} .

The automorphism σ of \mathcal{A} is induced by the shift $\alpha : i \in \mathbb{Z} \rightarrow i + 1$.

Then the crossed product $\mathcal{A} \rtimes_\sigma \mathbb{Z}$ acts faithfully on the Hilbert space

$$K = \sum_{i \in \mathbb{Z}} \bigoplus u^i \mathcal{H},$$

where u is the implementing unitary in $\mathcal{A} \rtimes_\sigma \mathbb{Z}$ for the automorphism σ of \mathcal{A} . Let p be the unit of $\mathcal{A}_0 \subset \mathcal{A} \rtimes_\sigma \mathbb{Z}$ and put

$$w = up.$$

We remark $w^j p = w^j$.

Then Cuntz algebra \mathcal{O}_n is reresented as $p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p$, which is the C^* subalgebra $C^*(\mathcal{A}_0, w)$ of $(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})$ generated by $\{\mathcal{A}_0, w\}$. There exists a conditional expectation E of $C^*(\mathcal{A}_0, w)$ onto \mathcal{A}_0 with $E(w^j) = 0$ for all $j = 1, 2, \dots$. The unique tracial state τ of \mathcal{A}_0 is extended to the state ϕ of $C^*(\mathcal{A}_0, w)$ by $\phi = \tau \cdot E$. Then ϕ is the unique log n -KMS state of $C^*(\mathcal{A}_0, w)$ ([**OP**]).

4.2. Since

$$\sigma^j(p)(\mathcal{H}) = \mathcal{H}_j, \quad j \in \mathbb{Z},$$

the algebra $p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p$ is acting faithfully on

$$pK = \sum_{i \in \mathbb{Z}} \bigoplus u^i \mathcal{H}_{-i}.$$

The restriction $\sigma|_{\mathcal{A}_0}$ of σ to \mathcal{A}_0 is the one sided non commutative Bernoulli shift. Cuntz's canonical inner endomorphism Φ of \mathcal{O}_n is nothing but the extension of $\sigma|_{\mathcal{A}_0}$ to the Cuntz algebra $C^*(\mathcal{A}_0, w)$ which maps

$$a \rightarrow \sigma(a), \quad (a \in \mathcal{A}_0), \quad \text{and} \quad w \rightarrow vw,$$

where

$$v = \sum_{j=1}^n e((j, 1), (1, j)) \in A_{[0,1]},$$

([**Cu2**], cf. [**Ch2**]).

4.3. Let $k, m \in \mathbb{N}$. We define

$$K(k, m) = \sum_{l=-k}^k \bigoplus u^l H_{[-l, -l+m]}$$

and we denote the orthogonal projection of K onto $K(k, m)$ by $Q(k, m)$. The set $\{u^j \delta(\mu) : -k \leq j \leq k, \mu \in W_{[-j, -j+m]}^n\}$ is an orthonormal basis of $K(k, m)$. We denote by $E((j, \mu), (l, \nu))$ the partial isometry in $B(K(k, m))$ such that

$$E((j, \mu), (l, \nu)) : u^l \delta(\nu) \rightarrow u^j \delta(\mu), \quad \left(\mu \in W_{[-j, -j+m]}^n, \nu \in W_{[-l, -l+m]}^n \right).$$

Then the set

$$\begin{aligned} & \mathcal{E}(k, m) \\ &= \left\{ E((j, \mu), (l, \nu)) : -k \leq j, l \leq k, \mu \in W_{[-j, -j+m]}^n, \nu \in W_{[-l, -l+m]}^n \right\} \end{aligned}$$

is a matrix units of $B(K(k, m))$.

4.4. Let $k, m \in \mathbb{N}$. We define the completely positive unital linear map

$$\varphi_{k,m} : p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p \rightarrow B(K(k, m))$$

by

$$\varphi_{k,m}(x) = Q(k, m)xQ(k, m)|_{K(k, m)}, \quad x \in p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p.$$

We remark that if $e(\mu, \nu)w^j \neq 0$ for ν in $W_{[0, b]}^n$, ($b \geq j$), then

$$\nu = (1, \dots, 1, \nu_j, \dots, \nu_b) \quad \text{and} \quad \delta(\nu) \in H_{[j, b]}.$$

For two integers a and b with $a < b$, we let

$$\omega_{a,b} = \left\{ e(\mu, \nu)w^j : 0 \leq j \leq a \quad \text{and} \quad \mu, \nu \in W_{[0, b]}^n \right\}.$$

Let $e(\mu, \nu)w^j \in \omega_{a,b}$ for $a, b \in \mathbb{N}$, ($a < b$) and $e(\mu, \nu)w^j \neq 0$. Since $\sigma^{-l}(p)\delta(\mu) = \delta(\mu)$ for $u^l\delta(\mu) \in K(k, m)$, we have that if $k \geq a$ and $m \geq b$ then

$$\varphi_{k,m}(e(\mu, \nu)w^j) = \sum_{l=-k}^{k-j} E((j+l, \alpha^{-(j+l)}(\mu) \cdot \beta_l), (l, \alpha^{-(j+l)}(\nu) \cdot \gamma_l)),$$

where

$$\begin{aligned} \beta_l &= (1, \dots, 1) \in W_{[-(j+l)+b+1, -(j+l)+m]}^n, \\ \gamma_l &= (1, \dots, 1) \in W_{[-l+b+1, -l+m]}^n. \end{aligned}$$

We remark that

$$\delta(\alpha^{-(j+l)}(\nu)) \in H_{[-l, -l+b+1]},$$

so that $E((j+l, \alpha^{-(j+l)}(\mu) \cdot \beta_l), (l, \alpha^{-(j+l)}(\nu) \cdot \gamma_l)) \in \mathcal{E}(k, m)$.

4.5. We define the linear map

$$\psi_{k,m} : B(K(k, m)) \rightarrow p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p$$

by

$$\psi_{k,m}(E((j, \mu), (l, \nu))) = \frac{1}{2k+1} p u^j e(\mu, \nu) u^{*l} p,$$

for $E((j, \mu), (l, \nu)) \in \mathcal{E}(k, m)$.

Let T_j , ($j \in \mathbb{Z}$) be the unitary operator on K defined by

$$T_j(u^i \delta(\mu)) = u^{i+j} \delta(\alpha^{-j}(\mu)), \quad i \in \mathbb{Z}, \quad \mu \in W_{\infty}^n.$$

Then we have

$$w - \lim_{r \rightarrow \infty} \sum_{i=-r}^r T_i E((j, \mu), (l, \nu)) T_i^* = u^j e(\mu, \nu) u^{*l}$$

for any $E((j, \mu), (l, \nu)) \in B(K(k, m))$. Here ω is a nontrivial ultrafilter on \mathbb{N} . Hence $\psi_{k,m}$ is a unital completely positive map from $B(K(k, m))$ to $p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p$. Since $u^j p = w^j$, we have

$$\psi_{k,m} \cdot \varphi_{k,m}(e(\mu, \nu)w^j) = \frac{2k-j+1}{2k+1}e(\mu, \nu)w^j,$$

for all $e(\mu, \nu)w^j \in \omega_{a,b}$, $a \leq k$ and $b \leq m$.

Theorem 4.6. *Let Φ be Cuntz's canonical inner endomorphism of \mathcal{O}_n . Then*

$$ht(\Phi) = \log n.$$

Proof. Let $e(\mu, \nu)w^j \in \omega_{a,b}$. Then we have, for $a \leq k$ and $b \leq m$, by §4.5

$$\|\psi_{k,m} \cdot \varphi_{k,m}(e(\mu, \nu)w^j) - e(\mu, \nu)w^j\| = \frac{j}{2k+1}\|e(\mu, \nu)w^j\| \leq \frac{a}{2k+1}$$

and we have for an $i \in \mathbb{N}$

$$\begin{aligned} \Phi^i(e(\mu, \nu)w^j) &= \sigma^i(e(\mu, \nu)) \sum_{s=1}^n e(\beta_s, \gamma_s)w^j \\ &= \sum_{s=1}^n e(\bar{\beta}_s \cdot \alpha^i(\mu), \gamma_s \cdot \nu_j)w^j. \end{aligned}$$

Here

$$\beta_s = (1, \dots, 1, \underset{i-1}{s}, 1, \dots, 1) \in W_{[0, j+i-1]}^n,$$

$$\gamma_s = (1, \dots, 1, s) \in W_{[0, j+i-1]}^n$$

and

$$\bar{\beta}_s = (1, \dots, 1, \underset{i-1}{s}) \in W_{[0, i-1]}^n, \quad \nu_j = (\nu_{j+i}, \dots, \nu_b) \in W_{[j+i, b+i]}^n.$$

Hence for $k \geq a$ and $m \geq b+i$ we have

$$\|\psi_{k,m} \cdot \varphi_{k,m}(\Phi^i(e(\mu, \nu)w^j)) - \Phi^i(e(\mu, \nu)w^j)\| \leq \frac{an}{2k+1}.$$

Therefore, we have for $N \in \mathbb{N}$

$$\begin{aligned} rcp \left(\bigcup_{i=0}^N \Phi^i \left(\omega_{a,b} \cup (\omega_{a,b})^* : \frac{an}{2k+1} \right) \right) &\leq \text{rank } B(K(k, N+b+1)) \\ &= (2k+1)n^{N+b+1}, \end{aligned}$$

where $(\omega_{a,b})^* = \{x^*; x \in \omega_{a,b}\}$. This implies that for all integers a, b with $a < b$,

$$ht \left(\Phi, \omega_{a,b} \cup (\omega_{a,b})^* ; \frac{an}{2k+1} \right) \leq \lim_{N \rightarrow \infty} \frac{1}{N} \log \left((2k+1)n^{N+b+1} \right) = \log n.$$

Increasing k , we have $ht(\Phi, \omega_{a,b} \cup (\omega_{a,b})^*) \leq \log n$, for all $a, b \in \mathbb{N}$ with $a < b$. Put $\omega_a = \omega_{a,2a} \cup (\omega_{a,2a})^*$, for $a \in \mathbb{N}$. Then the set $\{\omega_a : a \in \mathbb{N}\}$ is an increasing sequence of finite subsets of $p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p$ and the linear span of $\bigcup \{\omega_a : a \in \mathbb{N}\}$ is dense in $p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p$. Hence

$$ht(\Phi) = \sup_{a \in \mathbb{N}} ht(\Phi, \omega_a) \leq \log n.$$

On the other hand, the restriction $\Phi|_{\mathcal{A}_0}$ of Φ to \mathcal{A}_0 is $\sigma|_{\mathcal{A}_0} = \alpha_n|_{\mathcal{A}_0}$ and $h_{\tau}(\alpha_n|_{\mathcal{A}_0}) = h_{\tau}(\alpha_n) = \log n$. Since there exists a conditional expectation of \mathcal{O}_n onto \mathcal{A}_0 and $\tau \cdot \alpha_n|_{\mathcal{A}_0} = \tau$,

$$\log n = h_{\tau}(\alpha_n|_{\mathcal{A}_0}) \leq ht(\Phi|_{\mathcal{A}_0}) \leq ht(\Phi) \leq \log n$$

by the version for unital $*$ -endomorphisms of [V, Proposition 4.4]. Therefore, $ht(\Phi) = \log n$. \square

Corollary 4.7. *Let ϕ be the unique $\log n$ -KMS state of \mathcal{O}_n . Then*

$$h_{\phi}(\Phi) = \log n.$$

Proof. Let τ be the unique tracial state of \mathcal{A}_0 and E be the conditional expectation of $p(\mathcal{A} \rtimes_{\sigma} \mathbb{Z})p$ onto \mathcal{A}_0 , then $\phi = \tau \cdot E$. Hence $\phi \cdot \Phi = \phi$. This relation implies, by the endomorphism version of [V, Proposition 4.6],

$$\log n = h_{\tau}(\sigma|_{\mathcal{A}_0}) \leq h_{\phi}(\Phi) \leq ht(\Phi) = \log n.$$

Therefore $h_{\phi}(\Phi) = \log n$. \square

5. Entropy of Longo's canonical endomorphism.

In this section we apply the result in §4 to Longo's canonical endomorphism. We use the same notations as in §4.

5.1. Let τ_i be the tracial state of A_i , for $i \in \mathbb{N}$ and let

$$\tilde{A} = \bigotimes_{i=0}^{\infty} (A_i, \tau_i).$$

The \tilde{A} has the canonical trace $\bigotimes_{i=0}^{\infty} \tau_i$, which we denote by τ . The shift $\sigma|_{\mathcal{A}_0}$ is extended to the $*$ -endomorphism γ of the hyperfinite II_1 factor \tilde{A} . The γ is canonical in the sense of [Ch3]. Hence we have the extension algebra $\tilde{M} = \langle \tilde{A}, \sigma \rangle$, which is the injective type $\text{III}_{1/n}$ factor generated by \tilde{A} and an isometry W . Then γ is extended to the canonical $*$ -endomorphism Γ of \tilde{M} and

$$\Gamma(a) = \gamma(a), a \in \tilde{A}, \quad \text{and} \quad \Gamma(W) = \pi_{\phi}(vW).$$

The Γ is Longo's canonical endomorphism for the inclusion $\tilde{N} \subset \tilde{M}$ [Ch3, Theorem 6.10]. Here the subfactor \tilde{N} is obtained naturally in the step of

constructing \tilde{M} . The factor \tilde{M} is the von Neumann algebra generated by $\pi_\phi(\langle \mathcal{A}_0, \sigma|_{\mathcal{A}_0} \rangle)$ and the C^* -algebra $\langle \mathcal{A}_0, \sigma|_{\mathcal{A}_0} \rangle$ is \mathcal{O}_n . Hence Γ is the extension of Φ to \tilde{M} . Since Φ is ϕ -preserving, as an application of 4.7 Corollary, we have the following by [CNT, Theorem VII.2]:

Corollary 5.2. *Let M be the von Neumann algebra generated by $\pi_\phi(\mathcal{O}_n)$ and let Γ be the extension of Cuntz's canonical endomorphism Φ of \mathcal{O}_n to M . Then Γ is Longo's canonical endomorphism and*

$$h_\phi(\Gamma) = \log n.$$

References

- [Ch1] M. Choda, *Entropy for $*$ -endomorphisms and relative entropy for subalgebras*, J. Operator Theory, **25** (1991), 125-140.
- [Ch2] ———, *Canonical $*$ -endomorphisms and simple C^* -algebras*, J. Operator Theory, **33** (1995), 235-248.
- [Ch3] ———, *Extension algebras of II_1 factors via $*$ -endomorphisms*, Int. J. Math., **5** (1994), 635-655.
- [CNT] A. Connes, H. Narnhofer and W. Thirring, *Dynamical entropy of C^* algebras and von Neumann algebras*, Commun. Math. Phys., **112** (1987), 691-719.
- [CS] A. Connes and E. Størmer, *Entropy of II_1 von Neumann algebras*, Acta Math., **134** (1975), 289-306.
- [Cu1] J. Cuntz, *Simple C^* -algebras generated by isometries*, Commun. Math. Phys., **57** (1977), 173-185.
- [Cu2] ———, *Regular actions of Hopf algebras on the C^* -algebra generated by a Hilbert space*, in 'Operator algebras, mathematical physics, and low dimensional topology' (Istanbul, 1991), 87-100, Res. Notes Math., **5**, Wellesley, MA, (1993).
- [HS] U. Haagerup and E. Størmer, *Equivalence of normal states on von Neumann algebras and flow of weights*, Adv. Math., **83** (1990), 180-262.
- [Hu] T. Hudetz, *Topological entropy for appropriately approximated C^* -algebras*, J. Math. Phys., **35** (1994), 4303-4333.
- [I1] M. Izumi, *Canonical extension on factors, in Subfactors*, H. Araki et al. (eds.), World Scientific, Singapore, (1994), 129-138.
- [I2] ———, *Subalgebras of infinite C^* -algebras with finite indices, II*, Cuntz-Krieger algebras, preprint.
- [L1] R. Longo, *Index of subfactors and statistics of quantum fields, I*, Commun. Math. Phys., **126** (1989), 145-155.
- [L2] ———, *Duality for Hopf algebras and for subfactors*, Commun. Math. Phys., **159** (1994), 133-150.
- [LR] R. Longo and J.E. Roberts, *A theory of dimension*, preprint.
- [OP] D. Olesen and G.K. Pedersen, *Some C^* -dynamical systems with a single KMS state*, Math. Scand., **42** (1978), 111-118.

- [P] W. Paschke, *The crossed product of a C^* -algebra by an endomorphism*, Proc. Amer. Math. Soc., **80** (1980), 113-118.
- [R] M. Rørdam, *Classification of certain infinite simple C^* -algebras*, J. Funct. Anal., **131** (1995), 415-458.
- [S] E. Størmer, *Entropy in operator algebras*, preprint in University of Oslo, (1994).
- [T] K. Thomsen, *Topological entropy for endomorphisms of local C^* -algebras*, Comm. Math. Phys., **164** (1994), 181-193.
- [V] D. Voiculescu, *Dynamical approximation entropies and topological entropy in operator algebras*, Comm. Math. Phys., **170** (1995), 249-281.

Received July 1, 1997 and revised October 2, 1998.

OSAKA KYOIKU UNIVERSITY

KASHIWARA 582-8582

JAPAN

E-mail address: marie@cc.osaka-kyoiku.ac.jp