

Pacific Journal of Mathematics

APPROXIMATION OF RECURRENCE IN NEGATIVELY
CURVED METRIC SPACES

CHARALAMBOS CHARITOS AND GEORGIOS TSAPOGAS

Volume 195 No. 1

September 2000

APPROXIMATION OF RECURRENCE IN NEGATIVELY CURVED METRIC SPACES

CHARALAMBOS CHARITOS AND GEORGIOS TSAPOGAS

For metric spaces with curvature less than or equal to χ , $\chi < 0$, it is shown that a recurrent geodesic is approximated by closed geodesics. A counter example is provided for the converse.

1. Introduction and preliminaries.

In hyperbolic geometry it has been shown lately that many geometric properties are determined by the distance function on the space itself rather than the differential structure. It is shown in this work that, partially, this is the case with the notion of recurrence. For complete hyperbolic manifolds, a recent result of Aebischer, Hong and McCullough (see [1]) states that a geodesic is recurrent if and only if it is approximated by closed geodesics. We show that, in metric spaces with curvature less than or equal to χ , $\chi < 0$, recurrent geodesics are approximated by closed geodesics (see Theorem 2 below). The proof of the converse statement crucially depends on the manifold structure, in particular on the fact that two geodesics coincide if they do so on an open interval. Hence, the converse statement fails in our context due to the bifurcation property of geodesics. A counterexample exhibiting this failure is provided in Section 4 below. A geodesic γ is called *recurrent* if there exists a sequence $\{t_n\} \subset \mathbb{R}$, $t_n \rightarrow \infty$ such that $t_n\gamma \rightarrow \gamma$ as $t_n \rightarrow \infty$. Convergence in this definition is meant to be uniform convergence on compact sets which, in fact, induces the topology on the space GX consisting of all (local) isometries $\mathbb{R} \rightarrow X$ when X is (not) simply connected. \mathbb{R} acts on GX by right translations, namely, $(t, g) \rightarrow tg$, where $tg : \mathbb{R} \rightarrow X$ is the geodesic defined by $tg(s) = g(s + t)$, $s \in \mathbb{R}$. This action is simply the geodesic flow. The notion of convergence in the above definition is analogous to the tangential condition which defines recurrence in the manifold case. We use the notion of approximation given in Definition 6 below which was introduced in [1] in order to characterize recurrent geodesics in hyperbolic manifolds.

X will always denote a locally compact, complete, geodesic metric space with curvature less than or equal to χ , $\chi < 0$. Recall that a geodesic metric space is said to have curvature less than or equal to χ if each $x \in X$ has a neighborhood V_x such that every geodesic triangle of perimeter strictly

less than $\frac{2\pi}{\sqrt{\chi}}$ ($=+\infty$ when $\chi \leq 0$) contained in V_x satisfies $CAT - (\chi)$ inequality (see [11] for definitions and basic properties). We will denote the metric by $d(\cdot, \cdot)$ and will use the same letter to denote distance when the metric space to which we refer is understood. All curves are assumed to be parametrized by arclength. A *geodesic segment* in X is an isometry $c : I \rightarrow X$, where I is a closed interval in \mathbb{R} . A *geodesic* in X is a map $c : \mathbb{R} \rightarrow X$ such that for each closed interval $I \subset \mathbb{R}$, the map $c|_I : I \rightarrow X$ is a geodesic segment. A *local geodesic segment* (usually called *geodesic arc*) in X is a map $c : I \rightarrow X$ such that for each $t \in I$ there is an $\varepsilon > 0$ such that $c|_{[t-\varepsilon, t+\varepsilon] \cap I} : [t-\varepsilon, t+\varepsilon] \cap I \rightarrow X$ is a geodesic segment. Similarly, a *local geodesic* $\mathbb{R} \rightarrow X$ is defined. A *closed geodesic* in X is a local geodesic $c : \mathbb{R} \rightarrow X$ which is a periodic map.

Definition 1. An oriented geodesic g in X is said to be approximated by closed geodesics if, for every $\varepsilon > 0$ and every $x \in \text{Im } g$, there exists a closed oriented geodesic c such that for some point $y \in \text{Im } c$,

$$d(c(t+t_y), g(t+t_x)) < \varepsilon$$

for all $t \in [0, \text{period}(c)]$, where $t_x, t_y \in \mathbb{R}$ with $x = g(t_x)$ and $y = c(t_y)$.

The following theorem is the main result of this paper.

Theorem 2. *Let X be a locally compact, complete, geodesic metric space which has curvature less than or equal to χ , $\chi < 0$. If a geodesic or geodesic ray in X is recurrent, then it is approximated by closed geodesics.*

The proof of Theorem 2 uses the notion of quasi-geodesic and its stability properties. We will closely follow notation and terminology appearing in [8, Ch. 3] where we refer the reader for first definitions and basic properties of quasi-geodesics. Here we only recall the following definition.

Definition 3. Let $f : [a, b] \rightarrow X$ be a continuous map with $-\infty \leq a \leq b \leq +\infty$ and λ, κ, L real numbers with $\lambda \geq 1, \kappa \geq 0, L > 0$.

f is a (λ, κ, L) -quasi-geodesic if for every subinterval $[a', b']$ of $[a, b]$ satisfying

$$\text{length } f([a', b']) \leq L,$$

the following inequality holds

$$\text{length } f([a', b']) \leq \lambda d(f(a'), f(b')) + \kappa.$$

The next proposition is a well know fact for $CAT - (\chi)$ spaces. We include a short proof of it, since it is difficult to find exact reference (when X is a geometric polyhedron this result follows from [3, p. 403]).

Proposition 4. *Let M be a complete geodesic space satisfying $CAT - (\chi)$ inequality with $\chi < 0$. Every local geodesic segment in M is a geodesic segment.*

Proof. Let $\delta : [0, L] \rightarrow M$ be a local geodesic segment in M , $L > 0$. Set

$$l = \sup \{ t \in [0, L] \mid \delta|_{[0, t]} \text{ is a geodesic segment} \}.$$

Apparently, $l > 0$ and by completeness of M , $\delta|_{[0, l]}$ is a geodesic segment joining $\delta(0)$ with $\delta(l)$. Assuming the conclusion is not true, i.e., $l < L$, let ε be a positive number such that $\delta|_{[l-\varepsilon, l+\varepsilon]}$ is a geodesic segment. Denote by $[\delta(0), \delta(l+\varepsilon)]$ the geodesic segment in M joining $\delta(0)$ with $\delta(l+\varepsilon)$. Since $\delta|_{[0, l+\varepsilon]}$ is not the geodesic segment joining $\delta(0)$ with $\delta(l+\varepsilon)$,

$$(1) \quad d(\delta(0), \delta(l+\varepsilon)) < d(\delta(0), \delta(l)) + d(\delta(l), \delta(l+\varepsilon)).$$

The points $\delta(0)$, $\delta(l)$ and $\delta(l+\varepsilon)$ define a geodesic triangle in M . Denote by $\Delta = (\overline{\delta(0)}, \overline{\delta(l)}, \overline{\delta(l+\varepsilon)})$ the corresponding comparison triangle which is non-degenerate by inequality (1). Choose points B on $\delta|_{[0, l]}$ and B' on $\delta|_{[l, l+\varepsilon]}$ such that $d(B, \delta(l)) = d(B', \delta(l)) = \varepsilon' < \varepsilon$ and denote by \overline{B} and $\overline{B'}$ the corresponding points on the comparison triangle. Then by (1) the angle of Δ at $\overline{\delta(l)}$ is smaller than π and therefore

$$d(\overline{B}, \overline{B'}) < d(\overline{B}, \overline{\delta(l)}) + d(\overline{\delta(l)}, \overline{B'}) = 2\varepsilon'.$$

By comparison, $d(B, B') \leq d(\overline{B}, \overline{B'})$ so we obtain

$$d(B, B') < d(B, \delta(l)) + d(\delta(l), B').$$

This contradicts the fact that $\delta|_{[l-\varepsilon', l+\varepsilon']}$ is a geodesic segment. \square

Let \tilde{X} be the universal cover of X and $p : \tilde{X} \rightarrow X$ the projection map. \tilde{X} becomes a metric space as follows: Given $\tilde{x}, \tilde{y} \in \tilde{X}$ choose any curve $\tilde{c} : [a, b] \rightarrow \tilde{X}$ with $\tilde{c}(a) = \tilde{x}$ and $\tilde{c}(b) = \tilde{y}$ and define the distance from \tilde{x} to \tilde{y} to be the length of the unique length minimizing curve in the homotopy class of $p\tilde{c}$ with endpoints fixed. For the existence of the length minimizing curve see [10]. This distance function is a metric on \tilde{X} which inherits the properties of X , namely, \tilde{X} becomes a complete geodesic locally compact (hence, proper) metric space. $\pi_1(X)$ acts on \tilde{X} and the action commutes with p . As the projection p is a local isometry, it follows that $\pi_1(X)$ acts on \tilde{X} by local isometries. Using the fact that \tilde{X} is geodesic and Proposition 4, it is routine to show that $\pi_1(X)$ acts on \tilde{X} by isometries. In addition, \tilde{X} has curvature less than or equal to χ , $\chi < 0$ and, by a theorem of Gromov (see for example [11, p. 325]), \tilde{X} satisfies $CAT - (\chi)$ inequality.

GX is by definition the space of all local geodesics $\mathbb{R} \rightarrow X$ and, by Proposition 4 above, $G\tilde{X}$ is the space consisting of all global geodesics $\mathbb{R} \rightarrow \tilde{X}$. The topology on these spaces is uniform convergence on compact sets. The boundary $\partial\tilde{X}$ can be defined using either equivalence classes of sequences or, equivalence classes of geodesic rays. The local compactness assumption on X implies that \tilde{X} is proper and hence the two definitions coincide (see

[8, Ch. 2]). We will be using them interchangeably. For any two distinct points ξ, η in $\partial\tilde{X}$ there exists a unique, up to parametrization, (oriented) geodesic g with $g(-\infty) = \xi$ and $g(\infty) = \eta$ (see for example [5, Prop. 2]). We need the following lemma which asserts that the projection of a point onto a geodesic always exists.

Lemma 5. *Let g be a geodesic in $G\tilde{X}$ (or a geodesic segment) and x_0 a point in \tilde{X} . There exists a unique real number s such that $g(s)$ realizes the distance of x_0 from $\text{Im } g$, i.e., $\text{dist}(x_0, \text{Im } g) = d(x_0, g(s))$.*

Proof. We may assume that $x_0 \notin \text{Im } g$. Existence is apparent. Assume that $s \neq s'$ are two such numbers. The points $g(s), g(s')$ and x_0 define a non-degenerate geodesic triangle in \tilde{X} and denote by $\Delta = \left(\overline{g(s)}, \overline{g(s')}, \overline{x_0}\right)$ the corresponding comparison triangle. Δ is an equilateral triangle in the unique complete simply connected Riemannian 2-manifold of constant sectional curvature χ . Hence, the angles of Δ at $\overline{g(s)}$ and $\overline{g(s')}$ are each less than $\pi/2$. Therefore, there exists a point $\overline{g(t)}$ on the side of Δ opposite to x_0 such that $d(x_0, \overline{g(t)}) < d(x_0, \overline{g(s)}) = d(x_0, \overline{g(s')})$. By $CAT - (\chi)$ inequality, $d(x_0, g(t)) \leq d(x_0, g(s))$, a contradiction. \square

Remark 1. If $c \in GX$ is a closed geodesic and $x_0 \in X$, the same argument applied to a lifting \tilde{c} of c shows that there exists a unique point $B \in \text{Im } c$ such that $d(x_0, B) = \text{dist}(x_0, \text{Im } c)$.

Remark 2. Set $\partial^2\tilde{X} = \left\{(\xi, \eta) \in \partial\tilde{X} \times \partial\tilde{X} : \xi \neq \eta\right\}$ and let $\rho : G\tilde{X} \rightarrow \partial^2\tilde{X}$ be the fiber bundle given by $\rho(g) = (g(-\infty), g(+\infty))$. Since for any two distinct points ξ, η in $\partial\tilde{X}$ there exists a unique (oriented) geodesic g with $g(-\infty) = \xi$ and $g(\infty) = \eta$ (see for example [5, Prop. 2]), the fiber of ρ is \mathbb{R} . Moreover, this bundle is trivial (see for example [4, Th. 4.8]). To define a trivialization, let x_0 be a base point and let

$$(2) \quad H : G\tilde{X} \xrightarrow{\cong} \partial^2\tilde{X} \times \mathbb{R}$$

be the trivialization of ρ with respect to x_0 defined by

$$H(g) = (g(-\infty), g(+\infty), s)$$

where $-s$ is the real number provided by Lemma 5. Note that the composite of the geodesic flow $\mathbb{R} \times G\tilde{X} \rightarrow G\tilde{X}$ with H is given by the formula

$$(\xi_1, \xi_2, s) \longrightarrow (\xi_1, \xi_2, s + t)$$

for all $(\xi_1, \xi_2) \in \partial^2\tilde{X}$ and $s \in \mathbb{R}$.

2. Recurrent geodesics.

Definition 6. A geodesic γ in X is called recurrent if there exists a sequence $\{t_n\} \subset \mathbb{R}, t_n \rightarrow \infty$ such that $t_n \gamma \rightarrow \gamma$ as $t_n \rightarrow \infty$.

For a recurrent geodesic γ in X there exists a sequence of closed (in fact, piece-wise geodesic) curves $\{\gamma_n\}_{n \in \mathbb{N}}$, associated to γ as follows: Fix a convex neighborhood U of $\gamma(0)$, i.e., a neighborhood which satisfies the following property: For all $x, y \in U$ there exists a unique geodesic segment with endpoints x and y lying entirely in U . Such a neighborhood exists (see for example [2]). If $\{t_n\}$ is the sequence given by Definition 6 above and $\varepsilon_n = d(\gamma(0), \gamma(t_n))$, let $K \in \mathbb{N}$ such that $\gamma(t_n) \in U$ for all $n \geq K$. Define $\gamma_n, n \geq K$ to be the curve

$$(3) \quad \gamma_n : [0, t_n + \varepsilon_n] \rightarrow X$$

with $\gamma_n(t) = \gamma(t) \ \forall t \in [0, t_n]$ and $\gamma_n|_{[t_n, t_n + \varepsilon_n]}$ the unique geodesic segment in U joining $\gamma(t_n)$ with $\gamma(0)$. Note that $t_n + \varepsilon_n$ is the period of the closed curve γ_n . In the sequel, we will refer to these closed curves by writing $\gamma_n, n \in \mathbb{N}$ but it will always be implicit that n is large enough so that γ_n are defined.

Using the following lemma, we may assume that given a recurrent geodesic γ , the associated closed curves $\{\gamma_n\}_{n \in \mathbb{N}}$ are not homotopic to a point.

Lemma 7. *Given a recurrent geodesic γ there exists $M \in \mathbb{N}$ such that each closed curve $\gamma_n, n \in \mathbb{N}$ associated to γ is not homotopic to a point, provided $n \geq M$.*

Proof. Let $\tilde{\gamma}$ be a lift of γ to the universal cover \tilde{X} of X parametrized so that $\tilde{\gamma}(0)$ projects to $\gamma(0) = \gamma_n(0)$. The curve $\gamma_n|_{[0, t_n]}$ is a local geodesic segment and, by Proposition 4, its lift $\tilde{\gamma}_n|_{[0, t_n]}$ to \tilde{X} starting at $\tilde{\gamma}(0)$ is a geodesic segment. Moreover, $\gamma_n|_{[t_n, t_n + \varepsilon_n]}$ and its lift $\tilde{\gamma}_n|_{[t_n, t_n + \varepsilon_n]}$ to \tilde{X} starting at $\tilde{\gamma}_n(t_n)$ are both geodesic segments. We have

$$\begin{aligned} d(\tilde{\gamma}_n(t_n + \varepsilon_n), \tilde{\gamma}_n(0)) &\geq d(\tilde{\gamma}_n(t_n), \tilde{\gamma}_n(0)) - d(\tilde{\gamma}_n(t_n + \varepsilon_n), \tilde{\gamma}_n(t_n)) \\ &= t_n - \varepsilon_n. \end{aligned}$$

Since $\varepsilon_n \rightarrow 0$ and $t_n \rightarrow \infty$ as $n \rightarrow \infty$, we may choose $M \in \mathbb{N}$ such that $\tilde{\gamma}_n(t_n + \varepsilon_n), \tilde{\gamma}_n(0)$ are distinct for all $n \geq M$. Therefore, $\tilde{\gamma}_n|_{[0, t_n + \varepsilon_n]}$, which is the lift of the closed curve γ_n starting at $\tilde{\gamma}(0) = \tilde{\gamma}_n(0)$, has distinct endpoints and, therefore, $\gamma_n, n \geq M$ is not homotopic to a point. \square

The following proposition shows that the lifts (to the universal cover \tilde{X}) of the closed curves γ_n associated to a recurrent geodesic γ are, for n large enough, quasi-geodesics with arbitrarily large L . Recall that a $CAT - (\chi)$ space is a δ -hyperbolic space in the sense of Gromov (see for example [11,

Sec. 2]). This applies to the universal covering \tilde{X} , since it satisfies $CAT-(\chi)$ inequality globally. Let δ denote the hyperbolicity constant of the space \tilde{X} .

Proposition 8. *Let γ be a recurrent geodesic in X and $\{\gamma_n\}_{n \in \mathbb{N}}$ the associated closed curves. For every $L > 0$, there exists $N \in \mathbb{N}$ such that all lifts $\tilde{\gamma}_n : \mathbb{R} \rightarrow \tilde{X}$ of γ_n with $n \geq N$ are (λ, κ, L) -quasi-geodesics provided $\kappa > 16\delta$ and $\lambda = 1$, where δ is the hyperbolicity constant of \tilde{X} .*

Proof. Let γ be a recurrent geodesic and $L > 0$ be given. The sequence $\{t_n\}$ given by Definition 6 converges to infinity. Moreover, $\varepsilon_n = d(\gamma(0), \gamma(t_n)) \rightarrow 0$ and $t_n + \varepsilon_n = \text{period}(\gamma_n)$ also converges to infinity as $n \rightarrow \infty$. Hence, we may choose N such that

$$(4) \quad t_n + \varepsilon_n > L \quad \text{and} \quad \varepsilon_n < \frac{1}{2}(\kappa - 16\delta) \quad \text{for all } n \geq N.$$

Let now $[a, b]$ be any interval with $b - a < L$ (cf. Definition 3). For each $n \geq N$ there exists an integer k_n such that

$$(5) \quad \tilde{\gamma}_n([a, b]) \subset \tilde{\gamma}_n([(k_n - 1)(t_n + \varepsilon_n), k_n(t_n + \varepsilon_n) + t_n]).$$

Denote by $[\tilde{\gamma}_n(a), \tilde{\gamma}_n(b)]$ the unique geodesic segment in \tilde{X} joining $\tilde{\gamma}_n(a)$ with $\tilde{\gamma}_n(b)$ and set

$$\begin{aligned} A_n &:= \tilde{\gamma}_n(k_n(t_n + \varepsilon_n) - \varepsilon_n) \\ y_{k_n} &:= \tilde{\gamma}_n(k_n(t_n + \varepsilon_n)). \end{aligned}$$

The distance of any point on $[\tilde{\gamma}_n(a), \tilde{\gamma}_n(b)]$ from $\tilde{\gamma}_n([a, b])$ is bounded by a number which depends on the hyperbolicity constant δ of the space \tilde{X} and on the number of geodesic segments which constitute $\tilde{\gamma}_n([a, b])$, see [8, Lemma 1.5, p. 25]. In our case here, $\tilde{\gamma}_n([a, b])$ consists of at most three geodesic segments (since the right hand side of inclusion (5) above consists of 3 geodesic segments) and the bound is 8δ . Hence we have

$$(6) \quad d(y_{k_n}, [\tilde{\gamma}_n(a), \tilde{\gamma}_n(b)]) \leq \varepsilon_n.$$

By Lemma 5, let B_n be the point on $[\tilde{\gamma}_n(a), \tilde{\gamma}_n(b)]$ which realizes the distance in the left hand side of inequality 6. Assume that neither $\tilde{\gamma}_n(a)$ nor $\tilde{\gamma}_n(b)$ lies on the geodesic segment $[A_n, y_{k_n}]$. Then we have the following triangle inequalities

$$\begin{aligned} d(\tilde{\gamma}_n(a), A_n) &\leq d(\tilde{\gamma}_n(a), B_n) + d(B_n, y_{k_n}) + d(y_{k_n}, A_n) \\ d(y_{k_n}, \tilde{\gamma}_n(b)) &\leq d(y_{k_n}, B_n) + d(B_n, \tilde{\gamma}_n(b)) \end{aligned}$$

which, after employing the fact that $d(A_n, y_{k_n}) = \varepsilon_n$, become

$$\begin{aligned} \text{length } \tilde{\gamma}_n([a, b]) &= d(\tilde{\gamma}_n(a), A_n) + d(A_n, y_{k_n}) + d(y_{k_n}, \tilde{\gamma}_n(b)) \\ &\leq 2\varepsilon_n + 2d(y_{k_n}, B_n) + d(\tilde{\gamma}_n(a), \tilde{\gamma}_n(b)) \\ &\leq 2\varepsilon_n + 2 \cdot 8\delta + d(\tilde{\gamma}_n(a), \tilde{\gamma}_n(b)) \\ &\leq \kappa + d(\tilde{\gamma}_n(a), \tilde{\gamma}_n(b)). \end{aligned}$$

by inequality (6)
by inequality (4)

The case where $\widetilde{\gamma}_n(a)$ and/or $\widetilde{\gamma}_n(b)$ lies on $[A_n, y_{k_n}]$ is treated similarly. \square

Corollary 9. *For $n \in \mathbb{N}$ sufficiently large, the isometry of \widetilde{X} in $\pi_1(X)$ which corresponds to the homotopy class of the closed curve γ_n is hyperbolic.*

Proof. It suffices to show that each $\widetilde{\gamma}_n : \mathbb{R} \rightarrow \widetilde{X}$ determines exactly two boundary points $\widetilde{\gamma}_n(-\infty)$, $\widetilde{\gamma}_n(+\infty)$. By Lemma 8 there exists an $M \in \mathbb{N}$ such that $\widetilde{\gamma}_n$ is a quasi-isometry for all $n \geq M$. Each such $\widetilde{\gamma}_n$ induces a map $\partial\mathbb{R} \rightarrow \partial\widetilde{X}$ which is a homeomorphism onto its image, see [8, Th. 2.2, p. 35]. As $\partial\mathbb{R}$ consists of two distinct points, $\widetilde{\gamma}_n(-\infty)$, $\widetilde{\gamma}_n(+\infty) \in \partial\widetilde{X}$ are also distinct for all $n \geq M$. \square

It now follows that a recurrent geodesic γ in X as well as each of the (oriented) closed curves γ_n , $n \geq M$ (cf. Lemma 7 and Corollary 9 above) determine exactly two boundary points in $\partial\widetilde{X}$ denoted by $\widetilde{\gamma}(-\infty)$, $\widetilde{\gamma}(+\infty)$ and $\widetilde{\gamma}_n(-\infty)$, $\widetilde{\gamma}_n(+\infty)$ respectively. We need the following lemma concerning these boundary points. Recall that $\widetilde{X} \cup \partial\widetilde{X}$ is a compact space which is metrizable (see [8, p. 134]), and we will denote such metric by $d_{\widetilde{X} \cup \partial\widetilde{X}}$.

Lemma 10. *$\widetilde{\gamma}_n(-\infty) \rightarrow \widetilde{\gamma}(-\infty)$ and $\widetilde{\gamma}_n(+\infty) \rightarrow \widetilde{\gamma}(+\infty)$ as $n \rightarrow \infty$.*

Proof. As above, let $\varepsilon_n = \text{length}(\text{Im } \gamma_n) - t_n$ so that $t_n + \varepsilon_n$ is the period of γ_n . We first show that $\widetilde{\gamma}_n(+\infty) \rightarrow \widetilde{\gamma}(+\infty)$. Consider the sequence $\widetilde{\gamma}_n(k(t_n + \varepsilon_n))$, $k \in \mathbb{N}$ which converges to $\widetilde{\gamma}_n(+\infty)$ as $k \rightarrow \infty$. Thus, there exists $k_n \in \mathbb{N}$ such that

$$(7) \quad d_{\widetilde{X} \cup \partial\widetilde{X}}(\widetilde{\gamma}_n(k_n(t_n + \varepsilon_n)), \widetilde{\gamma}_n(+\infty)) < 1/n.$$

Now consider the sequences $y_n := \widetilde{\gamma}_n(k_n(t_n + \varepsilon_n))$ and $x_n := \widetilde{\gamma}(t_n)$, $n \in \mathbb{N}$. Since $x_n \rightarrow \widetilde{\gamma}(+\infty)$, by inequality (7) above it is enough to show that the sequences $\{x_n\}$ and $\{y_n\}$ represent the same element in $\partial\widetilde{X}$ or, in other words, that the hyperbolic product $(x_n, y_n)_{x_0}$ with respect to the base point $x_0 := \widetilde{\gamma}(0)$ converges to $+\infty$ as $n \rightarrow +\infty$. For the notion of hyperbolic product of sequences and their equivalence, see [8].

The stability property of quasi-geodesics states (see Corollary 1.10 of [8, p. 31]) that given any two numbers $\kappa \geq 0$ and $\lambda \geq 1$, there exists a constant C depending on λ, κ and on the hyperbolicity constant δ of the space such that if L is bigger than $2C$ then every (λ, κ, L) -quasi-geodesic $f : [a, b] \rightarrow \widetilde{X}$ lies within a C -neighborhood of the geodesic segment $[f(a), f(b)]$. By choosing $\lambda = 1$, $\kappa > 16\delta$ where δ is the hyperbolicity constant of the space \widetilde{X} and $L > 2C$ we obtain, by Proposition 8 above, a natural number N such that all $\widetilde{\gamma}_n : \mathbb{R} \rightarrow \widetilde{X}$ with $n \geq N$ are (λ, κ, L) -quasi-geodesics. In particular, $\widetilde{\gamma}_n : [0, k_n(t_n + \varepsilon_n)] \rightarrow \widetilde{X}$ are (λ, κ, L) -quasi-geodesics for all $n \geq N$. Therefore, by Corollary 1.10 of [8, p. 31] as explained above,

$$d(x_n, x'_n) < C \quad \forall n \geq N$$

where x'_n denotes the projection of x_n on the geodesic segment $[\widetilde{\gamma}(0), y_n]$ (cf. Lemma 5). Hence,

$$\begin{aligned} (x_n, y_n)_{x_0} &= \frac{1}{2} (d(x_n, x_0) + d(y_n, x_0) - d(x_n, y_n)) \\ &\geq \frac{1}{2} (d(x'_n, x_0) - C + d(y_n, x_0) - d(x'_n, y_n) - C) \\ &= (x'_n, y_n)_{x_0} - C \\ &= d(x_0, x'_n) - C. \end{aligned}$$

Apparently, $d(x_0, x'_n) \rightarrow \infty$ as $n \rightarrow +\infty$ and, hence, $(x_n, y_n)_{x_0} \rightarrow \infty$ as required.

In order to show that $\widetilde{\gamma}_n(-\infty) \rightarrow \widetilde{\gamma}(-\infty)$ we work in a similar manner: The sequence $\widetilde{\gamma}_n(-k(t_n + \varepsilon_n))$, $k \in \mathbb{N}$ converges to $\widetilde{\gamma}_n(-\infty)$ as $k \rightarrow \infty$. Hence, there exists $k_n \in \mathbb{N}$ such that $d_{\widetilde{X} \cup \partial \widetilde{X}}(\widetilde{\gamma}_n(-k_n(t_n + \varepsilon_n)), \widetilde{\gamma}_n(-\infty)) < 1/n$. As before, sequences $\{y_n\}$ and $\{x_n\}$ are defined by $y_n := \widetilde{\gamma}_n(-k_n(t_n + \varepsilon_n))$ and $x_n := \widetilde{\gamma}(-t_n)$, $n \in \mathbb{N}$. Then we use the same arguments to show that the hyperbolic product $(x_n, y_n)_{x_0}$ with respect to the base point $x_0 := \widetilde{\gamma}(0)$ converges to $+\infty$ as $n \rightarrow +\infty$. \square

3. Proof of main theorem.

Let γ be a recurrent geodesic, $\varepsilon > 0$ and $x \in \text{Im } \gamma$ be given. We may assume that $x = \gamma(0)$. Let $\{t_n\}$ be the sequence given by Definition 6 and $\{\gamma_n\}$ the sequence of the associated closed curves given by formula (3) above. For each $n \in \mathbb{N}$, there exists a unique closed geodesic c_n in the free homotopy class of γ_n . The number $t_n + \varepsilon_n$ is the period of γ_n and let s_n denote the period of c_n (apparently, $s_n < t_n + \varepsilon_n$). Let B_n be the projection of $\gamma(0)$ onto $\text{Im } c_n$, i.e., $d(\gamma(0), B_n) = d(\gamma(0), \text{Im } c_n)$. Such a point exists and is unique by Remark 1 following Lemma 5. Lift γ to an isometry $\widetilde{\gamma} : \mathbb{R} \rightarrow \widetilde{X}$ with a base point $\widetilde{\gamma}(0)$ satisfying $p(\widetilde{\gamma}(0)) = \gamma(0)$, where $p : \widetilde{X} \rightarrow X$ is the universal covering map. Lift each c_n to an isometry $\widetilde{c}_n : \mathbb{R} \rightarrow \widetilde{X}$ and parametrize it so that $\widetilde{c}_n(0)$ is a point \widetilde{B}_n satisfying

$$d(\widetilde{B}_n, \widetilde{\gamma}(0)) = d(B_n, \gamma(0)) \quad \text{and} \quad p(\widetilde{B}_n) = B_n.$$

For the reader's convenience, we have gathered all the above notation in Figure 1.

Since

$$p(\widetilde{\gamma}_n(t_n + \varepsilon_n)) = p(\widetilde{\gamma}(0)) = \gamma(0)$$

and γ_n, c_n are homotopic, the isometry ϕ_n of \widetilde{X} which translates \widetilde{c}_n (in the positive direction) satisfies

$$\phi_n(\widetilde{\gamma}(0)) = \widetilde{\gamma}_n(t_n + \varepsilon_n).$$




Figure 1.

Moreover,

$$\begin{aligned}
 (8) \quad d\left(\tilde{\gamma}_n(t_n + \epsilon_n), \phi_n\left(\widetilde{B}_n\right)\right) &= d\left(\phi_n\left(\tilde{\gamma}(0)\right), \phi_n\left(\widetilde{B}_n\right)\right) \\
 &= d\left(\tilde{\gamma}(0), \widetilde{B}_n\right).
 \end{aligned}$$

We now proceed to show that given $\varepsilon > 0$, there exists $N \in \mathbb{N}$ such that for all $n \geq N$

$$(9) \quad d(\tilde{\gamma}(s), \tilde{c}_n(s)) < \varepsilon \quad \forall s \in [0, s_n].$$

Recall that s_n is the period of c_n and $s_n < t_n + \varepsilon_n = \text{period}(\gamma_n)$. Using Lemma 10 and the fact that γ_n, c_n are homotopic for all n large enough, we have that $\tilde{c}_n(+\infty) \rightarrow \tilde{\gamma}(+\infty)$ and $\tilde{c}_n(-\infty) \rightarrow \tilde{\gamma}(-\infty)$. Let $H : G\tilde{X} \xrightarrow{\sim} \partial^2 \tilde{X} \times \mathbb{R}$ be the trivialization of the fiber bundle $G\tilde{X} \rightarrow \partial^2 \tilde{X}$ with respect to the base point $x_0 = \tilde{\gamma}(0)$. This homeomorphism was described in Remark 2 following Lemma 5. By the choice of parametrization for each \tilde{c}_n made above (i.e., $\tilde{c}_n(0) = \widetilde{B}_n$), we have that $H^{-1}(\tilde{c}_n(-\infty), \tilde{c}_n(+\infty), 0) = \tilde{c}_n$. Moreover, $H^{-1}(\tilde{\gamma}(-\infty), \tilde{\gamma}(+\infty), 0) = \tilde{\gamma}$ and, thus, $\tilde{c}_n \rightarrow \tilde{\gamma}$ uniformly on compact sets. Observe that such convergence is weaker than property (9). However, it implies, in particular, that $\text{dist}(\tilde{\gamma}(0), \text{Im } \tilde{c}_n) \rightarrow 0$ as $n \rightarrow \infty$. Hence, we may choose $N \in \mathbb{N}$ such that

$$(10) \quad d(\tilde{\gamma}(0), \widetilde{B}_n) < \varepsilon/5, \quad \text{for all } n \geq N.$$

Moreover, we may choose N such that, in addition, the following inequality is satisfied

$$(11) \quad \varepsilon_n = d(\tilde{\gamma}(t_n), \tilde{\gamma}(t_n + \varepsilon_n)) < \varepsilon/5, \quad \text{for all } n \geq N.$$

To show inequality (9), let $s \in [0, s_n]$ be arbitrary and let D_n (resp. F_n) be the point on the geodesic segment $[\tilde{\gamma}(0), \tilde{\gamma}_n(t_n + \varepsilon_n)]$

(resp. $\left[\tilde{\gamma}(0), \phi_n(\widetilde{B}_n) \right]$) whose distance from $\tilde{\gamma}(0)$ is equal to s . Then,

$$\begin{aligned} d(\tilde{\gamma}(s), \tilde{c}_n(s)) &\leq d(\tilde{\gamma}(s), D_n) + d(D_n, F_n) + d(F_n, \tilde{c}_n(s)) \\ &\leq d(\tilde{\gamma}(s), D_n) + d(D_n, F_n) + d(F_n, F'_n) + d(F'_n, \tilde{c}_n(s)) \end{aligned}$$

where F'_n is the point on $\left[\widetilde{B}_n, \phi_n(\widetilde{B}_n) \right]$ satisfying

$$d(F_n, \phi_n(\widetilde{B}_n)) = d(F'_n, \phi_n(\widetilde{B}_n)).$$

By comparison (see for example [12, Prop. 29]) we have

$$\begin{aligned} d(\tilde{\gamma}(s), D_n) &\leq d(\tilde{\gamma}(t_n + \varepsilon_n), \tilde{\gamma}_n(t_n + \varepsilon_n)) \leq 2\varepsilon_n \\ d(D_n, F_n) &\leq d(\tilde{\gamma}(t_n + \varepsilon_n), \phi_n(\widetilde{B}_n)) \\ d(F_n, F'_n) &\leq d(\tilde{\gamma}(0), \widetilde{B}_n) \\ d(F'_n, \tilde{c}_n(s)) &\leq \left| d(\tilde{\gamma}(0), \phi_n(\widetilde{B}_n)) - d(\widetilde{B}_n, \phi_n(\widetilde{B}_n)) \right| < d(\tilde{\gamma}(0), \widetilde{B}_n). \end{aligned}$$

Combining the above inequalities with inequalities (8), (10) and (11), we obtain property (9) which completes the proof of the existence of a sequence of closed geodesics approximating a given recurrent geodesic. \square

Remark. Let Γ be a discrete group of isometries of a locally compact, complete geodesic metric space Y satisfying $CAT - (\chi)$ inequality, $\chi < 0$. The notion of controlled concentration points in the limit set of Γ can be defined as follows. $\xi \in \partial\Gamma$ is a *controlled concentration point* if it admits a neighborhood U containing ξ with the following property: For every neighborhood V of ξ there exists an element $\gamma \in \Gamma$ such that $\gamma(U) \subset V$ and $\xi \in \gamma(V)$. Following [1], one can show that ξ is a controlled concentration point if and only if there exists a sequence of $\{\phi_n\}$ of distinct elements of Γ such that $\phi_n(\xi) \rightarrow \xi$ and $\phi_n(0) \rightarrow \eta$ with $\eta \neq \xi$. The proof in this more general setting is identical with the one provided in [1] except that the convergence property used there, namely, $\phi_n(x) \rightarrow \eta$ for all $x \in Y \cup \partial Y$, is provided in our case by Proposition 7.2 in [6, Ch. 1]. The latter property for ξ is equivalent to the existence of a recurrent geodesic γ with $\gamma(+\infty) = \xi$ and $\gamma(-\infty) = \eta$. Hence we obtain the following connection between recurrent geodesics and controlled concentration points which also holds for manifolds (see [1]).

Theorem 11. *Let Y be a locally compact, complete geodesic metric space Y satisfying $CAT - (\chi)$ inequality, $\chi < 0$ and Γ a discrete group of isometries of Y . A limit point $\xi \in \partial Y$ is a controlled concentration point if and only if $\gamma(+\infty) = \xi$ for some recurrent geodesic γ in Y .*

4. Construction of a counterexample.

As it was mentioned in the introduction, approximation by closed geodesics does not imply recurrence. The following example demonstrates the existence of a geodesic in a $CAT - (\chi)$, $\chi < 0$ space which is not recurrent but can be approximated by closed geodesics in the sense of Definition 1. Let X be the union of two hyperbolic cylinders identified along a (convex) geodesic strip bounded by two geodesic segments (see Figure 2). We may adjust the geometry of X so that the unique simple closed geodesic in each cylinder, denoted by c_1 and c_2 , have a common image in the geodesic strip, namely, the geodesic segment indicated by letters A and B in Figure 2. Using Cor. 5 of [2] and the fact that the geodesic strip is a convex closed subset it follows that X is a $CAT - (\chi)$ space with $\chi < 0$.




Figure 2.

Let ω_1 and ω_2 be the periods of c_1 and c_2 respectively and assume that c_1 and c_2 are parametrized so that $c_1(0) = c_2(0) = B$ and clockwise i.e., $c_1(s) = c_2(s)$ for all $s \in [0, d(A, B)]$. Define $\gamma : \mathbb{R} \rightarrow X$ as follows:

$$\begin{aligned} \gamma(t) &= c_1(t), & \text{for } t \in [0, \omega_1] \\ \gamma(t) &= c_2(t), & \text{for } t \in (-\infty, 0] \cup [\omega_1, +\infty). \end{aligned}$$

It is apparent that γ can be approximated by closed geodesics in the sense of Definition 1. We proceed to show that γ is not recurrent by showing that, γ and $s\gamma$ are not close in the compact open topology for any positive real s . For this it suffices to show that there exists $\varepsilon > 0$ and a compact $M \subset \mathbb{R}$

such that for any positive $s \in \mathbb{R}$,

$$(12) \quad d(s\gamma(t_0), \gamma(t_0)) \geq \varepsilon \quad \text{for some } t_0 \in M.$$

For simplicity, we may assume that $d(A, B) = \omega_1/2 = \omega_2/4$. Pick $\varepsilon < d(A, B)/2$ and choose a compact $M \subset \mathbb{R}$ containing the real numbers 0 and $3\omega_1/4$. Let s be arbitrary positive real. If

$$d(\gamma(s), \gamma(0)) = d(s\gamma(0), \gamma(0)) \geq \varepsilon$$

then Equation (12) is satisfied for the number $t_0 = 0$. If $d(\gamma(s), \gamma(0)) < \varepsilon$ then for $t_0 = 3\omega_1/4$ we have

$$d(s\gamma(t_0), \gamma(t_0)) = d\left(s\gamma\left(\frac{3\omega_1}{4}\right), c_1\left(\frac{3\omega_1}{4}\right)\right) > \frac{\omega_1}{4} = \frac{d(A, B)}{2} > \varepsilon.$$

This completes the proof that γ is not recurrent and, therefore, approximation by closed geodesics does not imply recurrence.

References

- [1] B. Aebicher, S. Hong and McCullough, *Recurrent geodesics and controlled concentration points*, Duke Math. J., **75**(3) (1994), 759-774.
- [2] W. Ballman, *Singular spaces of non-positive curvature*, in Sur les groupes hyperboliques d'après Gromov (Seminaire de Berne), édité par E. Ghys et P. de la Harpe, (a paraitre chez Birkhäuser), 1990.
- [3] M.R. Bridson, *Geodesics and geometry in metric simplicial complexes*, in Group Theory from a Geometrical Viewpoint, (ICTP, Trieste, Italy, March 26 - April 6, 1990), E. Ghys and A. Haefliger eds. (1991).
- [4] Ch. Champetier, *Petite simplification dans les groupes hyperboliques*, Ann. Fac. Sci. Toulouse, VI. Ser., Math., **3**(2) (1994), 161-221.
- [5] C. Charitos, *Closed geodesics in ideal polyhedra of dimension 2*, Rocky Mountain Journal of Mathematics, **26**(1) (1996), 507-521.
- [6] M. Coornaert, *Sur les groupes proprement discontinus d'isométries des espaces hyperboliques au sens de Gromov*, Thèse U.L.P., Publication de l'IRMA.
- [7] ———, *Measures de Patterson-Sullivan sur le bord d'un espace hyperbolique au sens de Gromov*, Pacific J. Math., **159**(2) (1993), 241-270.
- [8] M. Coornaert, T. Delzant and A. Papadopoulos, *Géométrie et théorie des groupes*, Lecture Notes in Mathematics, **1441**, Springer-Verlag, (1990).
- [9] M. Gromov, *Hyperbolic groups*, in 'Essays in Group Theory', MSRI Publ., **8**, Springer Verlag, (1987), 75-263.
- [10] ———, *Structures métriques pour les variétés riemanniennes*, written with J. Lafontaine and P. Pansu, Cedric Fernand Nathan, Paris, 1981.
- [11] F. Paulin, *Constructions of hyperbolic groups via hyperbolization of polyhedra*, in Group Theory from a Geometrical Viewpoint, (ICTP, Trieste, Italy, March 26 - April 6, 1990), E. Ghys and A. Haefliger eds. (1991).

- [12] M. Troyanov, *Espaces à courbure négative et groupés hyperboliques* in Sur les groupes hyperboliques d'après Gromov (Seminaire de Berne), édité par E. Ghys et P. de la Harpe, (a paraitre chez Birkhäuser), 1990.

Received September 23, 1998 and revised May 27, 1999.

AGRICULTURAL UNIVERSITY OF ATHENS
75 IERA ODOS
ATHENS 11855
GREECE
E-mail address: bakis@auadec.aua.gr

UNIVERSITY OF THE AEGEAN
KARLOVASSI
SAMOS 83200
GREECE
E-mail address: gtsap@aegean.gr